

World Journal of *Clinical Cases*

World J Clin Cases 2022 December 26; 10(36): 13148-13469



MINIREVIEWS

- 13148** Liver injury in COVID-19: Holds ferritinophagy-mediated ferroptosis accountable
Jia FJ, Han J
- 13157** Amebic liver abscess by *Entamoeba histolytica*
Usuda D, Tsuge S, Sakurai R, Kawai K, Matsubara S, Tanaka R, Suzuki M, Takano H, Shimozawa S, Hotchi Y, Tokunaga S, Osugi I, Katou R, Ito S, Mishima K, Kondo A, Mizuno K, Takami H, Komatsu T, Oba J, Nomura T, Sugita M
- 13167** Living with liver disease in the era of COVID-19-the impact of the epidemic and the threat to high-risk populations
Barve P, Choday P, Nguyen A, Ly T, Samreen I, Jhooty S, Umeh CA, Chaudhuri S
- 13179** Cortical bone trajectory screws in the treatment of lumbar degenerative disc disease in patients with osteoporosis
Guo S, Zhu K, Yan MJ, Li XH, Tan J
- 13189** Probiotics for preventing gestational diabetes in overweight or obese pregnant women: A review
Deng YF, Wu LP, Liu YP

ORIGINAL ARTICLE

Retrospective Cohort Study

- 13200** Effectiveness of microwave endometrial ablation combined with hysteroscopic transcervical resection in treating submucous uterine myomas
Kakinuma T, Kakinuma K, Shimizu A, Kaneko A, Kagimoto M, Okusa T, Suizu E, Saito K, Matsuda Y, Yanagida K, Takeshima N, Ohwada M
- 13208** Antibody and complement levels in patients with hypersplenism associated with cirrhotic portal hypertension and therapeutic principles
Zhang K, Zeng M, Li YJ, Wu HF, Wu JC, Zhang ZS, Zheng JF, Lv YF

Retrospective Study

- 13216** Case series in Indonesia: B.1.617.2 (delta) variant of SARS-CoV-2 infection after a second dose of vaccine
Karuniawati A, Syam AF, Achmadasyah A, Ibrahim F, Rosa Y, Sudarmono P, Fadilah F, Rasmin M
- 13227** Endobronchial ultrasound-guided transbronchial needle aspiration in intrathoracic lymphadenopathy with extrathoracic malignancy
Li SJ, Wu Q
- 13239** Analysis of the clinical efficacy of two-stage revision surgery in the treatment of periprosthetic joint infection in the knee: A retrospective study
Qiao YJ, Li F, Zhang LD, Yu XY, Zhang HQ, Yang WB, Song XY, Xu RL, Zhou SH

- 13250** Prognostic factors for disease-free survival in postoperative patients with hepatocellular carcinoma and construction of a nomogram model
Luo PQ, Ye ZH, Zhang LX, Song ED, Wei ZJ, Xu AM, Lu Z
- 13264** Oral higher dose prednisolone to prevent stenosis after endoscopic submucosal dissection for early esophageal cancer
Zhan SG, Wu BH, Li DF, Yao J, Xu ZL, Zhang DG, Shi RY, Tian YH, Wang LS
- 13274** Predictive value of the unplanned extubation risk assessment scale in hospitalized patients with tubes
Liu K, Liu Z, Li LQ, Zhang M, Deng XX, Zhu H
- 13284** Classification of rectal cancer according to recurrence types - comparison of Japanese guidelines and Western guidelines
Miyakita H, Kamei Y, Chan LF, Okada K, Kayano H, Yamamoto S
- 13293** Risk of critical limb ischemia in long-term uterine cancer survivors: A population-based study
Chen MC, Chang JJ, Chen MF, Wang TY, Huang CE, Lee KD, Chen CY
- 13304** Serum Spondin-2 expression, tumor invasion, and antitumor immune response in patients with cervical cancer
Zhang LL, Lin S, Zhang Y, Yao DM, Du X
- 13313** Thoracic para-aortic lymph node recurrence in patients with esophageal squamous cell carcinoma: A propensity score-matching analysis
Li XY, Huang LS, Yu SH, Xie D
- 13321** Anastomotic leakage in rectal cancer surgery: Retrospective analysis of risk factors
Brisinda G, Chiarello MM, Pepe G, Cariati M, Fico V, Mirco P, Bianchi V

META-ANALYSIS

- 13337** Successful outcomes of unilateral *vs* bilateral pedicle screw fixation for lumbar interbody fusion: A meta-analysis with evidence grading
Sun L, Tian AX, Ma JX, Ma XL

CASE REPORT

- 13349** Pregnancy-induced leukocytosis: A case report
Wang X, Zhang YY, Xu Y
- 13356** Acute moderate to severe ulcerative colitis treated by traditional Chinese medicine: A case report
Wu B
- 13364** Solitary hyoid plasmacytoma with unicentric Castleman disease: A case report and review of literature
Zhang YH, He YF, Yue H, Zhang YN, Shi L, Jin B, Dong P
- 13373** Recurrence of intratendinous ganglion due to incomplete excision of satellite lesion in the extensor digitorum brevis tendon: A case report
Park JJ, Seok HG, Yan H, Park CH

- 13381** Two methods of lung biopsy for histological confirmation of acute fibrinous and organizing pneumonia: A case report
Liu WJ, Zhou S, Li YX
- 13388** Application of 3D-printed prosthesis in revision surgery with large inflammatory pseudotumour and extensive bone defect: A case report
Wang HP, Wang MY, Lan YP, Tang ZD, Tao QF, Chen CY
- 13396** Undetected traumatic cardiac herniation like playing hide-and-seek-delayed incidental findings during surgical stabilization of flail chest: A case report
Yoon SY, Ye JB, Seok J
- 13402** Laparoscopic treatment of pyogenic liver abscess caused by fishbone puncture through the stomach wall and into the liver: A case report
Kadi A, Tuergan T, Abulaiti Y, Shalayiadang P, Tayier B, Abulizi A, Tuohuti M, Ahan A
- 13408** Hepatic sinusoidal obstruction syndrome induced by tacrolimus following liver transplantation: Three case reports
Jiang JY, Fu Y, Ou YJ, Zhang LD
- 13418** *Staphylococcus aureus* bacteremia and infective endocarditis in a patient with epidermolytic hyperkeratosis: A case report
Chen Y, Chen D, Liu H, Zhang CG, Song LL
- 13426** Compound heterozygous p.L483P and p.S310G mutations in GBA1 cause type 1 adult Gaucher disease: A case report
Wen XL, Wang YZ, Zhang XL, Tu JQ, Zhang ZJ, Liu XX, Lu HY, Hao GP, Wang XH, Yang LH, Zhang RJ
- 13435** Short-term prone positioning for severe acute respiratory distress syndrome after cardiopulmonary bypass: A case report and literature review
Yang JH, Wang S, Gan YX, Feng XY, Niu BL
- 13443** Congenital nephrogenic diabetes insipidus arginine vasopressin receptor 2 gene mutation at new site: A case report
Yang LL, Xu Y, Qiu JL, Zhao QY, Li MM, Shi H
- 13451** Development of dilated cardiomyopathy with a long latent period followed by viral fulminant myocarditis: A case report
Lee SD, Lee HJ, Kim HR, Kang MG, Kim K, Park JR
- 13458** Hoffa's fracture in a five-year-old child diagnosed and treated with the assistance of arthroscopy: A case report
Chen ZH, Wang HF, Wang HY, Li F, Bai XF, Ni JL, Shi ZB

LETTER TO THE EDITOR

- 13467** Precautions before starting tofacitinib in persons with rheumatoid arthritis
Swarnakar R, Yadav SL

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Janardhan Mydam, MD, Assistant Professor, Consultant Physician-Scientist, Statistician, Division of Neonatology, Department of Pediatrics, John H. Stroger, Jr. Hospital of Cook County 1969 W. Ogden, Chicago, IL 60612, United States. mydamj@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

December 26, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Oral higher dose prednisolone to prevent stenosis after endoscopic submucosal dissection for early esophageal cancer

Sheng-Gang Zhan, Ben-Hua Wu, De-Feng Li, Jun Yao, Zheng-Lei Xu, Ding-Guo Zhang, Rui-Yue Shi, Yan-Hui Tian, Li-Sheng Wang

Specialty type: Medicine, research and experimental

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C, C
Grade D (Fair): D
Grade E (Poor): E

P-Reviewer: Dilek ON, Turkey;

Lim KT, Singapore; Salimi M, Iran; Sumi K, Japan

Received: August 28, 2022

Peer-review started: August 28, 2022

First decision: October 5, 2022

Revised: October 22, 2022

Accepted: December 5, 2022

Article in press: December 5, 2022

Published online: December 26, 2022



Sheng-Gang Zhan, Ben-Hua Wu, De-Feng Li, Jun Yao, Zheng-Lei Xu, Ding-Guo Zhang, Rui-Yue Shi, Yan-Hui Tian, Li-Sheng Wang, Department of Gastroenterology, Second Clinical Medical College of Jinan University, Shenzhen People's Hospital, Shenzhen 518020, Guangdong Province, China

Corresponding author: Li-Sheng Wang, PhD, Doctor, Department of Gastroenterology, Second Clinical Medical College of Jinan University, Shenzhen People's Hospital, 1017 Dongmen North Road, Shenzhen 518020, Guangdong Province, China. wangls168@163.com

Abstract

BACKGROUND

Esophageal stenosis is one of the main complications of endoscopic submucosal dissection (ESD) for the treatment of large-area superficial esophageal squamous cell carcinoma and precancerous lesions ($\geq 3/4$ of the lumen). Oral prednisone is useful to prevent esophageal stenosis, but the curative effect remains controversial.

AIM

To share our experience of the precautions against esophageal stenosis after ESD to remove large superficial esophageal lesions.

METHODS

Between June 2019 and March 2022, we enrolled patients with large superficial esophageal squamous cell carcinoma and high-grade intraepithelial neoplasia experienced who underwent ESD. Prednisone (50 mg/d) was administered orally on the second morning after ESD for 1 mo, and tapered gradually (5 mg/wk) for 13 wk.

RESULTS

In total, 14 patients met the inclusion criteria. All patients received ESD without operation-related bleeding or perforation. There were 11 patients with $\geq 3/4$ and $< 7/8$ of lumen mucosal defects and 1 patient with $\geq 7/8$ of lumen mucosal defect and 2 patients with the entire circumferential mucosal defects due to ESD. The longitudinal extension of the esophageal mucosal defect was < 50 mm in 3 patients and ≥ 50 mm in 11 patients. The esophageal stenosis rate after ESD was 0% (0/14). One patient developed esophageal candida infection on the 30th d after

ESD, and completely recovered after 7 d of administration of oral fluconazole 100 mg/d. No other adverse events of oral steroids were found.

CONCLUSION

Oral prednisone (50 mg/d) and prolonged prednisone usage time may effectively prevent esophageal stricture after ESD without increasing the incidence of glucocorticoid-related adverse events. However, further investigation of larger samples is required to warrant feasibility and safety.

Key Words: Early esophageal cancer; Stenosis; Prednisone; Endoscopic submucosal dissection

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Esophageal stenosis is one of the main complications of endoscopic submucosal dissection (ESD) for the remedy of large-area superficial esophageal squamous cell carcinoma and precancerous lesions ($\geq 3/4$ of the lumen). Oral prednisone (30 mg/d) is one of the most commonly used treatment measures to prevent postoperative stenosis after esophageal ESD; however, several studies have drawn inconsistent conclusions. For the first time, we took a higher dose of prednisone (50 mg/d) orally to prevent esophageal stenosis after esophageal ESD and no stenosis occurred in 14 patients, meanwhile, no significant glucocorticoid-related adverse events occurred.

Citation: Zhan SG, Wu BH, Li DF, Yao J, Xu ZL, Zhang DG, Shi RY, Tian YH, Wang LS. Oral higher dose prednisone to prevent stenosis after endoscopic submucosal dissection for early esophageal cancer. *World J Clin Cases* 2022; 10(36): 13264-13273

URL: <https://www.wjgnet.com/2307-8960/full/v10/i36/13264.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v10.i36.13264>

INTRODUCTION

Endoscopic submucosal dissection (ESD) is one of main treatment measures for early esophageal cancer and esophageal high-grade intraepithelial neoplasia[1]. It is minimally invasive and permits *en bloc* resection of large esophageal lesions without esophagectomy. However, esophageal stenosis generally occurs after ESD resection of esophageal lesions, especially for lesions $\geq 3/4$ of the lumen. Multivariate analysis has shown that esophageal mucosal membrane defects over $3/4$ of the lumen is an important predictor of stenosis formation. Without prophylactic treatment, the occurrence rate of esophageal strictures can reach 83.3%-94.1%; especially when the lesion affects the whole circumference of the esophagus, the rate of esophageal strictures is even higher[2,3]. This often requires repeated endoscopic balloon dilatation (EBD) to alleviate symptoms; however, the benefit is limited[4].

Recently, it has been reported that hormones, as a preventive treatment, can reduce the occurrence of stricture after esophageal ESD[5,6]. Yamaguchi *et al*[5] first reported that oral prednisone 30 mg/d can effectively prevent esophageal stenosis after ESD, and the postoperative stenosis rate can be reduced to 5.3% (1/19). A study also showed that oral prednisone 30 mg /d is not effective in preventing esophageal stenosis in patients with an entire circumference or $\geq 7/8$ circumferential mucosal defects [7]. Meanwhile, one case reported that a patient with superficial esophageal squamous cell carcinoma received high-dose dexamethasone therapy (40 mg for 4 d) for multiple myeloma on the 9th d after ESD. After three courses of treatment, no esophageal stenosis was found in the follow-up gastroscopy, the histopathological evaluation showed that the submucosa became thinner, and the fibrosis degree of wound scar after ESD was the lowest[8]. In a prospective study by Nakamura, 11 patients with superficial esophageal squamous cell carcinoma with lesions $\geq 3/4$ of the circumference were treated with steroid pulse therapy from the next day after ESD (intravenous infusion of sodium methylprednisolone succinate 500 mg/d for 3 consecutive days from the next day after ESD)[9]. It was found that although steroid pulse therapy was safe, steroid pulse therapy had no preventive effect on esophageal stenosis after ESD. Therefore, oral hormone is an effective method to prevent esophageal stenosis after esophageal ESD, but the dose, use time, effectiveness, and safety of the hormone need to be further studied.

MATERIALS AND METHODS

Patients

Between June 2019 and March 2022, 74 patients with superficial esophageal squamous cell carcinoma or precancerous lesions of esophagus were *en bloc* resected with ESD at the Digestive Endoscopy Center of Shenzhen People's Hospital (Guangdong province, China). Of these, 18 patients accepted mucosal resection *via* ESD, and the mucosal defects involved a 3/4 or larger circumference of the esophageal lumen. However, 1 patient who had received additional chemoradiotherapy (CRT) and 3 patients who had undergone additional surgery were removed from our study. Ultimately, 14 patients were included in this study. The indication criteria were as follows: (1) Before esophageal ESD, the intrapapillary capillary of the lesion mucosa observed by narrow band imaging magnifying endoscopy was type B1; (2) The preoperative lesions of ESD were histologically confirmed as superficial esophageal squamous cell carcinoma or esophageal high-grade intraepithelial neoplasia; (3) Thoracoabdominal enhanced computed tomography (CT) showed no lymph node or distant metastasis; (4) Provision of written informed consent was given; (5) There was no achalasia; and (6) There was no corrosive injury of esophagus. The exclusion criteria were as follows: (1) Patients who could not be followed up for 6 mo or longer; (2) patients whose stenosis formed before esophageal ESD; (3) patients who had prior esophageal cancer with CRT; and (4) patients with additional CRT or additional esophagectomy after non-curative ESD.

ESD procedure

The ESD was operated in an operating room. The patients were endotracheal intubated and kept in the left lying position. An endoscope (GIF-Q260J; Olympus Co., Tokyo, Japan) was attached with forward water delivery function and was used with carbon dioxide insufflation. The head end of the endoscope was connected with a pellucid cap (diameter 12.4 mm, length 4 mm, D-201-11804; Olympus Co.). Iodine 0.75% staining was used for identifying the range of the lesion, and a dual knife (KD-650Q; Olympus Co.) marking dots 3 mm away from the edge of the lesion. An electronic surgical workstation (VIO 200D; ERBE, Tübingen, Germany) was used with an operating mode (Forced Coagulation model, effect 2, maximum power 20 W). The 0.9% saline solution containing 0.3% indigo carmine was used for adequate submucosal injection along the marked dots. After circumferential mucosal incision, submucosal dissection was operated by the dual knife (KD-650Q) from the oral side to the anal side of the lesion under an operating mode (Endocut I mode, Forced Coagulation model, effect 2, maximum power 50 W, VIO 200D; ERBE). Bleeding during the operation and visible blood vessels were handled with hemostatic forceps (FD-410LR; Olympus Co.) under a soft coagulating mode (Soft Coagulation mode, effect 4, maximum power 80 W, VIO 200D; ERBE).

Postoperative-related bleeding was defined as bleeding requiring blood transfusion or surgical intervention, or bleeding that resulted in a 2 g/dL decline in hemoglobin levels. Postoperative-related perforation was diagnosed by endoscopy or chest CT[7]. All patients treated with proton pump inhibitor, esomeprazole, with a dose of 20 mg, twice a day after ESD for 28 d[5,9]. After ESD, each patient received two pieces of calcium carbonate and vitamin D3 chewable tablets (each tablet contains 300 mg calcium and 60 IU vitamin D3) per day until the prednisone was stopped to prevent glucocorticoid-induced osteoporosis[10].

Management for esophageal stenosis prevention

Prednisolone was taken orally at a dose of 50 mg/d from the next morning after ESD for 1 mo, and then decreased gradually (45, 40, 35, 30, 25, 20, 15, 10, and 5 mg for 7 d each) until 13 wk later.

Follow-up

Regular endoscopy was examined at 1, 3, and 6 mo after ESD operation, and then annually thereafter. In addition, endoscopic examination was performed whenever the patient had dysphagia to determine whether there was esophageal stenosis. EBD was performed subsequently if esophageal stricture was identified. Any abnormal mucosa required biopsy for pathological evaluation of whether there was local tumor recurrence. Meanwhile, regular physical and blood examinations were carried out to evaluate the side effects of the steroid. Bone mineral density testing was performed before ESD treatment and 6 mo after ESD treatment.

Outcomes

The main outcome measure was incidence of esophageal stenosis. Esophageal stenosis was defined as the inability of 9.9 mm diameter gastroscope (GIF-Q260J; Olympus) to pass through the esophageal stenosis. Secondary observation indicators of glucocorticoid-related adverse events were observed at 1, 3, and 6 mo after ESD such as newly diagnosed diabetes or aggravation of diabetes, peptic ulcer, adrenocortical insufficiency, aggravation of osteoporosis or fracture, and corticosteroid-related mental disorders.

Table 1 Background characteristics of patients, *n* (%)

Characteristics	
Sex	
Male	9 (64)
Female	5 (36)
Age in yr, mean (range)	62.1 (45-75)
Tumor location	
Cervical esophagus	0 (0)
Upper thoracic esophagus	1 (7)
Middle thoracic esophagus	7 (50)
Lower thoracic esophagus	6 (43)
Macroscopic type	
0-IIa	13 (93)
0-IIc	1 (7)
Tumor size in mm, mean (range)	51.7 (43.8-60.7)
Resection size in mm, mean (range)	55.5 (47.5-65.0)
Transverse extension of mucosal defect	
≥ 3/4 and < 7/8 circumference	11 (79)
≥ 7/8 circumference	1 (7)
The entire circumference	2 (14)
Longitudinal extension of the mucosal defect	
< 50 mm	3 (21)
≥ 50 mm	11 (79)
Depth of tumor invasion	
M1 or M2	12 (86)
M3	2 (14)

M1: Tumor limited to the epithelium; M2: Tumor invading the lamina propria; M3: Tumor invading the muscularis mucosa.

End point: The follow-up was terminated if tumor recurrence and serious adverse events of glucocorticoid and procedure-related complications (procedure-related bleeding and procedure-related perforation) occurred.

Statistical analyses

Continuous variables are presented as the mean ± standard deviation or median (interquartile range, 25%-75%). Categorical variables are expressed by proportion. Data analyses were conducted using SPSS 23.0 software (version 23.0 for Mac).

RESULTS

Background characteristics of patients

After ESD surgery, there were 18 patients with mucosal defects more than 3/4 of the esophageal circumference. One patient received additional CRT treatment, and 3 patients received additional surgery and were removed from this study. Eventually, a total of 14 patients met the criteria. Patients and characteristics of lesions are shown in Table 1 and Table 2. Male patients accounted for 64%, with a mean age of 62.1 years (ranging from 45 to 75 years). According to the Paris endoscopic classification, 13 cases of endoscopic tumor morphology were classified as type 0-IIa and 1 case was type 0-IIc. The lesions were mainly located in the middle and lower esophagus, and 1 case was located in the upper esophagus. Each patient successfully received esophageal ESD treatment, and postoperative pathology confirmed

Table 2 The feasibility and safety of prophylactic treatment, *n* (%)

Outcome	
Stricture rate	0 (0)
Main complication	1 (7)
Procedure-related bleeding	0 (0)
Procedure-related perforation	0 (0)
Newly diagnosed diabetes or aggravation of diabetes	0 (0)
Pepticulcer	0 (0)
Adrenocortical insufficiency	0 (0)
Aggravation of osteoporosis or fracture	0 (0)
Infection	1 (7)
Corticosteroid related mental disorders	0 (0)
Follow-up period (mo), median (range)	13 (6-28)
Residual	0 (0)
Recurrence	0 (0)

that the lesion was completely removed. All patients had no procedure-related bleeding and procedure-related perforation after esophageal ESD. The mean resection size was 55.5 mm in diameter (ranging from 47.5 mm to 65.0 mm). According to the range of esophageal mucosal defect, 11 cases involved $\geq 3/4$ and $< 7/8$ circumference, 1 case involved $\geq 7/8$ circumference, and 2 cases involved the entire circumference. The longitudinal extension of mucosal defect was < 50 mm in 3 patients and ≥ 50 mm in 11 patients. In 12 cases, the depth of invasion of pathological tissues was limited within the epithelium and lamina propria mucosa, whereas 2 lesions were limited within muscularis mucosa without lymphovascular infiltration.

The shortest follow-up time of all cases was 6 mo, the longest follow-up time was 28 mo, and the median follow-up time was 13 mo. During this period, all patients were followed up by endoscopy regularly without dysphagia. The incidence of esophageal stenosis was 0% (0/14) (Table 3). Representative cases are shown in Figures 1 and 2.

Only 1 patient developed esophageal *Candida* infection on the 30th d after ESD and recovered completely after 7d of treatment with oral fluconazole 100 mg/d. Glucocorticoid-related adverse events were observed such as newly diagnosed diabetes or aggravation of diabetes, pepticulcer, adrenocortical insufficiency, aggravation of osteoporosis or fracture, and corticosteroid-related mental disorders.

DISCUSSION


In the current study, increasing the dose of oral hormone (prednisone acetate 50 mg/d) and prolonging the treatment time (13 wk) were effective to prevent esophageal stenosis in patients with mucosal defects $\geq 3/4$ circumference after ESD. Studies have shown that the occurrence of esophageal stricture after ESD is related to the infiltration of postoperative inflammatory cells and vascular proliferation[11, 12]. At the same time, epithelial cells proliferate and migrate from the edge of the wound after ESD, fibroblasts proliferate continuously, and finally fibrous scar is formed. This process is divided into three stages: acute inflammation, proliferation, and remodeling; however, the duration of this process is unknown. Through the dog model study, Honda found that within about 1 mo after esophageal EMR, the mucosal defect healed and was covered by squamous cells. Although the proper muscle layer was not damaged, muscle fiber atrophy still occurred in the 1st wk after operation, and finally fibrosis was formed[13]. Some clinical studies have also shown that esophageal stenosis mostly occurs within 2 to 4 wk after surgery[3,14], but this was confined to endoscopic observation. Glucocorticoid has a strong anti-inflammatory effect, which not only inhibits the synthesis of collagen but also promotes the decomposition of collagen to inhibit the formation of stenosis. In our study, all cases did not have esophageal stenosis after ESD. We believe that increasing the dose of prednisolone can enhance the anti-inflammatory effect in the acute inflammatory period, especially in the critical period of the 1st mo after ESD. Meanwhile, we speculate that the process of esophageal stenosis may last longer than expected, and prolonging the usage of prednisolone may inhibit the proliferation of fibroblasts steadily to prevent esophageal remodeling and the formation of esophageal stenosis.

Table 3 Clinical outcomes for 14 patients with early esophageal cancer after circumferential endoscopic submucosal dissection

	Case 1	Case 2	Case 3	Case 4	Case 5	Case 6	Case 7	Case 8	Case 9	Case 10	Case 11	Case 12	Case 13	Case 14
Sex	Male	Male	Male	Female	Female	Male	Male	Male	Female	Male	Female	Male	Male	Female
Age in yr	66	45	50	75	67	61	63	45	70	72	67	55	66	67
Tumor location	Lt	Ut	Lt	Mt	Mt	Mt	Lt	Lt	Mt	Mt	Mt	Lt	Lt	Mt
Macroscopic type	0-IIa	0-IIa	0-IIa	0-IIa	0-IIc	0-IIa	0-IIa	0-IIa	0-IIa	0-IIa	0-IIa	0-IIa	0-IIa	0-IIa
Tumor size in mm	50.7	49.5	48.4	43.8	47.2	53.3	60.7	49.2	59.8	46.5	59.5	45.2	51.8	58.6
Resection size in mm	56.1	52.6	52.7	47.5	50.8	57.6	65.0	53.2	64.7	50.1	63.5	47.1	54.6	61.2
Transverse extension of mucosal defect	≥ 3/4 and < 7/8	≥ 3/4 and < 7/8	≥ 3/4 and < 7/8	≥ 3/4 and < 7/8	≥ 3/4 and < 7/8	≥ 3/4 and < 7/8	Entire	≥ 3/4 and < 7/8	Entire	≥ 3/4 and < 7/8	≥ 7/8	≥ 3/4 and < 7/8	≥ 3/4 and < 7/8	≥ 3/4 and < 7/8
Longitudinal extension of the mucosal defect in mm	≥ 50	≥ 50	< 50	≥ 50	≥ 50	≥ 50	≥ 50	< 50	≥ 50	≥ 50	≥ 50	< 50	≥ 50	≥ 50
Depth of tumor invasion	M3	M2	M1	M1	M2	M3	M2	M1	M2	M1	M2	M1	M1	M2
Stricture	No	No	No	No	No	No	No	No	No	No	No	No	No	No
Follow up time in mo	9	12	12	12	12	12	15	18	28	28	6	6	7	8
Residual	No	No	No	No	No	No	No	No	No	No	No	No	No	No
Recurrence	No	No	No	No	No	No	No	No	No	No	No	No	No	No

Lt: Lower thoracic esophagus; M1: Tumor limited to the epithelium; M2: Tumor invading the lamina propria; M3: Tumor invading the muscularis mucosa; Mt: Middle thoracic esophagus; Ut: Upper thoracic esophagus.

There are several reports on the application of steroids to prevent stenosis after ESD operation for large-area superficial esophageal squamous cell carcinoma and precancerous lesions. Yamaguchi *et al*[5] reported the therapeutic effect of oral prednisolone after esophageal ESD for the first time. In their report, prednisone was taken on the 3rd d after ESD, with an initial dose of 30 mg/d, and then decreased gradually (30, 30, 25, 25, 20, 15, 10, and 5 mg for 1 wk each). The incidence of stenosis after semi-circumferential ESD resection and entire circumferential ESD resection were 6.3% (1/16) and 0% (0/3), respectively[5]. However, for the cases of circumferential esophageal mucosal defect after ESD, Sato *et al* [15] found that oral prednisone 30 mg/d could not reduce the incidence of postoperative esophageal stenosis, but could decrease the total number of EBD expansions required. In Kadota's[7] study, the stenosis rate of patients with less than entire circumferential ESD resection and with oral prednisone 30 mg/d administration was similar to the results by Yamaguchi *et al*[5], whereas patients with entire circumferential ESD resection showed higher stenosis rate (10/14) even with additional local submucosal steroid injections[7]. Meanwhile, two studies of submucosal injection of triamcinolone acetonide within the mucosal defects combined with oral prednisone in the prevention of esophageal stenosis post-ESD for lesions more than 3/4 circumference have obtained completely opposite results. Chu *et al*[16] reported that after treated with submucosal injection of triamcinolone acetonide within the mucosal defects combined with oral prednisone 30 mg/d, the incidence of esophageal stenosis was only 18.2% (2/11), including lesions with total circumferential resection. Surprisingly, in the Hanaoka *et al* [17] study of 12 cases with whole circumferential defect, the same steroid submucosal injection combined with oral prednisone 5 mg/d were used for post-ESD treatment; nevertheless, 11 patients failed to avoid postoperative stenosis. This discrepancy may be caused by different doses of orally-taken prednisolone in these studies. A study about short-term usage of oral prednisolone (30, 20, and 10 mg/d for 1 wk each) for mucosal defects ≥ 3/4 circumference, including 3 patients with total circumferential resection that showed a stenosis rate of 18% (3/17), and 1 of 3 patients with total circumferential resection withstood stenosis[18]. Accordingly, we speculate that the prevention of esophageal stenosis after esophageal ESD by oral prednisone was correlated with the dose and the use time. In our study, we increased the dose of prednisone to 50 mg/d and prolonged the treatment time to 13 wk. During our follow-up, 14 patients had no feedback of dysphagia symptoms, and no esophageal stenosis observed by endoscopic examination. Especially the 2 patients with entire circumference mucosal defects, although the esophageal wounds were fibrotic, the 9.9 mm diameter gastroscop (GIF-Q260J; Olympus)



DOI: 10.12998/wjcc.v10.i36.13264 Copyright ©The Author(s) 2022.

Figure 1 Representative case (case 1). A 67-yr-old female who underwent endoscopic resection for large superficial esophageal squamous cell carcinoma. A: Endoscopic view of the tumor after Lugol's staining. The tumor spread to more than 3/4 of the circumference of the esophageal lumen; B: Endoscopic view of the ulcer bed immediately after endoscopic submucosal dissection. The width of the mucosal defect was $\geq 3/4$ and less than the entire circumference. Then oral steroid was administered as a prophylactic treatment; C: Endoscopic view on the 30th d. The mucosal defect was still undergoing re-epithelialization, and a 9.9 mm diameter gastroscope (Olympus GIF-Q260J) could pass. D: Endoscopic view on the 180th d; Complete epithelialization is shown and a 9.9 mm diameter gastroscope (Olympus GIF-Q260J) could pass without dysphagia.


could pass, and we did not add EBD treatment. In addition, studies have shown that the injury of the intrinsic muscle layer was one of the risk factors for esophageal stenosis after ESD for early esophageal cancer and precancerous lesions[19,20]. Therefore, we paid more attention to avoid the injury of the intrinsic muscle layer as much as possible during ESD operation, which we think is also helpful for the prevention of postoperative esophageal stenosis.

Furthermore, systemic steroids are associated with adverse events, including newly diagnosed diabetes or aggravation of diabetes, peptic ulcer, adrenocortical insufficiency, aggravation of osteoporosis or fracture, and corticosteroid-related mental disorders. Stuck *et al*[21] showed that when the cumulative dose of oral prednisone exceeded 700 mg, the risk of infectious complications in patients taking prednisone increased with the increase of prednisone dosage. One study also found that even short-term steroid use is related to increased risks of adverse events[22]. However, in our protocol, the accumulated dose of oral steroids was 3075 mg, which was higher than that of other studies, and proton pump inhibitor, oral calcium, and vitamin D3 were taken simultaneously. One patient was found to have esophageal *Candida* infection on the 30th d after operation, and completely recovered after 7 d of oral fluconazole 100 mg/d therapy, and no patients experienced other adverse incidents related to orally-taken prednisolone. Therefore, we believe that the treatment scheme of increasing the dose of prednisone (50 mg/d) is safe, but still needs long-term follow-up and observation.

This study had several limitations. First, this study was a retrospective analysis in single-centered, and possible bias could not be avoided. Second, the follow-up time was insufficient and could not comprehensively evaluate the feasibility and safety of the hormone. Third, the number of subjects was relatively small, and the control group was lacking, so statistical difference analysis could not be conducted. Due to these limitations, prospective randomized controlled studies should be established to validate the efficacy and safety of prophylactic steroid therapy.

CONCLUSION

In conclusion, increasing the dose of oral prednisone (50 mg/d) and prolonging the usage time (total 13



DOI: 10.12998/wjcc.v10.i36.13264 Copyright ©The Author(s) 2022.

Figure 2 Representative case (case 3). A 70-yr-old female who underwent endoscopic resection for large superficial esophageal squamous cell carcinoma. A: Endoscopic view of the tumor after Lugol's staining. The tumor spread to about the entire circumference of the esophageal lumen; B: Endoscopic view of the ulcer bed immediately after endoscopic submucosal dissection. The width of the mucosal defect was the entire lumen circumference. Then oral steroid was administered as a prophylactic treatment; C: Endoscopic view 6 mo later. The mucosal defect underwent complete epithelialization, and an 9.9 mm diameter gastroscope (Olympus GIF-Q260J) could pass; D: Endoscopic view after 1 yr. The endoscope could pass without dysphagia.

wk) may effectively prevent esophageal stenosis after ESD to remove large-area superficial esophageal squamous cell carcinoma or precancerous lesions of esophagus, and does not increase the incidence of glucocorticoid-related adverse events.

ARTICLE HIGHLIGHTS

Research background

Esophageal stenosis is one of the main complications of endoscopic submucosal dissection (ESD) for the treatment of large-area superficial esophageal squamous cell carcinoma and precancerous lesions ($\geq 3/4$ of the lumen). Oral prednisone is useful to prevent esophageal stenosis, but the curative effect remains controversial.

Research motivation

Explore more effective methods to prevent esophageal stenosis after ESD for early esophageal cancer and precancerous lesions.

Research objectives

We shared our experience of the precautions against esophageal stenosis after ESD to remove large superficial esophageal lesions.

Research methods

Patients with large superficial esophageal squamous cell carcinoma and high-grade intraepithelial neoplasia experienced ESD were enrolled. Prednisone (50 mg/d) was administered orally on the 2nd d after ESD for 1 mo, and tapered gradually (5 mg/wk) for 13 wk.

Research results

According to the range of esophageal mucosal defect, 11 cases involved $\geq 3/4$ and $< 7/8$ circumference, 1 case involved $\geq 7/8$ circumference, and 2 cases involved the entire circumference. The incidence of esophageal stenosis was 0% (0/14), and only 1 patient developed esophageal *Candida* infection on the 30th d after ESD and recovered completely after 7d of treatment with oral fluconazole 100 mg/d.

Research conclusions

Further investigation of larger samples is required to warrant feasibility and safety.

Research perspectives

In conclusion, increasing the dose of oral prednisone (50 mg/d) and prolonging the usage time (total 13 wk) may effectively prevent esophageal stenosis after ESD removing large-area superficial esophageal squamous cell carcinoma or precancerous lesions of esophagus, and does not increase the incidence of glucocorticoid-related adverse events.

FOOTNOTES

Author contributions: Zhan SG and Wang LS were responsible for the design of the study and reviewed the manuscript; Zhan SG, Wu BH, Li DF, and Yao J extracted the data; Zhan SG, Xu ZL, Zhang DG, Shi RY, and Tian YH performed the data analysis; Zhan GS and Wang LS were responsible for revising the manuscript; All authors have read and approved the final manuscript.

Institutional review board statement: This study was approved by the Ethics Committee of Second Clinical Medical College of Jinan University, Shenzhen People's Hospital (Approval No. LL-KY-2022150-02).

Informed consent statement: Patients were not required to give informed consent to the study because the analysis used anonymous clinical data that were obtained after each patient agreed to treatment by written consent.

Conflict-of-interest statement: The authors have no conflicts of interest to declare.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Sheng-Gang Zhan 0000-0002-3318-2635; Ben-Hua Wu 0000-0002-9392-9470; De-Feng Li 0000-0003-3118-6840; Jun Yao 0000-0002-3472-1602; Zheng-Lei Xu 0000-0002-5413-7390; Ding-Guo Zhang 0000-0001-7728-9672; Rui-Yue Shi 0000-0001-9123-2103; Yan-Hui Tian 0000-0003-4671-2938; Li-Sheng Wang 0000-0002-7418-6114.

S-Editor: Liu GL

L-Editor: Filipodia

P-Editor: Liu GL

REFERENCES

- 1 **Oyama T**, Tomori A, Hotta K, Morita S, Kominato K, Tanaka M, Miyata Y. Endoscopic submucosal dissection of early esophageal cancer. *Clin Gastroenterol Hepatol* 2005; **3**: S67-S70 [PMID: 16013002 DOI: 10.1016/s1542-3565(05)00291-0]
- 2 **Shi Q**, Ju H, Yao LQ, Zhou PH, Xu MD, Chen T, Zhou JM, Chen TY, Zhong YS. Risk factors for postoperative stricture after endoscopic submucosal dissection for superficial esophageal carcinoma. *Endoscopy* 2014; **46**: 640-644 [PMID: 24830402 DOI: 10.1055/s-0034-1365648]
- 3 **Ono S**, Fujishiro M, Niimi K, Goto O, Kodashima S, Yamamichi N, Omata M. Predictors of postoperative stricture after esophageal endoscopic submucosal dissection for superficial squamous cell neoplasms. *Endoscopy* 2009; **41**: 661-665 [PMID: 19565442 DOI: 10.1055/s-0029-1214867]
- 4 **Yoda Y**, Yano T, Kaneko K, Tsuruta S, Oono Y, Kojima T, Minashi K, Ikematsu H, Ohtsu A. Endoscopic balloon dilatation for benign fibrotic strictures after curative nonsurgical treatment for esophageal cancer. *Surg Endosc* 2012; **26**: 2877-2883 [PMID: 22543993 DOI: 10.1007/s00464-012-2273-9]
- 5 **Yamaguchi N**, Isomoto H, Nakayama T, Hayashi T, Nishiyama H, Ohnita K, Takeshima F, Shikuwa S, Kohno S, Nakao K. Usefulness of oral prednisolone in the treatment of esophageal stricture after endoscopic submucosal dissection for

- superficial esophageal squamous cell carcinoma. *Gastrointest Endosc* 2011; **73**: 1115-1121 [PMID: [21492854](#) DOI: [10.1016/j.gie.2011.02.005](#)]
- 6 **Hashimoto S**, Kobayashi M, Takeuchi M, Sato Y, Narisawa R, Aoyagi Y. The efficacy of endoscopic triamcinolone injection for the prevention of esophageal stricture after endoscopic submucosal dissection. *Gastrointest Endosc* 2011; **74**: 1389-1393 [PMID: [22136782](#) DOI: [10.1016/j.gie.2011.07.070](#)]
 - 7 **Kadota T**, Yano T, Kato T, Imajoh M, Noguchi M, Morimoto H, Osera S, Yoda Y, Oono Y, Ikematsu H, Ohtsu A, Kaneko K. Prophylactic steroid administration for strictures after endoscopic resection of large superficial esophageal squamous cell carcinoma. *Endosc Int Open* 2016; **4**: E1267-E1274 [PMID: [28028531](#) DOI: [10.1055/s-0042-118291](#)]
 - 8 **Ono S**, Fujishiro M, Kodashima S, Minatsuki C, Hirano K, Niimi K, Goto O, Yamamichi N, Fukuda T, Seto Y, Koike K. High-dose dexamethasone may prevent esophageal stricture after endoscopic submucosal dissection. *Clin J Gastroenterol* 2010; **3**: 155-158 [PMID: [26190123](#) DOI: [10.1007/s12328-010-0147-z](#)]
 - 9 **Nakamura J**, Hikichi T, Watanabe K, Sato M, Obara K, Ohira H. Feasibility of Short-Period, High-Dose Intravenous Methylprednisolone for Preventing Stricture after Endoscopic Submucosal Dissection for Esophageal Cancer: A Preliminary Study. *Gastroenterol Res Pract* 2017; **2017**: 9312517 [PMID: [28828004](#) DOI: [10.1155/2017/9312517](#)]
 - 10 **Buckley L**, Guyatt G, Fink HA, Cannon M, Grossman J, Hansen KE, Humphrey MB, Lane NE, Magrey M, Miller M, Morrison L, Rao M, Robinson AB, Saha S, Wolver S, Bannuru RR, Vaysbrot E, Osani M, Turgunbaev M, Miller AS, McAlindon T. 2017 American College of Rheumatology Guideline for the Prevention and Treatment of Glucocorticoid-Induced Osteoporosis. *Arthritis Rheumatol* 2017; **69**: 1521-1537 [PMID: [28585373](#) DOI: [10.1002/art.40137](#)]
 - 11 **Wick G**, Grundtman C, Mayerl C, Wimpissinger TF, Feichtinger J, Zelger B, Sgonc R, Wolfram D. The immunology of fibrosis. *Annu Rev Immunol* 2013; **31**: 107-135 [PMID: [23516981](#) DOI: [10.1146/annurev-immunol-032712-095937](#)]
 - 12 **Duffield JS**, Lupher M, Thannickal VJ, Wynn TA. Host responses in tissue repair and fibrosis. *Annu Rev Pathol* 2013; **8**: 241-276 [PMID: [23092186](#) DOI: [10.1146/annurev-pathol-020712-163930](#)]
 - 13 **Honda M**, Nakamura T, Hori Y, Shionoya Y, Yamamoto K, Nishizawa Y, Kojima F, Shigeno K. Feasibility study of corticosteroid treatment for esophageal ulcer after EMR in a canine model. *J Gastroenterol* 2011; **46**: 866-872 [PMID: [21597933](#) DOI: [10.1007/s00535-011-0400-3](#)]
 - 14 **Mizuta H**, Nishimori I, Kuratani Y, Higashidani Y, Kohsaki T, Onishi S. Predictive factors for esophageal stenosis after endoscopic submucosal dissection for superficial esophageal cancer. *Dis Esophagus* 2009; **22**: 626-631 [PMID: [19302207](#) DOI: [10.1111/j.1442-2050.2009.00954.x](#)]
 - 15 **Sato H**, Inoue H, Kobayashi Y, Maselli R, Santi EG, Hayee B, Igarashi K, Yoshida A, Ikeda H, Onimaru M, Aoyagi Y, Kudo SE. Control of severe strictures after circumferential endoscopic submucosal dissection for esophageal carcinoma: oral steroid therapy with balloon dilation or balloon dilation alone. *Gastrointest Endosc* 2013; **78**: 250-257 [PMID: [23453294](#) DOI: [10.1016/j.gie.2013.01.008](#)]
 - 16 **Chu Y**, Chen T, Li H, Zhou P, Zhang Y, Chen W, Zhong Y, Yao L, Xu M. Long-term efficacy and safety of intralesional steroid injection plus oral steroid administration in preventing stricture after endoscopic submucosal dissection for esophageal epithelial neoplasms. *Surg Endosc* 2019; **33**: 1244-1251 [PMID: [30171398](#) DOI: [10.1007/s00464-018-6404-9](#)]
 - 17 **Hanaoka N**, Ishihara R, Uedo N, Takeuchi Y, Higashino K, Akasaka T, Kanesaka T, Matsuura N, Yamasaki Y, Hamada K, Iishi H. Refractory strictures despite steroid injection after esophageal endoscopic resection. *Endosc Int Open* 2016; **4**: E354-E359 [PMID: [27004256](#) DOI: [10.1055/s-0042-100903](#)]
 - 18 **Kataoka M**, Anzai S, Shirasaki T, Ikemiyagi H, Fujii T, Mabuchi K, Suzuki S, Yoshida M, Kawai T, Kitajima M. Efficacy of short period, low dose oral prednisolone for the prevention of stricture after circumferential endoscopic submucosal dissection (ESD) for esophageal cancer. *Endosc Int Open* 2015; **3**: E113-E117 [PMID: [26135649](#) DOI: [10.1055/s-0034-1390797](#)]
 - 19 **Ezoe Y**, Muto M, Horimatsu T, Morita S, Miyamoto S, Mochizuki S, Minashi K, Yano T, Ohtsu A, Chiba T. Efficacy of preventive endoscopic balloon dilation for esophageal stricture after endoscopic resection. *J Clin Gastroenterol* 2011; **45**: 222-227 [PMID: [20861798](#) DOI: [10.1097/MCG.0b013e3181f39f4e](#)]
 - 20 **Miwata T**, Oka S, Tanaka S, Kagemoto K, Sanomura Y, Urabe Y, Hiyama T, Chayama K. Risk factors for esophageal stenosis after entire circumferential endoscopic submucosal dissection for superficial esophageal squamous cell carcinoma. *Surg Endosc* 2016; **30**: 4049-4056 [PMID: [26703127](#) DOI: [10.1007/s00464-015-4719-3](#)]
 - 21 **Stuck AE**, Minder CE, Frey FJ. Risk of infectious complications in patients taking glucocorticosteroids. *Rev Infect Dis* 1989; **11**: 954-963 [PMID: [2690289](#) DOI: [10.1093/clinids/11.6.954](#)]
 - 22 **Waljee AK**, Rogers MA, Lin P, Singal AG, Stein JD, Marks RM, Ayanian JZ, Nallamotheu BK. Short term use of oral corticosteroids and related harms among adults in the United States: population based cohort study. *BMJ* 2017; **357**: j1415 [PMID: [28404617](#) DOI: [10.1136/bmj.j1415](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

