

# World Journal of *Clinical Cases*

*World J Clin Cases* 2022 December 26; 10(36): 13148-13469



### MINIREVIEWS

- 13148** Liver injury in COVID-19: Holds ferritinophagy-mediated ferroptosis accountable  
*Jia FJ, Han J*
- 13157** Amebic liver abscess by *Entamoeba histolytica*  
*Usuda D, Tsuge S, Sakurai R, Kawai K, Matsubara S, Tanaka R, Suzuki M, Takano H, Shimozawa S, Hotchi Y, Tokunaga S, Osugi I, Katou R, Ito S, Mishima K, Kondo A, Mizuno K, Takami H, Komatsu T, Oba J, Nomura T, Sugita M*
- 13167** Living with liver disease in the era of COVID-19-the impact of the epidemic and the threat to high-risk populations  
*Barve P, Choday P, Nguyen A, Ly T, Samreen I, Jhooty S, Umeh CA, Chaudhuri S*
- 13179** Cortical bone trajectory screws in the treatment of lumbar degenerative disc disease in patients with osteoporosis  
*Guo S, Zhu K, Yan MJ, Li XH, Tan J*
- 13189** Probiotics for preventing gestational diabetes in overweight or obese pregnant women: A review  
*Deng YF, Wu LP, Liu YP*

### ORIGINAL ARTICLE

#### Retrospective Cohort Study

- 13200** Effectiveness of microwave endometrial ablation combined with hysteroscopic transcervical resection in treating submucous uterine myomas  
*Kakinuma T, Kakinuma K, Shimizu A, Kaneko A, Kagimoto M, Okusa T, Suizu E, Saito K, Matsuda Y, Yanagida K, Takeshima N, Ohwada M*
- 13208** Antibody and complement levels in patients with hypersplenism associated with cirrhotic portal hypertension and therapeutic principles  
*Zhang K, Zeng M, Li YJ, Wu HF, Wu JC, Zhang ZS, Zheng JF, Lv YF*

#### Retrospective Study

- 13216** Case series in Indonesia: B.1.617.2 (delta) variant of SARS-CoV-2 infection after a second dose of vaccine  
*Karuniawati A, Syam AF, Achmadasyah A, Ibrahim F, Rosa Y, Sudarmono P, Fadilah F, Rasmin M*
- 13227** Endobronchial ultrasound-guided transbronchial needle aspiration in intrathoracic lymphadenopathy with extrathoracic malignancy  
*Li SJ, Wu Q*
- 13239** Analysis of the clinical efficacy of two-stage revision surgery in the treatment of periprosthetic joint infection in the knee: A retrospective study  
*Qiao YJ, Li F, Zhang LD, Yu XY, Zhang HQ, Yang WB, Song XY, Xu RL, Zhou SH*

- 13250** Prognostic factors for disease-free survival in postoperative patients with hepatocellular carcinoma and construction of a nomogram model  
*Luo PQ, Ye ZH, Zhang LX, Song ED, Wei ZJ, Xu AM, Lu Z*
- 13264** Oral higher dose prednisolone to prevent stenosis after endoscopic submucosal dissection for early esophageal cancer  
*Zhan SG, Wu BH, Li DF, Yao J, Xu ZL, Zhang DG, Shi RY, Tian YH, Wang LS*
- 13274** Predictive value of the unplanned extubation risk assessment scale in hospitalized patients with tubes  
*Liu K, Liu Z, Li LQ, Zhang M, Deng XX, Zhu H*
- 13284** Classification of rectal cancer according to recurrence types - comparison of Japanese guidelines and Western guidelines  
*Miyakita H, Kamei Y, Chan LF, Okada K, Kayano H, Yamamoto S*
- 13293** Risk of critical limb ischemia in long-term uterine cancer survivors: A population-based study  
*Chen MC, Chang JJ, Chen MF, Wang TY, Huang CE, Lee KD, Chen CY*
- 13304** Serum Spondin-2 expression, tumor invasion, and antitumor immune response in patients with cervical cancer  
*Zhang LL, Lin S, Zhang Y, Yao DM, Du X*
- 13313** Thoracic para-aortic lymph node recurrence in patients with esophageal squamous cell carcinoma: A propensity score-matching analysis  
*Li XY, Huang LS, Yu SH, Xie D*
- 13321** Anastomotic leakage in rectal cancer surgery: Retrospective analysis of risk factors  
*Brisinda G, Chiarello MM, Pepe G, Cariatì M, Fico V, Mirco P, Bianchi V*

**META-ANALYSIS**

- 13337** Successful outcomes of unilateral *vs* bilateral pedicle screw fixation for lumbar interbody fusion: A meta-analysis with evidence grading  
*Sun L, Tian AX, Ma JX, Ma XL*

**CASE REPORT**

- 13349** Pregnancy-induced leukocytosis: A case report  
*Wang X, Zhang YY, Xu Y*
- 13356** Acute moderate to severe ulcerative colitis treated by traditional Chinese medicine: A case report  
*Wu B*
- 13364** Solitary hyoid plasmacytoma with unicentric Castleman disease: A case report and review of literature  
*Zhang YH, He YF, Yue H, Zhang YN, Shi L, Jin B, Dong P*
- 13373** Recurrence of intratendinous ganglion due to incomplete excision of satellite lesion in the extensor digitorum brevis tendon: A case report  
*Park JJ, Seok HG, Yan H, Park CH*

- 13381** Two methods of lung biopsy for histological confirmation of acute fibrinous and organizing pneumonia: A case report  
*Liu WJ, Zhou S, Li YX*
- 13388** Application of 3D-printed prosthesis in revision surgery with large inflammatory pseudotumour and extensive bone defect: A case report  
*Wang HP, Wang MY, Lan YP, Tang ZD, Tao QF, Chen CY*
- 13396** Undetected traumatic cardiac herniation like playing hide-and-seek-delayed incidental findings during surgical stabilization of flail chest: A case report  
*Yoon SY, Ye JB, Seok J*
- 13402** Laparoscopic treatment of pyogenic liver abscess caused by fishbone puncture through the stomach wall and into the liver: A case report  
*Kadi A, Tuergan T, Abulaiti Y, Shalayiadang P, Tayier B, Abulizi A, Tuohuti M, Ahan A*
- 13408** Hepatic sinusoidal obstruction syndrome induced by tacrolimus following liver transplantation: Three case reports  
*Jiang JY, Fu Y, Ou YJ, Zhang LD*
- 13418** *Staphylococcus aureus* bacteremia and infective endocarditis in a patient with epidermolytic hyperkeratosis: A case report  
*Chen Y, Chen D, Liu H, Zhang CG, Song LL*
- 13426** Compound heterozygous p.L483P and p.S310G mutations in GBA1 cause type 1 adult Gaucher disease: A case report  
*Wen XL, Wang YZ, Zhang XL, Tu JQ, Zhang ZJ, Liu XX, Lu HY, Hao GP, Wang XH, Yang LH, Zhang RJ*
- 13435** Short-term prone positioning for severe acute respiratory distress syndrome after cardiopulmonary bypass: A case report and literature review  
*Yang JH, Wang S, Gan YX, Feng XY, Niu BL*
- 13443** Congenital nephrogenic diabetes insipidus arginine vasopressin receptor 2 gene mutation at new site: A case report  
*Yang LL, Xu Y, Qiu JL, Zhao QY, Li MM, Shi H*
- 13451** Development of dilated cardiomyopathy with a long latent period followed by viral fulminant myocarditis: A case report  
*Lee SD, Lee HJ, Kim HR, Kang MG, Kim K, Park JR*
- 13458** Hoffa's fracture in a five-year-old child diagnosed and treated with the assistance of arthroscopy: A case report  
*Chen ZH, Wang HF, Wang HY, Li F, Bai XF, Ni JL, Shi ZB*

**LETTER TO THE EDITOR**

- 13467** Precautions before starting tofacitinib in persons with rheumatoid arthritis  
*Swarnakar R, Yadav SL*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Janardhan Mydam, MD, Assistant Professor, Consultant Physician-Scientist, Statistician, Division of Neonatology, Department of Pediatrics, John H. Stroger, Jr. Hospital of Cook County 1969 W. Ogden, Chicago, IL 60612, United States. mydamj@gmail.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

December 26, 2022

**COPYRIGHT**

© 2022 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

# Staphylococcus aureus bacteremia and infective endocarditis in a patient with epidermolytic hyperkeratosis: A case report

Yu Chen, Dian Chen, Hao Liu, Chen-Guang Zhang, Lin-Lin Song

**Specialty type:** Emergency medicine

**Provenance and peer review:** Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B  
Grade C (Good): C  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Ghimire R, Nepal; Nakaji K, Japan

**Received:** September 16, 2022

**Peer-review started:** September 16, 2022

**First decision:** November 11, 2022

**Revised:** November 21, 2022

**Accepted:** December 5, 2022

**Article in press:** December 5, 2022

**Published online:** December 26, 2022



**Yu Chen, Chen-Guang Zhang, Lin-Lin Song**, Department of Emergency, Beijing Tsinghua Changgung Hospital, Beijing 102218, China

**Dian Chen**, Department of Dermatology, Beijing Tsinghua Changgung Hospital, Beijing 102218, China

**Hao Liu**, Department of Pathology, Beijing Tsinghua Changgung Hospital, Beijing 102218, China

**Corresponding author:** Lin-Lin Song, PhD, Chief Physician, Doctor, Department of Emergency, Beijing Tsinghua Changgung Hospital, No. 168 Litang Road, Changping District, Beijing 102218, China. [slla01139@btch.edu.cn](mailto:slla01139@btch.edu.cn)

## Abstract

### BACKGROUND

*Staphylococcus aureus* bacteraemia (SAB) is among the leading causes of bacteraemia and infectious endocarditis. The frequency of infectious endocarditis (IE) among SAB patients ranges from 5% to 10%-12%. In adults, the characteristics of epidermolytic hyperkeratosis (EHK) include hyperkeratosis, erosions, and blisters. Patients with inflammatory skin diseases and some diseases involving the epidermis tend to exhibit a disturbed skin barrier and tend to have poor cell-mediated immunity.

### CASE SUMMARY

We describe a case of SAB and infective endocarditis in a 43-year-old male who presented with fever of unknown origin and skin diseases. After genetic tests, the skin disease was diagnosed as EHK.

### CONCLUSION

A breached skin barrier secondary to EHK, coupled with inadequate sanitation, likely provided the opportunity for bacterial seeding, leading to IE and deep-seated abscess or organ abscess. EHK may be associated with skin infection and multiple risk factors for extracutaneous infections. Patients with EHK should be treated early to minimize their consequences. If patients with EHK present with prolonged fever of unknown origin, IE and organ abscesses should be ruled out, including metastatic spreads.

**Key Words:** *Staphylococcus aureus* bacteremia; Infective endocarditis; Epidermolytic

hyperkeratosis; Case report

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Emergency physicians often encounter patients with fever of unknown origin, some of who present with skin diseases. We hope the case can heighten awareness that, in patients with epidermolytic hyperkeratosis or other skin diseases presented with prolonged pyrexia, infectious endocarditis, *Staphylococcus aureus* bacteraemia and organ abscess could be identified and treated early to minimize the consequences and avoid further life-threatening episodes.

**Citation:** Chen Y, Chen D, Liu H, Zhang CG, Song LL. *Staphylococcus aureus* bacteremia and infective endocarditis in a patient with epidermolytic hyperkeratosis: A case report. *World J Clin Cases* 2022; 10(36): 13418-13425

**URL:** <https://www.wjgnet.com/2307-8960/full/v10/i36/13418.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v10.i36.13418>

## INTRODUCTION

Epidermolytic hyperkeratosis (EHK), originally termed bullous congenital ichthyosiform erythroderma, is a rare autosomal dominant disorder caused by mutations in keratins 1 and 10. It has characteristics of erythema, blistering, erosions and skin denudation present at birth and develop marked hyperkeratosis in adults[1,2]. EHK is easy to distinguish from other congenital ichthyoses through its pathological features[2]. These manifestations are due to mutations in genes mostly involved in skin barrier formation. When the skin barrier function is significantly impaired, it can lead to hypernatremic dehydration, impaired thermoregulation, increased risk for electrolyte imbalances and infection. A breached skin barrier secondary to EHK is likely to provide the opportunity for bacterial seeding and invasion, leading to *Staphylococcus aureus* bacteraemia (SAB) and infective endocarditis, as in our case. We discuss a case of SAB and infective endocarditis in a 43-year-old man with EHK and no history of drug addiction.

## CASE PRESENTATION

### Chief complaints

A 43-year-old male presented with a 10-d history of pyrexia, fatigue, diarrhoea, and weight loss.

### History of present illness

Upon presentation, he was pyrexial at 39°C, accompanied by fatigue, confusion, and diarrhoea.

### History of past illness

He regarded his skin disease as psoriasis since childhood, with recurrently diffuse palmoplantar hyperkeratosis, erythema and hypernatremia on the flexor surfaces of both arms. He was not on regular medications and had never sought medical advice about his hyperkeratosis. He had never smoked, did not drink alcohol, and had no history of drug addiction. The patient had never received professional dermatological treatment.

### Personal and family history

The patient had no significant prior family history.

### Physical examination

The patient was pyrexial at 39°C with a blood pressure of 132/78 mmHg on admission. His hyperkeratosis affected his upper and lower limbs as diffuse palmoplantar hyperkeratosis, erythema, and scales on the flexor surfaces of both arms (Figure 1). He had a systolic murmur over the mitral area but did not exhibit the meningeal irritation sign or cardiopulmonary or abdominal issues.

### Laboratory examinations

White cell counts and C-reactive protein levels were elevated to  $17.07 \times 10^9$  cells/L and above 250 mg/L, respectively. His procalcitonin levels were 3.38 ng/mL. His tests for human immunodeficiency virus





DOI: 10.12998/wjcc.v10.i36.13418 Copyright ©The Author(s) 2022.

**Figure 1 Images of the skin lesions.** Diffuse palmoplantar hyperkeratosis, erythema, and scales on the flexor surfaces of both arms and lower limbs. A: Palms; B: Opisthenar; C: Arms; D: Planta pedis; E: Dorsum of the foot.

(HIV) and syphilis antibodies were negative. His electrocardiogram showed sinus tachycardia. Blood cultures isolated methicillin-sensitive *Staphylococcus aureus* (MSSA) 2 and 4 d after admission to the hospital. Transthoracic echocardiography indicated mitral valve disease with severe insufficiency.

### Imaging examinations

Cranial computed tomography revealed multiple low-density shadows in the brain (Figure 2). The computed tomography (CT) results of the chest were normal. Abdominal enhanced CT showed multiple low-density shadows in the spleen, which were considered to be splenic abscesses with subcapsular effusion (Figure 2). Given the persistent fever, bacteraemia, abscesses, and mitral valve insufficiency, we conducted transoesophageal echocardiography (Figure 3), which revealed a wart in the mitral valve with prolapse along with severe insufficiency. He was diagnosed with definitive infective endocarditis, clinically suspected to be attributable to his skin lesions. He was urgently referred to a dermatologist, who advised treatment with tretinoin, urea cream for external use, skin biopsy (Figure 4), and genetic tests. Later, a pathological examination was conducted on his skin, which showed epidermal hyperkeratosis, acantholysis, and lymphocytes infiltrating the superficial dermis around the blood vessels and adjuncts. Genetic tests revealed mutations in the keratin 1 (*KRT1*) gene (Figure 4).

## FINAL DIAGNOSIS

The final diagnosis of the presented case was EHK combined with infective endocarditis (IE).


## TREATMENT

He was started on intravenous 0.5 g levofloxacin once a day (QD). Four days later, levofloxacin was escalated to vancomycin 500 mg Q12 h due to a positive blood culture for Gram-positive coccal bacteraemia and persistent fever. When infective endocarditis and brain abscesses are diagnosed, the selection of antibiotics depends on drug sensitivity experiments; based on the results, his treatment was changed to 2 g ceftriaxone QD.

## OUTCOME AND FOLLOW-UP


Cardiac surgery was planned for approximately 6-8 wk after completion of the course of antibiotics. Because of the use of antibiotics, his white cell counts and C-reactive protein levels remained within the






DOI: 10.12998/wjcc.v10.i36.13418 Copyright ©The Author(s) 2022.

**Figure 2 Computed tomography.** A and B: Cranial computed tomography showing multiple low-density shadows (blue arrow, A and B) in the brain; C and D: Abdominal enhanced CT scan showing multiple low-density shadows (green arrow, C and D) in the spleen, considered a splenic abscess with subcapsular effusion.



DOI: 10.12998/wjcc.v10.i36.13418 Copyright ©The Author(s) 2022.

**Figure 3 Transesophageal echocardiography.** A and B: The anterior leaflet of the mitral valve was detected with a wart (white arrow) of a cord-like medium echoic substance about 10mm and the posterior leaflet was detected with the wart (white arrow) of a medium echoic substance about 7 mm × 7 mm; C and D: There was severe mitral regurgitation and the regurgitation bundle was distributed along the posterior leaflet of the mitral valve. There was mild tricuspid valve and the aortic valve regurgitation.



**Figure 4 Histopathological, and genetic features of the patient.** A and B: The c.539A>G mutation was detected in KRT1; C: The pathological examination indicated epidermal hyperkeratosis, acantholysis and lymphocytes infiltrating the superficial dermis around blood vessels and adjuncts (H&E stain, original magnification 100×); D: c.539A>G detection using Next Generation Sequencing and Sanger sequencing.

normal range. Subsequent echocardiograms still showed a wart in the mitral valve with prolapse and severe insufficiency, but the valve excrescences appeared to be getting smaller. CT of the chest was repeated, which revealed a small pleural effusion. Cranial CT was also repeated, which indicated shrinking lesions. Blood cultures were repeated and were negative, and he remained afebrile. Because of his medical insurance policy, the patient decided to return to his local hospital for further treatment.

**DISCUSSION**

The clinical manifestations, histopathological findings, and mutations in the KRT1 gene observed in this patient were consistent with the diagnostic criteria of EHK[3,4], a rare genodermatosis with autosomal-dominant inheritance[5]. EHK is associated with erythema, blistering, erosions and skin denudation present at birth, and develop marked hyperkeratosis in adult. The KRT1 mutation c.539A>G: p. E180G was found; this has been reported previously[6].

Patients with skin diseases often exhibit a disturbed skin barrier and tend to have poor cell-mediated immunity, such as atopic dermatitis, irritant contact dermatitis, ichthyosis, rosacea, and acne[7,8]. Patients with EHK, especially with subtypes of ichthyosis, are prone to repeated episodes of skin infections, including bacterial infection and fungal infection, but the mechanism is still elusive[4,9]. Several factors have been proposed, including disruption of skin barrier function, hyperkeratotic plaques and defective cell-mediated immunity[10-12]. *Staphylococcus aureus* is a leading cause of human bacterial infection, and the most common site of *Staphylococcus aureus* infection is the skin[13]. Skin and soft tissue infections with *Staphylococcus* remain a dominant cause of bacteraemia and IE[14,15].

The definition of SAB is as presence of  $\geq 1$  positive blood cultures for *Staphylococcus aureus*[16]. The most frequent site is the anterior nares, but the skin, axilla, oropharynx, perineum, and vagina may also be colonized. These colonized sites may serve as reservoirs for future infections. Colonization is relatively higher among patients with insulin-dependent diabetes, skin damage, HIV infection, and in patients undergoing maintenance haemodialysis[16]. SAB can lead to seeding and invasion of virtually any body site and associated complications[17]. The case fatality rates for SAB have only slightly improved in recent decades[18]. Patients with SAB can develop all kinds of complications, such as IE, epidural abscess, vertebral osteomyelitis, brain abscess, and discitis, which may be difficult to identify and can lead to high rates of disability and mortality[17,19].

SAB is a leading cause of bacteraemia and infective endocarditis. Only a minority of bacteraemic patients will show involvement of the heart valves. The frequency of endocarditis among SAB patients ranges from 5% to 10%-12%[20]. IE is a severe complication in patients with nosocomial SAB. It has a low incidence but high mortality. Therefore, prevention and early detection of IE are important. The incidence is highest in those with a previous history of IE or who have prosthetic, repaired valves, intravenous drug use or rheumatic heart disease[21,22]. Our patient had no history of these risk factors except EHK.

## CONCLUSION

A case of atopic dermatitis, IE and multiple cerebral infarctions has been reported[23], but there are no previous reports of EHK and IE, such as in our case. Some factors have been speculated to increase the risk of infection, such as including disruption of skin barrier integrity, defective cell-mediated immunity, and delayed keratin scaling. In our case, a breached skin barrier secondary to EHK, coupled with inadequate skin sanitation, likely provided the opportunity for bacterial seeding by MSSA, triggering IE and abscesses. EHK may be associated with skin infection and multiple risk factors for extracutaneous infections. Patients with EHK should be treated early to minimize the consequences. If patients with EHK present with prolonged fever, IE and organ abscesses should be suspected or checked, including metastatic spreads. In conclusion, we highlight that in the absence of thorough treatment, clinicians should recognize that patients with EHK are susceptible to bacterial infections owing to disruption of the skin barrier.

## ACKNOWLEDGEMENTS

We would like to thank the patient and his family. We extend our thanks to the dermatology, echocardiography, radiology, and pathology departments of the Beijing Tsinghua Changgung Hospital for facilitating the acquisition of the relevant materials.

## FOOTNOTES

**Author contributions:** Chen Y and Song LL analyzed the data and wrote the manuscript; Liu H and Chen D contributed new pathological datum and analytic tools; Zhang CG designed the research study; All authors have read and approve the final manuscript.

**Informed consent statement:** All study participants provided informed written consent prior to study enrollment.

**Conflict-of-interest statement:** We declare that we have no financial and personal relationships with other people or organizations that can inappropriately influence our work, there is no professional or other personal interest of any nature or kind in any product, service and/or company that could be construed as influencing the position presented in, or the review of, the manuscript entitled.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by

external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Lin-Lin Song 0000-0001-6398-6725.

**S-Editor:** Liu JH

**L-Editor:** A

**P-Editor:** Liu JH

## REFERENCES

- Miao H, Dong R, Zhang S, Yang L, Liu Y, Wang T. Inherited ichthyosis and fungal infection: an update on pathogenesis and treatment strategies. *J Dtsch Dermatol Ges* 2021; **19**: 341-350 [PMID: 33448147 DOI: 10.1111/ddg.14389]
- Rice AS, Crane JS. Epidermolytic Hyperkeratosis. [Updated 2022 Aug 1]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK544323/>
- Kurosawa M, Takagi A, Tamakoshi A, Kawamura T, Inaba Y, Yokoyama K, Kitajima Y, Aoyama Y, Iwatsuki K, Ikeda S. Epidemiology and clinical characteristics of bullous congenital ichthyosiform erythroderma (keratinolytic ichthyosis) in Japan: results from a nationwide survey. *J Am Acad Dermatol* 2013; **68**: 278-283 [PMID: 23182068 DOI: 10.1016/j.jaad.2012.06.044]
- Peter Rout D, Nair A, Gupta A, Kumar P. Epidermolytic hyperkeratosis: clinical update. *Clin Cosmet Investig Dermatol* 2019; **12**: 333-344 [PMID: 31190940 DOI: 10.2147/CCID.S166849]
- Hayashida MT, Mitsui GL, Reis NI, Fantinato G, Jordão Neto D, Mercante AM. Epidermolytic Hyperkeratosis--case report. *An Bras Dermatol* 2015; **90**: 888-891 [PMID: 26734873 DOI: 10.1590/abd1806-4841.20153966]
- Sashikawa M, Tsuda H, Komine M, Ohtsuki M. Novel missense mutation c.539A>G; p.Glu180Gly in keratin 1 causing epidermolytic ichthyosis. *J Dermatol* 2021; **48**: e579-e580 [PMID: 34486163 DOI: 10.1111/1346-8138.16142]
- Proksch E. pH in nature, humans and skin. *J Dermatol* 2018; **45**: 1044-1052 [PMID: 29863755 DOI: 10.1111/1346-8138.14489]
- Takahashi M, Hagiya H, Tanaka S, Yamamoto K, Honda H, Hasegawa K, Otsuka F. Persistent methicillin-resistant *Staphylococcus aureus* bacteremia in an adult patient with Netherton's syndrome: A case report. *J Infect Chemother* 2022; **28**: 978-981 [PMID: 35277342 DOI: 10.1016/j.jiac.2022.02.021]
- DiGiovanna JJ, Bale SJ. Clinical heterogeneity in epidermolytic hyperkeratosis. *Arch Dermatol* 1994; **130**: 1026-1035 [PMID: 8053700]
- Vyas NS, Kannan S, N Jahnke M, Hu RH, Choate KA, Shwayder TA. Congenital Ichthyosiform Erythroderma Superimposed with Chronic Dermatophytosis: A Report of Three Siblings. *Pediatr Dermatol* 2016; **33**: e6-e9 [PMID: 26645853 DOI: 10.1111/pde.12728]
- Hoetzenecker W, Schanz S, Schaller M, Fierlbeck G. Generalized tinea corporis due to *Trichophyton rubrum* in ichthyosis vulgaris. *J Eur Acad Dermatol Venereol* 2007; **21**: 1129-1131 [PMID: 17714151 DOI: 10.1111/j.1468-3083.2006.02117.x]
- Schmuth M, Yosipovitch G, Williams ML, Weber F, Hintner H, Ortiz-Urda S, Rappersberger K, Crumrine D, Feingold KR, Elias PM. Pathogenesis of the permeability barrier abnormality in epidermolytic hyperkeratosis. *J Invest Dermatol* 2001; **117**: 837-847 [PMID: 11676820 DOI: 10.1046/j.0022-202x.2001.01471.x]
- Becker RE, Bubeck Wardenburg J. *Staphylococcus aureus* and the skin: a longstanding and complex interaction. *Skinmed* 2015; **13**: 111-9; quiz 120 [PMID: 26137737]
- Al-Bayati A, Alshami A, AlAzzawi M, Al Hillan A, Hossain M. Metastatic Osteoarticular Infective Endocarditis by Methicillin-sensitive *Staphylococcus Aureus*. *Cureus* 2020; **12**: e8124 [PMID: 32542174 DOI: 10.7759/cureus.8124]
- Salvador VB, Chapagain B, Joshi A, Brennessel DJ. Clinical Risk Factors for Infective Endocarditis in *Staphylococcus aureus* Bacteremia. *Tex Heart Inst J* 2017; **44**: 10-15 [PMID: 28265207 DOI: 10.14503/THIJ-15-5359]
- Rongpharpi SR, Duggal S, Kalita H, Duggal AK. *Staphylococcus aureus* bacteremia: targeting the source. *Postgrad Med* 2014; **126**: 167-175 [PMID: 25295661 DOI: 10.3810/pgm.2014.09.2811]
- Keynan Y, Rubinstein E. *Staphylococcus aureus* bacteremia, risk factors, complications, and management. *Crit Care Clin* 2013; **29**: 547-562 [PMID: 23830653 DOI: 10.1016/j.ccc.2013.03.008]
- Holland TL, Arnold C, Fowler VG Jr. Clinical management of *Staphylococcus aureus* bacteremia: a review. *JAMA* 2014; **312**: 1330-1341 [PMID: 25268440 DOI: 10.1001/jama.2014.9743]
- Veronese C, Pellegrini M, Maiolo C, Morara M, Armstrong GW, Ciardella AP. Multimodal ophthalmic imaging of *staphylococcus aureus* bacteremia associated with chorioretinitis, endocarditis, and multifocal brain abscesses. *Am J Ophthalmol Case Rep* 2020; **17**: 100577 [PMID: 31886435 DOI: 10.1016/j.ajoc.2019.100577]
- Kern WV. Management of *Staphylococcus aureus* bacteremia and endocarditis: progresses and challenges. *Curr Opin Infect Dis* 2010; **23**: 346-358 [PMID: 20592532 DOI: 10.1097/QCO.0b013e32833bccc8a]
- Thornhill MH, Jones S, Prendergast B, Baddour LM, Chambers JB, Lockhart PB, Dayer MJ. Quantifying infective endocarditis risk in patients with predisposing cardiac conditions. *Eur Heart J* 2018; **39**: 586-595 [PMID: 29161405 DOI: 10.1093/eurheartj/ehx655]
- Cahill TJ, Prendergast BD. Infective endocarditis. *Lancet* 2016; **387**: 882-893 [PMID: 26341945 DOI: 10.1016/S0140-6736(15)00067-7]

- 23 **Furumura Y**, Nishida M, Imanishi A, Maekawa N, Fukai K. Infectious endocarditis with multiple cerebral infarctions in a patient with severe atopic dermatitis. *J Dermatol* 2019; **46**: e353-e354 [PMID: [31090225](#) DOI: [10.1111/1346-8138.14924](#)]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

