

World Journal of *Clinical Cases*

World J Clin Cases 2023 April 6; 11(10): 2123-2362



Contents

Thrice Monthly Volume 11 Number 10 April 6, 2023

EVIDENCE REVIEW

- 2123 Fractional flow reserve and non-hyperemic indices: Essential tools for percutaneous coronary interventions

Boutaleb AM, Ghafari C, Ungureanu C, Carlier S

REVIEW

- 2140 Diagnosis, treatment protocols, and outcomes of liver transplant recipients infected with COVID-19

Hashem M, El-Kassas M

MINIREVIEWS

- 2160 Treatment of stellate ganglion block in diseases: Its role and application prospect

Deng JJ, Zhang CL, Liu DW, Huang T, Xu J, Liu QY, Zhang YN

- 2168 Clinical application of SARS-CoV-2 antibody detection and monoclonal antibody therapies against COVID-19

Sun J, Yang ZD, Xie X, Li L, Zeng HS, Gong B, Xu JQ, Wu JH, Qu BB, Song GW

- 2181 Cheesy material on macroscopic on-site evaluation after endoscopic ultrasound-guided fine-needle biopsy: Don't miss the tuberculosis

Delsa H, Bellahammou K, Okasha HH, Ghalim F

- 2189 Liver manifestations in COVID-19 patients: A review article

Helou M, Nasr J, El Osta N, Jabbour E, Husni R

- 2201 Breast reconstruction: Review of current autologous and implant-based techniques and long-term oncologic outcome

Malekpour M, Malekpour F, Wang HTH

- 2213 Update on the current management of persistent and recurrent primary hyperparathyroidism after parathyroidectomy

Pavlidis ET, Pavlidis TE

ORIGINAL ARTICLE

Retrospective Study

- 2226 Hepatobiliary system and intestinal injury in new coronavirus infection (COVID-19): A retrospective study

Kozlov KV, Zhdanov KV, Ratnikova AK, Ratnikov VA, Tishkov AV, Grinevich V, Kravchuk YA, Miklush PI, Nikiforova PO, Gordienko VV, Popov AF, Andryukov BG

- 2237 Impact of lockdown policies during the COVID-19 outbreak on a trauma center of a tertiary hospital in China

Shen BS, Cheng WY, Liang ZR, Tang Q, Li KY

Observational Study

- 2246** Interaction between the left ventricular ejection fraction and left ventricular strain and its relationship with coronary stenosis
Gui HY, Liu SW, Zhu DF

CASE REPORT

- 2254** Neonatal hyperinsulinism with an *ABCC8* mutation: A case report
Liu MT, Yang HX
- 2260** Unilateral contrast-induced encephalopathy with contrast medium exudation: A case report
Zhang ZY, Lv H, Wang PJ, Zhao DY, Zhang LY, Wang JY, Hao JH
- 2267** Diagnosis and treatment of primary seminoma of the prostate: A case report and review of literature
Cao ZL, Lian BJ, Chen WY, Fang XD, Jin HY, Zhang K, Qi XP
- 2276** Primary intra-abdominal paraganglioma: A case report
Guo W, Li WW, Chen MJ, Hu LY, Wang XG
- 2282** Successful surgical treatment of bronchopleural fistula caused by severe pulmonary tuberculosis: A case report and review of literature
Shen L, Jiang YH, Dai XY
- 2290** Clinical and genetic features of Kenny-Caffey syndrome type 2 with multiple electrolyte disturbances: A case report
Yuan N, Lu L, Xing XP, Wang O, Jiang Y, Wu J, He MH, Wang XJ, Cao LW
- 2301** Dupilumab for treatment of severe atopic dermatitis accompanied by lichenoid amyloidosis in adults: Two case reports
Zhao XQ, Zhu WJ, Mou Y, Xu M, Xia JX
- 2308** Reabsorption of intervertebral disc prolapse after conservative treatment with traditional Chinese medicine: A case report
Wang CA, Zhao HF, Ju J, Kong L, Sun CJ, Zheng YK, Zhang F, Hou GJ, Guo CC, Cao SN, Wang DD, Shi B
- 2315** Development of subdural empyema from subdural effusion after suppurative encephalitis: A case report
Yang RX, Chen B, Zhang Y, Yang Y, Xie S, He L, Shi J
- 2321** Treatment of periprosthetic knee infection and coexistent periprosthetic fracture: A case report and literature review
Hao LJ, Wen PF, Zhang YM, Song W, Chen J, Ma T
- 2329** Formation of a rare curve-shaped thoracolith documented on serial chest computed tomography images: A case report
Hsu FC, Huang TW, Pu TW
- 2336** Neurofibromatosis type 1 with multiple gastrointestinal stromal tumors: A case report
Yao MQ, Jiang YP, Yi BH, Yang Y, Sun DZ, Fan JX

- 2343** Coexisting cytomegalovirus colitis in an immunocompetent patient with *Clostridioides difficile* colitis: A case report
Kim JH, Kim HS, Jeong HW
- 2349** Paradoxical vocal fold motion masquerading as post-anesthetic respiratory distress: A case report
Baek J, Jee DL, Choi YS, Kim SW, Choi EK
- 2355** Full neurological recovery from severe nonexertional heat stroke with multiple organ dysfunction: A case report
Du F, Zheng JW, Zhao YB, Yang K, Li HN

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Abdullah Gul, MD, PhD, Associate Professor, Department of Urology, University of Health Sciences Turkey, Bursa Yuksek Ihtisas Training and Research Hospital, Bursa 16310, Turkey. dr_abdullahgul@hotmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Xu Guo*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

April 6, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Primary intra-abdominal paraganglioma: A case report

Wei Guo, Wei-Wei Li, Min-Jie Chen, Ling-Yu Hu, Xiao-Guang Wang

Specialty type: Surgery

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B, B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Lucke-Wold B, United States; Maurea S, Italy

Received: December 26, 2022

Peer-review started: December 26, 2022

First decision: January 20, 2023

Revised: January 29, 2023

Accepted: March 3, 2023

Article in press: March 3, 2023

Published online: April 6, 2023



Wei Guo, Graduate School, Zhejiang Chinese Medicine University, Hangzhou 310053, Zhejiang Province, China

Wei Guo, Department of General Surgery, Jiaying The Second Hospital, Jiaying 314000, Zhejiang Province, China

Wei-Wei Li, Min-Jie Chen, Ling-Yu Hu, Xiao-Guang Wang, Department of Hepatobiliary Surgery, Jiaying The Second Hospital, Jiaying 314000, Zhejiang Province, China

Corresponding author: Xiao-Guang Wang, Doctor, Academic Editor, Doctor, Department of Hepatobiliary Surgery, Jiaying The Second Hospital, No. 1518 Huancheng North Road, Nanhu District, Jiaying 314000, Zhejiang Province, China. xiaoguangwangs@163.com

Abstract

BACKGROUND

Paragangliomas are rare neuroendocrine tumors. We hereby report a case of a localized paraganglioma found in the abdominal cavity, and review the relevant literature to improve the understanding of this disease.

CASE SUMMARY

A 29-year-old Chinese female patient was referred to our hospital due to an abdominal mass found on physical examination. Imaging revealed a mass in the left upper abdomen, suggestive of either a benign stromal tumor or an ectopic accessory spleen. Laparoscopic radical resection was subsequently performed, and histopathological analysis confirmed the diagnosis of a paraganglioma. The patient was followed up 3 months post-operation, and reported good recovery with no metastasis.

CONCLUSION

Radical resection can effectively treat intra-abdominal paragangliomas, with few side effects and low recurrence risk. In addition, early and accurate diagnosis and timely intervention are essential for the prognosis of this disease.

Key Words: Abdominal cavity; Paraganglioma; Surgical treatment; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Intraperitoneal paraganglioma is clinically rare without characteristic imaging findings, and many clinicians may never encounter it. However, clinicians must remain vigilant to suspect, identify, locate, and remove the tumor, as the associated symptoms and hypertension can be cured by surgical resection. If the tumor is not diagnosed and removed, there is a risk of death and heart disease. Therefore, due to the small number of cases, the lack of understanding of its clinical features and imaging signs, especially easy to miss diagnosis.

Citation: Guo W, Li WW, Chen MJ, Hu LY, Wang XG. Primary intra-abdominal paraganglioma: A case report. *World J Clin Cases* 2023; 11(10): 2276-2281

URL: <https://www.wjgnet.com/2307-8960/full/v11/i10/2276.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i10.2276>

INTRODUCTION

Paragangliomas are rare neuroendocrine tumors that can occur at any age. It represents an important cause of secretory hypertension as it is often characterized by the excessive production of catecholamines. This study aimed to report a case of an intra-abdominal paraganglioma, and improve the understanding of this disease by elucidating the clinical features of the patient, while reviewing the relevant literature.

CASE PRESENTATION

Chief complaints

A 29-year-old woman was referred to the Department of General Surgery of the Second Hospital of Jiaxing on July 20, 2022 following findings of an abdominal mass 5 d ago.

History of present illness

Initial abdominal computed tomography (CT) revealed soft tissue nodules with central calcification in the left upper abdomen, which warranted further imaging with contrast-enhanced CT.

History of past illness

The patient had no surgical or tumor history. Besides a 2-year history of Hashimoto's thyroiditis and hyperthyroidism, no other underlying diseases such as hypertension and diabetes were reported.

Personal and family history

The patient denied any family history of cancer.

Physical examination

Physical examination was grossly unremarkable. No gastrointestinal symptoms such as abdominal pain, distension, nausea and vomiting, jaundice, abnormal bowel movements, or abnormal stool forms were noted. Urinary symptoms such as gross hematuria, frequency, urgency, and dysuria were not noted as well. The patient was not pyrexia, and denied any chest tightness or shortness of breath.


Laboratory examinations

Laboratory results were as follows: Thyroid stimulating hormone, 0.003 μ IU/mL; anti-thyroglobulin, 282.92 IU/mL, anti-thyroid peroxidase antibody, 149.91 IU/mL (reference range, 0–34 IU/mL); and CA19-9, 2.0 U/mL (reference range 0–37 U/mL). The remaining laboratory test results were otherwise unremarkable.

Pathological examination of the lesion suggested a paraganglioma (Figure 1). Immunohistochemical staining was subsequently performed, which demonstrated the following: Syn (-), CgA (-), CD56 (-), Ki-67 (+ 5%), CD10 (-), S-100 (+), NES (+), AE1/AE3 (-), SOX10 (-), and P53 (wild type). A diagnosis of an intra-abdominal paraganglioma was thus confirmed.

Imaging examinations

Contrast-enhanced CT confirmed a mass in the left upper abdomen. At this stage, a benign stromal tumor of mesenchymal origin or an ectopic accessory spleen was considered (Figure 2).



DOI: 10.12998/wjcc.v11.i10.2276 Copyright ©The Author(s) 2023.

Figure 1 Pathological result suggestive of a paraganglioma (HE × 400).



DOI: 10.12998/wjcc.v11.i10.2276 Copyright ©The Author(s) 2023.

Figure 2 Contrast-enhanced computed tomography showing a mass in the left upper abdomen, indicating either a benign mesenchymal tumor or an ectopic accessory spleen.

FINAL DIAGNOSIS

The patient was diagnosed with a primary intra-abdominal paraganglioma.

TREATMENT

Laparoscopic radical resection of the abdominal mass was indicated. During the operation, the mass was found adhered to the surrounding omentum in the left upper abdominal region, in close proximity to the spleen, stomach, small intestine, and colon. The lesion was approximately 22 mm × 26 mm in size. Postoperative chemotherapy was not indicated.

OUTCOME AND FOLLOW-UP

The patient recovered well without any discomfort. At 3 mo follow-up, no complications such as recurrence or metastasis were observed.

DISCUSSION

Paragangliomas are rare neuroendocrine tumors originating from either the adrenal medulla, or the extra-adrenal sympathetic and parasympathetic ganglia. Symptoms involve the classic triad of headache, palpitation, and sweating. Adrenal and extra-adrenal sympathetic paragangliomas are often characterized by the excessive production of catecholamines, which can not only result in hypertension, but also associate with the risk of acute cardiovascular complications. Diagnosis is often based on plasma or urinary metanephrine measurements and nuclear imaging. Moreover, normal catecholamine levels have been reported to virtually exclude the presence of a sympathetic paraganglioma[1]. Intra-abdominal paragangliomas are particularly rare, with incidence of approximately 1 in 500000.

Parasympathetic paragangliomas are usually located in the head and neck, while sympathetic paragangliomas are more commonly located in the abdomen, followed by the chest and pelvis[2]. And the tumor that may present with cranial neuropathies when located along the skull base[3]. Head and neck paragangliomas are usually painless and slow growing, and are thus often an incidental clinical finding. They mainly manifest as carotid body tumors and vagal paragangliomas. As parasympathetic paragangliomas are non-secretory, symptoms are usually secondary to mass effects. These may include neck pain and dysphagia, conductive hearing loss and pulsatile tinnitus in cervical tympanic paragangliomas, as well as lower cranial nerve defects in advanced tumors[4]. Paragangliomas located outside of the head and neck may also be non-secretory, and are often accompanied by mild symptoms[5].

Due to the catecholamine-secreting nature of abdominal paragangliomas, the common clinical symptoms may include malignant hypertension, palpitation, headache, dizziness, anxiety, metabolic disorder syndrome, orthopnea, oliguria, anuria, and hepatic encephalopathy, among others. In rare cases, patients may experience paraganglioma crises, an endocrine emergency resulting in life-threatening hemodynamic instability and end-organ damage[6]. This can often be misdiagnosed as septic shock, heart failure, thyroid storm, and malignant hyperthermia[7,8]. Given its mortality rate of approximately 15%, early recognition of the signs and symptoms of paragangliomas is thereby critical[7,9]. It was found that in patients with paraganglioma, the metaiodobenzylguanidine (MIBG) uptake intensity ratio was significantly higher in malignant lesions than in benign lesions. Therefore, iodine-131 MIBG uptake was able to distinguish between benign and malignant diseases, which was not helped by magnetic resonance imaging[10]. Furthermore, one study used radio-labeled MIBG and somatostatin analogues for scintillation imaging for correct localization. The results showed that MIBG was more accurate in imaging pheochromocytoma than somatostatin analogues. But somatostatin analogues are more accurate than MIBGs in detecting neuroendocrine tumors[11]. While symptoms are often paroxysmal, the clinical manifestations of paragangliomas can vary based on catecholamine subtypes, and may range from asymptomatic to life-threatening[12]. As exemplified in our case, the lack of obvious clinical abnormalities such as abdominal pain, abdominal distension, hypertension, and dizziness could have led to misdiagnosis or missed diagnosis.

Paragangliomas are mostly benign in nature, with surgical resection being the main treatment of choice. In contrast, malignant paragangliomas often warrant a multidisciplinary approach, involving endocrinology, oncology, surgery, nuclear medicine, radiotherapy, interventional radiology, and histology. So far, there are no standardized treatment regimens for metastatic diseases. Current treatment measures mainly involve beta-blockers and catecholamine synthesis inhibitors (A-methyl-p-tyrosine) to prevent tumor progression and minimize catecholamine-induced symptoms[13].

The diagnosis of paraganglioma in our case was mainly based on pathology, and was confirmed upon findings of Syn (-) and S-100 (+) on immunohistochemical staining. While morphologically similar to malignant perivascular epithelioid cell and stromal tumors, S-100 played a role as a differentiating factor. As a neurogenic index, the positive expression of S-100 observed in our patient was in keeping with the origin of paragangliomas from chromaffin cells of the neural crest. In contrast, perivascular epithelioid cell tumors are mostly malignant in nature, and are characterized by positive HMB45, SMA, and Desmin, the latter 2 of which are myogenic indices. Stromal tumors are also commonly malignant, and are distinguished by the expression of CD117. In our case, the low expression of Ki-67 indicated a benign tumor.

CONCLUSION

In conclusion, intra-abdominal paragangliomas are clinically rare with no characteristic imaging findings, and are, as such, easily missed. However, surgical resection can associate with good clinical prognosis.

ACKNOWLEDGEMENTS

We thank the patient for his contribution to this case report. We also thank the Department of General Surgery of the Second Affiliated Hospital of Jiaxing University for their support with the treatment of

this case.

FOOTNOTES

Author contributions: Li WW and Chen MJ contributed to the treatment of case and data collection; Guo W, Hu LY, and Wang XG contributed to the writing and revising the manuscript; All authors have read and approve the final version of manuscript.

Supported by Jiaxing Science and Technology Planning Project, No. 2020AY30017.

Informed consent statement: Informed written consent was obtained from the patient for publication of this report and any accompanying images.

Conflict-of-interest statement: All the authors declare that they have no conflict of interest.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Wei Guo 0000-0003-4479-4124; Ling-Yu Hu 0000-0002-5698-7700; Xiao-Guang Wang 0000-0002-5698-8800.

S-Editor: Liu JH

L-Editor: A

P-Editor: Liu JH

REFERENCES

- 1 Cornu E, Belmihoub I, Burnichon N, Grataloup C, Zinzindohoué F, Baron S, Billaud E, Azizi M, Gimenez-Roqueplo AP, Amar L. [Pheochromocytoma and paraganglioma]. *Rev Med Interne* 2019; **40**: 733-741 [PMID: 31493938 DOI: 10.1016/j.revmed.2019.07.008]
- 2 Lenders JW, Duh QY, Eisenhofer G, Gimenez-Roqueplo AP, Grebe SK, Murad MH, Naruse M, Pacak K, Young WF Jr; Endocrine Society. Pheochromocytoma and paraganglioma: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab* 2014; **99**: 1915-1942 [PMID: 24893135 DOI: 10.1210/jc.2014-1498]
- 3 Moor R, Goutnik M, Lucke-Wold B, Laurent D, Chen S, Friedman W, Rahman M, Allen N, Rivera-Zengotita M, Koch M. Clival Paraganglioma, Case Report and Literature Review. *OBM Neurobiol* 2022; **6** [PMID: 35844205 DOI: 10.21926/obm.neurobiol.2203128]
- 4 Smith JD, Harvey RN, Darr OA, Prince ME, Bradford CR, Wolf GT, Else T, Basura GJ. Head and neck paragangliomas: A two-decade institutional experience and algorithm for management. *Laryngoscope Invest Otolaryngol* 2017; **2**: 380-389 [PMID: 29299512 DOI: 10.1002/lto2.122]
- 5 Fishbein L. Pheochromocytoma/Paraganglioma: Is This a Genetic Disorder? *Curr Cardiol Rep* 2019; **21**: 104 [PMID: 31367972 DOI: 10.1007/s11886-019-1184-y]
- 6 Meijs AC, Snel M, Corssmit EPM. Pheochromocytoma/paraganglioma crisis: case series from a tertiary referral center for pheochromocytomas and paragangliomas. *Hormones (Athens)* 2021; **20**: 395-403 [PMID: 33575936 DOI: 10.1007/s42000-021-00274-6]
- 7 Whitelaw BC, Prague JK, Mustafa OG, Schulte KM, Hopkins PA, Gilbert JA, McGregor AM, Aylwin SJ. Pheochromocytoma [corrected] crisis. *Clin Endocrinol (Oxf)* 2014; **80**: 13-22 [PMID: 24102156 DOI: 10.1111/cen.12324]
- 8 Juszczak K, Drewa T. Adrenergic crisis due to pheochromocytoma - practical aspects. A short review. *Cent European J Urol* 2014; **67**: 153-155 [DOI: 10.5173/cej.2014.02.art7]
- 9 Martucci VL, Pacak K. Pheochromocytoma and paraganglioma: diagnosis, genetics, management, and treatment. *Curr Probl Cancer* 2014; **38**: 7-41 [PMID: 24636754 DOI: 10.1016/j.crrproblecancer.2014.01.001]
- 10 Maurea S, Cuocolo A, Imbriaco M, Pellegrino T, Fusari M, Cuocolo R, Liuzzi R, Salvatore M. Imaging characterization of benign and malignant pheochromocytoma or paraganglioma: comparison between MIBG uptake and MR signal intensity ratio. *Ann Nucl Med* 2012; **26**: 670-675 [PMID: 22752959 DOI: 10.1007/s12149-012-0624-1]
- 11 Lastoria S, Maurea S, Vergara E, Acampa W, Varrella P, Klain M, Muto P, Bernardy JD, Salvatore M. Comparison of labeled MIBG and somatostatin analogs in imaging neuroendocrine tumors. *Q J Nucl Med* 1995; **39**: 145-149 [PMID: 9002775]
- 12 Petrák O, Rosa J, Holaj R, Štrauch B, Krátká Z, Kvasnička J, Klímová J, Waldauf P, Hamplová B, Markvartová A, Novák

- K, Michalský D, Widimský J, Zelinka T. Blood Pressure Profile, Catecholamine Phenotype, and Target Organ Damage in Pheochromocytoma/Paraganglioma. *J Clin Endocrinol Metab* 2019; **104**: 5170-5180 [PMID: 31009053 DOI: 10.1210/jc.2018-02644]
- 13 Corssmit EPM, Snel M, Kapiteijn E. Malignant pheochromocytoma and paraganglioma: management options. *Curr Opin Oncol* 2020; **32**: 20-26 [PMID: 31599769 DOI: 10.1097/CCO.000000000000589]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

