World Journal of *Clinical Cases*

World J Clin Cases 2023 April 26; 11(12): 2582-2854





Published by Baishideng Publishing Group Inc

W J C C World Journal of Clinical Cases

Contents

Thrice Monthly Volume 11 Number 12 April 26, 2023

REVIEW

2582	Controversies in the management of acute pancreatitis: An update	
	Manrai M, Dawra S, Singh AK, Jha DK, Kochhar R	
2604	Classification of osteogenesis imperfecta: Importance for prophylaxis and genetic counseling	

Panzaru MC, Florea A, Caba L, Gorduza EV

MINIREVIEWS

- 2621 Emerging role of dual biologic therapy for the treatment of inflammatory bowel disease McCormack MD, Wahedna NA, Aldulaimi D, Hawker P
- 2631 Pancreatic cancer and depression Michoglou K, Ravinthiranathan A, San Ti S, Dolly S, Thillai K
- Mediastinal lesions in children 2637 Çinar HG, Gulmez AO, Üner Ç, Aydin S
- Exit strategies in inflammatory bowel disease: Looking beyond anti-tumor necrosis factors 2657 Crispino F, Michielan A, Grova M, Tieppo C, Mazza M, Rogger TM, Armelao F
- 2670 Medicinal cannabis products for the treatment of acute pain Fiore M, Alfieri A, Di Franco S, Petrou S, Damiani G, Pace MC
- 2677 Role of in vitamin D in irritable bowel syndrome Yu XL, Wu QQ, He LP, Zheng YF

ORIGINAL ARTICLE

Retrospective Cohort Study

2684 Analysis of oxidative stress and antioxidative potential in premature ovarian insufficiency Kakinuma K, Kakinuma T

Retrospective Study

- 2694 Surgical management of pituitary adenoma during pregnancy Jia XY, Guo XP, Yao Y, Deng K, Lian W, Xing B
- 2708 Role of pre-existing incomplete intestinal metaplasia in gastric adenocarcinoma: A retrospective case series analysis

Bogdanova I, Polaka I, Aleksandraviča I, Dzērve Z, Anarkulova L, Novika V, Tolmanis I, Leja M



Carl	World Journal of Clinical Cases
Contents Thrice Monthly Volume 11 Number 12 April 26, 20	
	Observational Study
2716	Severe/critical COVID-19 early warning system based on machine learning algorithms using novel imaging scores
	Li QY, An ZY, Pan ZH, Wang ZZ, Wang YR, Zhang XG, Shen N
2729	Mediating effect of mindfulness level on the relationship between marital quality and postpartum depression among primiparas
	Yang J, Lin XZ, Guo QW, Wang CL, Yang RY, Zhang JW, Zeng Y
2740	Ferric carboxymaltose for anemia in Crohn's disease patients at a tertiary center: A retrospective observational cohort study
	Siqueira NSN, Pascoal LB, Rodrigues BL, de Castro MM, Martins ASC, Araújo DOS, Gomes LEM, Camargo MG, Ayrizono MLS, Leal RF
	META-ANALYSIS
2753	Is metaphyseal ulnar shortening osteotomy superior to diaphyseal ulnar shortening osteotomy in the treatment of ulnar impaction syndrome? A meta-analysis
	Deng HL, Lu ML, Tang ZM, Mao QL, Zhao JM
2766	Relationship between body mass index and short-term postoperative prognosis in patients undergoing colorectal cancer surgery
	Li Y, Deng JJ, Jiang J
	CASE REPORT
2780	Cardiac amyloidosis presenting as pulmonary arterial hypertension: A case report
	Gao M, Zhang WH, Zhang ZG, Yang N, Tong Q, Chen LP
2788	Short-term outcome of total knee replacement in a patient with hemophilia: A case report and review of literature
	Yin DL, Lin JM, Li YH, Chen P, Zeng MD
2796	Modified inferior oblique anterior transposition for dissociated vertical deviation combined with superior oblique palsy: A case report
	Zong Y, Wang Z, Jiang WL, Yang X
2803	Treatment of talipes equinovarus after triceps surae intramuscular hemangioma surgery by Ilizarov technology in adults: A case report
	Chen ZX, Wang MY, Zhang C, Ding ZQ, Chen W
2811	Open surgery: Still a great option to treat patients with post-traumatic arteriovenous fistulas: A case report
	Kalinin R, Suchkov I, Mzhavanadze N, Borisova Y, Panin I

2817 Recovery from Bell's palsy after treatment using uncultured umbilical cord-derived mesenchymal stem cells: A case report

Ahn H, Jung WJ, Lee SY, Lee KH



Conton	World Journal of Clinical Cases
conten	Thrice Monthly Volume 11 Number 12 April 26, 2023
2825	Pancreatic neuroendocrine tumor detected by technetium-99m methoxy-2-isobutylisonitrile single photon emission computed tomography/computed tomography: A case report
	Liu CJ, Yang HJ, Peng YC, Huang DY
2832	Furazolidone-induced pulmonary toxicity in Helicobacter pylori infection: Two case reports
	Ye Y, Shi ZL, Ren ZC, Sun YL
2839	Efficacy of anlotinib combined with radioiodine to treat scalp metastasis of papillary thyroid cancer: A case report and review of literature
	Zhang LY, Cai SJ, Liang BY, Yan SY, Wang B, Li MY, Zhao WX
2848	Endoscopic ultrasound-guided transrectal drainage of a pelvic abscess after Hinchey II sigmoid colon diverticulitis: A case report
	Drnovšek J, Čebron Ž, Grosek J, Janež J



Contents

Thrice Monthly Volume 11 Number 12 April 26, 2023

ABOUT COVER

Editorial Board Member of World Journal of Clinical Cases, Mohamed Eltayeb Abdelrahman Naiem, MBBS, MD, Assistant Professor, Surgeon, Department of Surgery, Faculty of Medicine, University of Khartoum, Khartoum 102, Sudan. m-altayeb@live.com

AIMS AND SCOPE

The primary aim of World Journal of Clinical Cases (WJCC, World J Clin Cases) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Ying-Yi Yuan; Production Department Director: Xu Guo; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS
World Journal of Clinical Cases	https://www.wjgnet.com/bpg/gerinfo/204
ISSN	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 2307-8960 (online)	https://www.wjgnet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
April 16, 2013	https://www.wjgnet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Thrice Monthly	https://www.wjgnet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku	PUBLICATION MISCONDUCT https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
https://www.wjgnet.com/2307-8960/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS
April 26, 2023	https://www.wjgnet.com/bpg/GerInfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2023 Baishideng Publishing Group Inc	https://www.f6publishing.com

© 2023 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: bpgoffice@wjgnet.com https://www.wjgnet.com



W J C C World Journal of Clinical Cases

Submit a Manuscript: https://www.f6publishing.com

World J Clin Cases 2023 April 26; 11(12): 2796-2802

DOI: 10.12998/wjcc.v11.i12.2796

ISSN 2307-8960 (online)

CASE REPORT

Modified inferior oblique anterior transposition for dissociated vertical deviation combined with superior oblique palsy: A case report

Yao Zong, Ze Wang, Wen-Lan Jiang, Xian Yang

Specialty type: Medicine, research and experimental

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): 0 Grade C (Good): C, C Grade D (Fair): 0 Grade E (Poor): 0

P-Reviewer: Bernardes A, Portugal; Nambi G, Saudi Arabia

Received: November 15, 2022 Peer-review started: November 15, 2022 First decision: February 14, 2023 Revised: March 3, 2023 Accepted: March 22, 2023 Article in press: March 22, 2023 Published online: April 26, 2023



Yao Zong, Wen-Lan Jiang, Xian Yang, Department of Ophthalmology, The Affiliated Hospital of Qingdao University, Qingdao 266003, Shandong Province, China

Ze Wang, Department of Ophthalmology, Nanjing South East Eye Hospital, Nanjing 210007, Jiangsu Province, China

Corresponding author: Xian Yang, MD, PhD, Professor, Department of Ophthalmology, The Affiliated Hospital of Qingdao University, No. 16 Jiangsu Road, Shinan District, Qingdao 266003, Shandong Province, China. yangxian_zhao@qdu.edu.cn

Abstract

BACKGROUND

Inferior oblique anterior transposition (IOAT) has emerged as an effective surgery in the management of dissociated vertical deviation (DVD) combined with superior oblique palsy (SOP). Traditional IOAT usually provides satisfactory primary position alignment and simultaneously restricts the superior floating phenomenon. However, it also increases the risk of the anti-elevation syndrome and narrowing of the palpebral fissure in straight-ahead gaze, especially after the unilateral operation.

CASE SUMMARY

We report the outcomes of the modified unilateral IOAT in two patients with unilateral DVD combined with SOP. The anterior-nasal fibers of the inferior oblique muscle were attached at 9 mm posterior to the corneal limbus along the temporal board of the inferior rectus muscle, the other fibers were attached a further 5 mm temporal to the anterior-nasal fibers. Postoperatively, both hypertropia and floating were improved, and no obvious complications occurred.

CONCLUSION

In these cases, the modified unilateral IOAT was an effective and safe surgical method for treating DVD with SOP.

Key Words: Anterior transposition; Inferior oblique muscles; Dissociated vertical deviation; Superior oblique palsy; Anti-elevation syndrome; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.



WJCC https://www.wjgnet.com

Core Tip: Inferior oblique anterior transposition (IOAT) is effective for dissociated vertical deviation combined with superior oblique palsy; however, unilateral IOAT also increases the risk of the antielevation syndrome. This modified unilateral IOAT involves a more backward new insertion and the new insertion line perpendicular to the inferior rectus muscle axis, which provides satisfactory primary position alignment and restricts the superior floating phenomenon without obvious postoperative complications.

Citation: Zong Y, Wang Z, Jiang WL, Yang X. Modified inferior oblique anterior transposition for dissociated vertical deviation combined with superior oblique palsy: A case report. World J Clin Cases 2023; 11(12): 2796-2802

URL: https://www.wjgnet.com/2307-8960/full/v11/i12/2796.htm DOI: https://dx.doi.org/10.12998/wjcc.v11.i12.2796

INTRODUCTION

The clinical features of dissociated vertical deviation (DVD) with superior oblique palsy (SOP) are variable and include unstable vertical strabismus after covering in the primary position due to significant DVD; spontaneous upward drifting of one or both eyes when binocularity is blocked after neutralizing the hypertropia; and superior oblique muscle (SO) paralysis[1,2]. The goals of surgery are improvement of primary position hypertropia and reduction of upward floating. Inferior oblique anterior transposition (IOAT) has been recognized as the standard surgical procedure [1,2]. The complications after this procedure remain a difficult problem, such as the anti-elevation syndrome (AES)[3] and eyelid fission narrowing in straight-ahead gaze^[4], especially after unilateral surgery. We here report the results of a modified unilateral IOAT in two patients with DVD and SOP.

IOAT was first proposed by Gobin^[5] in 1964 and used in the treatment of esotropia with V sign, the inferior oblique muscle (IO) was transposed to the equatorial sclera (about 14 mm from the corneal limbus) and approximately 50% of patients were orthotropic after the procedure. The application of IOAT in the treatment of inferior oblique overaction (IOOA) and DVD was first proposed by Elliott and Nankin^[6] in 1981. IOAT is a more effective treatment for IOOA and DVD than simple IO recession, with the anterior fibers of the IO fixed temporally at the inferior rectus (IR) insertion (Elliott's point). It was reported that 67% (14/21) of patients' hypertropia was corrected, but it was easily overcorrected, resulting in restricted elevation in 24% of the patients.

This novel unilateral IOAT method involves a more backward reinsertion line perpendicular to the IR axis with J-deformity. The anterior-nasal fibers of the IO were attached at 9 mm posterior to the corneal limbus along the temporal board of the IR, and the posterior fibers were attached a further 5 mm temporal to the first suture.

This procedure was performed by a single surgeon (YX) for DVD with SOP in the two patients between June and August 2020. One day after surgery, they were orthotropic in primary gaze without diplopia and remained orthotropic one year after the procedure without obvious AES. This approach ensured surgical effectiveness and prevented the onset of AES, especially after unilateral surgery.

This study was approved by the Medical Ethics Committee of Qingdao University. This report complies with all local laws and the principles of the Declaration of Helsinki.

CASE PRESENTATION

Chief complaints

Case 1: An 11-year-old Chinese girl presented with persistent exotropia after a strabismus procedure five years earlier.

Case 2: A 17-year-old Chinese girl presented with intermittent exotropia of the left eye. She had been tilting her head to the left since infancy.

History of present illness

Case 1: The patient had been tilting her head to the right since infancy and, in 2015, was diagnosed with constant exotropia. She underwent lateral rectus recession (6 mm) with medial rectus resection (5 mm) in her right eye, but residual exotropia and abnormal head position persisted postoperatively through to the current presentation.

Case 2: The patient presented with intermittent exotropia of the left eye. She had been tilting her head to the right since infancy without any prior treatment.



WJCC | https://www.wjgnet.com

History of past illness

The patients had no previous medical history.

Personal and family history

Case 1: The patient's parents and sister are healthy and have no family history of similar disease.

Case 2: The patient's parents, sister and brother are healthy and have no family history of similar disease.

Physical examination

Case 1: The patient's temperature was 36.3 °C, heart rate was 92 bpm, respiratory rate was 23 breaths/ min, and blood pressure was 94/54 mmHg.

Case 2: The patient's temperature was 36.5 °C, heart rate was 98 bpm, respiratory rate was 20 breaths/ min, and blood pressure was 104/67 mmHg.

Laboratory examinations

Routine blood and biochemical tests were normal in both cases.

Imaging examinations

Ultrasonography did not indicate intraocular disease in both cases.

Ophthalmological examination

Case 1: On examination, the patient's best-corrected visual acuity (BCVA) was 1.0 in both eyes. She had primary position exotropia of 40Δ at both distance and near, and a hypertropia of 10Δ in the left eye by alternate prism cover test. DVD was observed in both eyes, although more pronounced in the left, and only appearing when covering one eye. There was bilateral IOOA (+3) and a significant V-pattern. A positive Bielschowsky tilt test was found in both sides[7] (Figure 1A).

Case 2: The patient's BCVA was 1.0 in both eyes, with refractions of -3.00 for the right and $+3.00 + 0.25 \times 65^{\circ}$ for the left eye. Exodeviation and hypertropia were highly variable due to significant dissociated horizontal deviation[8] in the left eye and DVD in the right. Exotropia of $10-20\Delta$ and right hypertropia of $15-30\Delta$ were observed on alternate prism cover test in the primary position at distance and near. She had latent nystagmus in both eyes and an anomalous head posture. There was unilateral superior oblique underaction (-2) along with IOOA (+2) in the right eye, and a positive Bielschowsky tilt test on the right-hand side (Figure 2A).

FINAL DIAGNOSIS

The final diagnosis for both cases were DVD with SOP.

TREATMENT

Case 1

Levofloxacin eye drops were applied to prevent infection in both eyes 2 d before the surgery. With the patient and her mother's consent, she underwent the modified unilateral IOAT with lateral rectus recession (5 mm) and medial rectus resection (4 mm) in her left eye under general anesthesia.

The anterior-nasal fibers of the left IO were attached at 9 mm posterior to the corneal limbus along the temporal board of the left IR, and the posterior fibers were attached a further 5 mm temporal to the first suture using an absorbable 6/0 suture. The left lateral rectus (LR) was backward to 5 mm from the original insertion site and the left medial rectus was resected about 4 mm and re-sutured to the original insertion site using an absorbable 6/0 sutures. At the end of the surgery, the surgeon applied tobramycin dexamethasone ointment to the left eye, bandaged it and waited for the patient to wake up before sending the patient back to the ward. After the surgery, the patient was instructed to apply levofloxacin eye drops and tobramycin dexamethasone drops to the left eye.

Case 2

Levofloxacin eye drops were applied to prevent infection in both eyes 2 d before the surgery. With the patient and her mother's consent, she underwent the modified unilateral IOAT in her right eye with lateral rectus recession 7 mm in her left eye under general anesthesia.

Raisbidena® WJCC | https://www.wjgnet.com



DOI: 10.12998/wjcc.v11.i12.2796 Copyright ©The Author(s) 2023.

Figure 1 Clinical photographs of Case 1. A: Preoperative images showing primary position exotropia and hypertropia in the left eye, and positive Bielschowsky tilt tests demonstrating superior oblique palsy in both eyes; B and C: Postoperative images showing primary position orthophoria without obvious antielevation syndrome in the left eye at the 1-d and 12-mo follow-up.

> The anterior-nasal fibers of the right IO were attached at 9 mm posterior to the corneal limbus along the temporal board of the right IR, and the posterior fibers were attached a further 5 mm temporal to the first suture using an absorbable 6/0 suture. The left LR was backward to 7 mm from the original insertion site using an absorbable 6/0 suture. At the end of the surgery, the surgeon applied tobramycin dexamethasone ointment to both eyes, bandaged them and waited for the patient to wake up before sending the patient back to the ward. After the surgery, the patient was instructed to apply antibiotic eye drops to the both eyes.

OUTCOME AND FOLLOW-UP

Case 1

One day after surgery, primary gaze orthophoria was achieved, without head turn or diplopia (Figure 1B), and one year after surgery, she remained orthophoric in primary gaze and without obvious AES and eyelid fission narrowing (Figure 1C).

Case 2

One day after surgery she achieved orthophoria in primary gaze, without head turn and diplopia (Figure 2B). Twelve months after surgery, she remained orthophoric in primary gaze without limitation of elevation in both eyes (Figure 2C).

DISCUSSION

SOP is the most common paralysis of single cyclovertical muscle and is the fourth cranial nerve (trochlear nerve) palsy, which involves the SO[9]. The incidence of SOP is approximately 12.9 (95% confidence interval 9.0-16.9) per 100000 adolescents younger than 19 years of age[10] and 54.1 (95% confidence interval 50.2–58.0) per 100000 adults[11]. In SOP, there is hypertropia of the paralytic eye





DOI: 10.12998/wjcc.v11.i12.2796 Copyright ©The Author(s) 2023.

Figure 2 Clinical photographs of Case 2. A: Preoperative images showing primary position exotropia and hypertropia in the right eye, and a positive Bielschowsky tilt test demonstrating superior oblique palsy in the right eye; B and C: Postoperative images showing primary position orthophoria without obvious antielevation syndrome in the right eye at the 2-wk and 12-mo follow-up.

> with abnormal head position. Examination of version shows underaction of the SO and overaction of the antagonist IO. The main surgical methods are to weaken the antagonist muscle of the paralytic muscle; to weaken the yoke muscle of the another eye; to strengthen the paralytic muscle; and to weaken the superior rectus (SR) of the paralytic eye[9]. DVD is an innervation disorder which occurs in more than 50% of infantile strabismus[12]. The main clinical feature is either eye slowly drifting upward and outward with extortion when occluded. Traditional surgical treatment for DVD focuses on the SR recession, and for patients with IOOA, IOAT is also considered[13].

> By displacing the IO insertion to anterior to the temporal insertion site of the IR (Elliott point) in a crossed-swords manner, IOAT was more effective improvement of IOOA and DVD compared with the standard recession of IO[6]. This IOAT has been widely employed; however, the risks of overaction and AES were high, especially in unilateral IOAT (more than half). AES is an elevation restriction in abduction, that might be contributed by both suturing the IO anterior to the IR insertion, and spreading the posterior lateral corner of the IO out temporally[14].

> Stager[15,16] established that the neurovascular bundle of the IO was an ancillary origin of the posterior-temporal fibers of the IO following IOAT for limiting eye upturn, and the neurofibrovascular bundle attaches to the IO 2 mm temporal to the IR, 12 mm from the insertion of the IO. Based on the autopsy results from cadaveric eyes by Apt L[17], the temporal length of the IO was about 14 mm, so the Elliott point might increase the risk of excessive muscle tension of temporal IO fibers. Kushner[18] suggested that AES might be an "innervational restriction," as opposed to a mechanical restriction and appears to be caused by tautness of the lateral fibers of the transposed IO. Therefore, they sutured the anterior nasal corner of the IO to the temporal corner of the IR insertion and let the posterior temporal corner hang back. This approach is better for preventing AES, but less powerful for reducing DVD.

> To avoid AES, some have suggested that the insertion line should be oriented parallel to the IR axis, the anterior IO nasal fibers should be at or just (≤ 2 mm) anterior to the temporal IR insertion in a "crossed-swords" manner, and the posterior temporal fibers should be folded and buried under the



WJCC | https://www.wjgnet.com

fixed anterior IO nasal fibers^[19]. This transposition has been seen to weaken mild to moderate IOOA and correct small primary position hypertropia. Guemes and Wright[20] suggested that the bilateral graded IOAT was effective in normalizing versions and correcting vertical deviations in primary position, and the new insertion (the temporal insertion site of the IR or 1-2 mm behind) line perpendicular to the IR axis with a "J" deformity for the IOOA combined with DVD (6 to 15 PD), without longterm observation of complications.

Thus, the reattachment points and attachment line directions are the key factors affecting operative outcomes and complications. Based on the above study, we chose a new anterior transposition to 9 mm posterior to the corneal limbus along the temporal board of the IR to reduce the risk of AES, with the wider insertion (about 5 mm) perpendicular to the IR axis to enhance the orthotic effects on vertical strabismus and spontaneous upward drifting.

The basic idea of IO transposition surgery according to the degree of IOOA is gradually moving the insertion of the IO anteriorly; if 1-mm anterior IR insertion still does not sufficiently improve the vertical strabismus and spontaneous upward drifting, a new line of attachment perpendicular to the IR insertion will be used. This surgical approach chose the new insertion line perpendicular to the IR axis to decline the upward drift of the DVD and vertical strabismus, weakening the effect of the anterior transposition of the new insertion of the IOAT, thus preventing the onset of AES. This approach ensured surgical effectiveness and prevented the onset of AES, especially after unilateral surgery.

Other advantages to this version of the IOAT procedure is that using the limbus as a reference point is more accurate and reasonable than the IR, and nasal transposition is not necessarily required which reduces operating difficulty.

CONCLUSION

In the cases presented here, the modified unilateral IOAT improved the patients' primary position hypertropia and superior floating phenomenon, with no obvious postoperative anti-elevation syndrome and no significant effect on lower lid configuration and function for 1 year. Therefore, this procedure is an effective and safe surgical methodfor t reating DVD with SOP. Certainly, more data are required to corroborate these findings and a controlled comparative study is necessary to draw definitive conclusions. In addition, a cohort study is in progress to better assess the procedure.

ACKNOWLEDGEMENTS

Thanks to Professor Ze Wang for his valuable advice on surgical design and article writing.

FOOTNOTES

Author contributions: Zong Y and Yang X were the patient's surgeons; Zong Y contributed to the conception, manuscript writing, revision, and final approval of the manuscript; Jiang WL contributed to the provision of study materials; Yang X and Wang Z contributed to the design, manuscript writing and revision; all authors have read and approved the final manuscript.

Supported by The Natural Science Foundation of Shandong Province, No. ZR2018BH013; The China Postdoctoral Science Foundation, No. 2017M612214.

Informed consent statement: Informed written consent was obtained from the patients for publication of this report and any accompanying images.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is noncommercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: China

ORCID number: Yao Zong 0000-0003-2300-1527; Wen-Lan Jiang 0000-0002-4393-4570; Xian Yang 0000-0002-6821-7930.



S-Editor: Hu YR L-Editor: A P-Editor: Cai YX

REFERENCES

- Yang SH. Dissociated vertical deviation and paralytic vertical strabismus. Zhongguo Xieshi Yu Xiaoer Yanke Zazhi2010 [DOI: 10.3969/j.issn.1005-328X.2010.01.001]
- Yang HK, Kim JH, Hwang JM. Congenital superior oblique palsy and trochlear nerve absence: a clinical and radiological 2 study. Ophthalmology 2012; 119: 170-177 [PMID: 21924501 DOI: 10.1016/j.ophtha.2011.06.038]
- 3 Cho YA, Kim JH, Kim S. Antielevation syndrome after unilateral anteriorization of the inferior oblique muscle. Korean J Ophthalmol 2006; 20: 118-123 [PMID: 16892649 DOI: 10.3341/kjo.2006.20.2.118]
- 4 Kushner BJ. The effect of anterior transposition of the inferior oblique muscle on the palpebral fissure. Arch Ophthalmol 2000; 118: 1542-1546 [PMID: 11074811 DOI: 10.1001/archopht.118.11.1542]
- Gobin MH. Anteroposion of the inferior oblique muscle in V-esotropia. Ophthalmologica 1964; 148: 325-341 [PMID: 5 14270337 DOI: 10.1159/000304704]
- Elliott RL, Nankin SJ. Anterior transposition of the inferior oblique. J Pediatr Ophthalmol Strabismus 1981; 18: 35-38 6 [PMID: 7264849 DOI: 10.3928/0191-3913-19810501-08]
- Wright KW, Yi N. Torticollis, Nystagmus and Incomitant Strabismus. In: Color Atlas Of Strabismus Surgery: Color 7 Atlas Of Strabismus Surgery. New York: Springer New York, 2015: 52-55
- Brodsky MC. Dissociated horizontal deviation: clinical spectrum, pathogenesis, evolutionary underpinnings, diagnosis, 8 treatment, and potential role in the development of infantile esotropia (an American Ophthalmological Society thesis). Trans Am Ophthalmol Soc 2007; 105: 272-293 [PMID: 18427617]
- 9 Chang MY, Coleman AL, Tseng VL, Demer JL. Surgical interventions for vertical strabismus in superior oblique palsy. Cochrane Database Syst Rev 2017; 11: CD012447 [PMID: 29178265 DOI: 10.1002/14651858.CD012447.pub2]
- 10 Tollefson MM, Mohney BG, Diehl NN, Burke JP. Incidence and types of childhood hypertropia: a population-based study. Ophthalmology 2006; 113: 1142-1145 [PMID: 16647125 DOI: 10.1016/j.ophtha.2006.01.038]
- Martinez-Thompson JM, Diehl NN, Holmes JM, Mohney BG. Incidence, types, and lifetime risk of adult-onset strabismus. Ophthalmology 2014; 121: 877-882 [PMID: 24321142 DOI: 10.1016/j.ophtha.2013.10.030]
- The American Academy of Ophthalmology. Section 06: Pediatric Ophthalmology and Strabismus. In 2018-2019 Basic 12 and Clinical Science Course (BCSC). San Francisco: The American Academy of Ophthalmology, 2019: 129-130
- 13 Hatt SR. Wang X. Holmes JM. Interventions for dissociated vertical deviation. Cochrane Database Syst Rev 2015; 2015: CD010868 [PMID: 26587695 DOI: 10.1002/14651858.CD010868.pub2]
- 14 Kushner BJ. Restriction of elevation in abduction after inferior oblique anteriorization. J AAPOS 1997; 1: 55-62 [PMID: 10530986 DOI: 10.1016/s1091-8531(97)90024-0]
- 15 Stager DR. Costenbader lecture. Anatomy and surgery of the inferior oblique muscle: recent findings. J AAPOS 2001; 5: 203-208 [PMID: 11507578 DOI: 10.1067/mpa.2001.116273]
- 16 Stager DR. The neurofibrovascular bundle of the inferior oblique muscle as the ancillary origin of that muscle. J AAPOS 1997; 1: 216-225 [PMID: 10532767 DOI: 10.1016/s1091-8531(97)90041-0]
- Apt L, Call NB. Inferior oblique muscle recession. Am J Ophthalmol 1978; 85: 95-100 [PMID: 619690 DOI: 17 10.1016/s0002-9394(14)76672-3]
- Kushner BJ. How to Perform Superior Surgery on the Inferior Oblique and Avoid Inferior Surgery on the Superior 18 Oblique. In: Strabismus: Practical Pearls You Won't Find in Textbooks. Switzerland: Springer Nature, 2017: 221-231
- Jeon H, Kwon H, Choi HY. Modified anterior transposition of the inferior oblique muscle. J AAPOS 2018; 22: 361-365.e1 19 [PMID: 30227248 DOI: 10.1016/j.jaapos.2018.06.008]
- Guemes A, Wright KW. Effect of graded anterior transposition of the inferior oblique muscle on versions and vertical 20 deviation in primary position. J AAPOS 1998; 2: 201-206 [PMID: 10532737 DOI: 10.1016/s1091-8531(98)90053-2]



WJCC | https://www.wjgnet.com



Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: bpgoffice@wjgnet.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

