

World Journal of *Clinical Cases*

World J Clin Cases 2023 April 26; 11(12): 2582-2854



Contents

Thrice Monthly Volume 11 Number 12 April 26, 2023

REVIEW

- 2582** Controversies in the management of acute pancreatitis: An update
Manrai M, Dawra S, Singh AK, Jha DK, Kochhar R
- 2604** Classification of osteogenesis imperfecta: Importance for prophylaxis and genetic counseling
Panzaru MC, Florea A, Caba L, Gorduza EV

MINIREVIEWS

- 2621** Emerging role of dual biologic therapy for the treatment of inflammatory bowel disease
McCormack MD, Wahedna NA, Aldulaimi D, Hawker P
- 2631** Pancreatic cancer and depression
Michoglou K, Ravinthiranathan A, San Ti S, Dolly S, Thillai K
- 2637** Mediastinal lesions in children
Çinar HG, Gulmez AO, Üner Ç, Aydin S
- 2657** Exit strategies in inflammatory bowel disease: Looking beyond anti-tumor necrosis factors
Crispino F, Michielan A, Grova M, Tieppo C, Mazza M, Rogger TM, Armelao F
- 2670** Medicinal cannabis products for the treatment of acute pain
Fiore M, Alfieri A, Di Franco S, Petrou S, Damiani G, Pace MC
- 2677** Role of in vitamin D in irritable bowel syndrome
Yu XL, Wu QQ, He LP, Zheng YF

ORIGINAL ARTICLE

Retrospective Cohort Study

- 2684** Analysis of oxidative stress and antioxidative potential in premature ovarian insufficiency
Kakinuma K, Kakinuma T

Retrospective Study

- 2694** Surgical management of pituitary adenoma during pregnancy
Jia XY, Guo XP, Yao Y, Deng K, Lian W, Xing B
- 2708** Role of pre-existing incomplete intestinal metaplasia in gastric adenocarcinoma: A retrospective case series analysis
Bogdanova I, Polaka I, Aleksandraviča I, Dzērve Z, Anarkulova L, Novika V, Tolmanis I, Leja M

Observational Study

- 2716** Severe/critical COVID-19 early warning system based on machine learning algorithms using novel imaging scores
Li QY, An ZY, Pan ZH, Wang ZZ, Wang YR, Zhang XG, Shen N
- 2729** Mediating effect of mindfulness level on the relationship between marital quality and postpartum depression among primiparas
Yang J, Lin XZ, Guo QW, Wang CL, Yang RY, Zhang JW, Zeng Y
- 2740** Ferric carboxymaltose for anemia in Crohn's disease patients at a tertiary center: A retrospective observational cohort study
Siqueira NSN, Pascoal LB, Rodrigues BL, de Castro MM, Martins ASC, Araiho DOS, Gomes LEM, Camargo MG, Ayrizono MLS, Leal RF

META-ANALYSIS

- 2753** Is metaphyseal ulnar shortening osteotomy superior to diaphyseal ulnar shortening osteotomy in the treatment of ulnar impaction syndrome? A meta-analysis
Deng HL, Lu ML, Tang ZM, Mao QL, Zhao JM
- 2766** Relationship between body mass index and short-term postoperative prognosis in patients undergoing colorectal cancer surgery
Li Y, Deng JJ, Jiang J

CASE REPORT

- 2780** Cardiac amyloidosis presenting as pulmonary arterial hypertension: A case report
Gao M, Zhang WH, Zhang ZG, Yang N, Tong Q, Chen LP
- 2788** Short-term outcome of total knee replacement in a patient with hemophilia: A case report and review of literature
Yin DL, Lin JM, Li YH, Chen P, Zeng MD
- 2796** Modified inferior oblique anterior transposition for dissociated vertical deviation combined with superior oblique palsy: A case report
Zong Y, Wang Z, Jiang WL, Yang X
- 2803** Treatment of talipes equinovarus after triceps surae intramuscular hemangioma surgery by Ilizarov technology in adults: A case report
Chen ZX, Wang MY, Zhang C, Ding ZQ, Chen W
- 2811** Open surgery: Still a great option to treat patients with post-traumatic arteriovenous fistulas: A case report
Kalinin R, Suchkov I, Mzhavanadze N, Borisova Y, Panin I
- 2817** Recovery from Bell's palsy after treatment using uncultured umbilical cord-derived mesenchymal stem cells: A case report
Ahn H, Jung WJ, Lee SY, Lee KH

- 2825** Pancreatic neuroendocrine tumor detected by technetium-99m methoxy-2-isobutylisonitrile single photon emission computed tomography/computed tomography: A case report
Liu CJ, Yang HJ, Peng YC, Huang DY
- 2832** Furazolidone-induced pulmonary toxicity in *Helicobacter pylori* infection: Two case reports
Ye Y, Shi ZL, Ren ZC, Sun YL
- 2839** Efficacy of anlotinib combined with radioiodine to treat scalp metastasis of papillary thyroid cancer: A case report and review of literature
Zhang LY, Cai SJ, Liang BY, Yan SY, Wang B, Li MY, Zhao WX
- 2848** Endoscopic ultrasound-guided transrectal drainage of a pelvic abscess after Hinchey II sigmoid colon diverticulitis: A case report
Drnovšek J, Čebren Ž, Grosek J, Janež J

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Mohamed Eltayeb Abdelrahman Naiem, MBBS, MD, Assistant Professor, Surgeon, Department of Surgery, Faculty of Medicine, University of Khartoum, Khartoum 102, Sudan. m-altayeb@live.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

April 26, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Endoscopic ultrasound-guided transrectal drainage of a pelvic abscess after Hinchey II sigmoid colon diverticulitis: A case report

Jan Drnovšek, Žan Čebren, Jan Grosek, Jurij Janež

Specialty type: Medicine, research and experimental

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B, B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Bains L, India; Bal'afif F, Indonesia; Liang P, China

Received: January 24, 2023

Peer-review started: January 24, 2023

First decision: February 17, 2023

Revised: February 20, 2023

Accepted: March 24, 2023

Article in press: March 24, 2023

Published online: April 26, 2023



Jan Drnovšek, Department of Gastroenterology, University Medical Centre Ljubljana, Ljubljana 1000, Slovenia

Jan Drnovšek, Žan Čebren, Jan Grosek, Jurij Janež, Faculty of Medicine, University of Ljubljana, Ljubljana 1000, Slovenia

Žan Čebren, Jan Grosek, Jurij Janež, Department of Abdominal Surgery, University Medical Centre Ljubljana, Ljubljana 1000, Slovenia

Corresponding author: Jurij Janež, MD, PhD, Assistant Professor, Surgeon, Surgical Oncologist, Department of Abdominal Surgery, University Medical Centre Ljubljana, No. 7 Zaloška cesta, Ljubljana 1000, Slovenia. jurij.janez@kclj.si

Abstract

BACKGROUND

Acute diverticulitis is one of the most prevalent complications of diverticular disease and may result in abscess formation, perforation, fistula formation, obstruction, or bleeding. Diverticular abscesses may be initially treated with antibiotics and/or percutaneous drainage and/or surgery. Endoscopic ultrasound (EUS)-guided drainage techniques are increasingly used as a minimally invasive alternative to percutaneous or surgical approaches, as they are associated with better treatment outcomes, shorter recovery time and duration of hospitalization.

CASE SUMMARY

A 57-year-old female presented to the emergency department on account of abdominal pain and fever. Clinical examination revealed tenderness in the left lower abdominal quadrant, with elevated inflammatory markers in laboratory tests. Abdominal computed tomography (CT) revealed an 8 cm × 8 cm × 5 cm well-encapsulated abscess of the sigmoid colon, surrounded by numerous diverticula. A diagnosis of Hinchey II diverticular abscess was made, and the patient was admitted and commenced on appropriate antibiotic treatment. A transrectal EUS showed a fluid collection in direct contact with the sigmoid colon. Transluminal drainage was performed, and a lumen-apposing metal stent was inserted into the abscess collection. A follow-up CT scan showed a regression of the collection. The patient's general condition improved, and the stent was removed during a follow-up transrectal EUS that revealed no visible collection.

CONCLUSION

We report the first successful management of a pelvic abscess in patient with

Hinchey II acute diverticulitis using EUS-guided transluminal drainage in Slovenia. The technique appears effective for well-encapsulated intra-abdominal abscesses larger than 4 cm in direct contact with the intestinal wall of left colon.

Key Words: Acute diverticulitis; Diverticular disease; Pelvic abscess; Endoscopic drainage; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The incidence of colonic diverticulosis and its complications is rising in developed countries. An abscess due to acute diverticulitis may be initially treated with antibiotics and/or percutaneous drainage and/or surgery. Since percutaneous drainage of an abscess is not always feasible, endoscopic ultrasound-guided transluminal drainage seems to be an effective minimally invasive alternative for well-encapsulated intra-abdominal abscesses lying in direct contact with the intestinal wall, which reduces the need for surgery and stoma formation in selected patients. However, given the limitations of the supporting evidence, the optimal treatment strategy should be determined on a case-by-case basis by a multidisciplinary team.

Citation: Drnovšek J, Čebren Ž, Grosek J, Janež J. Endoscopic ultrasound-guided transrectal drainage of a pelvic abscess after Hinchey II sigmoid colon diverticulitis: A case report. *World J Clin Cases* 2023; 11(12): 2848-2854

URL: <https://www.wjgnet.com/2307-8960/full/v11/i12/2848.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i12.2848>

INTRODUCTION

Diverticulosis of the colon is defined as the existence of false diverticula or outpouchings of the mucosa and serosa through openings in the muscular layer of the bowel known as *locus minoris resistentiae*. The condition is remarkably prevalent in Western countries, affecting a substantial proportion of the middle-aged and elderly populations. Inflammation of a colonic diverticulum, or acute diverticulitis, is one of the most prevalent complications of diverticular disease and represents an abdominal emergency[1]. The incidence of acute diverticulitis is rising, especially among individuals under the age of 50. This is likely attributable to factors such as a low-fibre diet, obesity, smoking, physical inactivity, and a diet high in red meat[2-4]. Approximately 5% of patients with known diverticula are estimated to develop an episode of acute diverticulitis[5].

Numerous classifications and modifications describe the various stages of diverticular disease. The first widely used classification by Hinchey was intended as an intra-operative stratification of perforated diverticulitis with abscess or peritonitis, enabling surgeons to adjust the surgical approach. The treatment of acute diverticulitis depends on whether the presentation is uncomplicated or complicated. Complicated presentation occurs in approximately 12% of patients. An intraabdominal abscess (Hinchey Ib or II) can be managed non-operatively, but an abscess larger than 4 cm in diameter must be drained[3,6,7].

In recent years, minimally invasive methods have replaced classical surgical techniques where possible. Drainage is usually performed percutaneously by an interventional radiologist. However, percutaneous drainage is not always feasible, and endoscopic ultrasound (EUS)-guided drainage represents a compelling alternative that eliminates the need for an invasive surgical approach.

In the present report, we describe the clinical presentation, management, and outcome of EUS-guided drainage of a pelvic abscess due to complicated Hinchey II acute sigmoid diverticulitis. The management approach is discussed in the context of available literature.

CASE PRESENTATION

Chief complaints

A 57-year-old otherwise healthy female was admitted to the abdominal emergency department on account of abdominal pain and a fever of up to 38 °C since the preceding five days.

History of past illness

A patient's history of past illness is unremarkable.

Personal and family history

A patient's personal and family history is unremarkable.

Physical examination

Clinical examination revealed localized tenderness in the left lower abdominal quadrant without clinical signs of diffuse peritonitis.

Laboratory examinations

Laboratory tests at admission showed elevated inflammatory markers with a C-reactive protein (CRP) value of 353 mg/L and a leucocyte count of 12.4×10^9 /L.

Imaging examinations

An urgent abdominal computed tomography (CT) scan revealed a well-encapsulated pelvic abscess measuring 8 cm × 8 cm × 5 cm, located in the proximity of a very long sigmoid colon surrounded by numerous diverticula (Figure 1). There was only a small amount of free fluid in the lesser pelvis with no other pathological findings.

FINAL DIAGNOSIS

Pelvic abscess due to acute sigmoid diverticulitis (Hinchey II) without evidence of diffuse peritonitis.

TREATMENT

The patient was hospitalized at the department of abdominal surgery, and empirical antibiotic treatment with amoxicillin and clavulanic acid was commenced, in line with national guidelines for acute complicated diverticulitis. Additionally, in the first 48 h she was put on nil by mouth and received intravenous fluids and analgesics. Due to the size of the abscess and the elevated inflammatory markers despite antibiotic treatment, an interventional radiologist was consulted to perform percutaneous drainage. However, the collection was not safely approachable *via* the percutaneous route. After consulting a gastroenterologist, a decision was jointly made to attempt draining the collection transrectally under EUS guidance.


On the third day of admission, a transrectal EUS scan was performed, which showed an encapsulated fluid collection lying in direct contact with the wall of the sigmoid colon. Transluminal drainage of the collection was performed by placing a lumen-apposing metal stent (LAMS) into the collection (Figure 2). The procedure was completed without complications, and purulent content was observed leaking through the open stent. The patient's condition gradually improved over the first few days after the procedure; she started with enteral nutrition, abdominal pain diminished, and defecation was normal. The previously elevated inflammatory parameters also began to decline.

A follow-up abdominal CT scan showed an appropriate stent position and regression of the collection with only some residual gas at the site (Figure 3). As the patient was asymptomatic, the antibiotic treatment was switched from intravenous to enteral route by the 9th day after admission. She was temporarily discharged home with the prescribed antibiotic and care instructions.

The patient returned as scheduled on the 16th day after the primary admission and repeat laboratory studies showed complete normalization of inflammatory parameters (CRP 6 mg/L, leukocytes 6.6×10^9 /L). She reported no complaints and was discharged from the hospital on the same day. The antibiotic treatment was also discontinued after 16 d of administration (9 d intravenously and 8 d enterally).


OUTCOME AND FOLLOW-UP

Thirty days after the stent insertion, an elective follow-up transrectal EUS was performed at the outpatient clinic. The endoscopic view demonstrated almost complete closure of the abscess cavity, surrounded by granulation tissue; accordingly, the LAMS was endoscopically removed during the same procedure (Figure 4). The procedure was uneventful, and at a subsequent follow-up clinical review, the patient remained well without any complaints. Three months after the LAMS removal, we additionally performed a colonoscopy and noted the disappearance of the fistular canal at the previous puncture site and no other remarkable findings.




DOI: 10.12998/wjcc.v11.i12.2848 Copyright ©The Author(s) 2023.

Figure 1 Abdominal computed tomography scan in axial and coronal view. The well-encapsulated pelvic abscess measuring 8 cm × 8 cm × 5 cm is seen as a dense air-fluid collection in the proximity of the sigmoid colon. A: Axial view; B: Coronal view.



DOI: 10.12998/wjcc.v11.i12.2848 Copyright ©The Author(s) 2023.

Figure 2 Endoscopic ultrasound scan during lumen-apposing metal stent insertion. A: An Endoscopic ultrasound image of the inserted lumen-apposing metal stent (LAMS), connecting the bowel lumen and the abscess collection; B: An endoscopic view of in-situ LAMS, draining the pus from the abscess collection.



DOI: 10.12998/wjcc.v11.i12.2848 Copyright ©The Author(s) 2023.

Figure 3 Follow-up abdominal computed tomography scan in axial and coronal views. The lumen-apposing metal stent position is appropriate and lies transluminal in the sigmoid wall. Regression of the collection is seen with only some gas remaining at the site. A: Axial view; B: Coronal view.

DISCUSSION

We present a case of successful EUS-guided drainage of a pelvic abscess due to complicated Hinchey II acute sigmoid diverticulitis. Because the percutaneous route of abscess drainage was not safely accessible, the decision was made to use this alternative option after interdisciplinary consultations and discussions. The EUS-guided transrectal drainage with LAMS insertion was performed and, in




Figure 4 The findings during the removal of the lumen-apposing metal stent. A: The lumen-apposing metal stent (LAMS) is seen in situ with regression of the abscess cavity; B: The remaining fistular canal is covered with granulation tissue after the removal of LAMS.

combination with antibiotic treatment, proved effective for managing the diverticular abscess. LAMS removal was performed 30 d after the insertion uneventfully.

There are several classification systems for describing acute diverticulitis, but the modified Hinchey classification, introduced in 1978 and revised by Wasvary *et al*[8] in 1999, remains the most commonly used in the international literature[8-12]. In general, the management of acute diverticulitis is well-established in the guidelines[6,7,12,13]. For the Hinchey stages 0 and Ia, antibiotic treatment is usually sufficient; stages Ib and II require additional drainage if an abscess is larger than 4 cm[6,7,13]. The drainage may be performed surgically, percutaneously or endoscopically[11,13].

However, the most appropriate drainage method for the patient may not always be accurately determined, and treatment choice must be individualized. The surgical treatment being the most invasive and associated with the highest morbidity and mortality, should be performed as a last resort and only when other methods fail. The most widely accepted clinical practice is to perform percutaneous drainage *via* transabdominal anterior or transgluteal posterior approach; however, percutaneous drainage is not always feasible[11-14]. Additionally, even though it is considered safe, the complication rate of percutaneous drainage is 2.5% (ranging from 0 to 12.5%), and possible complications include enterocutaneous fistulas, small bowel lesions, and sciatic nerve injuries[15].

Deep pelvic abscesses often present a clinical challenge, since they are surrounded by intra-abdominal organs and other structures, which may interfere with safe percutaneous drainage route. Over the past few decades, endoscopic drainage has emerged as a compelling and feasible option when percutaneous drainage cannot be performed safely, reducing the need for surgery and stoma formation in selected patients[11]. However, endoscopic drainage of diverticular abscess is not yet reflected in the major guidelines, and its supporting evidence is currently limited to only several retrospective case studies or case reports[6,12-17]. In contrast, the experience with endoscopic drainage of intraabdominal abscesses is rich. Besides acute diverticulitis, an intraabdominal abscess may also be associated with other inflammatory gastrointestinal or urogenital conditions or may occur as a complication after surgical procedures. Appropriate drainage is of vital importance for a successful treatment of an intra-abdominal abscess[18,19]. Extensive and positive experiences with LAMS in treating peripancreatic collections also open new possibilities for treating other types of fluid collections. Their characteristic shape prevents stent dislocation, and the diameter of the lumen allows efficient drainage and further endoscopic necrosectomy *via* an open stent. EUS-guided transluminal drainage might be a safer procedure in comparison to surgical or percutaneous drainage techniques as the risk of vessel injury and a puncture site leakage are neglectable. Interestingly, fecal contamination in the abscess cavity is unlikely as the intrinsic negative luminal pressure ensures adequate drainage. Thus, EUS-guided transluminal drainage is a suitable drainage method of all pelvic abscesses, regardless of their etiology.

The crucial abscess characteristics for a safe EUS drainage are size larger than 4 cm, a well-formed capsule and the position in close contact with the intestinal wall of the left colon[14,19]. As already noted, although percutaneous drainage is considered the method of choice in the existing guidelines, it is not always feasible. In the event of a failed or impractical percutaneous approach, an attempt at endoscopic drainage should be considered in any patient without evidence of sepsis before initiating surgical therapy. Meanwhile, more case reports or larger prospective studies are needed to evaluate and compare both drainage methods. Because EUS-guided drainage is not included in the guidelines, the exact role of endoscopic drainage in the treatment of diverticular abscesses remains unclear.

CONCLUSION

Since percutaneous drainage of diverticular abscess is not always feasible, EUS-guided transluminal drainage appears to be a promising and practical alternative for managing well-encapsulated intra-abdominal abscesses that lie in direct contact with the intestinal wall of left colon, reducing the need for

surgery and stoma formation. However, given the limited evidence, a careful interdisciplinary review of each clinical case by the abdominal surgeon, gastroenterologist, and interventional radiologist is required to determine the optimal treatment strategy.

FOOTNOTES

Author contributions: Drnovšek J and Janež J conceived and designed the study; Čebon Ž contributed to the collection, analysis and interpretation of data; Čebon Ž and Drnovšek J drafted the manuscript; Grosek J and Janež J contributed to the critical revision; all authors issued final approval for the version to be submitted.

Informed consent statement: Our institution does not require ethical approval for publishing a case report. Written informed consent was obtained from the patient for the publication of this case report and accompanying images.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Slovenia

ORCID number: Jan Drnovšek 0000-0003-1831-1432; Žan Čebon 0000-0003-1896-7167; Jan Grosek 0000-0001-9832-4596; Jurij Janež 0000-0003-2543-5003.

S-Editor: Zhao S

L-Editor: A

P-Editor: Zhao S

REFERENCES

- 1 Wilkins T, Embry K, George R. Diagnosis and management of acute diverticulitis. *Am Fam Physician* 2013; **87**: 612-620 [PMID: 23668524]
- 2 Swanson SM, Strate LL. Acute Colonic Diverticulitis. *Ann Intern Med* 2018; **168**: ITC65-ITC80 [PMID: 29710265 DOI: 10.7326/AITC201805010]
- 3 Strate LL, Morris AM. Epidemiology, Pathophysiology, and Treatment of Diverticulitis. *Gastroenterology* 2019; **156**: 1282-1298.e1 [PMID: 30660732 DOI: 10.1053/j.gastro.2018.12.033]
- 4 Hawkins AT, Wise PE, Chan T, Lee JT, Glyn T, Wood V, Eglinton T, Frizelle F, Khan A, Hall J, Ilyas MIM, Michailidou M, Nfonam VN, Cowan ML, Williams J, Steele SR, Alavi K, Ellis CT, Collins D, Winter DC, Zaghiyan K, Gallo G, Carvello M, Spinelli A, Lightner AL. Diverticulitis: An Update From the Age Old Paradigm. *Curr Probl Surg* 2020; **57**: 100862 [PMID: 33077029 DOI: 10.1016/j.cpsurg.2020.100862]
- 5 Piscopo N, Ellul P. Diverticular Disease: A Review on Pathophysiology and Recent Evidence. *Ulster Med J* 2020; **89**: 83-88 [PMID: 33093692]
- 6 Francis NK, Sylla P, Abou-Khalil M, Arolo S, Berler D, Curtis NJ, Dolejs SC, Garfinkle R, Gorter-Stam M, Hashimoto DA, Hassinger TE, Molenaar CJL, Pucher PH, Schuermans V, Arezzo A, Agresta F, Antoniou SA, Arulampalam T, Boutros M, Bouvy N, Campbell K, Francione T, Haggerty SP, Hedrick TL, Stefanidis D, Truitt MS, Kelly J, Ket H, Dunkin BJ, Pietrabissa A. EAES and SAGES 2018 consensus conference on acute diverticulitis management: evidence-based recommendations for clinical practice. *Surg Endosc* 2019; **33**: 2726-2741 [PMID: 31250244 DOI: 10.1007/s00464-019-06882-z]
- 7 Schultz JK, Azhar N, Binda GA, Barbara G, Biondo S, Boermeester MA, Chabok A, Consten ECJ, van Dijk ST, Johansen A, Kruis W, Lambrechts D, Post S, Ris F, Rockall TA, Samuelsson A, Di Saverio S, Tartaglia D, Thorisson A, Winter DC, Bemelman W, Angenete E. European Society of Coloproctology: guidelines for the management of diverticular disease of the colon. *Colorectal Dis* 2020; **22** Suppl 2: 5-28 [PMID: 32638537 DOI: 10.1111/codi.15140]
- 8 Wasvary H, Turfah F, Kadro O, Beauregard W. Same hospitalization resection for acute diverticulitis. *Am Surg* 1999; **65**: 632-5; discussion 636 [PMID: 10399971]
- 9 Hinchey EJ, Schaaf PG, Richards GK. Treatment of perforated diverticular disease of the colon. *Adv Surg* 1978; **12**: 85-109 [PMID: 735943]
- 10 Klarenbeek BR, de Korte N, van der Peet DL, Cuesta MA. Review of current classifications for diverticular disease and a translation into clinical practice. *Int J Colorectal Dis* 2012; **27**: 207-214 [PMID: 21928041 DOI: 10.1007/s00384-011-1314-5]
- 11 Zaborowski AM, Winter DC. Evidence-based treatment strategies for acute diverticulitis. *Int J Colorectal Dis* 2021; **36**:

- 467-475 [PMID: [33156365](#) DOI: [10.1007/s00384-020-03788-4](#)]
- 12 **Qaseem A**, Etzeandia-Ikobaltzeta I, Lin JS, Fitterman N, Shamliyan T, Wilt TJ; Clinical Guidelines Committee of the American College of Physicians*, Crandall CJ, Cooney TG, Cross JT Jr, Hicks LA, Maroto M, Mustafa RA, Obbley AJ, Owens DK, Tice J, Williams JW Jr; Clinical Guidelines Committee of the American College of Physicians. Diagnosis and Management of Acute Left-Sided Colonic Diverticulitis: A Clinical Guideline From the American College of Physicians. *Ann Intern Med* 2022; **175**: 399-415 [PMID: [35038273](#) DOI: [10.7326/M21-2710](#)]
- 13 **Sartelli M**, Weber DG, Kluger Y, Ansaloni L, Coccolini F, Abu-Zidan F, Augustin G, Ben-Ishay O, Biffl WL, Bouliaris K, Catena R, Ceresoli M, Chiara O, Chiarugi M, Coimbra R, Cortese F, Cui Y, Damaskos D, De' Angelis GL, Delibegovic S, Demetrasvili Z, De Simone B, Di Marzo F, Di Saverio S, Duane TM, Faro MP, Fraga GP, Gkiokas G, Gomes CA, Hardcastle TC, Hecker A, Karamarkovic A, Kashuk J, Khokha V, Kirkpatrick AW, Kok KYY, Inaba K, Isik A, Labricciosa FM, Latifi R, Leppäniemi A, Litvin A, Mazuski JE, Maier RV, Marwah S, McFarlane M, Moore EE, Moore FA, Negroi I, Pagani L, Rasa K, Rubio-Perez I, Sakakushev B, Sato N, Sganga G, Siquini W, Tarasconi A, Tolonen M, Ulrych J, Zachariah SK, Catena F. 2020 update of the WSES guidelines for the management of acute colonic diverticulitis in the emergency setting. *World J Emerg Surg* 2020; **15**: 32 [PMID: [32381121](#) DOI: [10.1186/s13017-020-00313-4](#)]
- 14 **Lisotti A**, Cominardi A, Bacchilega I, Linguetti R, Fusaroli P. EUS-guided transrectal drainage of pelvic fluid collections using electrocautery-enhanced lumen-apposing metal stents: a case series. *VideoGIE* 2020; **5**: 380-385 [PMID: [32821872](#) DOI: [10.1016/j.vgie.2020.04.014](#)]
- 15 **Gregersen R**, Mortensen LQ, Burchard J, Pommergaard HC, Rosenberg J. Treatment of patients with acute colonic diverticulitis complicated by abscess formation: A systematic review. *Int J Surg* 2016; **35**: 201-208 [PMID: [27741423](#) DOI: [10.1016/j.ijsu.2016.10.006](#)]
- 16 **Donatelli G**, Cereatti F, Fazi M, Ceci V, Dhumane P. Endoscopic ultrasound-guided drainage of intra-abdominal diverticular abscess. A case series. *J Minim Access Surg* 2021; **17**: 513-518 [PMID: [34558427](#) DOI: [10.4103/jmas.JMAS_184_20](#)]
- 17 **Hadithi M**, Bruno MJ. Endoscopic ultrasound-guided drainage of pelvic abscess: A case series of 8 patients. *World J Gastrointest Endosc* 2014; **6**: 373-378 [PMID: [25132921](#) DOI: [10.4253/wjge.v6.i8.373](#)]
- 18 **Dawod E**, Nieto JM. Endoscopic Ultrasound-Guided Drainage of Pelvic, Intra-abdominal, and Mediastinal Abscesses. In: *Interventional Endoscopic Ultrasound*. Springer 2019 [DOI: [10.1007/978-3-319-97376-0_17](#)]
- 19 **Dhindsa BS**, Naga Y, Saghir SM, Dhaliwal A, Ramai D, Cross C, Singh S, Bhat I, Adler DG. EUS-guided pelvic drainage: A systematic review and meta-analysis. *Endosc Ultrasound* 2021; **10**: 185-190 [PMID: [33463556](#) DOI: [10.4103/eus.eus_71_20](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

