

# World Journal of *Clinical Cases*

*World J Clin Cases* 2023 May 6; 11(13): 2855-3113



## Contents

Thrice Monthly Volume 11 Number 13 May 6, 2023

## OPINION REVIEW

- 2855 Long-term implications of fetal growth restriction  
*D'Agostin M, Di Sipio Morgia C, Vento G, Nobile S*

## REVIEW

- 2864 Appraisal of gastric stump carcinoma and current state of affairs  
*Shukla A, Kalayarasan R, Gnanasekaran S, Pottakkat B*
- 2874 Burden of severe infections due to carbapenem-resistant pathogens in intensive care unit  
*Pace MC, Corrente A, Passavanti MB, Sansone P, Petrou S, Leone S, Fiore M*

## MINIREVIEWS

- 2890 Individualized diabetes care: Lessons from the real-world experience  
*Khor XY, Pappachan JM, Jeeyavudeen MS*
- 2903 Clinical management of dural defects: A review  
*Dong RP, Zhang Q, Yang LL, Cheng XL, Zhao JW*
- 2916 Potential impact of music interventions in managing diabetic conditions  
*Eseadi C, Amedu AN*
- 2925 Implications of obesity and adiposopathy on respiratory infections; focus on emerging challenges  
*Lempesis IG, Georgakopoulou VE*

## ORIGINAL ARTICLE

## Case Control Study

- 2934 Association of C-reactive protein and complement factor H gene polymorphisms with risk of lupus nephritis in Chinese population  
*Li QY, Lv JM, Liu XL, Li HY, Yu F*

## Retrospective Study

- 2945 Comparison of the application value of transvaginal ultrasound and transabdominal ultrasound in the diagnosis of ectopic pregnancy  
*Hu HJ, Sun J, Feng R, Yu L*

## Observational Study

- 2956 Assessment of knowledge, cultural beliefs, and behavior regarding medication safety among residents in Harbin, China  
*Liu XT, Wang N, Zhu LQ, Wu YB*

## SYSTEMATIC REVIEWS

- 2966** Palliative oral care in terminal cancer patients: Integrated review  
*Silva ARP, Bodanezi AV, Chrun ES, Lisboa ML, de Camargo AR, Munhoz EA*

## META-ANALYSIS

- 2981** Effect of preoperative inspiratory muscle training on postoperative outcomes in patients undergoing cardiac surgery: A systematic review and meta-analysis  
*Wang J, Wang YQ, Shi J, Yu PM, Guo YQ*
- 2992** Efficacy and safety of intravenous tranexamic acid in total shoulder arthroplasty: A meta-analysis  
*Deng HM*

## CASE REPORT

- 3002** Awake laparoscopic cholecystectomy: A case report and review of literature  
*Mazzone C, Sofia M, Sarv  I, Litrico G, Di Stefano AML, La Greca G, Latteri S*
- 3010** Bilateral malignant glaucoma with bullous keratopathy: A case report  
*Ma YB, Dang YL*
- 3017** Finger compartment syndrome due to a high-pressure washer injury: A case report  
*Choi JH, Choi SY, Hwang JH, Kim KS, Lee SY*
- 3022** Primary dedifferentiated chondrosarcoma of the lung with a 4-year history of breast cancer: A case report  
*Wen H, Gong FJ, Xi JM*
- 3029** Importance of proper ventilator support and pulmonary rehabilitation in obese patients with heart failure: Two case reports  
*Lim EH, Park SH, Won YH*
- 3038** Multiple flexor tendon ruptures due to osteochondroma of the hamate: A case report  
*Kwon TY, Lee YK*
- 3045** Fractional flow reserve measured *via* left internal mammary artery after coronary artery bypass grafting: Two case reports  
*Zhang LY, Gan YR, Wang YZ, Xie DX, Kou ZK, Kou XQ, Zhang YL, Li B, Mao R, Liang TX, Xie J, Jin JJ, Yang JM*
- 3052** Uterine artery embolization combined with percutaneous microwave ablation for the treatment of prolapsed uterine submucosal leiomyoma: A case report  
*Zhang HL, Yu SY, Cao CW, Zhu JE, Li JX, Sun LP, Xu HX*
- 3062** Metachronous urothelial carcinoma in the renal pelvis, bladder, and urethra: A case report  
*Zhang JQ, Duan Y, Wang K, Zhang XL, Jiang KH*
- 3070** Unusual phenomenon-“polyp” arising from a diverticulum: A case report  
*Liew JJJ, Lim WS, Koh FH*

- 3076** Idiopathic steno-occlusive disease with bilateral internal carotid artery occlusion: A Case Report  
*Hamed SA, Yousef HA*
- 3086** Solitary acral persistent papular mucinosis nodule: A case report and summary of eight Korean cases  
*Park YJ, Shin HY, Choi WK, Lee AY, Lee SH, Hong JS*
- 3092** Eosinophilic fasciitis difficult to differentiate from scleroderma: A case report  
*Lan TY, Wang ZH, Kong WP, Wang JP, Zhang N, Jin DE, Luo J, Tao QW, Yan ZR*
- 3099** Misdiagnosis of scalp angiosarcoma: A case report  
*Yan ZH, li ZL, Chen XW, Lian YW, Liu LX, Duan HY*
- 3105** Discrepancy among microsatellite instability detection methodologies in non-colorectal cancer: Report of 3 cases  
*Şenocak Taşçı E, Yıldız İ, Erdamar S, Özer L*



**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Jina Yun, MD, PhD, Assistant Professor, Division of Hematology-Oncology, Department of Internal Medicine, Soonchunhyang University Bucheon Hospital, Soonchunhyang University School of Medicine, Bucheon 14584, South Korea. 19983233@schmc.ac.kr

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Si Zhao; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

May 6, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

# Bilateral malignant glaucoma with bullous keratopathy: A case report

Ya-Bin Ma, Ya-Long Dang

**Specialty type:** Medicine, research and experimental

**Provenance and peer review:** Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): 0  
Grade C (Good): C, C  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Tanabe S, Japan; Wang WW, China

**Received:** October 7, 2022

**Peer-review started:** October 7, 2022

**First decision:** December 26, 2022

**Revised:** January 22, 2023

**Accepted:** March 31, 2023

**Article in press:** March 31, 2023

**Published online:** May 6, 2023



**Ya-Bin Ma, Ya-Long Dang**, Department of Ophthalmology, Sanmenxia Central Hospital, Sanmenxia 472000, Henan Province, China

**Corresponding author:** Ya-Long Dang, MD, PhD, Associate Chief Physician, President, Surgeon, Department of Ophthalmology, Sanmenxia Central Hospital, Wenhua Street, Sanmenxia 472000, Henan Province, China. [dangyalong@126.com](mailto:dangyalong@126.com)

## Abstract

### BACKGROUND

Malignant glaucoma, caused by aqueous misdirection, is a challenging post-surgical complication presented with normal/high intraocular pressure and shallowing of the central and peripheral anterior chambers. Its incidence is about 0.6%-4.0%. It can be secondary to filtering surgeries, laser iridotomy, and cataract surgery. Short axial length and a history of angle closure glaucoma are its main risk factors. Here, we report a bilateral malignant glaucoma with bullous keratopathy in the patient's left eye.

### CASE SUMMARY

We present a case of bilateral malignant glaucoma. The cause of malignant glaucoma for each eye of this patient was different. Hence, the management strategy and selection of surgical methods were also different. However, the normal anterior chamber was ultimately maintained, and maximum visual function was preserved. Even though the left eye received multiple surgeries and corneal endothelial decompensation occurred, the formation of a retroendothelial fibrous membrane partially compensated for the function of the corneal endothelium.

### CONCLUSION

The formation of a retroendothelial fibrous membrane partially compensated for the function of the corneal endothelium.

**Key Words:** Malignant glaucoma; Corneal epithelial cell; Lens epithelial cell; Bullous keratopathy; Retroendothelial fibrous membrane; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Malignant glaucoma, caused by aqueous misdirection, is a challenging post-surgical complication presented with normal/high intraocular pressure and shallowing of the central and peripheral anterior chambers. Its incidence is about 0.6%-4.0%. It can be secondary to filtering surgeries, laser iridotomy, and cataract surgery. Short axial length and a history of angle closure glaucoma are its main risk factors. Here, we report bilateral malignant glaucoma with bullous keratopathy in the patient's left eye. Interestingly, the best corrected visual acuity of the left eye improved to 20/70, and bullous keratopathy was relieved after the migration and implant of lens epithelial cells into the corneal endothelium.

**Citation:** Ma YB, Dang YL. Bilateral malignant glaucoma with bullous keratopathy: A case report. *World J Clin Cases* 2023; 11(13): 3010-3016

**URL:** <https://www.wjgnet.com/2307-8960/full/v11/i13/3010.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v11.i13.3010>

## INTRODUCTION

Malignant glaucoma, caused by aqueous misdirection, is a challenging post-surgical complication presented with normal/high intraocular pressure (IOP) and shallowing of the central and peripheral anterior chambers. Its incidence is about 0.6%-4.0% [1-3]. It can be secondary to filtering surgeries, laser iridotomy, and cataract surgery. Short axial length and a history of angle closure glaucoma are its main risk factors [4,5]. Here, we reported bilateral malignant glaucoma with bullous keratopathy in the patient's left eye. Interestingly, the best corrected visual acuity (BCVA) of her left eye improved to 20/70, and bullous keratopathy was relieved after the migration and implant of lens epithelial cells into the corneal endothelium.

## CASE PRESENTATION

### Chief complaints

A 36-year-old woman complained of "blurred vision in both eyes" and was diagnosed with chronic angle-closure glaucoma.

### History of present illness

She received bilateral laser peripheral iridotomy and was prescribed with timolol, brimonidine, and brinzolamide twice per day and latanoprost daily. Her IOP remained high (38 mmHg-45 mmHg), and visual field worsened.

### History of past illness

The patient did not report any past illnesses.

### Physical examination


The BCVA of both eyes was 20/20. The central and peripheral anterior chambers were shallow. The anterior chamber angles were partially closed. The anterior chamber depth was 1.86 mm (OD) and 1.61 mm (OS). The pupil diameter was 6.5 mm (OD) and 2.8 mm (OS). The pupillary light reflex in her right eye was delayed (Figure 1). The lenses were transparent. Fundoscopy revealed that the right cup-disc ratio was 0.9, and the disc border was narrow. The left optic disc boundaries were clear and pale red in color. No apparent abnormalities were observed in either retina.

### Laboratory examinations

The right eye showed a tubular visual field. The contrast sensitivity in the left eye was decreased and accompanied with a visual field defect. The central corneal thicknesses were 535  $\mu$ m (OD) and 556  $\mu$ m (OS), and the IOPs were 45 mmHg (OD) and 40 mmHg (OS). The axial lengths were 19.55 mm (OD) and 19.59 mm (OS). The corneal endothelial cell density was 2488/mm<sup>2</sup> (OD) and 2890/mm<sup>2</sup> (OS).

### Imaging examinations

OCT revealed that the thickness of retinal nerve fiber layer in her right eye reduced (Figure 2A). No abnormality was observed in the left eye (Figure 2B).



DOI: 10.12998/wjcc.v11.i13.3010 Copyright ©The Author(s) 2023.

**Figure 1** Anterior segment photographs and pentacam analysis. A and B: Bilateral corneas show transparency. Pupil diameter: OD 6.5 mm and OS 2.8 mm. The hole of iridotomy was marked by a black arrow; B: Different perspectives; C and D: Central and peripheral anterior chambers were shallow, and partial angle closure was present in both eyes.

## FINAL DIAGNOSIS


Bilateral chronic angle-closure glaucoma

## TREATMENT

For her right eye, we performed cataract phacoemulsification combined with intraocular lens implantation. Postoperatively, the patient had a grade II shallow anterior chamber, intraocular lens protrusion, and IOP of 42 mmHg. We considered malignant glaucoma in her right eye. Therefore, periorbital injection of methylprednisolone and atropine mydriasis were continued for 3 d; however, there was no improvement. Subsequently, the patient underwent a capsulotomy and an anterior vitrectomy. Postsurgery, there was significant anterior chamber deepening with a central ACD of 2.89 mm and IOP fluctuating between 25 and 35 mmHg. Considering that this might be an instance of chronic angle closure in the right eye, we implanted an Ahmed drainage valve. Post-surgery, the IOP stabilized at 13 mmHg-19 mmHg, and BCVA was 20/25. The surgery had achieved the expected outcomes.

Due to the short axial length in her left eye and malignant glaucoma in her right eye, we performed cataract phacoemulsification and intraocular lens implantation, combined with anterior vitrectomy for her left eye. Post-surgery, the ACD was deepened at approximately 2.51 mm, but the IOP remained at 30 mmHg. One week later, considering the angle closure, we implanted an Ahmed valve. Postsurgery, the patient's anterior chamber gradually disappeared, and a grade III shallow anterior chamber was detected. There was significant corneal edema. IOP was 20 mmHg, and the Seidel test was negative. We considered the malignant glaucoma in her left eye. We administered periorbital injections of methylprednisolone and atropine eye drops twice per day. However, these treatments were ineffective. A viscoelastic agent was injected into the anterior chamber, and laser posterior capsulotomy was performed. However, a grade III shallow anterior chamber recurred. A peripheral iridectomy with zonulo-capsulo-hyaloidotomy was performed through the pars plana route. Tension sutures were used to reform the anterior chamber. On day 2 postsurgery, the ACD was significantly increased to 2.75 mm. The IOP was 15 mmHg. One-week post-surgery, corneal edema was still significant, and bullous changes could be seen in the epithelium (Figure 3A). The OCT and slit lamp examination suggested that nearly one-third of the cornea endothelial detached, and a translucent membrane attached and stretched the endothelial layer (Figures 3B and 4A and B). The slit lamp revealed a bundle of white fibers that proliferated behind the corneal endothelium near the pupillary limbus at the 10 o'clock position and were pulling on the intraocular lens surface. Yttrium-aluminum-garnet (YAG) laser was immediately carried out to remove the proliferative fibers. We saw that fiber bundles had high tension and significant rebound.






DOI: 10.12998/wjcc.v11.i13.3010 Copyright ©The Author(s) 2023.


**Figure 2** Optical coherence tomography of the optic nerve head. A: The retinal nerve fiber layer in various quadrants of the right eye appeared to have thinned; B: No apparent abnormalities were observed in the retinal nerve fiber layer of the left eye.

## OUTCOME AND FOLLOW-UP

Two weeks after the YAG laser, the corneal edema had abated, the corneal bullae had disappeared, and the peripheral cornea was transparent (Figure 3C). The OCT revealed that the corneal endothelial detachment gradually resolved, and a highly reflective membranous mass attached to the medial endothelial layer (Figures 3D and 4C and D). At the 1-year follow-up, the IOP was 17 mmHg, and ACD was normal. The peripheral cornea was partially transparent. The corneal edema persisted (central corneal thickness: 756  $\mu$ m), and a fibrous proliferative membrane could be seen in the epithelium. The corneal epithelial bullae disappeared, and BCVA was 20/70.



**Figure 3 Anterior segment photography and optical coherence tomography of the left eye.** A: After peripheral iridectomy with zonulo-capsulo-hyaloidotomy and tension suture fixation, the anterior chamber was reconstructed. However, a significant corneal edema and bullous changes were seen; B: Optical coherence tomography showed that extensive corneal endothelial detachment involving approximately half of the cornea and a translucent membrane attached and stretched the endothelial layer; C: Two weeks after yttrium-aluminum-garnet laser, the corneal edema had abated, the corneal bullae had disappeared, and the peripheral cornea was transparent. D: Optical coherence tomography showed the corneal endothelial detachment gradually resolved. A highly reflective membranous mass attached to the medial endothelial layer.



**Figure 4 Enlarged figure of corneal optical coherence tomography.** A and B: A corneal endothelial detachment (orange arrows) involving approximately half of the cornea and a translucent membrane (green arrows) attached and stretched the endothelial layer (orange arrows); B: Enlarged image; C and D: Two weeks after removal of the fibrous bundle, corneal endothelial detachment gradually disappeared. A highly reflective membranous mass (green arrows) was attached to the medial endothelial layer (orange arrows).

## DISCUSSION

We presented a case of bilateral malignant glaucoma after phacoemulsification and intraocular lens implantation combined with anterior vitrectomy and Ahmed drainage valve implantation. The patient's right anterior chamber and IOP fully recovered after anterior vitrectomy. Although we realized that the incidence of malignant glaucoma in her left eye was extremely high and carried out anterior vitrectomy during stage I, grade III shallow anterior chamber and corneal lens contact still occurred, resulting in corneal endothelial decompensation. Fortunately, the anterior chamber reformed, and the IOP stabilized after peripheral iridectomy with zonulo-capsulo-hyaloidotomy and a tension suture fixation in the anterior chamber. It is worth noting that 1 wk after the left intraocular lens was disengaged from the cornea, a bundle of tense fiber formed between the lens surface and the corneal endothelium. This was suspected to be caused by epithelial cells migrating and transdifferentiating into myofibroblasts in the corneal endothelium. These myofibroblasts were attached to the surface of the corneal endothelium, blocking the entry of aqueous humor into the subcorneal epithelium and thereby preventing bullous

keratopathy.

Short axial length and anterior segment crowding are risk factors for malignant glaucoma[6]. Wang *et al*[6] analyzed 1183 patients with angle-closure glaucoma and found that the axial length and ACD of those patients were significantly lower than of those with primary angle-closure glaucoma (axial length:  $21.44 \pm 1.18$  mm *vs*  $22.17 \pm 0.97$  mm, ACD:  $2.12 \pm 0.41$  mm *vs*  $2.49 \pm 0.48$  mm). As malignant glaucoma often involves both eyes[7,8], we performed a three-step lens resection combined with vitrectomy on the patient's left eye[9]. First, the central vitreous was removed *via* the pars plana vitrectomy to relieve crowding of the anterior segment. Second, lens phacoemulsification, combined with intraocular lens implantation, was carried out. Finally, anterior vitrectomy was performed. Postsurgery, the anterior chamber was stable and deep, and aqueous misdirection did not occur. However, as the patient had chronic angle closure, the IOP remained high. Considering that trabeculectomy has poor results after vitrectomy, we opted for Ahmed drainage valve implantation[10].

Postsurgery, the patient had a grade III shallow anterior chamber and intraocular lens protrusion, with the intraocular lens coming into contact with the corneal endothelium. Although the IOP did not exceed 21 mmHg, aqueous misdirection had occurred. The diagnosis of malignant glaucoma was confirmed. The reasons for this were: (1) Excessive aqueous humor drainage after valve implantation; and (2) Postoperative inflammation promotes the proliferation, adhesion, and transformation of lens epithelial cells and ciliary body edema, resulting in a blockage of normal aqueous humor circulation [11]. Therefore, we carried out a peripheral iridectomy with zonulo-capsulo-hyaloidotomy at the 6 o'clock position, combined with tension suture fixation in the anterior chamber[12]. The postoperative anterior chamber reformation was good.

After the malignant glaucoma occurred, the patient's left eye developed a grade III shallow anterior chamber, the intraocular lens was in contact with the corneal endothelium, and there was moderate-severe inflammation[11]. Inflammatory responses can promote epithelial-mesenchymal transformation of the lens epithelial cell[13]. In this patient, a fibrous bundle could be seen between the corneal endothelium and surface of the intraocular lens at the 10 o'clock 1 wk after anterior chamber reformation had been successfully performed.

The fibrous bundle had tension between the corneal endothelium and the intraocular lens. It pulled the corneal endothelium, resulting in partial detachment. After the fibrous bundle was resected with YAG laser, the corneal endothelium and fibrous membrane gradually attached to the posterior surface of the cornea. Even though the density of corneal endothelial cells was low (797 cells/mm<sup>2</sup>) and the function was poor (6A cells accounted for 20%), the corneal bullae gradually disappeared, together with the formation of the fibrous membrane behind the corneal endothelium. This suggests that fibrous membrane derived from lens epithelial transformation is somewhat impermeable to water.

## CONCLUSION

We presented a case of bilateral malignant glaucoma. The cause of malignant glaucoma for each eye of this patient was different. Hence, the management strategy and selection of surgical methods were also different. However, the normal anterior chamber was ultimately maintained, and maximum visual function was preserved. Even though the left eye received multiple surgeries and corneal endothelial decompensation occurred, the formation of a retroendothelial fibrous membrane partially compensated for the function of the corneal endothelium.

## FOOTNOTES

**Author contributions:** Dang YL contributed to conception of the study; Dang YL performed the surgery; Ma YB and Dang YL contributed significantly to analysis and manuscript preparation; all authors have read and approved the final manuscript.

**Supported by** The Central Guiding Local Science & Technology Projects, No. Z20221341047; The Major Scientific and Technological Project of Sanmenxia 2022, No. 2022001007

**Informed consent statement:** Written informed consent was obtained from the patient for the publication of this case report.

**Conflict-of-interest statement:** All authors report no relevant conflicts of interest for this article.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-



NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Ya-Long Dang [0000-0003-2194-6375](https://orcid.org/0000-0003-2194-6375).

**S-Editor:** Liu GL

**L-Editor:** Filipodia

**P-Editor:** Liu GL

## REFERENCES

- 1 **Chandler PA**, Simmons RJ, Grant WM. Malignant glaucoma. Medical and surgical treatment. *Am J Ophthalmol* 1968; **66**: 495-502 [PMID: [5676357](#) DOI: [10.1016/0002-9394\(68\)91535-3](#)]
- 2 **Simmons RJ**. Malignant glaucoma. *Br J Ophthalmol* 1972; **56**: 263-272 [PMID: [5032764](#) DOI: [10.1136/bjo.56.3.263](#)]
- 3 **Lowe RF**. Clinical types of primary angle closure glaucoma. *Aust N Z J Ophthalmol* 1988; **16**: 245-250 [PMID: [3179051](#) DOI: [10.1111/j.1442-9071.1988.tb01216.x](#)]
- 4 **Jacoby B**, Reed JW, Cashwell LF. Malignant glaucoma in a patient with Down's syndrome and corneal hydrops. *Am J Ophthalmol* 1990; **110**: 434-435 [PMID: [2145771](#) DOI: [10.1016/s0002-9394\(14\)77033-3](#)]
- 5 **Cohen RG**, Wu HK, Schuman JS. Glaucoma with inflammatory precipitates on the trabecular meshwork: a report of Grant's syndrome with ultrasound biomicroscopy of precipitates. *J Glaucoma* 1996; **5**: 266-270 [PMID: [8795772](#)]
- 6 **Wang M**, Tan Q, Jiang H, Xia X, Wang P, Jiang J, Liu D. [Clinical analysis of malignant glaucoma after glaucoma surgery]. *Zhongnan Daxue Xuebao Yixueban* 2015; **40**: 543-548 [PMID: [26032082](#) DOI: [10.11817/j.issn.1672-7347.2015.05.015](#)]
- 7 **Saunders PP**, Douglas GR, Feldman F, Stein RM. Bilateral malignant glaucoma. *Can J Ophthalmol* 1992; **27**: 19-21 [PMID: [1555130](#) DOI: [10.1001/archophth.1964.00970020601004](#)]
- 8 **Stan C**. Glaucom malign bilateral--prezentare de caz. *Conexiuni Medicale* 2007; 45-47 [DOI: [10.33311/conmed.2007.3.3.10](#)]
- 9 **Sharma A**, Sii F, Shah P, Kirkby GR. Vitrectomy-phacoemulsification-vitrectomy for the management of aqueous misdirection syndromes in phakic eyes. *Ophthalmology* 2006; **113**: 1968-1973 [PMID: [17074562](#) DOI: [10.1016/j.ophtha.2006.04.031](#)]
- 10 **Wang MH**, Li QM, Dong HT, Dong SQ, Li Y, Zheng CY. Ahmed Valves vs Trabeculectomy Combined with Pans Plana Vitrectomy for Neovascular Glaucoma with Vitreous Hemorrhage. *Eur J Ophthalmol* 2017; **27**: 774-780 [PMID: [28430329](#) DOI: [10.5301/ejo.5000973](#)]
- 11 **Martínez-de-la-Casa JM**, García-Feijó J, Castillo A, Polo V, Larrosa JM, Pablo L, García-Sánchez J. Glaucoma maligno tras cirugía combinada de implante valvular de Ahmed y facoemulsificación en el tratamiento del glaucoma crónico por cierre angular. *Arch Soc Esp Oftalmol* 2005; **80**: 11 [DOI: [10.4321/s0365-66912005001100009](#)]
- 12 **Balekudaru S**, Choudhari NS, Rewri P, George R, Bhende PS, Bhende M, Lingam V, Lingam G. Surgical management of malignant glaucoma: a retrospective analysis of fifty eight eyes. *Eye (Lond)* 2017; **31**: 947-955 [PMID: [28257136](#) DOI: [10.1038/eye.2017.32](#)]
- 13 **Gao L**, Zhou Y, Li N, Liu X, Cong F. Lens Epithelial Mesenchymal Transition and Long-Term Chronic Inflammation After Cataract Surgery. *Ocul Immunol Inflamm* 2022; 1-3 [PMID: [35708475](#) DOI: [10.1080/092737948.2022.2088567](#)]





Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

