

World Journal of *Clinical Cases*

World J Clin Cases 2023 May 26; 11(15): 3369-3663



Contents

Thrice Monthly Volume 11 Number 15 May 26, 2023

REVIEW

- 3369 Superior mesenteric artery syndrome: Diagnosis and management
Oka A, Awoniyi M, Hasegawa N, Yoshida Y, Tobita H, Ishimura N, Ishihara S

MINIREVIEWS

- 3385 Astrocytes in the central nervous system and their functions in health and disease: A review
Gradisnik L, Velnar T
- 3395 Progress in diagnosis and treatment of acute injury to the anterior talofibular ligament
Chen RP, Wang QH, Li MY, Su XF, Wang DY, Liu XH, Li ZL
- 3408 Synchronous manifestation of colorectal cancer and intraductal papillary mucinous neoplasms
Mirchev MB, Boeva I, Peshevska-Sekulovska M, Stoitsov V, Peruhova M
- 3418 Clinical infections in neurosurgical oncology: An overview
Velnar T, Kocivnik N, Bosnjak R
- 3434 Effectiveness and safety of subthreshold vibration over suprathreshold vibration in treatment of muscle fatigue in elderly people
Mohamed AA, Khaled E, Hesham A, Khalf A

ORIGINAL ARTICLE

Clinical and Translational Research

- 3444 Establishment of a prognostic model related to tregs and natural killer cells infiltration in bladder cancer
Yang YJ, Xu XQ, Zhang YC, Hu PC, Yang WX

Retrospective Study

- 3457 New native tissue repair for pelvic organ prolapse: Medium-term outcomes of laparoscopic vaginal stump-round ligament fixation
Kakinuma T, Kaneko A, Kakinuma K, Imai K, Takeshima N, Ohwada M
- 3464 Demographic characteristics of patients who underwent anterior cruciate ligament reconstruction at a tertiary care hospital in India
Malv SK, Mahmood A, Vatsya P, Garika SS, Mittal R, Nagar M
- 3471 Usefulness of transcatheter arterial embolization for eighty-three patients with secondary postpartum hemorrhage: Focusing on difference in angiographic findings
Kim BM, Jeon GS, Choi MJ, Hong NS
- 3481 Chronic otitis media and middle ear variants: Is there relation?
Gökharman FD, Şenbil DC, Aydın S, Karavaş E, Özdemir Ö, Yalçın AG, Koşar PN

Observational Study

- 3491** Observation of the effect of angiojet to treat acute lower extremity arterial embolization
Meng XH, Xie XP, Liu YC, Huang CP, Wang LJ, Liu HY, Fang X, Zhang GH
- 3502** Outbreak of methanol-induced optic neuropathy in early COVID-19 era; effectiveness of erythropoietin and methylprednisolone therapy
Tabatabaei SA, Amini M, Haydar AA, Soleimani M, Cheraqpour K, Shahriari M, Hassanian-Moghaddam H, Zamani N, Akbari MR

META-ANALYSIS

- 3511** Impact of heart failure on outcomes in patients with sepsis: A systematic review and meta-analysis
Zhu MY, Tang XK, Gao Y, Xu JJ, Gong YQ

CASE REPORT

- 3522** New clinical application of digital intraoral scanning technology in occlusal reconstruction: A case report
Hou C, Zhu HZ, Xue B, Song HJ, Yang YB, Wang XX, Sun HQ
- 3533** Rare adult neuronal ceroid lipofuscinosis associated with *CLN6* gene mutations: A case report
Wang XQ, Chen CB, Zhao WJ, Fu GB, Zhai Y
- 3542** Enzyme replacement therapy in two patients with classic Fabry disease from the same family tree: Two case reports
Harigane Y, Morimoto I, Suzuki O, Temmoku J, Sakamoto T, Nakamura K, Machii K, Miyata M
- 3552** Immune-mediated necrotizing myopathy: Report of two cases
Chen BH, Zhu XM, Xie L, Hu HQ
- 3560** Retroperitoneal cavernous hemangioma misdiagnosed as lymphatic cyst: A case report and review of the literature
Hou XF, Zhao ZX, Liu LX, Zhang H
- 3571** Malignant melanoma resection and reconstruction with the first manifestation of lumbar metastasis: A case report
Guo ZX, Zhao XL, Zhao ZY, Zhu QY, Wang ZY, Xu M
- 3578** Promising way to address massive intragastric clotting in patients with acute upper gastrointestinal bleeding: A case report
Liu SX, Shi B, Liu YF, Shan JY, Sun B
- 3583** Pyogenic spondylitis caused by *Escherichia coli*: A case report and literature review
Zou LC, Qian J, Bian ZY, Wang XP, Xie T
- 3592** Primary ovarian choriocarcinoma occurring in a postmenopausal woman: A case report
Dai GL, Tang FR, Wang DQ

- 3599** Treatment of severe open bite and mandibular condyle anterior displacement by mini-screws and four second molars extraction: A case report
Huang ZW, Yang R, Gong C, Zhang CX, Wen J, Li H
- 3612** Application of apical negative pressure irrigation in the nonsurgical treatment of radicular cysts: A case report
Chen GP, Zhang YZ, Ling DH
- 3619** Treatment of postherpetic neuralgia by bone marrow aspirate injection: A case report
Honda Pazili T
- 3625** Non-target lung embolization during portal vein embolization due to an unrecognized portosystemic venous fistula: A case report
Alharbi SR, Bin Nasif M, Alwaily HB
- 3631** Acute abdomen caused by spontaneous rupture of degenerative hysteromyoma during pregnancy: A case report
Xu Y, Shen X, Pan XY, Gao S
- 3637** Atypical progress of frozen shoulder after COVID-19 vaccination: A case report
Jo HS, Kim HM, Han JY, Park HK
- 3643** Co-existing squamous cell carcinoma and chronic myelomonocytic leukemia with *ASXL1* and *EZH2* gene mutations: A case report
Deng LJ, Dong Y, Li MM, Sun CG
- 3651** Diagnosis based on electromagnetic navigational bronchoscopy-guided biopsied peripheral lung lesions in a 10-year-old girl: A case report
Meng FZ, Chen QH, Gao M, Zeng L, Lin JR, Zheng JY
- 3658** Relationship between intralobar pulmonary sequestration and type A aortic dissection: A case report
Wang YJ, Chen YY, Lin GH

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Gulali Aktas, MD, Professor, Department of Internal Medicine, Abant Izzet Baysal University Hospital, Bolu 14030, Turkey. draliaktas@yahoo.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

May 26, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



New clinical application of digital intraoral scanning technology in occlusal reconstruction: A case report

Chao Hou, Hua-Zhang Zhu, Bai Xue, Hong-Jie Song, Ying-Bei Yang, Xiao-Xue Wang, Hui-Qiang Sun

Specialty type: Dentistry, oral surgery and medicine

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): B, B

Grade C (Good): 0

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Galiatsatos A, Greece; Nagamine T, Japan

Received: December 24, 2022

Peer-review started: December 24, 2022

First decision: February 17, 2023

Revised: February 28, 2023

Accepted: April 13, 2023

Article in press: April 13, 2023

Published online: May 26, 2023



Chao Hou, Hua-Zhang Zhu, Ying-Bei Yang, Xiao-Xue Wang, Department of Prosthodontics, School and Hospital of Stomatology, Cheeloo College of Medicine, Shandong University, Jinan 250010, Shandong Province, China

Chao Hou, Department of Stomatology, Zaozhuang Municipal Hospital, Zaozhuang 277100, Shandong Province, China

Bai Xue, Department of Audit, Shandong Youth Political College, Jinan 250103, Shandong Province, China

Hong-Jie Song, Department of Stomatology, Chengdu Second People's Hospital, Chengdu 610021, Sichuan Province, China

Hui-Qiang Sun, Department of Prosthodontics, School and Hospital of Stomatology, Cheeloo College of Medicine, Shandong University & Shandong Key Laboratory of Oral Tissue Regeneration & Shandong Engineering Laboratory for Dental Materials and Oral Tissue Regeneration & Shandong Provincial Clinical Research Center for Oral Diseases, Jinan 250012, Shandong Province, China

Corresponding author: Hui-Qiang Sun, Doctor, MD, Chief Doctor, Department of Prosthodontics, School and Hospital of Stomatology, Cheeloo College of Medicine, Shandong University & Shandong Key Laboratory of Oral Tissue Regeneration & Shandong Engineering Laboratory for Dental Materials and Oral Tissue Regeneration & Shandong Provincial Clinical Research Center for Oral Diseases, No. 44-1 Wenhuxilu Road, Jinan 250012, Shandong Province, China. whitedove69@163.com

Abstract

BACKGROUND

Digital intraoral scanning, although developing rapidly, is rarely used in occlusal reconstruction. To compensate for the technical drawbacks of current occlusal reconstruction techniques, such as time consumption and high technical requirements, digital intraoral scanning can be used in clinics. This report aims to provide a way of selecting the most suitable maxillo-mandibular relationship (MMR) during recovery.

CASE SUMMARY

A 68-year-old man with severely worn posterior teeth underwent occlusal reconstruction with fixed prosthesis using digital intraoral scanning. A series of digital models in different stages of treatment were obtained, subsequently

compared, and selected using digital intraoral scanning together with traditional measurements, such as cone beam computed tomography, joint imaging, and clinical examination. Using digital intraoral scanning, the MMR in different stages of treatment was accurately recorded, which provided feasibility for deciding the best occlusal reconstruction treatment, made the treatment process easier, and improved patient satisfaction.

CONCLUSION

This case report highlights the clarity, recordability, repeatability, and selectivity of digital intraoral scanning to replicate and transfer the MMR during occlusal reconstruction, expanding new perspectives for its design, fabrication, and postoperative evaluation.

Key Words: Occlusal reconstruction; Digital intraoral scanning; Maxillo-mandibular relationship; Cone beam computed tomography; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Selection of the most appropriate maxillo-mandibular relationship and transfer of occlusal information are crucial steps in occlusal reconstruction. The traditional method is complicated, has high technical sensitivity, and may cause the accumulation of errors. In this case, a digital occlusal relation transfer method was established using digital intraoral scanning to improve the accuracy and efficiency of occlusal reconstruction. This method not only prevents the loss of the optimal maxillo-mandibular relationship but also directly transfers the facial morphology of the temporary prosthesis to the final prosthesis, thus achieving a good transfer of aesthetic and functional information. Digital intraoral scanning makes recording the jaw position relationship between the maxilla and mandible, which is the most important but challenging aspect in the treatment process, clear, recordable, reproducible, and selectable.

Citation: Hou C, Zhu HZ, Xue B, Song HJ, Yang YB, Wang XX, Sun HQ. New clinical application of digital intraoral scanning technology in occlusal reconstruction: A case report. *World J Clin Cases* 2023; 11(15): 3522-3532

URL: <https://www.wjgnet.com/2307-8960/full/v11/i15/3522.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i15.3522>

INTRODUCTION

Occlusal reconstruction refers to all treatment methods that can change the existing abnormal occlusal contact relationship to restore the patient's normal occlusal function[1]. The key to occlusal reconstruction is the use of occlusal pads, mockups, temporary crowns, and other diagnostic temporary restorations, from noninvasive, minimally invasive to invasive procedures, and reversible to irreversible treatments, to verify, adjust, and run in the diagnostic occlusal relationship established in the step by step occlusal design process, and finally obtain the final occlusal relationship of the prosthesis that is most suitable for patients[2,3]. In traditional occlusal reconstruction, acquiring the maxillo-mandibular relationship (MMR) primarily depends on the doctors' experience and patients' subjective feedback and thus lacks an objective evaluation. In recent years, digital workflow, computer-aided design (CAD), and computer-aided manufacturing (CAM) have become indispensable components of prosthodontics[4]. Digital intraoral scanning is an innovative technology that promotes the development of dental prostheses. Digital intraoral scanning technology uses a small probe, an optical scanning head, to directly obtain the three-dimensional morphology and information of the color and texture of the surface of soft and hard tissues, such as the teeth, gums, and mucosa, in the patient's mouth, to help restore the three-dimensional morphology and color texture of the oral tissues to the greatest extent. Digital intraoral scanning allows the direct transformation of the scanned information of the intraoral dentition to a digital dental model. This method reduces the steps of making impressions and models; avoids the influence of human operation and materials on accuracy; and avoids errors in making, transferring, and pouring plaster models. Compared with traditional methods, digital intraoral scanning technology has the advantages of rapid speed, high precision, and patient comfort. Dentists are increasingly inclined to use digital intraoral scanning instead of traditional impressions[5]. Currently, there are a few reports on the application of digital intraoral scanning technology in occlusal reconstruction. With the help of digital tools, we accurately recorded the MMR in each stage, combined with cone beam computed tomography (CBCT), joint imaging, and clinical examination, to guide

clinicians in selecting a comfortable occlusal relationship at any stage for the final reconstruction. These records are intended to provide new ideas for the design of occlusal reconstruction, prosthesis production, and postoperative evaluation of fixed prostheses.

CASE PRESENTATION

Chief complaints

A 68-year-old man visited our hospital for crown restoration because of root canal treatment of the upper right posterior teeth 2 wk previously.

History of present illness

The patient's teeth were worn to varying degrees. One month prior, the patient's upper right posterior teeth developed symptoms of pulpitis due to severe enamel wear and dentin exposure for a long time. He was treated with root canal therapy. He returned to see a dentist for crown restoration.

History of past illness

The patient denied a history of systemic diseases, such as hypertension, coronary heart disease, and diabetes. He reported no history of food or drug allergy, trauma, surgery, or blood transfusion, and his vaccination history was unknown.

Personal and family history

No individual or family history of mental illness was reported.

Physical examination

The patient's facial morphology was left-right symmetrical, the lower 1/3 height of the face was shorter, the jaw space at rest was normal, the degree of opening was normal, the opening type was biased at the initial stage of closure, and there was no clicking or tenderness in either joint. The impact force of the bilateral condyles was large, according to the external auditory canal digital examination. All the teeth were worn to varying degrees, overbite and overjet were normal, and the incisal edge was slightly worn. The enamel of 15-17, 26, 27, 36, 37, and 44-47 was worn in large areas, the mesial lingual tips of 16 and 26 were worn to 1 mm below the gum, and most teeth were probe examined (+/-), cold stimulated (-), and percussed (-). Dental plaque 1°, dental calculus (+), pigment segment (+), and localized gingival congestion and swelling were observed. There were no loose teeth, and the periodontal probing depth was 2-4 mm (Figure 1).

Imaging examinations

Initial temporomandibular joint CBCT: The posterior space of both temporomandibular joints was narrow, the anterior space increased, the upper space increased slightly, and the condyle moved backward and upward (Figure 2A).

CBCT of the temporomandibular joint when wearing the occlusal pad: The anterior space of the joint decreased, the posterior space increased, the upper space slightly increased, and the condyle moved forward and downward. The left joint space was normal, and the right posterior joint space improved compared with when not wearing the occlusal pad (Figure 2B).

CBCT of the temporomandibular joint 1 mo after wearing the temporary crowns: The left joint space was normal, and the right posterior joint space was slightly wider than when wearing the jaw pad (Figure 2C).

CBCT of the temporomandibular joint after bonding the final denture: The left joint space was normal, the right posterior joint space was wider than before, and the bilateral temporomandibular joint space was improved (Figure 2D).


FINAL DIAGNOSIS

Heavy abrasion.

TREATMENT

Preparation before restoration

CBCT was performed before restoration (Figure 2A). Upper and lower alginate impressions were taken, the occlusal relationship was recorded, facial arch was transferred to the articulator, the lifting distance was analyzed and determined, and a dental diagnostic wax-up of 14-17, 24-27, 34-37, and 44-47 was



DOI: 10.12998/wjcc.v11.i15.3522 Copyright ©The Author(s) 2023.

Figure 1 Photo taken before occlusion reconstruction. A and B: Initial intraoral photographs; C and D: Initial face photograph.

made (Figure 3A), including the occlusal relationship, cusp and fossa relationship, and aesthetic adjustment. Simultaneously, the jaw pad was made according to the same jaw position relationship.

Preparation of the first digital model

The patient wore a maxillary occlusal pad and was instructed to visit once a week (Figure 3B). After wearing an occlusal pad for 3 mo, the patient reported no discomfort, and CBCT of the temporomandibular joint was performed (Figure 2B). At that time, the patient wore an occlusal pad for a digital intraoral scan (Figure 4A).

Fabrication a temporary prosthesis


The occlusal pad was divided into two at the central line for standby (Figure 3C). According to the principle of segmented preparation, tooth preparation was first performed. Referring to the Zhao *et al*[6] method of facial arch transfer to a cross-mounting articulator, the adapted jaw position and vertical distance records determined by the transitional treatment were transferred to the articulator, and temporary restorations for 14-17, 24-27, 34-37, and 44-47 for routine adjustments were made (Figure 5).

Preparation of the second digital model

After using the temporary crowns for 1 mo, CBCT of the temporomandibular joint was performed (Figure 2C). Joints, masticatory muscles, occlusal contact, and periodontal health of the teeth were re-evaluated, and formal restoration after full communication with the patient was commenced. Digital intraoral scanning models were then taken with the temporary crowns (Figure 4B).

Selection of the final occlusal relationship

Before the final restoration was made, the most suitable MMR to meet the requirements of the patient was determined. After completion of the above steps, two occlusal reconstruction models were obtained and retained through digital intraoral scanning (Figure 4). To select the most suitable model for occlusal reconstruction of the patient, comparisons were made using Shape 3D viewer software. With the image of the upper jaw used as the reference coordinate for the comparisons, the models overlapped each other in the same spatial position (Figure 6A). The yellow model was the jaw position relationship obtained with the occlusal pad, while the blue model was obtained after wearing the temporary crowns. A locally amplified image (Figure 6B) illustrated the difference in the MMR; the anterior edge of the lateral labial margin of the mandibular central incisor was shifted upward and medially by approximately 0.861 mm in the MMR with the temporary crowns compared with that with the occlusal pad. In addition, at the end of the mock-up phase, the occlusal reconstruction vertical distance decreased, overjet decreased slightly, and the patient had a more comfortable bite (Figure 7). Finally, we




DOI: 10.12998/wjcc.v11.i15.3522 Copyright ©The Author(s) 2023.

Figure 2 Cone beam computed tomography imaging. A: Cone beam computed tomography (CBCT) at the beginning; B: CBCT after wearing an occlusal pad for 3 mo; C: CBCT 1 mo after wearing the temporary crown; D: CBCT rechecked 2 wk after wearing the final denture. 1 and 2 are the coronal and sagittal planes of the right temporomandibular joint, respectively; 3 and 4 are the coronal and sagittal planes of the left temporomandibular joint, respectively.

determined that the occlusal relationship of the second oral scan was most suitable for the final restoration and reconstruction for the patient.


Trying on the final restoration

After removing the temporary crowns, 16 altron all-ceramic crowns were used. In addition, other aspects such as decreased overjet and better condylar position resulted in a more comfortable bite and a pleasing appearance.



DOI: 10.12998/wjcc.v11.i15.3522 Copyright ©The Author(s) 2023.

Figure 3 Preparation before restoration. A: Diagnostic wax pattern; B: Intraoral photographs of patients wearing maxillary occlusal pad; C: Divide the occlusal pad at the center line for standby.



DOI: 10.12998/wjcc.v11.i15.3522 Copyright ©The Author(s) 2023.

Figure 4 Occlusal reconstruction models obtained through digital intraoral scanning. A: Digital intraoral scanning with occlusal pad; B: Digital intraoral scanning after fine tooth preparation.


OUTCOME AND FOLLOW-UP

Eventually, after wearing the all-ceramic crowns for 2 wk, the patient felt beautiful and comfortable, without obvious discomfort or temporomandibular joint-related symptoms, and underwent CBCT (Figure 2D). At 1 mo follow-up visit, the patient's face shape was normal, the muscle strength of both sides of the maxillofacial region was symmetrical, and the occlusal relationship was excellent during oral examination. The jaw position did not slide during clenching compared with the jaw position during light bite. The patient reported an improved quality of life after treatment. A re-evaluation visit was in 3 mo; if there were no obvious symptoms, the patient could visit yearly.




DOI: 10.12998/wjcc.v11.i15.3522 Copyright ©The Author(s) 2023.

Figure 5 Intraoral photographs of temporary crowns worn on the preliminary prepared posterior teeth.



DOI: 10.12998/wjcc.v11.i15.3522 Copyright ©The Author(s) 2023.

Figure 6 Selection of the final occlusal relationship. A: Comparison of maxillary position between the end of occlusal pad and the end of mock up; B: A local magnified image showing that at the end of the mock up phase, compared to when wearing the occlusal pad, the most anterior end of the lateral labial margin of the mandibular central incisor is shifted upward and medially by approximately 0.861 mm.



DOI: 10.12998/wjcc.v11.i15.3522 Copyright ©The Author(s) 2023.

Figure 7 Comparison of anterior tooth overjet photographs. A: Anterior tooth overjet photographs at occlusal pad end-stage; B: Anterior tooth overjet photographs at temporary coronal end-stage.

DISCUSSION

Digital technology of stomatology refers to the technology that helps dentists realize an accurate, efficient, automatic, and intelligent diagnosis and treatment of oral diseases with the help of digital hardware or software. Digital hardware or software includes but is not limited to intraoral scanning, CBCT, digital diagnosis and design, digital occlusion analysis based on virtual articulators, artificial

intelligence, navigation, robotics, and related materials. Dental CAD and CAM are typical applications of digital stomatology. The process includes digital impression, denture CAD, and CAM[7]. In traditional occlusal reconstruction, the acquisition of jaw position relationship largely depends on the doctors' experience and patients' subjective feedback, and therefore lacks an objective evaluation. Digital tools enable an objective evaluation of the treatment to help verify whether the reconstructed occlusion is coordinated with the functional state of the oral and maxillofacial system. The application of digital technology can help simulate and predict the effect of reconstructed occlusion and the accuracy of replication and transfer of occlusal relationships before invasive procedures. Treatment that was originally highly dependent on clinical experience has become procedural and reproducible with digitalization, and the technical threshold for doctors is relatively low. Digitalization can help achieve personalized medical treatment that conforms to patients' anatomical and physiological states and improve the pathological state, patient satisfaction with treatment, and quality and efficiency of communication between doctors and patients and between doctors and technicians. Digitalization in prosthodontics has been widely accepted by peers and is regarded as an irreversible trend. There is also consensus to build a complete process of digital prosthodontics – digital diagnosis, digital designing, digital manufacturing, and digital treatment"[8,9].

When the patient adapts to the new occlusal relationship formed using the occlusal pad and has no obvious subjective discomfort, objective assessment is performed by palpation with both hands to check the strength of contraction of the temporal muscle and CBCT to check whether the condylar position is normal. Then, the occlusal pad is debugged, and the first digital intraoral scanning is performed to collect the digital information of the MMR, to obtain the first type of occlusal relationship. After a certain period of temporary crown treatment with varying degrees of adjustment, running-in diagnosis, and verification, depending on the adaptability of patients, digital intraoral scanning is used once again to acquire another type of occlusal relationship.

In the widely used traditional method, the key to achieving occlusal reconstruction lies in the clinical experience of doctors to guarantee the accuracy of MMR in the entire process of copying, transferring from the occlusal pad to the temporary crowns, and making the final restorations. Moreover, with the restriction of its mock-up stage, the MMR available is, simply, the one acquired in the final coronal phase and is, therefore, comparatively limited. In this case report, digital intraoral scanning accurately recorded a series of digital information of the MMR during occlusal pad commissioning and temporary crown wearing. These data can help us find the best position for occlusal reconstruction, thus minimizing the errors generated in the tedious process of recording the best MMR relationship.

In traditional occlusal reconstruction, adaptive grinding of diagnostic temporary restorations, such as occlusal pads and temporary crowns, has advantages and disadvantages. Adjusting the occlusal relationship in the horizontal direction can better reduce the overbite of the anterior teeth, which is conducive to the condyle sliding upward and forward to achieve a satisfying central position relationship. The vertical distance will decrease due to wear, thus reducing the increased vertical distance in the process of occlusal reconstruction. Digital intraoral scanning technology can not only prevent the loss of optimal jaw relationship caused by various errors in the occlusal pad commissioning stage and temporary crown fabrication and grinding process but also provide a chance to choose the appropriate MMR in the clinic. The application of digital intraoral scanning in the occlusal reconstruction of fixed prosthesis makes recording the jaw position relationship between the maxilla and mandible, which is the most essential but fuzzy aspect in the treatment process, clear, recordable, reproducible, and selectable.

We can select either the occlusal relationship of the first oral scan (Figure 4A) or the second (Figure 4B) for the final reconstruction. In the digital software system, at the end of the mock-up phase, compared to when wearing the occlusal pad, the most anterior end of the lateral labial margin of the mandibular central incisor was shifted upward and medially by approximately 0.861 mm. Compared with digitally recorded information on the occlusal relationship between the end of the temporary crown and the end of the occlusal pad wearing, the occlusal reconstruction vertical distance decreased, and the overjet decreased slightly. There were supernumerary teeth between the maxillary central incisors of this patient; however, he refused extraction of the supernumerary teeth and restoration of the occlusal relationship in the anterior tooth area. The occlusal relationship of the temporary crown stage, with a slightly smaller overjet, was used for posterior teeth occlusal reconstruction, which helped retain the guiding function of the anterior teeth to a greater extent and was more conducive to the comfort of the patient's temporomandibular joint. By comparing the occlusal relationship of the patient's two oral scans, we determined that the occlusal relationship of the second oral scan was most suitable for the final restoration and reconstruction. Digital intraoral scanning has two advantages in occlusal reconstruction: (1) The occlusal relationship can be accurately reproduced and transferred stepwise from the occlusal pad to the temporary crown to the final restoration; and (2) It can establish a perfect occlusion that includes correct lateral occlusal guidance, uniform contact during centric occlusion, and a reasonable opening angle during non-centric motion.

In addition, the final occlusal relationship replication and transfer of diagnostic temporary restorations in traditional full-mouth occlusal reconstruction often use cross-mounting articulators. The traditional steps in this method include taking multiple impressions, pouring models, and recording multiple occlusions. Each step has potential for errors, and they cannot be quantified. These complex

steps and potential errors can affect the occlusal accuracy of the final prosthesis and often requires considerable time to adjust in the clinic[10-12]. Digital intraoral scanning can minimize these errors and simplify the process. Digital intraoral scanning allows the direct transformation of the scanned information of the intraoral dentition to a digital dental model, three-dimensional prediction of the repair effect, and transfer the treatment design to the final prosthesis through digital design and production technology to enable the "what you see is what you get" treatment effect[13,14]. Using digital dental scanning equipment, the morphology of temporary restorations after functional adaptive wear, occlusal contact relationship, and refined abutment preparation can be recorded and the digital model established and matched. Digital intraoral scanning has become an important dental operation technology. Based on the data obtained, crown restoration and the production of implant superstructures, orthodontic devices, removable partial denture brackets, and other structures, are carried out [15-19]. The rigidity of the digital model can avoid transmission errors caused by the variety of mandibular movements and the errors while making occlusal records in the restoration process. It can also shorten the clinical operation time and steps while improving accuracy[20]. Treatment costs are expected to decline with the continuous development of digital technology. We believe that intraoral scanning will become popular in the next few years[21].

Owing to the close relationship between digitalization and artificial intelligence (AI), it is believed that with the rapid development of AI technology in the future, the application of digitalization technology in clinical practice will become increasingly more common and operation will become increasingly simple and standardized; this in turn will provide a strong support for the development of AI medicine.

CBCT has been widely used in various fields of stomatology, including orthodontic three-dimensional cephalometry, orthodontic treatment plan design, implant surgery planning and guide plate design, maxillofacial surgery diagnosis and surgery design, maxillofacial surgery navigation implementation and guide plate design, temporomandibular joint diagnosis, periodontal diagnosis, root canal apical morphology and root fracture diagnosis, sleep apnea syndrome, and soft palate and pharyngeal morphology diagnosis. It is the most widely used and promising digital technology in stomatology. In addition to routine clinical examination, the examination methods in the occlusal reconstruction process of the fixed prosthesis also include the use of imaging to evaluate the structural changes of the oral and maxillofacial system, evaluate the motor functions of the oral and maxillofacial system, detect bruits of the temporomandibular joint, and examine the electrophysiological function of the masticatory muscles[22,23]. In this case, CBCT was used for occlusal reconstruction in a patient with severe abrasion. By a combination of the clinical symptoms of the patient before repair and 1 mo after permanent repair and the morphological changes of the temporomandibular joint space and condylar position shown by CBCT, the clinical treatment effect of fixed denture occlusal reconstruction was objectively evaluated. Researchers believe that long-term dentition loss or serious wear leads to reduction in the vertical distance, and the condyle then shifts backward[24]. Consistent with the results of previous studies, in this patient, CBCT showed that the anterior space of the TMJ was significantly larger than the posterior space before occlusal reconstruction and that the condyle was displaced posteriorly. After digital intraoral scan-assisted occlusal reconstruction, CBCT showed that the left joint space was normal, the right posterior joint space was wider than before, and the space of the bilateral temporomandibular space was better. The patient's temporomandibular joint had not undergone pathological changes; therefore, only the jaw position and movement adapted to the physiological state of the joint muscles needed to be determined by digital equipment. Using this case as a reference, restoration with an accurate occlusal relationship can be achieved.

CONCLUSION

This clinical report showed that the application of digital intraoral scanning for occlusal reconstruction of a fixed prosthesis can accurately record the occlusal relationship of jaw position in different stages to avoid the loss of an optimal jaw relationship. Digital intraoral scanning makes recording the jaw position relationship between the maxilla and mandible, which is the most important but challenging aspect in the treatment process, clear, recordable, reproducible, and selectable. We hope that this clinical report will provide a new clinical reference for the design, fabrication, and postoperative evaluation of occlusal reconstruction of fixed prostheses. These conclusions should be verified with the treatment of more patients in the future.

ACKNOWLEDGEMENTS

The authors are grateful to Mr. Teng K for technical support.

FOOTNOTES

Author contributions: Sun HQ designed the research study; Hou C, Yang YB, and Wang XX performed the research; Xue B and Song HJ contributed the analytic tools and analyzed the data; Hou C and Zhu HZ wrote the manuscript; all authors have read and approved the final manuscript.

Informed consent statement: Informed written consent was obtained from the patient for publication of this report and any accompanying images.

Conflict-of-interest statement: The authors declare that they have no conflict of interest to disclose.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Hong-Jie Song 0000-0003-3784-9211; Hui-Qiang Sun 0000-0002-8878-4676.

S-Editor: Yan JP

L-Editor: A

P-Editor: Yuan YY

REFERENCES

- 1 **Edelhoff D**, Probst F, Ehrenfeld M, Prandtner O, Schweiger J, Liebermann A. Interdisciplinary full-mouth rehabilitation for redefining esthetics, function, and orofacial harmony. *J Esthet Restor Dent* 2019; **31**: 179-189 [PMID: 30609191 DOI: 10.1111/jerd.12455]
- 2 **Li DL**, Tan JG. Creating esthetic provisional restorations: step by step. *Zhonghua Kou Qiang Yi Xue Za Zhi* 2021; **56**: 226-230 [PMID: 33557512]
- 3 **Fabbri G**, Cannistraro G, Pulcini C, Sorrentino R. The full-mouth mock-up: a dynamic diagnostic approach (DDA) to test function and esthetics in complex rehabilitations with increased vertical dimension of occlusion. *Int J Esthet Dent* 2018; **13**: 460-474 [PMID: 30302437]
- 4 **Gao H**, Liu X, Liu M, Yang X, Tan J. Accuracy of three digital scanning methods for complete-arch tooth preparation: An in vitro comparison. *J Prosthet Dent* 2022; **128**: 1001-1008 [PMID: 33736864 DOI: 10.1016/j.prosdent.2021.01.029]
- 5 **Christensen GJ**. Will digital impressions eliminate the current problems with conventional impressions? *J Am Dent Assoc* 2008; **139**: 761-763 [PMID: 18520000 DOI: 10.14219/jada.archive.2008.0258]
- 6 **Zhao YW**, Gao R, Sun HQ. The Protocol of Fixed Reconstruction for Severely Worn Teeth Combined with Anterior Deep Bite. *Case Rep Dent* 2017; **2017**: 9378091 [PMID: 28912979 DOI: 10.1155/2017/9378091]
- 7 **Duret F**, Blouin JL, Duret B. CAD-CAM in dentistry. *J Am Dent Assoc* 1988; **117**: 715-720 [PMID: 3058771 DOI: 10.14219/jada.archive.1988.0096]
- 8 **van Noort R**. The future of dental devices is digital. *Dent Mater* 2012; **28**: 3-12 [PMID: 22119539 DOI: 10.1016/j.dental.2011.10.014]
- 9 **Rekow ED**. Digital dentistry: The new state of the art - Is it disruptive or destructive? *Dent Mater* 2020; **36**: 9-24 [PMID: 31526522 DOI: 10.1016/j.dental.2019.08.103]
- 10 **Liu Y**. Several important issues concerning occlusal reconstruction. *Hua Xi Kou Qiang Yi Xue Za Zhi* 2020; **38**: 357-363 [PMID: 32865351 DOI: 10.7518/hxkq.2020.04.001]
- 11 **Kordass B**, Gärtner C, Söhnel A, Bisler A, Voss G, Bockholt U, Seipel S. The virtual articulator in dentistry: concept and development. *Dent Clin North Am* 2002; **46**: 493-506, vi [PMID: 12222093 DOI: 10.1016/s0011-8532(02)00006-x]
- 12 **Lee W**, Kwon HB. Vertical repositioning accuracy of magnetic mounting systems on 4 articulator models. *J Prosthet Dent* 2018; **119**: 446-449 [PMID: 28689914 DOI: 10.1016/j.prosdent.2017.04.006]
- 13 **Liu YS**, Ye HQ, Gu M, Lv LW, Sun YC, Zhao YJ, Zhou YS. [Application of patient-participated digital design in esthetic rehabilitation of anterior teeth]. *Beijing Da Xue Xue Bao Yi Xue Ban* 2014; **46**: 90-94 [PMID: 24535356]
- 14 **Coachman C**, Georg R, Bohner L, Rigo LC, Sesma N. Chairside 3D digital design and trial restoration workflow. *J Prosthet Dent* 2020; **124**: 514-520 [PMID: 31924341 DOI: 10.1016/j.prosdent.2019.10.015]
- 15 **Miyazaki T**, Hotta Y, Kunii J, Kuriyama S, Tamaki Y. A review of dental CAD/CAM: current status and future perspectives from 20 years of experience. *Dent Mater J* 2009; **28**: 44-56 [PMID: 19280967 DOI: 10.4012/dmj.28.44]
- 16 **Goo CL**, Tan KB. Fabricating CAD/CAM Implant-Retained Mandibular Bar Overdentures: A Clinical and Technical Overview. *Case Rep Dent* 2017; **2017**: 9373818 [PMID: 28396807 DOI: 10.1155/2017/9373818]
- 17 **Alghazzawi TF**. Advancements in CAD/CAM technology: Options for practical implementation. *J Prosthodont Res* 2016; **60**: 72-84 [PMID: 26935333 DOI: 10.1016/j.jpor.2016.01.003]

- 18 **Kattadiyil MT**, Mursic Z, AlRumaih H, Goodacre CJ. Intraoral scanning of hard and soft tissues for partial removable dental prosthesis fabrication. *J Prosthet Dent* 2014; **112**: 444-448 [PMID: 24882595 DOI: 10.1016/j.prosdent.2014.03.022]
- 19 **Williams RJ**, Bibb R, Eggbeer D, Collis J. Use of CAD/CAM technology to fabricate a removable partial denture framework. *J Prosthet Dent* 2006; **96**: 96-99 [PMID: 16911885 DOI: 10.1016/j.prosdent.2006.05.029]
- 20 **Li Z**, Xia Y, Chen K, Zhao H, Liu Y. Maintenance of the Maxillomandibular Position with Digital Workflow in Oral Rehabilitation: A Technical Note. *Int J Prosthodont* 2018; **31**: 280-282 [PMID: 29723325 DOI: 10.11607/ijp.5663]
- 21 **Ho CT**, Lin HH, Lo LJ. Intraoral Scanning and Setting Up the Digital Final Occlusion in Three-Dimensional Planning of Orthognathic Surgery: Its Comparison with the Dental Model Approach. *Plast Reconstr Surg* 2019; **143**: 1027e-1036e [PMID: 31033828 DOI: 10.1097/PRS.0000000000005556]
- 22 **Hanchi Z**, Tingting L, Bilu X, Zhongjie L, Yang L. Features of condylography curves. *Hua Xi Kou Qiang Yi Xue Za Zhi* 2017; **35**: 555-560 [PMID: 29188656 DOI: 10.7518/hxkq.2017.05.021]
- 23 **Sharma S**, Crow HC, Kartha K, McCall WD Jr, Gonzalez YM. Reliability and diagnostic validity of a joint vibration analysis device. *BMC Oral Health* 2017; **17**: 56 [PMID: 28209141 DOI: 10.1186/s12903-017-0346-9]
- 24 **Krishnamoorthy B**, Mamatha N, Kumar VA. TMJ imaging by CBCT: Current scenario. *Ann Maxillofac Surg* 2013; **3**: 80-83 [PMID: 23662265 DOI: 10.4103/2231-0746.110069]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

