

# World Journal of *Clinical Cases*

*World J Clin Cases* 2023 May 26; 11(15): 3369-3663



## REVIEW

- 3369 Superior mesenteric artery syndrome: Diagnosis and management  
*Oka A, Awoniyi M, Hasegawa N, Yoshida Y, Tobita H, Ishimura N, Ishihara S*

## MINIREVIEWS

- 3385 Astrocytes in the central nervous system and their functions in health and disease: A review  
*Gradisnik L, Velnar T*
- 3395 Progress in diagnosis and treatment of acute injury to the anterior talofibular ligament  
*Chen RP, Wang QH, Li MY, Su XF, Wang DY, Liu XH, Li ZL*
- 3408 Synchronous manifestation of colorectal cancer and intraductal papillary mucinous neoplasms  
*Mirchev MB, Boeva I, Peshevska-Sekulovska M, Stoitsov V, Peruhova M*
- 3418 Clinical infections in neurosurgical oncology: An overview  
*Velnar T, Kocivnik N, Bosnjak R*
- 3434 Effectiveness and safety of subthreshold vibration over suprathreshold vibration in treatment of muscle fatigue in elderly people  
*Mohamed AA, Khaled E, Hesham A, Khalf A*

## ORIGINAL ARTICLE

## Clinical and Translational Research

- 3444 Establishment of a prognostic model related to tregs and natural killer cells infiltration in bladder cancer  
*Yang YJ, Xu XQ, Zhang YC, Hu PC, Yang WX*

## Retrospective Study

- 3457 New native tissue repair for pelvic organ prolapse: Medium-term outcomes of laparoscopic vaginal stump-round ligament fixation  
*Kakinuma T, Kaneko A, Kakinuma K, Imai K, Takeshima N, Ohwada M*
- 3464 Demographic characteristics of patients who underwent anterior cruciate ligament reconstruction at a tertiary care hospital in India  
*Malv SK, Mahmood A, Vatsya P, Garika SS, Mittal R, Nagar M*
- 3471 Usefulness of transcatheter arterial embolization for eighty-three patients with secondary postpartum hemorrhage: Focusing on difference in angiographic findings  
*Kim BM, Jeon GS, Choi MJ, Hong NS*
- 3481 Chronic otitis media and middle ear variants: Is there relation?  
*Gökharman FD, Şenbil DC, Aydın S, Karavaş E, Özdemir Ö, Yalçın AG, Koşar PN*

**Observational Study**

- 3491** Observation of the effect of angiojet to treat acute lower extremity arterial embolization  
*Meng XH, Xie XP, Liu YC, Huang CP, Wang LJ, Liu HY, Fang X, Zhang GH*
- 3502** Outbreak of methanol-induced optic neuropathy in early COVID-19 era; effectiveness of erythropoietin and methylprednisolone therapy  
*Tabatabaei SA, Amini M, Haydar AA, Soleimani M, Cheraqpour K, Shahriari M, Hassanian-Moghaddam H, Zamani N, Akbari MR*

**META-ANALYSIS**

- 3511** Impact of heart failure on outcomes in patients with sepsis: A systematic review and meta-analysis  
*Zhu MY, Tang XK, Gao Y, Xu JJ, Gong YQ*

**CASE REPORT**

- 3522** New clinical application of digital intraoral scanning technology in occlusal reconstruction: A case report  
*Hou C, Zhu HZ, Xue B, Song HJ, Yang YB, Wang XX, Sun HQ*
- 3533** Rare adult neuronal ceroid lipofuscinosis associated with *CLN6* gene mutations: A case report  
*Wang XQ, Chen CB, Zhao WJ, Fu GB, Zhai Y*
- 3542** Enzyme replacement therapy in two patients with classic Fabry disease from the same family tree: Two case reports  
*Harigane Y, Morimoto I, Suzuki O, Temmoku J, Sakamoto T, Nakamura K, Machii K, Miyata M*
- 3552** Immune-mediated necrotizing myopathy: Report of two cases  
*Chen BH, Zhu XM, Xie L, Hu HQ*
- 3560** Retroperitoneal cavernous hemangioma misdiagnosed as lymphatic cyst: A case report and review of the literature  
*Hou XF, Zhao ZX, Liu LX, Zhang H*
- 3571** Malignant melanoma resection and reconstruction with the first manifestation of lumbar metastasis: A case report  
*Guo ZX, Zhao XL, Zhao ZY, Zhu QY, Wang ZY, Xu M*
- 3578** Promising way to address massive intragastric clotting in patients with acute upper gastrointestinal bleeding: A case report  
*Liu SX, Shi B, Liu YF, Shan JY, Sun B*
- 3583** Pyogenic spondylitis caused by *Escherichia coli*: A case report and literature review  
*Zou LC, Qian J, Bian ZY, Wang XP, Xie T*
- 3592** Primary ovarian choriocarcinoma occurring in a postmenopausal woman: A case report  
*Dai GL, Tang FR, Wang DQ*

- 3599** Treatment of severe open bite and mandibular condyle anterior displacement by mini-screws and four second molars extraction: A case report  
*Huang ZW, Yang R, Gong C, Zhang CX, Wen J, Li H*
- 3612** Application of apical negative pressure irrigation in the nonsurgical treatment of radicular cysts: A case report  
*Chen GP, Zhang YZ, Ling DH*
- 3619** Treatment of postherpetic neuralgia by bone marrow aspirate injection: A case report  
*Honda Pazili T*
- 3625** Non-target lung embolization during portal vein embolization due to an unrecognized portosystemic venous fistula: A case report  
*Alharbi SR, Bin Nasif M, Alwaily HB*
- 3631** Acute abdomen caused by spontaneous rupture of degenerative hysteromyoma during pregnancy: A case report  
*Xu Y, Shen X, Pan XY, Gao S*
- 3637** Atypical progress of frozen shoulder after COVID-19 vaccination: A case report  
*Jo HS, Kim HM, Han JY, Park HK*
- 3643** Co-existing squamous cell carcinoma and chronic myelomonocytic leukemia with *ASXL1* and *EZH2* gene mutations: A case report  
*Deng LJ, Dong Y, Li MM, Sun CG*
- 3651** Diagnosis based on electromagnetic navigational bronchoscopy-guided biopsied peripheral lung lesions in a 10-year-old girl: A case report  
*Meng FZ, Chen QH, Gao M, Zeng L, Lin JR, Zheng JY*
- 3658** Relationship between intralobar pulmonary sequestration and type A aortic dissection: A case report  
*Wang YJ, Chen YY, Lin GH*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Gulali Aktas, MD, Professor, Department of Internal Medicine, Abant Izzet Baysal University Hospital, Bolu 14030, Turkey. draliaktas@yahoo.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

May 26, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

# Diagnosis based on electromagnetic navigational bronchoscopy-guided biopsied peripheral lung lesions in a 10-year-old girl: A case report

Fan-Zheng Meng, Qiong-Hua Chen, Man Gao, Li'e Zeng, Jie-Ru Lin, Jing-Yang Zheng

**Specialty type:** Pediatrics

**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B  
Grade C (Good): C  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Kwiecień I, Poland;  
Zeng C, United States

**Received:** February 21, 2023

**Peer-review started:** February 21, 2023

**First decision:** April 10, 2023

**Revised:** April 13, 2023

**Accepted:** April 19, 2023

**Article in press:** April 19, 2023

**Published online:** May 26, 2023



**Fan-Zheng Meng, Man Gao,** Department of Pediatrics, The First Hospital of Jilin University, Changchun 130021, Jilin Province, China

**Qiong-Hua Chen,** Department of Respiratory Medicine, The Third Clinical Medical College, Fujian Medical University, Quanzhou Women's and Children's Hospital, Quanzhou 362000, Fujian Province, China

**Li'e Zeng, Jie-Ru Lin,** Department of Respiratory Medicine, Quanzhou Women's and Children's Hospital, Quanzhou 362000, Fujian Province, China

**Jing-Yang Zheng,** Department of Respiratory Medicine, Quanzhou Women's and Children's Hospital, The Third Clinical Medical College, Fujian Medical University, Quanzhou 362000, Fujian Province, China

**Corresponding author:** Jing-Yang Zheng, MD, Professor, Department of Respiratory Medicine, Quanzhou Women's and Children's Hospital, The Third Clinical Medical College, Fujian Medical University, No. 700 Fengze Street, Fengze District, Quanzhou 362000, Fujian Province, China. [279277481@qq.com](mailto:279277481@qq.com)

## Abstract

### BACKGROUND

Electromagnetic navigational bronchoscopy (ENB) is an emerging diagnostic tool that enables practitioners to biopsy peripheral lung tissues that were previously only accessible under computed tomography (CT) guidance. However, few studies have investigated ENB use in children. Here, we report a case of a 10-year-old girl with peripheral lung lesions who complained of a 7-d persistent fever. She was diagnosed with *Streptococcus parasanguinis* infection based on findings obtained using ENB-guided transbronchial lung biopsy (TBLB).

### CASE SUMMARY

A 10-year-old girl presented with constitutional symptoms of cough and fever of 7 days' duration. Chest CT scans detected peripheral lung lesions and no endobronchial lesions. TBLB performed under the guidance of an ENB Lungpro navigation system was safe, well-tolerated, and effective for biopsying peripheral lung lesions. Examination of biopsied samples indicated the patient had a pulmonary *Streptococcus parasanguinis* infection, which was treated with antibiotics instead of more invasive treatment interventions. The patient's

symptoms resolved after she received a 3-wk course of oral linezolid. Comparisons of pre-treatment and post-treatment CT scans revealed absorption of some lung lesions within 7 mo of hospital discharge.

# CONCLUSION

ENB-guided TBLB biopsying of peripheral lung lesions in this child is a safe, well-tolerated, and effective alternative to conventional interventions.

**Key Words:** Electromagnetic navigational bronchoscopy; Lungpro; Transbronchial lung biopsy; Child; Diagnosis; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Electromagnetic navigational bronchoscopy-guided transbronchial lung biopsy is used to diagnose and treat pulmonary disorders in adults worldwide, but very few were reported in children. Moreover, diagnosing patients with peripheral lung lesions is still challenging, due to difficulties related to existing biopsy procedures. Here we present a case study conducted in China that demonstrated that electromagnetic navigational bronchoscopy-guided transbronchial lung biopsy is safe and effective when used to diagnose pediatric patients afflicted with peripheral pulmonary lesion-inducing disorders. Ultimately, clinicians adopting this procedure will be able to diagnose pulmonary disease cases without subjecting patients to invasive and risky surgical interventions.

**Citation:** Meng FZ, Chen QH, Gao M, Zeng L, Lin JR, Zheng JY. Diagnosis based on electromagnetic navigational bronchoscopy-guided biopsied peripheral lung lesions in a 10-year-old girl: A case report. *World J Clin Cases* 2023; 11(15): 3651-3657

**URL:** <https://www.wjgnet.com/2307-8960/full/v11/i15/3651.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v11.i15.3651>

# INTRODUCTION

With the development of auxiliary navigation equipment, such as the Lungpro navigation system, electromagnetic navigational bronchoscopy (ENB) has been increasingly applied in pulmonary clinical practice. Importantly, ENB can be used to guide transbronchial lung biopsying (TBLB) of peripheral lung tissues that were previously only accessible by computed tomography (CT)-guided conventional bronchoscopy. To perform this function, ENB systems are used to virtually plan the lung bronchoscopic path in advance and to navigationally guide the bronchoscope in real time to ensure accurate localization of the disease focus during the biopsy. Although ENB has been shown to improve success rates of TBLB in adults with pulmonary diseases[1], this technology has been infrequently used to diagnose pediatric pulmonary diseases. In fact, a search of medical literature[2,3] revealed only one report of a Chinese pediatric patient with pulmonary lesions who was successfully diagnosed using ENB-guided TBLB, which permitted physicians to safely obtain biopsied tissues to distinguish between lung infection and lung tumor metastasis when diagnosing the patient[3]. Here, we report the clinical application of ENB for diagnosing a second Chinese pediatric patient with pulmonary lesions.

# CASE PRESENTATION

## Chief complaints

A 7-d history of fever and cough.

## History of present illness

A ten-year old girl presented to our hospital with a fever and cough of 7 days' duration. Her temperature peaked at 39.2 °C. She had been initially admitted to a local hospital, where she was diagnosed with community-acquired pneumonia and treated with intravenous antibiotics that did not resolve her fever and cough. Due to lingering symptoms, she sought further treatment at our hospital.

## History of past illness

Her medical history was unremarkable.



### Personal and family history

The patient had no relevant personal or family history.

### Physical examination

At admission, the patient's weight was 34.0 kg and her vital signs were as follows: Body temperature, 37.0 °C; blood pressure, 90/62 mmHg; heart rate, 130 beats/min; respiratory rate, 20 breaths/min; oxygen saturation, 98% in room air. At the time of admission, moderate and fine moist rales were heard in both lungs with no wheezing observed.

### Laboratory examinations

Infection index white blood cell count,  $17.52 \times 10^9/L$ ; platelet count,  $256 \times 10^9/L$ ; erythrocyte sedimentation rate, 30 mm/h; C-reactive protein, 60.65 mg/L (0.00-5.00 mg/L) (December 16, 2021). Immune function was normal. Liver function: Alanine aminotransferase, 29 U/L (0-40 U/L); aspartate aminotransferase, 33U/L (0-40 U/L). A positive culture result for *Streptococcus parasanguinis* was obtained, with no fungal growth detected.

### Imaging examinations

During her first stay at our hospital, the patient underwent electronic lung bronchoscopy, which detected endobronchial inflammation in the left upper lobe proper segment and left lower lobe basal segment (Figure 1A). Results of CT retesting obtained 7 d later (Figure 1B) revealed growing, dense masses within lung tissues, as evidenced by darker shadows on the scan coinciding with the position of the outer basal segment of the lower lobe of the left lung that indicated the presence of multiple cystic translucent lesions. Moreover, various lesions of different sizes, multilocular thin-walled air-filled cavities, and patchy and highly dense shadows were observed that lacked clear boundaries and had approximate dimensions of 72 mm × 60 mm × 42 mm. The child was subsequently discharged with oral amoxicillin clavulanate potassium and advised to undergo an enhanced pulmonary CT scan 2 wk after discharge. The CT results revealed that the previously noted multiple cystic lesions within the lower left lung lobe and the dense mass within the basal segment outside that lobe remained unchanged in appearance. In contrast, the inflammatory lesions within the middle and lower lobes of the right lung were larger than before (Figure 1C). Due to the fact that the pathological nature of the peripheral lung lesion could not be determined from these findings, the patient underwent a TBLB procedure to ensure that appropriate treatment would be administered to treat the disorder.

The entire procedure was conducted using the following steps. Prior to TBLB, inspiratory and expiratory CT images were uploaded to the Lungpro system (LungPoint VBN, version 3.4, Broncus Medical Inc, CA, United States), which was used to construct a virtual airway leading to the above-mentioned peripheral lesions. Next, picture archiving and communication system-based integration was conducted to build a virtual three-dimensional image of the patient's lungs. Then, a bronchial centerline was used as a frame of reference to help the bronchoscope operator plan a path to the airway wall puncture point or focus point.

After the patient was prepared for the procedure, she was administered local anesthesia and intravenous midazolam, a mild sedative. Endobronchial occlusion was then performed using a bronchoscope (BF-P260, Olympus Ltd.). The patient was placed in the supine position and was administered oxygen through a nasal catheter. Using a nasal point of entry, the optimal navigational path to pulmonary lesions was selected, and the reconstructed three-dimensional image was superimposed onto the image as viewed under the bronchoscope. During real-time navigation, the position of the bronchoscope and the distance from the pleura and the focus were displayed.

Finally, Lungpro-guided TBLB was performed using a bronchoscope with a working channel diameter of 2.8 mm (Figure 2, Figure 3A). Biopsy samples taken from the posterior basal segment of the left lower lobe were sent to the hospital pathology department for microscopic examination (Figure 3B). The patient experienced no complications after the procedure.


## FINAL DIAGNOSIS

*Streptococcus parasanguinis* infection of the lung.

## TREATMENT


During the patient's first stay at our hospital, her fever decreased after intravenous antibiotic therapy (intravenous amoxicillin clavulanate potassium, oral azithromycin) and anti-inflammatory therapy (intravenous methylprednisolone). Results of ENB-guided TBLB performance (which utilized the Lungpro navigation system) indicated lung infection with *Streptococcus parasanguinis*. Based on this finding, the patient was treated with a 3-wk course of oral linezolid.





DOI: 10.12998/wjcc.v11.i15.3651 Copyright ©The Author(s) 2023.

**Figure 1** Bronchoscopy and computed tomography images at different stages of treatment. A: Basal segment of the lower lobe of the left lung under conventional bronchoscopy during the first hospitalization; B: Computed tomography (CT) scan showed a lesion in the outer basal segment of the lower lobe of the left lung grew bigger during the first hospitalization; C: CT scan showed a lesion in the outer basal segment of the lower lobe of the left lung grew bigger after the first hospitalization.



DOI: 10.12998/wjcc.v11.i15.3651 Copyright ©The Author(s) 2023.


**Figure 2** Three-dimensional reconstruction of thoracic computed tomography with the bronchial tree, tunnel path, and peripheral lung lesions location. A: Virtual bronchoscopic navigation images on LungPro indicate a bronchial route; B: Cross-sectional view; C: Coronal reconstruction view; D: Sagittal view; E: Three-dimensional view.

## OUTCOME AND FOLLOW-UP

The patient's symptoms disappeared after a 3-wk oral linezolid treatment, after which the patient's health status was monitored through regular follow-up visits. A second chest CT scan obtained at seven months post-treatment revealed increased transparency of the basal segment of the left lower lobe, multiple lesions of unequal size, and multilocular thin-walled air-filled cavities. A comparison of the first and second chest CT scans suggested that some lesions were absorbed before the 7-mo follow-up visit (Figure 3C).

## DISCUSSION

Modern imaging techniques have led to increased detection of peripheral lung lesions, of which the vast majority often require tissue biopsies for diagnosis. Barber *et al*[4] retrospectively analyzed etiologies of 88 pediatric cases with pulmonary nodules. They found that most nodules were caused by infections, while etiologies of the others included autoimmune diseases, hematologic diseases, vasculitis, and Crohn's disease. Nevertheless, diagnosing patients with peripheral lung lesions is still challenging due to difficulties related to existing biopsy procedures. The child described in this case report presented with cough and fever symptoms and rales. Chest high-resolution computed tomography (HRCT)



DOI: 10.12998/wjcc.v11.i15.3651 Copyright ©The Author(s) 2023.

**Figure 3** Transbronchial lung biopsy, postoperative pathological examination and computed tomography image results after treatment. A: Transbronchial lung biopsy under bronchoscopy; B: Endotracheal biopsy showing bacterial colonization, 200× magnification; C: High-resolution computed tomography (HRCT) obtained after treatment. Chest HRCT scans show the transparency of the basal segment of the left lower lobe increased, multiple lesions of unequal size, multilocular thin-walled air bag cavity can be seen. Compared with the former chest CT, some lesions are absorbed.

imaging revealed a dense mass, as evidenced by a dark area on the HRCT scan after the child had received regular antibiotic therapy. However, the lesion was located within the anterior segment of the left lower lobe in an area close to the anterior artery and thus would be inaccessible by conventional electronic bronchoscopy and require CT-guided percutaneous lung puncture treatment. As a major disadvantage, this procedure can lead to severe lung tissue damage, increasing the risk of intraoperative and postoperative bleeding and pneumothorax and leaving local skin wounds. To avoid these issues, we performed ENB-guided TBLB to obtain an accurate diagnosis for our patient.

Several bronchoscopic approaches are currently adopted to biopsy tissues in order to obtain diagnostic tissue samples, with pros and cons associated with each approach. Importantly, Bronchoscopy-guided modalities are increasingly being used to biopsy peripheral lung lesions, due to greater safety and superior diagnostic results obtained using these methods over results obtained using alternative biopsy techniques, such as percutaneous lung puncture, radial endobronchial ultrasound (EBUS), CT-guide flexible bronchoscopy, and combinations of these techniques[5,6]. In fact, a review of papers related to this topic suggests that diagnostic yields of peripheral lung nodule biopsies performed using conventional bronchoscopy range from 43% to 65%, while diagnostic yields of procedures used to biopsy smaller peripheral lung lesions are lower, from 14% to 31%[7]. Although bronchoscopy with navigation provides the lowest overall diagnostic yield of the three procedures (70%), this procedure has the best safety profile, as evidenced by its low rate of pneumothorax requiring a chest tube (1%)[8].

CT is currently considered the gold standard imaging technique for guiding bronchoscopic biopsies of pulmonary lesions, due to its effectiveness as a diagnostic tool and ability to target a bronchoscope to a selected cytologic sampling site accurately. Nevertheless, CT-guided transthoracic biopsy has several disadvantages, such as exposure to ionizing radiation, the need to move the patient to a radiology unit, and prolonged execution time. Moreover, this modality requires an accessible path for biopsy needle insertion and is associated with increased risks of pneumothorax, significant hemorrhaging, and respiratory failure.

Alternatively, radial EBUS is a current method to precisely locate peripheral lung lesions that are not visible by fluoroscopy, while it also provides clues regarding their internal structures. However, the effective use of this method requires careful patient selection, since the ultrasound wave does not easily penetrate aerated pulmonary parenchyma. Therefore, EBUS is used to locate peripheral pulmonary nodules that are not separated by areas of aerated pulmonary parenchyma, which would induce excessive acoustic impedance that would reflect the ultrasound wave. In addition, ultrasound is technically applicable for imaging peripheral pulmonary nodules that adhere to visceral pleura, detecting direct pleural infiltration, and detecting atelectasis involving subpleural lung parenchyma. Nonetheless, EBUS cannot target specific types of peripheral lesions that may be targeted using a navigational system.

TBLB, an important method used to clinically diagnose pulmonary lesion-associated disorders, offers advantages over conventional bronchoscopic biopsy methods that provide low diagnostic yields due to inaccurate targeting of lesions with biopsy forceps[9]. However, during TBLB operation, misalignments (offsets) between actual and virtual bronchoscopic images can occur when the bronchoscope is rotated as it advances to the target site, which can lead to lesion-targeting errors. Furthermore, rapid realignment of the virtual bronchoscopic image to actual bronchial branch points is not easily accomplished after the bronchoscope moves unexpectedly (*e.g.*, when the patient moves or coughs)[10]. To address these issues[10,11], the Lungpro system was recently developed to serve as a virtual

navigation device that can display virtual bronchoscopic images adjacent to actual video images (Video 1), thus enabling the operator to overlay an electronic pathway onto the endoscopic image and maintain their alignment in real-time. In practice, the identified lesion shape is initially overlaid onto the airway wall near the target location (as indicated on the aligned actual and virtual bronchoscopic images) to provide further biopsy site localization guidance. Importantly, this method can be applied to quickly and accurately guide the bronchoscope to the focus without puncturing blood vessels, thus making it possible to more effectively biopsy lung lesions present outside the airway[10,12]. Notably, when this procedure was used to biopsy our patient, no adverse events, such as pneumothorax or hemorrhage, occurred during or after the procedure.

Ultimately, the biopsy results obtained for our pediatric patient suggested that the child's lungs were infected with *S. parasanguinis*, a gram-positive, non-motile, catalase-negative organism belonging to the sanguinis group of viridans streptococci. Although this organism is not viewed as an important human pathogen, it has been isolated from human blood, urine, and throat samples[13], is a primary colonizer of teeth that supports biofilm formation on tooth surfaces[14], and is a common cause of native and prosthetic valve endocarditis. *S. parasanguinis*-induced endocarditis has been shown to begin with bacterial colonization on heart valve surfaces that is followed by biofilm formation that leads to colonization in valves with small surface aggregates of bacteria. These aggregates grow into rounded bacterial colonies that promote the deposition of a layer of fibrin and platelets on their surfaces that ultimately supports biofilm formation. This biofilm then shields the bacteria from antibiotics, thus making their eradication difficult despite the fact that the organisms are sensitive to vancomycin, linezolid, daptomycin, quinupristin-dalfopristin, clindamycin, and levofloxacin[15]. Notably, an analysis of 50 cancer patients with streptococcal infections revealed that 5 patients harbored *S. parasanguinis* with intermediate resistance to penicillin and azithromycin and susceptibility to vancomycin, linezolid, quinupristin-dalfopristin, and ceftriaxone[16]. These results were consistent with *S. parasanguinis* antibiotic responsiveness as observed in the current case study. In our patient, inflammatory lesions increased in size after treatment with amoxicillin clavulanate potassium and subsequently shrank significantly after 3 wk of oral linezolid treatment. Our results thus highlight the bacteriological and therapeutic challenges associated with pneumonia caused by *Streptococcus parasanguinis* infection.

## CONCLUSION

Worldwide, ENB-guided TBLB is used to diagnose and treat pulmonary disorders in adults but has been infrequently performed in children as its effectiveness and safety in pediatric patients remains unknown. Meanwhile, ENB technology is expensive, and ENB systems are not available in all countries of regions. Here we present a case study conducted in China that demonstrated that ENB-guided transbronchial lung biopsy is safe and effective when used to diagnose pediatric patients afflicted with peripheral pulmonary lesion-inducing disorders. Ultimately, clinicians adopting this procedure may help diagnose pulmonary disease cases without subjecting patients to invasive and risky surgical interventions.

## FOOTNOTES

**Author contributions:** Meng FZ and Chen QH were the first authors of this manuscript; Meng FZ and Zheng JY performed the operation; Chen QH, Zeng L and Lin JR collected clinical data and data interpretation; Meng FZ, Gao M, and Zheng JY helped revise the manuscript; Chen QH drafted the manuscript; all authors have read and approved the final manuscript.

**Supported by** The Natural Science Foundation of Jilin Province, No. 20200201475JC.

**Informed consent statement:** All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

**Conflict-of-interest statement:** The authors declare they have no competing interests.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>



Country/Territory of origin: China

ORCID number: Fan-Zheng Meng 0000-0001-8267-353X; Qiong-Hua Chen 0000-0002-5684-3379; Man Gao 0000-0002-2524-5034; Jing-Yang Zheng 0000-0002-5202-0121.

S-Editor: Ma YJ

L-Editor: A

P-Editor: Yu HG

## REFERENCES

- 1 Tamiya M, Okamoto N, Sasada S, Shiroyama T, Morishita N, Suzuki H, Yoshida E, Hirashima T, Kawahara K, Kawase I. Diagnostic yield of combined bronchoscopy and endobronchial ultrasonography, under LungPoint guidance for small peripheral pulmonary lesions. *Respirology* 2013; **18**: 834-839 [PMID: 23586738 DOI: 10.1111/resp.12095]
- 2 Piccione J, Hysinger EB, Vicencio AG. Pediatric advanced diagnostic and interventional bronchoscopy. *Semin Pediatr Surg* 2021; **30**: 151065 [PMID: 34172210 DOI: 10.1016/j.sempedsurg.2021.151065]
- 3 Yang H, Turgon E, Pan Y, Wen X, Zhang X, Shen Y, Wang F. Pilot study of archimedes virtual bronchoscopic navigation system-guided biopsy to diagnose lung nodules in children. *Front Pediatr* 2022; **10**: 1053289 [PMID: 36819191 DOI: 10.3389/fped.2022.1053289]
- 4 Barber A, Passarelli P, Dworsky ZD, Gatcliffe C, Ryu J, Lesser DJ. Clinical implications of pulmonary nodules detected in children. *Pediatr Pulmonol* 2021; **56**: 203-210 [PMID: 33118698 DOI: 10.1002/ppul.25146]
- 5 Ishida T, Asano F, Yamazaki K, Shinagawa N, Oizumi S, Moriya H, Munakata M, Nishimura M; Virtual Navigation in Japan Trial Group. Virtual bronchoscopic navigation combined with endobronchial ultrasound to diagnose small peripheral pulmonary lesions: a randomised trial. *Thorax* 2011; **66**: 1072-1077 [PMID: 21749984 DOI: 10.1136/thx.2010.145490]
- 6 Eberhardt R, Anantham D, Ernst A, Feller-Kopman D, Herth F. Multimodality bronchoscopic diagnosis of peripheral lung lesions: a randomized controlled trial. *Am J Respir Crit Care Med* 2007; **176**: 36-41 [PMID: 17379850 DOI: 10.1164/rccm.200612-1866oc]
- 7 Kalanjeri S, Holladay RC, Gildea TR. State-of-the-Art Modalities for Peripheral Lung Nodule Biopsy. *Clin Chest Med* 2018; **39**: 125-138 [PMID: 29433709 DOI: 10.1016/j.ccm.2017.11.007]
- 8 Wang Memoli JS, Nietert PJ, Silvestri GA. Meta-analysis of guided bronchoscopy for the evaluation of the pulmonary nodule. *Chest* 2012; **142**: 385-393 [PMID: 21980059 DOI: 10.1378/chest.11-1764]
- 9 Han Y, Kim HJ, Kong KA, Kim SJ, Lee SH, Ryu YJ, Lee JH, Kim Y, Shim SS, Chang JH. Diagnosis of small pulmonary lesions by transbronchial lung biopsy with radial endobronchial ultrasound and virtual bronchoscopic navigation versus CT-guided transthoracic needle biopsy: A systematic review and meta-analysis. *PLoS One* 2018; **13**: e0191590 [PMID: 29357388 DOI: 10.1371/journal.pone.0191590]
- 10 Eberhardt R, Kahn N, Gompelmann D, Schumann M, Heussel CP, Herth FJ. LungPoint--a new approach to peripheral lesions. *J Thorac Oncol* 2010; **5**: 1559-1563 [PMID: 20802352 DOI: 10.1097/JTO.0b013e3181e8b308]
- 11 Merritt SA, Gibbs JD, Yu KC, Patel V, Rai L, Cornish DC, Bascom R, Higgins WE. Image-guided bronchoscopy for peripheral lung lesions: a phantom study. *Chest* 2008; **134**: 1017-1026 [PMID: 18583513 DOI: 10.1378/chest.08-0603]
- 12 Silvestri GA, Herth FJ, Keast T, Rai L, Gibbs J, Wibowo H, Sterman DH. Feasibility and safety of bronchoscopictransparenchymal nodule access in canines: a new real-time image-guided approach to lung lesions. *Chest* 2014; **145**: 833-838 [PMID: 24202737 DOI: 10.1378/chest.13-1971]
- 13 Fernández-Garayzábal JF, Fernández E, Las Heras A, Pascual C, Collins MD, Domínguez L. Streptococcus parasanguis: new pathogen associated with asymptomatic mastitis in sheep. *Emerg Infect Dis* 1998; **4**: 645-647 [PMID: 9866743 DOI: 10.3201/eid0404.980417]
- 14 Chen Q, Wu H, Fives-Taylor PM. Investigating the role of secA2 in secretion and glycosylation of a fimbrial adhesin in Streptococcus parasanguis FW213. *Mol Microbiol* 2004; **53**: 843-856 [PMID: 15255897 DOI: 10.1111/j.1365-2958.2004.04116.x]
- 15 Sader HS, Streit JM, Fritsche TR, Jones RN. Antimicrobial susceptibility of gram-positive bacteria isolated from European medical centres: results of the Daptomycin Surveillance Programme (2002-2004). *Clin Microbiol Infect* 2006; **12**: 844-852 [PMID: 16882289 DOI: 10.1111/j.1469-0691.2006.01550.x]
- 16 Sadjadi SA, Ali H. Streptococcus parasanguis peritonitis: report of a case and review of the literature. *Perit Dial Int* 2011; **31**: 603-604 [PMID: 21976479 DOI: 10.3747/pdi.2011.00011]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

