

# World Journal of *Clinical Cases*

*World J Clin Cases* 2023 June 6; 11(16): 3664-3931



## Contents

Thrice Monthly Volume 11 Number 16 June 6, 2023

## REVIEW

- 3664 Kikuchi-Fujimoto disease: A comprehensive review  
*Mahajan VK, Sharma V, Sharma N, Rani R*
- 3680 Current diagnostic tools and treatment modalities for rectal prolapse  
*Oruc M, Erol T*

## MINIREVIEWS

- 3694 Application of laparoscopic surgery in gallbladder carcinoma  
*Wu X, Li BL, Zheng CJ*
- 3706 Current research of idiopathic normal pressure hydrocephalus: Pathogenesis, diagnosis and treatment  
*Ishida T, Murayama T, Kobayashi S*
- 3714 *Helicobacter pylori* plays a key role in gastric adenocarcinoma induced by spasmolytic polypeptide-expressing metaplasia  
*Li ML, Hong XX, Zhang WJ, Liang YZ, Cai TT, Xu YF, Pan HF, Kang JY, Guo SJ, Li HW*
- 3725 Review of deep learning and artificial intelligence models in fetal brain magnetic resonance imaging  
*Vahedifard F, Adepoju JO, Supanich M, Ai HA, Liu X, Kocak M, Marathu KK, Byrd SE*
- 3736 Diabetes more than retinopathy, it's effect on the anterior segment of eye  
*Morya AK, Ramesh PV, Kaur K, Gurnani B, Heda A, Bhatia K, Sinha A*

## ORIGINAL ARTICLE

## Retrospective Cohort Study

- 3750 Long term outcomes of Cohen's cross trigonal reimplantation for primary vesicoureteral reflux in poorly functioning kidney  
*Ansari MS, Banthia R, Jain S, Kaushik VN, Danish N, Yadav P*

## Retrospective Study

- 3756 Dexmedetomidine-induced anesthesia in elderly patients undergoing hip replacement surgery  
*Li JQ, Yuan H, Wang XQ, Yang M*

## Observational Study

- 3765 Hypoperfusion context as a predictor of 28-d all-cause mortality in septic shock patients: A comparative observational study  
*Kataria S, Singh O, Juneja D, Goel A, Bhide M, Yadav D*

- 3780** Psychological review of hemodialysis patients and kidney transplant recipients during the COVID-19 pandemic

*Gundogmus AG, Oguz EG, Guler-Cimen S, Kocyigit Y, Dogan AE, Ayli MD*

- 3791** Incidence and peri-operative risk factors for development of acute kidney injury in patients after cardiac surgery: A prospective observational study

*Dimopoulos S, Zagkotsis G, Kinti C, Rouvali N, Georgopoulou M, Mavraki M, Tasouli A, Lyberopoulou E, Roussakis A, Vasileiadis I, Nanas S, Karabinis A*

### Randomized Controlled Trial

- 3802** Coaxial radiography guided puncture technique for percutaneous transforaminal endoscopic lumbar discectomy: A randomized control trial

*Chen LP, Wen BS, Xu H, Lu Z, Yan LJ, Deng H, Fu HB, Yuan HJ, Hu PP*

### CASE REPORT

- 3813** Blood typing and transfusion therapy in a patient with A2 subtype acute myeloid leukemia M2: A case report

*Kuang XC, Zhang SH, Cen YJ, Zhang JB, Liu YS*

- 3822** Valve repair after infective endocarditis secondary to perforation caused by *Streptococcus gordonii*: A case report

*Qu YF, Yang J, Wang JY, Wei B, Ye XH, Li YX, Han SL*

- 3830** *Prevotella oris*-caused meningitis and spinal canal infection: A case report

*Zhang WW, Ai C, Mao CT, Liu DK, Guo Y*

- 3837** Severe liver trauma with complex portal and common bile duct avulsion: A case report and review of the literature

*Mitricof B, Kraft A, Anton F, Barcu A, Barzan D, Haiducu C, Brasoveanu V, Popescu I, Moldovan CA, Botea F*

- 3847** TACC diagnosed by transoesophageal endoscopic ultrasonography: A case report

*Pu XX, Xu QW, Liu BY*

- 3852** Ruptured teratoma mimicking a pelvic inflammatory disease and ovarian malignancy: A case report

*Lai PH, Ding DC*

- 3858** Purpura annularis telangiectodes of Majocchi: A case report

*Pu YJ, Jiang HJ, Zhang L*

- 3864** Giant cyst in heterotopic pregnancy: A case report

*Kong YY, Chanda K, Ying XY*

- 3870** High doses of dextromethorphan induced shock and convulsions in a 19-year-old female: A case report

*Shimozawa S, Usuda D, Sasaki T, Tsuge S, Sakurai R, Kawai K, Matsubara S, Tanaka R, Suzuki M, Hotchi Y, Tokunaga S, Osugi I, Katou R, Ito S, Asako S, Mishima K, Kondo A, Mizuno K, Takami H, Komatsu T, Oba J, Nomura T, Sugita M*

- 3877** Postpartum ovarian vein thrombosis after cesarean section and vaginal delivery: Two case reports

*Zhu HD, Shen W, Wu HL, Sang X, Chen Y, Geng LS, Zhou T*

- 3885** Traumatic pancreatic ductal injury treated by endoscopic stenting in a 9-year-old boy: A case report  
*Kwon HJ, Jung MK, Park J*
- 3891** Novel mutation c.2090\_2091del in neurodevelopmental-craniofacial syndrome with variable renal and cardiac abnormalities in an 18.5-mo-old boy: A case report  
*Li Y, Zhou Z, Xu Y, Wang ZR*
- 3899** Reading impairment after neonatal hypoglycemia with parieto-temporo-occipital injury without cortical blindness: A case report  
*Kurahashi N, Ogaya S, Maki Y, Nonobe N, Kumai S, Hosokawa Y, Ogawa C, Yamada K, Maruyama K, Miura K, Nakamura M*
- 3907** Unusual clinical presentation of oral pyogenic granuloma with severe alveolar bone loss: A case report and review of literature  
*Lomeli Martínez SM, Bocanegra Morando D, Mercado González AE, Gómez Sandoval JR*
- 3915** Intraoperative photodynamic therapy for tracheal mass in non-small cell lung cancer: A case report  
*Jung HS, Kim HJ, Kim KW*
- 3921** Coexistence of urinary tuberculosis and urothelial carcinoma: A case report  
*Tsai YC, Li CC, Chen BT, Wang CY*

**LETTER TO THE EDITOR**

- 3929** Symmetric DWI hyperintensities in CMT1X patients after SARS-CoV-2 vaccination should not be classified as stroke-like lesions  
*Finsterer J*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Ashraf F Hefny, MD, MSc, Associate Professor, Surgeon, Department of Surgery, College of Medicine and Health Sciences, UAE University, Al Ain 00000, United Arab Emirates. ahefny@uaeu.ac.ae

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Si Zhao; Production Department Director: Xu Guo; Editorial Office Director: Jin-Lei Wang.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

June 6, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Retrospective Cohort Study

## Long term outcomes of Cohen's cross trigonal reimplantation for primary vesicoureteral reflux in poorly functioning kidney

Mohd Sualeh Ansari, Ravi Banthia, Shrey Jain, Vinay N Kaushik, Nayab Danish, Priyank Yadav

**Specialty type:** Urology and nephrology**Provenance and peer review:** Invited article; Externally peer reviewed.**Peer-review model:** Single blind**Peer-review report's scientific quality classification**Grade A (Excellent): 0  
Grade B (Very good): 0  
Grade C (Good): C, C, C  
Grade D (Fair): 0  
Grade E (Poor): 0**P-Reviewer:** He G, China; Sarier M, Turkey**Received:** February 15, 2023**Peer-review started:** February 15, 2023**First decision:** April 11, 2023**Revised:** April 19, 2023**Accepted:** May 6, 2023**Article in press:** May 6, 2023**Published online:** June 6, 2023**Mohd Sualeh Ansari, Ravi Banthia, Shrey Jain, Vinay N Kaushik, Nayab Danish, Priyank Yadav,** Department of Urology and Renal Transplantation, Sanjay Gandhi Postgraduate Institute of Medical Sciences, Lucknow 226014, Uttar Pradesh, India**Corresponding author:** Priyank Yadav, MCh, Assistant Professor, Department of Urology and Renal Transplantation, Sanjay Gandhi Postgraduate Institute of Medical Sciences, Raebareli Road, Lucknow 226014, Uttar Pradesh, India. [priyankmama@gmail.com](mailto:priyankmama@gmail.com)

## Abstract

## BACKGROUND

Open ureteric reimplantation by cross trigonal technique described by Cohen is considered a common surgical option for correction of vesicoureteral reflux (VUR). There is a lack of evidence in literature though for what happens to such kidneys, in the long run, particularly those which are poorly functioning.

## AIM

To assess the long-term outcomes of ureteric reimplantation in poorly functioning kidneys in children with unilateral primary VUR.

## METHODS

Children with unilateral primary VUR and a relative renal function of less than 35% who underwent open or laparoscopic ureteric reimplantation between January 2005 and January 2017 were included in the study. Patients who had a follow up of less than five years were excluded. Preoperative evaluation consisted of a voiding cystourethrogram and Dimercaptosuccinic acid (DMSA) scan. In the follow-up period, patients underwent a diuretic scan at 6 weeks and 6 months. Follow up ultrasound was done for change in grade of hydronephrosis and retrovesical ureteric diameter. Subsequent follow up was done at 6 monthly intervals with evaluation for proteinuria and hypertension and any recurrent urinary tract infection (UTI). For assessment of cortical function, DMSA was repeated annually for 5 years after surgery. A paired-samples *t*-test was used to test the mean difference of DMSA between pre-post observations.

## RESULTS

During this period, 36 children underwent ureteric reimplantation for unilateral primary VUR. After excluding those with insufficient follow-up, 31 were included in the analysis. Most of the patients were males ( $n = 26/31$ , 83.8%). Patient's age (mean  $\pm$  SD, range) was  $5.21 \pm 3.71$ , 1-18 years. The grades of VUR were grade II (1

patient), grade III (8 patients), grade IV (10 patients), and grade V (12 patients). The pre and postoperative DMSA was  $24.064 \pm 12.02$  and  $24.06 \pm 10.93$ , which was almost the same (statistically equal, paired-samples *t*-test:  $P = 0.873$ ). The median (range) follow-up duration was 82 (60-120) mo. One patient had persistent reflux after surgery (preoperative: grade IV, postoperative: grade III), and the very same patient developed recurrent UTI. The difference in the preoperative and postoperative DRF was less than 10% in 29 patients. In one patient, the DRF decreased by 17% (22% to 05%) while in another patient, the DRF increased by 12% (25% to 37%) after surgery. None of the patients had an increase in scarring after surgery. 15% of patients were hypertensive before surgery and all of them continued to be hypertensive after surgery while none developed hypertension after surgery. None of the patients had significant proteinuria ( $> 150$  mg/d) during the follow-up period.

### CONCLUSION

Children with unilateral primary VUR and poorly functioning kidney maintain the renal function over the long term in most cases. Hypertension and proteinuria do not progress over time in these patients.

**Key Words:** Vesicoureteral reflux; Ureteric reimplantation; Relative renal function; Poorly functioning kidney; Unilateral; Long term

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** This article presents the outcomes of reimplantation of ureters in patients with vesicoureteral reflux in patients without any secondary causes of reflux. In this study, we found that children with unilateral primary vesicoureteral reflux and poorly functioning kidney maintain the renal function over the long term in most cases. All the patients who were hypertensive before surgery continued to be hypertensive after surgery while no patient developed new hypertension after surgery.

**Citation:** Ansari MS, Banthia R, Jain S, Kaushik VN, Danish N, Yadav P. Long term outcomes of Cohen's cross trigonal reimplantation for primary vesicoureteral reflux in poorly functioning kidney. *World J Clin Cases* 2023; 11(16): 3750-3755

**URL:** <https://www.wjgnet.com/2307-8960/full/v11/i16/3750.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v11.i16.3750>

## INTRODUCTION

Vesicoureteral reflux (VUR) is defined as the retrograde flow of urine from the bladder to the upper urinary tracts[1]. Its clinical challenges arise from the fact that it is usually asymptomatic and even in asymptomatic cases it is responsible for pyelonephritic scarring and can be associated with congenital renal dysmorphism. In most cases, VUR resolve spontaneously; 20% to 30% will have further infections, and few of them will experience long-term sequelae like renal scarring[2]. Extensive renal scarring decreases renal function and can result in renal insufficiency, hypertension, or end-stage renal disease [3]. A dilemma exists whether to opt for nephrectomy or ureteric reimplantation for the poorly functional kidney. Open ureteric reimplantation by cross trigonal technique described by Cohen is amongst the most common options for surgical correction of VUR, but there is a dearth of evidence in the literature for what happens to such kidneys, in the long run, particularly those which are poorly functioning. We at our institute conducted a retrospective study to assess the long-term outcomes of unilateral ureteric reimplantation in poorly functioning kidneys in children with unilateral primary VUR. The primary objective of our study was to determine if a kidney with significant unilateral reflux nephropathy maintains relative renal function after ureteric reimplantation on long term follow up (5 years or more).

## MATERIALS AND METHODS

We retrospectively reviewed data of 168 patients who underwent ureteric reimplantation from 2003-2019. Children with unilateral primary VUR and a relative renal function of less than 35% who underwent open or laparoscopic ureteric reimplantation between January 2005 and December 2017 were included in the study. Patients who had a follow up of less than five years, patients older than 18



years, having a neurogenic bladder, posterior urethral valves, bladder exstrophy and bilateral VUR were excluded. Data was collected by two independent researchers to minimize bias. Preoperative evaluation consisted of a voiding cystourethrogram and Dimercaptosuccinic acid (DMSA) scan, EC scan, and S. Creatinine. In the follow-up period, patients underwent an EC scan at 6 wk and 6 mo. Ultrasounds for hydronephrosis and retrovesical ureteral diameter were done at 6 mo interval. Subsequent follow up was done at 6-monthly intervals with evaluation for proteinuria and hypertension. For assessment of cortical function, DMSA was repeated 1 year after surgery and annually thereafter. A paired-samples *t*-test was used to test the mean difference between pre-post observations. Only the patients with complete data records and regular follow up were included in the study.

## RESULTS

During this period, 36 children underwent ureteric reimplantation for unilateral primary VUR. After excluding those with insufficient follow-up (*i.e.* follow up less than 5 years), 31 were included in the analysis. Most of the patients were males ( $n = 26/31$ , 83.8%). Patient's age (mean  $\pm$  SD, range) was  $5.21 \pm 3.71$ , 1-18 years. The grades of VUR were grade- II (1 patient), grade- III (8 patients), grade- IV (10 patients), and grade- V (12 patients)[4]. The renal function estimated by pre and postoperative DMSA scan was  $24.06 \pm 12.02$  and  $24.06 \pm 10.93$ , which was almost the same (statistically equal, paired-samples *t*-test:  $P = 0.873$ ). The median (range) follow-up duration was 82 (60-120) months. One patient had persistent reflux after surgery (preoperative: Grade IV, postoperative: grade III). One patient had recurrent UTI. The difference in the preoperative and postoperative DRF was less than 10% in 26 patients. In one patient, the DRF decreased by 17% (22% to 05%) while in another patient, the DRF increased by 12% (25% to 37%) after surgery. There was no significant change in Serum creatinine in almost all patients, as mean serum creatinine preoperatively and postoperatively was  $0.866 \pm 0.68$  and  $0.811 \pm 0.54$  respectively (statistically equal, paired - sample *t*-test:  $P = 0.583$ ). None of the patients had any increase in scarring after surgery or any development of new scars. A total of 15% of patients were hypertensive before surgery and all of them continued to be hypertensive after surgery, while no new patients developed hypertension. None of the patients showed significant proteinuria ( $> 150$  mg/d) during the follow-up period. No patients presented an increase in hydronephrosis or retrovesical ureteric diameter after surgery.

## DISCUSSION

Primary VUR is still a significant health problem for many children and young adolescents, as these patients are at high risk of developing reflux nephropathy[5]. The association of nephropathy with UTI and VUR is well established[6]. Reflux nephropathy is irreversible and may lead to renal insufficiency, renin-mediated hypertension, chronic renal failure, decreased somatic growth and morbidity during pregnancy. To avoid renal scarring these patients must be tested appropriately and promptly treated with antibiotics. In the present study, we found that the majority of children who have primary VUR and a poorly functioning kidney, maintain their renal function for a long time. In addition, hypertension and proteinuria in these patients usually do not progress with time.

Earlier studies used IVP for follow up after ureteric reimplantation to look for new scar formation and most of them reported a substantial rate of new scar formation (5% to 31%) and scar progression (12% to 20%)[7-10]. But soon the accuracy of IVP to assess reflux related renal scars was questioned and renal scintigraphy became a well-established method to quantitatively measure renal damage in kidneys with reflux and accuracy of this modality is well established now. A study performed by Choi *et al*[11] was the first of its kind which used dimercaptosuccinic acid renal scintigraphy instead of IVP. This study reported 0% new scar formation and 2.6% progression of renal scarring. Our study results are similar to study by Choi *et al*[11] and we also report 0% new scar formation and 0% progression of renal scarring.

The observation of our study is not in accordance with studies conducted by Atwell *et al*[12], Scott *et al*[13], and Carson *et al*[14] who reported accelerated renal growth after surgical correction of VUR. Conversely, large series reported by Rushton *et al*[15] and Olbing *et al*[16] concluded that there was no significant difference in renal size after surgical correction in children with primary VUR which is like the observations of our study.

Renal functional impairment is one of the most devastating complications associated with VUR and is rare fortunately. The outcome of relative and overall renal function after ureteric reimplantation in children with unilateral primary VUR with significantly diminished relative renal function was assessed. A study by Schiepers *et al*[17]. Found that DMSA uptake did not change significantly during the 5 years interval in 89% of children after surgical correction of VUR, although this study had only 13 children with diminished RRF. A previous study by Nepple *et al*[18] in 2005, which included 32 children showed that relative renal function is maintained in such patients, although the mean follow-up period in their study was 3.7 years (0.3 to 12.9 years). In our series we observed 31 children who had strictly



minimum 5 year follow up with mean follow up of 82 mo (60-120 mo) and most of our patients maintained their RRF in the long term. In this study we identified only one patient who had significantly decreased renal function and one patient also had significantly improved renal function.

Serum creatinine was measured preoperatively and was measured on follow up along with DMSA scan. Preoperative and post-operative serum creatinine show insignificant difference.

Ureteral reflux resolution rate was 96.4% which is similar to the success rate of 92%-99% reported by others. In a study by Nepple *et al*[18] reflux persisted in 4 of 32 children, which was treated with the Glenn-Anderson ureteral advancement procedure. Reflux persisted in 2 of 51 in the study by Grossklaus *et al*[19]. In the present study reflux persisted in 1 of 31 children (preoperative: grade IV, postoperative: grade III) which was treated by endoscopic method, and we continue to use this modality in patients with recurrence with a lower grade of reflux and surgical treatment is reserved for those with a higher or similar grade of reflux with UTI.

Hypertension is a known complication of reflux for many years and was present in 15% of the patients, while no patient developed newly detected hypertension after surgery. Wallace *et al*[20] conducted a study 12 years after correcting reflux, found hypertension in 12.8% of the patients (18.5% among children with bilateral scarring and 11.3% among those with unilateral scars). On the contrary, Belloli *et al*[21] detected hypertension in only 4.4% of patients who had corrective surgery. In our study we followed up patients with a mean period of 82 mo (60-120 mo) and none of the patients developed new-onset hypertension or proteinuria.

A similar study was conducted by Mor *et al*[22] showed that even patients who were treated successfully by ureteric reimplantation during childhood are prone to recurrent UTI, progressive renal scarring, hypertension, and complications during pregnancy. In that study only 31% (100) of the potential original study group of 322 patients could be fully evaluated. It was assumed that patients who had an uneventful did not enrolled themselves for the study and therefore showed the above results. This was the drawback of the study which reflected negatively biased results.

There are limitations to this retrospective study. The finding of no change in relative renal function may be the result of low statistical power from a small sample size. Although the follow-up period was good enough to determine that these patients maintain overall relative renal function. Since most of the patients in our study were males, we did not compare the outcomes between males and females. Future studies can take this limitation into account. Further, we did not explore the effect of bilateral VUR on renal function and thus, similar renal dynamics may not be observed with bilateral reflux nephropathy.

## CONCLUSION

Even poorly functional kidneys maintain their function in the long term and there is need to establish a protocol for the long-term follow-up of patients who have had ureteric reimplantation during childhood. Management with ureteric reimplantation rather than nephrectomy may be warranted even in children with significantly reduced RRF based on the continued stability of relative renal function of the affected kidney. Hypertension and proteinuria do not progress over time in these patients.

## ARTICLE HIGHLIGHTS

### Research background

Vesicoureteral reflux (VUR) is when urine flows backwards from the bladder to the upper urinary tracts. It often has no symptoms but can cause kidney damage and scarring, leading to renal insufficiency, hypertension, or end-stage renal disease. Surgical options like ureteric reimplantation may be considered for poorly functioning kidneys, but there is limited evidence on long-term outcomes.

### Research motivation

The motivation behind the research was to address the lack of evidence in the literature regarding the long-term outcomes of open ureteric reimplantation for poorly functioning kidneys with VUR. This research was conducted to provide better insights and guidance on the optimal management of VUR with poorly functioning kidneys.

### Research objectives

The aim of the study was to determine if poorly functioning kidneys with VUR maintain relative renal function after the surgery on long-term follow-up of 5 years or more.

### Research methods

The study involved a retrospective review of medical data for 168 patients who underwent ureteric reimplantation between 2003-2019. The research focused on children with unilateral primary VUR and a

relative renal function of less than 35% who underwent open or laparoscopic ureteric reimplantation between January 2005 and December 2017. Data was collected by two independent researchers, and patients were excluded based on specific criteria. Preoperative evaluation included a voiding cystourethrogram and Dimercaptosuccinic acid (DMSA) scan, EC scan, and S. Creatinine. Follow-up was done at 6-month intervals, and data was collected through various tests to assess the patient's cortical function. A paired-samples t-test was used to compare pre- and post-surgery observations. Only patients with complete data records and regular follow-up were included in the study.

### Research results

The study included 31 children who underwent ureteric reimplantation for unilateral primary VUR. The patients were predominantly male, and their mean age was  $5.21 \pm 3.71$  years. The pre- and postoperative renal function, as measured by DMSA scan, remained statistically equal in most patients, and there was no significant change in serum creatinine. Only one patient had persistent reflux after surgery, and one had a recurrent UTI. None of the patients showed an increase in scarring, proteinuria, or hydro-nephrosis after surgery, and there were no new cases of hypertension.

### Research conclusions

The study found that even poorly functioning kidneys maintain their function in the long term after undergoing ureteric reimplantation, suggesting that this procedure may be a better option than nephrectomy. The results also indicate the need to establish a protocol for long-term follow-up of patients who have undergone this procedure, as hypertension and proteinuria do not progress over time in these patients.

### Research perspectives

The study findings suggest the need for further research on the long-term outcomes of ureteric reimplantation in poorly functioning kidneys in children with VUR. Future studies could investigate the factors that influence the success of ureteric reimplantation, such as age of the patient, severity of VUR, and degree of renal scarring. Additionally, studies could explore alternative treatments for VUR and their long-term outcomes, such as endoscopic injection of bulking agents or laparoscopic ureteric reimplantation.

---

## FOOTNOTES

**Author contributions:** Ansari MS and Yadav P designed the research study; Banthia R, Jain S, and Kaushik VN performed the research; Danish N and Banthia R analyzed the data and wrote the manuscript; Ansari MS and Yadav P edited the manuscript; All authors have read and approve the final manuscript.

**Institutional review board statement:** The study was reviewed and approved by the Institutional Ethics Committee of Sanjay Gandhi Postgraduate Institute of Medical Sciences (Approval No. 2020-185-IP-EXP-24).

**Informed consent statement:** This is to inform that consent waiver has been obtained from the Institutional Ethics Committee for the research article.

**Conflict-of-interest statement:** All the authors declare that there are no conflicts of interest regarding the publication of this study.

**Data sharing statement:** No additional data are available.

**STROBE statement:** The authors have read the STROBE Statement – checklist of items, and the manuscript was prepared and revised according to the STROBE Statement – checklist of items.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** India

**ORCID number:** Priyank Yadav 0000-0002-2509-0116.

**Corresponding Author's Membership in Professional Societies:** American Urological Association, No. 883547.

**S-Editor:** Liu JH

**L-Editor:** A

P-Editor: Liu JH

## REFERENCES

- 1 Nino F, Ilari M, Noviello C, Santoro L, Räscher IM, Martino A, Cobellis G. Genetics of Vesicoureteral Reflux. *Curr Genomics* 2016; **17**: 70-79 [PMID: 27013925 DOI: 10.2174/1389202916666151014223507]
- 2 Williams G, Fletcher JT, Alexander SI, Craig JC. Vesicoureteral reflux. *J Am Soc Nephrol* 2008; **19**: 847-862 [PMID: 18322164 DOI: 10.1681/ASN.2007020245]
- 3 Elder JS, Peters CA, Arant BS Jr, Ewalt DH, Hawtrey CE, Hurwitz RS, Parrott TS, Snyder HM 3rd, Weiss RA, Woolf SH, Hasselblad V. Pediatric Vesicoureteral Reflux Guidelines Panel summary report on the management of primary vesicoureteral reflux in children. *J Urol* 1997; **157**: 1846-1851 [PMID: 9112544 DOI: 10.1016/S0022-5347(01)64882-1]
- 4 Lebowitz RL, Olbing H, Parkkulainen KV, Smellie JM, Tamminen-Möbius TE. International system of radiographic grading of vesicoureteric reflux. International Reflux Study in Children. *Pediatr Radiol* 1985; **15**: 105-109 [PMID: 3975102 DOI: 10.1007/BF02388714]
- 5 Risdon RA, Yeung CK, Ransley PG. Reflux nephropathy in children submitted to unilateral nephrectomy: a clinicopathological study. *Clin Nephrol* 1993; **40**: 308-314 [PMID: 8299237]
- 6 Bailey RR. The relationship of vesico-ureteric reflux to urinary tract infection and chronic pyelonephritis-reflux nephropathy. *Clin Nephrol* 1973; **1**: 132-141 [PMID: 4783715]
- 7 Elo J, Tallgren LG, Alfthan O, Sarna S. Character of urinary tract infections and pyelonephritic renal scarring after antireflux surgery. *J Urol* 1983; **129**: 343-346 [PMID: 6834504 DOI: 10.1016/s0022-5347(17)52089-3]
- 8 Prospective trial of operative versus non-operative treatment of severe vesicoureteric reflux in children: five years' observation. Birmingham Reflux Study Group. *Br Med J (Clin Res Ed)* 1987; **295**: 237-241 [PMID: 2888509 DOI: 10.1136/bmj.295.6592.237]
- 9 Olbing H, Claesson I, Ebel KD, Seppänen U, Smellie JM, Tamminen-Möbius T, Wikstad I. Renal scars and parenchymal thinning in children with vesicoureteral reflux: a 5-year report of the International Reflux Study in Children (European branch). *J Urol* 1992; **148**: 1653-1656 [PMID: 1433582 DOI: 10.1016/s0022-5347(17)36995-1]
- 10 Weiss R, Duckett J, Spitzer A. Results of a randomized clinical trial of medical versus surgical management of infants and children with grades III and IV primary vesicoureteral reflux (United States). The International Reflux Study in Children. *J Urol* 1992; **148**: 1667-1673 [PMID: 1433585 DOI: 10.1016/s0022-5347(17)36998-7]
- 11 Choi H, Oh SJ, So Y, Lee DS, Lee A, Kim KM. No further development of renal scarring after antireflux surgery in children with primary vesicoureteral reflux: review of the results of 99mtechnetium dimercapto-succinic acid renal scan. *J Urol* 1999; **162**: 1189-1192 [PMID: 10458463 DOI: 10.1016/S0022-5347(01)68126-6]
- 12 Atwell JD, Vijay MR. Renal growth following reimplantation of the ureters for reflux. *Br J Urol* 1978; **50**: 367-370 [PMID: 709057 DOI: 10.1111/j.1464-410x.1978.tb04211.x]
- 13 Scott JE, Stansfeld JM. Ureteric reflux and kidney scarring in children. *Arch Dis Child* 1968; **43**: 468-470 [PMID: 5661876 DOI: 10.1136/adc.43.230.468]
- 14 Carson CC 3rd, Kelalis PP, Hoffman AD. Renal growth in small kidneys after ureteroneocystostomy. *J Urol* 1982; **127**: 1146-1148 [PMID: 7087024 DOI: 10.1016/s0022-5347(17)54272-x]
- 15 Rushton HG, Majd M, Jantusch B, Wiedermann BL, Belman AB. Renal scarring following reflux and nonreflux pyelonephritis in children: evaluation with 99mtechnetium-dimercaptosuccinic acid scintigraphy. *J Urol* 1992; **147**: 1327-1332 [PMID: 1314912 DOI: 10.1016/s0022-5347(17)37555-9]
- 16 Olbing H, Hirche H, Koskimies O, Lax H, Seppänen U, Smellie JM, Tamminen-Möbius T, Wikstad I. Renal growth in children with severe vesicoureteral reflux: 10-year prospective study of medical and surgical treatment: the International Reflux Study in Children (European branch). *Radiology* 2000; **216**: 731-737 [PMID: 10966703 DOI: 10.1148/radiology.216.3.r00au35731]
- 17 Schiepers C, Mesotten L, Proesmans W, Vereecken R, Verbruggen A, de Roo M. Surgical correction of vesicoureteral reflux: 5-year follow-up with 99Tcm-DMSA scintigraphy. *Nucl Med Commun* 2001; **22**: 217-224 [PMID: 11258409 DOI: 10.1097/00006231-200102000-00014]
- 18 Nepple KG, Austin JC, Hawtrey CE, Cooper CS. Kidneys with reflux nephropathy maintain relative renal function after ureteral reimplantation. *J Urol* 2005; **174**: 1606-1608 [PMID: 16148663 DOI: 10.1097/01.ju.0000183882.09237.29]
- 19 Grossklaus DJ, Pope JC, Adams MC, Brock JW. Is postoperative cystography necessary after ureteral reimplantation? *Urology* 2001; **58**: 1041-1045 [PMID: 11744485 DOI: 10.1016/s0090-4295(01)01467-4]
- 20 Wallace DM, Rothwell DL, Williams DI. The long-term follow-up of surgically treated vesicoureteric reflux. *Br J Urol* 1978; **50**: 479-484 [PMID: 753497 DOI: 10.1111/j.1464-410x.1978.tb06195.x]
- 21 Belloli G, Bedogni L, Salano F, Biscuola G, Meschi V. [Long-term evaluation of renal damage in primary vesico-renal reflux after corrective surgery]. *Pediatr Med Chir* 1985; **7**: 643-652 [PMID: 3837230]
- 22 Mor Y, Leibovitch I, Zalts R, Lotan D, Jonas P, Ramon J. Analysis of the long-term outcome of surgically corrected vesico-ureteric reflux. *BJU Int* 2003; **92**: 97-100 [PMID: 12823390 DOI: 10.1046/j.1464-410x.2003.04264.x]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

