

World Journal of *Clinical Cases*

World J Clin Cases 2023 June 6; 11(16): 3664-3931



Contents

Thrice Monthly Volume 11 Number 16 June 6, 2023

REVIEW

- 3664 Kikuchi-Fujimoto disease: A comprehensive review
Mahajan VK, Sharma V, Sharma N, Rani R
- 3680 Current diagnostic tools and treatment modalities for rectal prolapse
Oruc M, Erol T

MINIREVIEWS

- 3694 Application of laparoscopic surgery in gallbladder carcinoma
Wu X, Li BL, Zheng CJ
- 3706 Current research of idiopathic normal pressure hydrocephalus: Pathogenesis, diagnosis and treatment
Ishida T, Murayama T, Kobayashi S
- 3714 *Helicobacter pylori* plays a key role in gastric adenocarcinoma induced by spasmolytic polypeptide-expressing metaplasia
Li ML, Hong XX, Zhang WJ, Liang YZ, Cai TT, Xu YF, Pan HF, Kang JY, Guo SJ, Li HW
- 3725 Review of deep learning and artificial intelligence models in fetal brain magnetic resonance imaging
Vahedifard F, Adepoju JO, Supanich M, Ai HA, Liu X, Kocak M, Marathu KK, Byrd SE
- 3736 Diabetes more than retinopathy, it's effect on the anterior segment of eye
Morya AK, Ramesh PV, Kaur K, Gurnani B, Heda A, Bhatia K, Sinha A

ORIGINAL ARTICLE

Retrospective Cohort Study

- 3750 Long term outcomes of Cohen's cross trigonal reimplantation for primary vesicoureteral reflux in poorly functioning kidney
Ansari MS, Banthia R, Jain S, Kaushik VN, Danish N, Yadav P

Retrospective Study

- 3756 Dexmedetomidine-induced anesthesia in elderly patients undergoing hip replacement surgery
Li JQ, Yuan H, Wang XQ, Yang M

Observational Study

- 3765 Hypoperfusion context as a predictor of 28-d all-cause mortality in septic shock patients: A comparative observational study
Kataria S, Singh O, Juneja D, Goel A, Bhide M, Yadav D

- 3780** Psychological review of hemodialysis patients and kidney transplant recipients during the COVID-19 pandemic

Gundogmus AG, Oguz EG, Guler-Cimen S, Kocyigit Y, Dogan AE, Ayli MD

- 3791** Incidence and peri-operative risk factors for development of acute kidney injury in patients after cardiac surgery: A prospective observational study

Dimopoulos S, Zagkotsis G, Kinti C, Rouvali N, Georgopoulou M, Mavraki M, Tasouli A, Lyberopoulou E, Roussakis A, Vasileiadis I, Nanas S, Karabinis A

Randomized Controlled Trial

- 3802** Coaxial radiography guided puncture technique for percutaneous transforaminal endoscopic lumbar discectomy: A randomized control trial

Chen LP, Wen BS, Xu H, Lu Z, Yan LJ, Deng H, Fu HB, Yuan HJ, Hu PP

CASE REPORT

- 3813** Blood typing and transfusion therapy in a patient with A2 subtype acute myeloid leukemia M2: A case report

Kuang XC, Zhang SH, Cen YJ, Zhang JB, Liu YS

- 3822** Valve repair after infective endocarditis secondary to perforation caused by *Streptococcus gordonii*: A case report

Qu YF, Yang J, Wang JY, Wei B, Ye XH, Li YX, Han SL

- 3830** *Prevotella oris*-caused meningitis and spinal canal infection: A case report

Zhang WW, Ai C, Mao CT, Liu DK, Guo Y

- 3837** Severe liver trauma with complex portal and common bile duct avulsion: A case report and review of the literature

Mitricof B, Kraft A, Anton F, Barcu A, Barzan D, Haiducu C, Brasoveanu V, Popescu I, Moldovan CA, Botea F

- 3847** TACC diagnosed by transoesophageal endoscopic ultrasonography: A case report

Pu XX, Xu QW, Liu BY

- 3852** Ruptured teratoma mimicking a pelvic inflammatory disease and ovarian malignancy: A case report

Lai PH, Ding DC

- 3858** Purpura annularis telangiectodes of Majocchi: A case report

Pu YJ, Jiang HJ, Zhang L

- 3864** Giant cyst in heterotopic pregnancy: A case report

Kong YY, Chanda K, Ying XY

- 3870** High doses of dextromethorphan induced shock and convulsions in a 19-year-old female: A case report

Shimozawa S, Usuda D, Sasaki T, Tsuge S, Sakurai R, Kawai K, Matsubara S, Tanaka R, Suzuki M, Hotchi Y, Tokunaga S, Osugi I, Katou R, Ito S, Asako S, Mishima K, Kondo A, Mizuno K, Takami H, Komatsu T, Oba J, Nomura T, Sugita M

- 3877** Postpartum ovarian vein thrombosis after cesarean section and vaginal delivery: Two case reports

Zhu HD, Shen W, Wu HL, Sang X, Chen Y, Geng LS, Zhou T

- 3885** Traumatic pancreatic ductal injury treated by endoscopic stenting in a 9-year-old boy: A case report
Kwon HJ, Jung MK, Park J
- 3891** Novel mutation c.2090_2091del in neurodevelopmental-craniofacial syndrome with variable renal and cardiac abnormalities in an 18.5-mo-old boy: A case report
Li Y, Zhou Z, Xu Y, Wang ZR
- 3899** Reading impairment after neonatal hypoglycemia with parieto-temporo-occipital injury without cortical blindness: A case report
Kurahashi N, Ogaya S, Maki Y, Nonobe N, Kumai S, Hosokawa Y, Ogawa C, Yamada K, Maruyama K, Miura K, Nakamura M
- 3907** Unusual clinical presentation of oral pyogenic granuloma with severe alveolar bone loss: A case report and review of literature
Lomeli Martínez SM, Bocanegra Morando D, Mercado González AE, Gómez Sandoval JR
- 3915** Intraoperative photodynamic therapy for tracheal mass in non-small cell lung cancer: A case report
Jung HS, Kim HJ, Kim KW
- 3921** Coexistence of urinary tuberculosis and urothelial carcinoma: A case report
Tsai YC, Li CC, Chen BT, Wang CY

LETTER TO THE EDITOR

- 3929** Symmetric DWI hyperintensities in CMT1X patients after SARS-CoV-2 vaccination should not be classified as stroke-like lesions
Finsterer J

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Ashraf F Hefny, MD, MSc, Associate Professor, Surgeon, Department of Surgery, College of Medicine and Health Sciences, UAE University, Al Ain 00000, United Arab Emirates. ahefny@uaeu.ac.ae

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Si Zhao; Production Department Director: Xu Guo; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

June 6, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Retrospective Study

Dexmedetomidine-induced anesthesia in elderly patients undergoing hip replacement surgery

Jun-Qing Li, Hao Yuan, Xiao-Qiang Wang, Meng Yang

Specialty type: Medicine, research and experimental**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind**Peer-review report's scientific quality classification**Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0**P-Reviewer:** Roh DJ, United States;
Shelfbine SJ, United States**Received:** March 13, 2023**Peer-review started:** March 13, 2023**First decision:** March 28, 2023**Revised:** April 13, 2023**Accepted:** April 24, 2023**Article in press:** April 24, 2023**Published online:** June 6, 2023**Jun-Qing Li, Hao Yuan, Xiao-Qiang Wang, Meng Yang**, Department of Anesthesiology, Fuyang Hospital Affiliated to Anhui Medical University, Fuyang 236000, Anhui Province, China**Corresponding author:** Jun-Qing Li, MD, Attending Doctor, Department of Anesthesiology, Fuyang Hospital of Anhui Medical University, No. 99 Huangshan Road, Fuyang Hefei Modern Industrial Park, Yingzhou District, Fuyang 236000, Anhui Province, China.lijunqing005@126.com

Abstract

BACKGROUND

A femoral neck fracture is a common and frequently reported issue in orthopedics, with a greater rate of incidence among the elderly. Due to their advanced age and the presence of some primary diseases, both anesthesia and surgery are increasingly difficult in elderly patients with a femoral neck fractures. In fact, general anesthesia can easily induce complications such as cognitive dysfunction, which is not conducive to postoperative recovery.

AIM

To analyze the efficacy of dexmedetomidine in inducing anesthesia for elderly patients undergoing hip replacement surgery.

METHODS

A total of 98 elderly patients undergoing hip replacement in our hospital from June 2020 to June 2021 were randomly divided into control group (49 cases) and observation group (49 cases). The control group was given general anesthesia, and the observation group was combined with dexmedetomidine for anesthesia on the basis of the control group. Both groups were observed until the patients were discharged. The vital signs, serum inflammatory factors and renal function indexes of the two groups were compared before, during and 6 h after operation. The postoperative recovery and adverse events of the two groups were statistically analyzed.

RESULTS

Compared with the mean arterial pressure of the two groups, the intraoperative and postoperative 6 h was higher than that before the operation, the intraoperative was lower than the postoperative 6 h ($P < 0.05$); the blood oxygen saturation of the two groups was higher than that before operation and 6 h after operation, and the observation group was higher than the control group 6 h after

operation ($P < 0.05$). The heart rate of the two groups was lower during and 6 h after operation than that before operation, and higher at 6 h after operation than that during operation ($P < 0.05$). The levels of serum C-reactive protein, tumor necrosis factor- α , interleukin-1 β and kidney injury molecule-1 in the two groups were higher during operation and 6 h after operation than those before operation ($P < 0.05$). The level of serum urea nitrogen in the two groups was higher than that before operation, and that in the observation group was lower than that in the control group ($P < 0.05$). During hospitalization, the first time of getting out of bed, recovery time of grade II muscle strength, recovery time of grade III muscle strength and hospitalization time in the observation group were shorter than those in the control group ($P < 0.05$).

CONCLUSION

Dexmedetomidine can effectively improve the vital signs of elderly patients undergoing hip replacement surgery, reduce the body's inflammatory response and renal function damage, and promote postoperative recovery. Meanwhile, dexmedetomidine showcased a good safety profile and a good anesthetic outcome.

Key Words: Hip replacement; Old age; Dexmedetomidine; Anesthetic effect; Vital signs

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Dexmedetomidine is a type of drug that selectively activates the α_2 -adrenergic receptors. It stimulates the postsynaptic membrane receptors, inhibits sympathetic nerves, and effectively maintains the hemodynamic fluctuations in the body to achieve the desired sedative effects.

Citation: Li JQ, Yuan H, Wang XQ, Yang M. Dexmedetomidine-induced anesthesia in elderly patients undergoing hip replacement surgery. *World J Clin Cases* 2023; 11(16): 3756-3764

URL: <https://www.wjgnet.com/2307-8960/full/v11/i16/3756.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i16.3756>

INTRODUCTION

Recent years have witnessed an increase in the aging population of China and hence in its population structure. Because the hip bone of the elderly bears a huge load and is subjected to a relatively greater extent of activity, it is more prone to fracture. Hip replacement is the main approach for the treatment of the hip fracture. Due to the decline in the dopaminergic, cholinergic, and other neuroregulatory system functions in elderly patients, the extent of postoperative inflammatory response is significantly increased, which can easily cause multiple organ dysfunction. This factor is detrimental to the postoperative healing of elderly patients. Recent studies have revealed that maintaining the function of important organs in elderly patients undergoing hip replacement surgery during anesthesia and reducing any potential bodily damage is conducive to postoperative recovery[1-4]. Midazolam and propofol are commonly applied anesthetics in elderly patients during surgery. The combination of these two agents has a good anesthetic outcome, but it cannot effectively protect organ functions[4,5]. Dexmedetomidine is a highly selective α_2 receptor agonist, which has a good effect in reducing the body's stress response and in stabilizing the patient's intraoperative hemodynamics. Hence, it has a good application prospect in perioperative organ protection[6,7]. However, the applicability of dexmedetomidine as anesthesia in elderly patients with hip replacement surgery is not established yet. This study intended to analyze the anesthetic outcome of dexmedetomidine in elderly patients with hip replacement.

MATERIALS AND METHODS

General information

A total of 98 elderly patients scheduled for hip replacement surgery who were admitted to our hospital during June 2020-2021, were randomly assigned to two groups: Control ($n = 49$) and observation ($n = 49$). In the observation group: There were 23 women and 26 men with body mass index (BMI) of 20-26 kg/m² (average: 24.02 \pm 0.41 kg/m²) and average age 67.69 \pm 2.79 years (age range: 62-74 years). The study included patients with injuries ranging from 18 to 65 years old, with 10 cases of injury resulting from crush injuries, 24 from traffic accidents, and 15 from falls. In the control group, there were 22

women and 27 men of BMI 20–27 kg/m² (average: 23.67 ± 0.37 kg/m²). The average age of the control subjects is 68.55 ± 2.25 years (age range: 63–73 years). In these subjects, 25 cases of injury were caused by traffic accidents, 9 by squeezing, and 15 by falling. The general data of the two groups were comparable ($P > 0.05$).

Inclusion and exclusion criteria

Inclusion criteria: The diagnostic criteria of hip fracture met the “Expert consensus on the diagnosis and treatment of hip fracture in the elderly (2017)” [8], which was confirmed by preoperative X-ray examination: Age ≥ 60 years; those who met the surgical indications of hip replacement; traumatic fracture admission; complete clinical data; informed consent of this study. Exclusion criteria: Preoperative severe malnutrition; patients with severe infectious diseases; patients with severe organ dysfunction and malignant tumors.

Anesthesia method

The control group was administered the general anesthesia, as follows: Anesthesia induction: Midazolam (Jiangsu Enhua Pharmaceutical Co., Ltd., Sinopharm H19990027, 1 mL: 5 mg) 0.03–0.05 mg/kg, sufentanil (Yichang Renfu Pharmaceutical Co., Ltd., Sinopharm H20054171, 1 mL: 50 µg) 0.3–0.5 µg/kg, etomidate (Jiangsu Enhua Pharmaceutical Co., Ltd., Sinopharm H20020511, 10 mL: 20 mg) 0.2–0.3 mg/kg, Rocuronium bromide (N. V. Organon, import drug registration number H20140847, 5 mL: 50 mg) 0.6–1.0 mg/kg. A tracheal catheter or I-GEL laryngeal mask was inserted after 2–5 min of muscle relaxation. Anesthesia maintenance was performed as follows: Sevoflurane (Albert Pharmaceutical Trading Co., Ltd., import drug registration number H20150020, 250 mL) inhalation concentration 1%–2%, propofol (Xi'an Libang Pharmaceutical Co., Ltd., Sinopharm approval number: H19990282, 20 mL: 0.2 g) 2–5 mg/(kg·h), remifentanyl (Yichang Renfu Pharmaceutical Co., Ltd., Sinopharm approval number H20030197, 1 mg) 0.05–0.2 µg/(kg·min). Intermittent intravenous injection of cisatracurium (Zhejiang Xianju Pharmaceutical Co., Ltd., Sinopharm H20090202, 5 mg) 3–4 mg to maintain muscle relaxation. Sufentanil was added, with the total amount not exceeding 1 µg/kg, depending on the patient's response to anesthesia. The parameters of mechanical ventilation were set as follows: The recommended respiratory parameters were a tidal volume of 6–10 mL/kg, a respiratory rate of 8–12 breaths/min, and an inspiratory-to-expiratory ratio of 1:2. The mean arterial pressure (MAP) and heart rate (HR) were maintained within the basic value of ± 20%, and vasoactive drugs were administered based on the blood pressure of the patient during the surgery. Postoperative patient-controlled intravenous analgesia included sufentanil 200 µg, dezocine (Yangzijiang Pharmaceutical Group Co., Ltd., Sinopharm H20184150, 1 mL: 5 mg) 10 mg, tropisetron (Hainan Lingkan Pharmaceutical Co., Ltd., Sinopharm H20060287, 5 mg) 10 mg, normal saline 144 mL, total 150 mL; duration of 75 h, intravenous self-control 2 mL/h, load 0–2 mL, continuous volume 1.5–2 mL/h, and a lock time of 30 min.

Based on the anesthesia scheme of the control group, the observation group was treated with dexmedetomidine (Yangzijiang Pharmaceutical Group Co., Ltd., Chinese Pharmacopoeia: H20183219, 2 mL: 0.2 mg). Anesthesia induction and maintenance were performed consistently between the two groups. Before anesthesia induction, the patients were intravenously administered 4 µg/mL dexmedetomidine for 10 min, which was then changed to 0.2 µg/(kg·h). Both groups were observed until the patient was discharged from the hospital.

Observation indicators

Vital signs Multifunctional ECG monitor [MX550, Philips (China) Investment Co., Ltd.] was used to measure the MAP, HR, and blood oxygen saturation (SpO₂) of the two groups before, during, and 6 h after the surgery.

The levels of serum inflammatory factors C-reactive protein (CRP), tumor necrosis factor-α (TNF-α), and interleukin-1β (IL-1β) were determined by enzyme-linked immunosorbent assay (ELISA) before, during, and 6 h after the surgery for the two study groups (Shanghai Enzyme-linked Biotechnology Co., Ltd.).

To examine the renal functions, the levels of serum kidney injury molecule-1 (KIM-1), urea nitrogen (BUN), and creatinine (Cr) were detected by ELISA before, during, and 6 h after the surgery.

The following postoperative recovery factors were assessed and recorded: The first time getting out of bed, the recovery time of grade II muscle strength, the recovery time of grade III muscle strength, and the time of hospitalization.

The incidence of the following adverse events was recorded for the two groups of elderly patients, intraoperative hypotension, bradycardia, hypoxemia, and general body movement-related adverse events.

Statistical analysis

SPSS 21.0 statistical software was used for data analysis, with $P < 0.05$ considered to indicate a statistically significant difference. The measurement data included vital signs, serum inflammatory factor levels, renal function, and postoperative recovery. The mean ± SD represented the use of repeated measures analysis of variance for comparison. If the football symmetry test failed, the G-G correction

Table 1 Comparison of vital signs between the two groups (mean \pm SD)

Group	<i>n</i>	MAP (mmHg)	SpO ₂ (%)	HR (beat/min)
Preoperative				
Control group	49	90.02 \pm 3.39	96.24 \pm 3.07	88.12 \pm 4.16
Observation group	49	91.12 \pm 3.37	97.29 \pm 2.45	87.14 \pm 4.11
<i>F</i> value	-	2.136	1.871	1.389
<i>P</i> value	-	0.147	0.064	0.242
Intraoperative				
Control group	49	75.61 \pm 4.49 ^a	99.63 \pm 0.49 ^a	81.59 \pm 3.21 ^a
Observation group	49	79.61 \pm 3.41 ^a	99.80 \pm 0.41 ^a	79.12 \pm 3.26 ^a
<i>F</i> value	-	23.995	1.863	3.779
<i>P</i> value	-	< 0.001	0.066	< 0.001
6 h after operation				
Control group	49	82.78 \pm 3.74 ^{a,d}	98.49 \pm 0.68 ^{a,d}	86.73 \pm 4.28 ^{a,d}
Observation group	49	88.49 \pm 4.67 ^{a,d}	98.92 \pm 0.28 ^{a,d}	83.43 \pm 3.45 ^{a,d}
<i>F</i> value	-	46.040	10.001	4.202
<i>P</i> value	-	< 0.001	0.056	< 0.001

^a*P* < 0.05 *vs* preoperative.^d*P* < 0.05 *vs* intraoperative.MAP: Mean arterial pressure; HR: Heart rate; SpO₂: Blood oxygen saturation.

method was applied. Under the interaction, the separation effect between the population and time was analyzed further. The adverse events were counted as data, expressed as cases (percent) [*n* (%)], and passed through χ^2 tests for comparison.

RESULTS

Comparison of the vital signs between the two groups of patients

The vital signs of MAP, SpO₂, and HR data of the patients in the control and observation groups met the spherical symmetry test. The MAP, SpO₂ between the groups, time, and the interaction between the groups and time should be subjected to detailed analysis to understand the individual effect. The MAP of the two groups was lower than that before and after 6 h of the surgery; the MAP value was higher in the observation group than in the control group (*P* < 0.05). The SpO₂ standards in both groups were greater than that before and 6 h after the surgery. The HR in the observation group was higher than that in the control group (*P* < 0.05). In contrast with the HR of the two groups, the HR during and 6 h after the surgery was below that recorded before surgery, while the HR at 6 h after surgery was greater than that during surgery (*P* < 0.05). During and 6 h after surgery, the HR in the observation group was lower than that in the control group (*P* < 0.01) (Table 1).

Comparison of serum inflammatory factor levels between the two groups

The levels of serum CRP, TNF- α , and IL-1 β between the two groups were not satisfied with the results of the football symmetry test and the G-G correction method. The interaction among the groups, time, and between the groups and time was found to be significant, after which the individual effects were analyzed. The serum levels of CRP and TNF- α were recorded at 6 h after surgery in both two groups. The levels of TNF- α and IL-1 β were higher than those before and during surgery, but lower than those of the control group (*P* < 0.05) (Table 2).

Comparison of renal function indexes between the two groups

The renal function satisfied the spherical symmetry test. The group, time, and interaction between the group and time of KIM-1 and BUN have significant further analysis of the individual effect. As shown in Table 3, there was no interaction between the Cr index time and group (*P* > 0.05). Comparison of the serum KIM-1 Levels between the two groups showed that the levels during and 6 h after the surgery were higher than that before the surgery, those at 6 h after the surgery were higher than those during

Table 2 Comparison of serum inflammatory factor levels between the two groups (mean \pm SD, ng/L)

Group	<i>n</i>	CRP	TNF- α	IL-1 β
Preoperative				
Control group	49	3.33 \pm 1.21	15.11 \pm 4.20	173.55 \pm 23.52
Observation group	49	3.15 \pm 1.20	15.50 \pm 4.03	173.15 \pm 22.50
<i>F</i> value	-	0.562	0.220	0.086
<i>P</i> value	-	0.455	0.640	0.932
Intraoperative				
Control group	49	15.53 \pm 4.22 ^a	53.23 \pm 8.12 ^a	199.25 \pm 22.10 ^a
Observation group	49	7.25 \pm 3.20 ^a	31.20 \pm 5.21 ^a	190.23 \pm 21.33 ^a
<i>F</i> value	-	119.785	255.597	4.222
<i>P</i> value	-	< 0.001	< 0.001	0.043
6 h after operation				
Control group	49	18.25 \pm 4.13 ^{a,d}	60.27 \pm 9.23 ^{a,d}	293.25 \pm 11.50 ^{a,d}
Observation group	49	11.73 \pm 2.50 ^{a,d}	38.25 \pm 8.20 ^{a,d}	203.19 \pm 12.37 ^{a,d}
<i>F</i> value	-	89.449	156.076	1393.312
<i>P</i> value	-	< 0.001	< 0.001	< 0.001

^a*P* < 0.05 *vs* preoperative.^d*P* < 0.05 *vs* intraoperative.CRP: C-reactive protein; TNF- α : Tumor necrosis factor-alpha; IL-1 β : Interleukin-1 β .**Table 3 Comparison of renal function between two groups (mean \pm SD)**

Group	<i>n</i>	KIM-1 (ng/L)	BUN (ng/L)	Cr (μ mol/L)
Preoperative				
Control group	49	16.27 \pm 5.11	5.20 \pm 1.29	76.15 \pm 11.09
Observation group	49	16.83 \pm 4.07	5.29 \pm 1.26	75.56 \pm 12.59
<i>t</i> value	-	0.361	0.130	0.062
<i>P</i> value	-	0.549	0.720	0.804
Intraoperative				
Control group	49	28.22 \pm 6.55 ^a	6.98 \pm 1.22 ^a	95.52 \pm 11.51 ^a
Observation group	49	21.09 \pm 5.55 ^a	6.07 \pm 1.13 ^a	91.12 \pm 13.30 ^a
<i>t</i> value	-	33.745	14.779	3.976
<i>P</i> value	-	< 0.001	< 0.001	0.049
6 h after operation				
Control group	49	89.55 \pm 8.55 ^{a,d}	6.08 \pm 1.23	86.05 \pm 10.34 ^{a,d}
Observation group	49	52.50 \pm 6.76 ^{a,d}	5.98 \pm 1.13	81.25 \pm 11.36 ^{a,d}
<i>F</i> value	-	566.402	0.197	4.787
<i>P</i> value	-	< 0.001	0.658	0.031

^a*P* < 0.05 *vs* preoperative.^d*P* < 0.05 *vs* intraoperative.

KIM-1: Serum kidney injury molecule-1; BUN: Urea nitrogen; Cr: Creatinine.

the surgery, during and 6 h after surgery, the observation group was lower than the control group ($P < 0.05$). The serum BUN levels after the surgery were elevated in both groups when compared to that before surgery. The observation group showed significantly lower levels than the control group ($P < 0.05$) for the measured parameter. The comparison of serum Cr levels between the two groups revealed that the levels during and 6 h after surgery were higher than those before surgery. It is lower during and 6 h after surgery. **Table 3** depicts that the levels of the measured parameter in the observation group during and 6 h after surgery were significantly lower compared to those in the control group ($P < 0.05$).

Comparison of postoperative recovery between the two groups

The factors of the first time of getting out of bed, the recovery time of grade II muscle strength, grade III muscle strength, and hospitalization time were shorter in the observation group than in the control group ($P < 0.05$; **Table 4**).

Comparison of adverse events between the two groups of patients

During hospitalization, the rate of incidence of adverse events in the control group was higher than that in the observation group ($P < 0.05$; **Table 5**).

DISCUSSION

Hip fracture is the most common type of fracture occurring in the elderly. Hip fractures are commonly treated in clinics through total hip arthroplasty as one of the main preferred methods for the elderly. Due to the reduction of all aspects of the body in elderly patients undergoing hip arthroplasty, the stress response and inflammatory response induced by surgery can aggravate the damage to important organs of the patients, which is not conducive to the recovery of the patients[9-11]. During the surgery, the use of certain anesthetic drugs can maintain the hemodynamic stability of the patient and reduce the damage to the patient's organs. Midazolam and propofol are common anesthetics applied in clinical practice, which have proven good sedative and analgesic effects, but they are not effective in protecting the functions of important organs of patients and need to be combined with other anesthetics for anesthesia[12,13]. In elderly patients undergoing hip replacement surgery, changes in hemodynamics, respiratory functions, and elevated levels of the inflammatory response can lead to short-term postoperative cognitive impairment[14-16]. Based on the current findings, the MAP of the observation group was higher than that of the control group during surgery. In addition, the HR of the observation group was lower than that of the control group during and after surgery. At 6 h after surgery, the MAP and SpO₂ levels as well as the serum CRP and TNF- α levels were higher for the patients in the observation group than for those in the control group. The levels of IL-1 β and dexmedetomidine were significantly lower in the observation group than in the control group. This finding suggests that dexmedetomidine administration may be an effective intervention for improving vital signs and reducing inflammation in elderly patients with hip joint disorders. Dexmedetomidine can facilitate reaching the depth of anesthesia as soon as possible and thereby speed up the surgery process. Meanwhile, it can stabilize the hemodynamics of patients by inhibiting the secretion of norepinephrine and the activity of sympathetic nerves. In addition, it can maintain the patient's respiratory drive and improve the levels of MAP, HR, and SpO₂[17]. Dexmedetomidine can alleviate the synthesis of corticosteroids and glucocorticoids by enhancing parasympathetic activity and inhibiting sympathetic activity, which can effectively reduce the body's stress response and inflammatory response in elderly hip-replacement patients[18]. In elderly patients undergoing hip replacement surgery, increased levels of inflammatory response and hemodynamic changes may cause kidney damage. KIM-1 is a trans-membrane protein that is not expressed during normal renal functions, but appears after renal injury, thereby serving as a marker of early renal injury. BUN and Cr are the traditional indicators of renal functions, with elevated levels of renal functions. The present results revealed that the average serum levels of KIM-1, BUN, and Cr in the observation group were lower than those in the control group during and 6 h after the surgery, thereby indicating that dexmedetomidine can effectively alleviate renal injury in elderly hip-replacement patients and protect renal functions. Dexmedetomidine can effectively reduce the inflammatory response and damage caused by them to the renal cells in elderly patients undergoing hip-replacement surgery. On the other hand, dexmedetomidine can effectively inhibit the release of norepinephrine, reduce sympathetic nerve excitability, and increase renal blood flow, which cumulatively protects renal functions. The present results also suggest that the postoperative recovery of the observation group was better than that of the control group. Moreover, the total frequency of adverse events was lesser than that of the control group, which signifies that dexmedetomidine can effectively enhance the recovery of articulation coxae in the elderly. We thus found that dexmedetomidine could reduce the body's inflammatory response and kidney damage and stabilize the patient's vital signs, thereby indicating a good safety profile. Meanwhile, its sedative effect is non-anesthetic, rapidly metabolized in the body, and shows a strong dose dependency. The low-dose dexmedetomidine infusion used in this study led to quick awakening and restoration of consciousness in elderly patients after anesthesia, implying its safety and efficacy.

Table 4 Comparison of postoperative recovery between the two groups (mean \pm SD)

Group	<i>n</i>	The first time to get out of bed (h)	II grade muscle strength recovery time (h)	III grade muscle strength recovery time (h)	Hospitalization time (d)
Control group	49	26.84 \pm 8.73	20.41 \pm 7.72	36.41 \pm 8.71	7.94 \pm 3.72
Observation group	49	20.67 \pm 8.78	12.82 \pm 6.48	27.61 \pm 7.85	5.92 \pm 3.62
<i>t</i> value	-	3.485	5.274	5.253	2.725
<i>P</i> value	-	< 0.001	< 0.001	< 0.001	< 0.001

Table 5 Comparison of adverse reactions between the two groups, *n* (%)

Group	<i>n</i>	Hypotension	Bradycardia	Hypoxemia	General body movement	Total incidence of adverse reactions
Control group	49	3 (6.12)	1 (2.04)	2 (4.08)	2 (4.08)	8 (16.33)
Observation group	49	1 (2.04)	1 (2.04)	0 (0.00)	0 (0.00)	2 (4.08)
χ^2 value	-	-	-	-	-	4.009
<i>P</i> value	-	-	-	-	-	0.045

CONCLUSION

In summary, dexmedetomidine could effectively improve the vital signs of elderly patients undergoing hip replacement surgery, reduce the body's inflammatory response and renal function damage, and promote postoperative recovery. Meanwhile, it demonstrated good safety and anesthetic outcomes. This finding has significant clinical implications and merits further investigations for optimized application. The relevant indicators included in this study only compared the data of three different time points, before, during, and 6 h after the surgery, because of which the conclusion may be biased. In addition, as this study was conducted on a small sample size from a single center, the results may not apply to different populations. Subsequent research should include large sample size, a diverse subject population, and different age-group patients. We believe that a follow-up study on the changes of indicators at multiple time points and further analyses of the effect of dexmedetomidine in anesthesia during hip replacement surgery in the elderly are warranted to supplement the present findings.

ARTICLE HIGHLIGHTS

Research background

A femoral neck fracture is a common and frequently-occurring disease in orthopedics, with a high incidence in the elderly population. For elderly patients with a femoral neck fracture, due to their older age and more basic diseases, the difficulty of anesthesia and surgery is increased. General anesthesia can easily induce complications such as cognitive dysfunction, which is not conducive to postoperative recovery.

Research motivation

This study analyzed the anesthetic effect of dexmedetomidine in elderly patients undergoing hip replacement.

Research objectives

This study aimed to provide a reference for the prognosis and anesthesia of clinical-related surgeries.

Research methods

A total of 98 elderly patients undergoing hip replacement were randomly divided into control group and observation group. Both groups were observed until the patients were discharged. The vital signs, serum inflammatory factors and renal function indexes of the two groups were compared before, during and 6 h after operation.

Research results

Dexmedetomidine has good safety and good anesthetic effect.

Research conclusions

Dexmedetomidine can effectively improve the vital signs of elderly patients undergoing hip replacement, reduce the body's inflammatory response, reduce renal function damage, and promote postoperative recovery. At the same time, it has good safety and good anesthetic effect.

Research perspectives

This study indicates that diazepam has good application prospects in elderly patients undergoing hip replacement surgery. Future research can further explore the anesthesia effect and safety of diazepam, search for more optimized anesthesia plans, and improve the success rate of hip replacement surgery in elderly patients.

FOOTNOTES

Author contributions: Li JQ and Yuan H proposed concepts for this study; Wang XQ and Yang M collected data; Li JQ, Yuan H, and Wang XQ contributed to formal analysis; Li JQ and Yang M contributed to the survey; Li JQ, Yuan H, and Yang M contributed to this method; Li JQ, Yuan H, Wang XQ, and Yang M supervised the study; Li JQ validated this study; Yuan H and Wang XQ contributed to the visualization of research; Li JQ and Yuan H initially drafted this manuscript; Li JQ, Yuan H, Wang XQ, and Yang M reviewed and edited the manuscript.

Institutional review board statement: The study was reviewed and approved by the Institutional Review Committee of the First Hospital Affiliated to Anhui Medical University.

Informed consent statement: All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

Conflict-of-interest statement: We declare that there are no conflicts of interest.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Jun-Qing Li 0000-0002-6550-7860.

S-Editor: Wang JL

L-Editor: A

P-Editor: Fan JR

REFERENCES

- 1 **Chen H**, Li F. Effect of Dexmedetomidine with Different Anesthetic Dosage on Neurocognitive Function in Elderly Patients After Operation Based on Neural Network Model. *World Neurosurg* 2020; **138**: 688-695 [PMID: 31931250 DOI: 10.1016/j.wneu.2020.01.012]
- 2 **Lin C**, Tu H, Jie Z, Zhou X, Li C. Effect of Dexmedetomidine on Delirium in Elderly Surgical Patients: A Meta-analysis of Randomized Controlled Trials. *Ann Pharmacother* 2021; **55**: 624-636 [PMID: 32885993 DOI: 10.1177/1060028020951954]
- 3 **Prathapadas U**, Hrishi AP, Appavoo A, Vimala S, Sethuraman M. Effect of Low-Dose Dexmedetomidine on the Anesthetic and Recovery Profile of Sevoflurane-Based Anesthesia in Patients Presenting for Supratentorial Neurosurgeries: A Randomized Double-Blind Placebo-Controlled Trial. *J Neurosci Rural Pract* 2020; **11**: 267-273 [PMID: 32367982 DOI: 10.1055/s-0040-1703968]
- 4 **Li Y**, Wang C, Bi M, Gao J, Zhang X, Tian H. Effect of dexmedetomidine on brain function and hemodynamics in patients undergoing lung cancer resection. *Oncol Lett* 2020; **20**: 1077-1082 [PMID: 32724346 DOI: 10.3892/ol.2020.11675]
- 5 **Song C**, Lu Q. Effect of dexmedetomidine supplementation for thoracoscopic surgery: a meta-analysis of randomized controlled trials. *J Cardiothorac Surg* 2022; **17**: 70 [PMID: 35382835 DOI: 10.1186/s13019-022-01803-z]
- 6 **Learmonth ID**, Young C, Rorabeck C. The operation of the century: total hip replacement. *Lancet* 2007; **370**: 1508-1519 [PMID: 17964352 DOI: 10.1016/S0140-6736(07)60457-7]
- 7 **Gupta N**, Das S, Chatterjee N, Munjal M. A Retrospective Study of Ultrasound-Guided Pericapsular Nerve Group Block With Dexamethasone: An Excellent Option for Early Mobility Following Total Hip Replacement Surgery. *Cureus* 2022;

- 14: e32515 [PMID: 36654572 DOI: 10.7759/cureus.32515]
- 8 **Mei B**, Meng G, Xu G, Cheng X, Chen S, Zhang Y, Zhang M, Liu X, Gu E. Intraoperative Sedation With Dexmedetomidine is Superior to Propofol for Elderly Patients Undergoing Hip Arthroplasty: A Prospective Randomized Controlled Study. *Clin J Pain* 2018; **34**: 811-817 [PMID: 29528863 DOI: 10.1097/AJP.0000000000000605]
- 9 **Uusalo P**, Jätinvuori H, Löytyniemi E, Kosola J, Saari TI. Intranasal Low-Dose Dexmedetomidine Reduces Postoperative Opioid Requirement in Patients Undergoing Hip Arthroplasty Under General Anesthesia. *J Arthroplasty* 2019; **34**: 686-692.e2 [PMID: 30733071 DOI: 10.1016/j.arth.2018.12.036]
- 10 **Park JW**, Kim EK, Lee HT, Park S, Do SH. The Effects of Propofol or Dexmedetomidine Sedation on Postoperative Recovery in Elderly Patients Receiving Lower Limb Surgery under Spinal Anesthesia: A Retrospective Propensity Score-Matched Analysis. *J Clin Med* 2021; **10** [PMID: 33401537 DOI: 10.3390/jcm10010135]
- 11 **Xu T**, Chen X, Li X, Wang M. Effect of dexmedetomidine-assisted ultrasound-guided lower extremity nerve block on postoperative cognitive function in elderly patients undergoing hip surgery. *Am J Transl Res* 2022; **14**: 7977-7984 [PMID: 36505284]
- 12 **Kim DJ**, Kim SH, So KY, Jung KT. Effects of dexmedetomidine on smooth emergence from anaesthesia in elderly patients undergoing orthopaedic surgery. *BMC Anesthesiol* 2015; **15**: 139 [PMID: 26446479 DOI: 10.1186/s12871-015-0127-4]
- 13 **Shin HJ**, Woo Nam S, Kim H, Yim S, Han SH, Hwang JW, Do SH, Na HS. Postoperative Delirium after Dexmedetomidine versus Propofol Sedation in Healthy Older Adults Undergoing Orthopedic Lower Limb Surgery with Spinal Anesthesia: A Randomized Controlled Trial. *Anesthesiology* 2023; **138**: 164-171 [PMID: 36534899 DOI: 10.1097/ALN.0000000000004438]
- 14 **Kim JE**, Kim NY, Lee HS, Kil HK. Effects of intrathecal dexmedetomidine on low-dose bupivacaine spinal anesthesia in elderly patients undergoing transurethral prostatectomy. *Biol Pharm Bull* 2013; **36**: 959-965 [PMID: 23727917 DOI: 10.1248/bpb.b12-01067]
- 15 **Du X**, Yu J, Mi W. The effect of dexmedetomidine on the perioperative hemodynamics and postoperative cognitive function of elderly patients with hypertension: Study protocol for a randomized controlled trial. *Medicine (Baltimore)* 2018; **97**: e12851 [PMID: 30412077 DOI: 10.1097/MD.00000000000012851]
- 16 **Liu T**, Liu FC, Xia Y, Luo TJ, Wang F, Chen B, Wang C, Gao GK, Zhai WT, Liu W. Effect of dexmedetomidine on the Montreal Cognitive Assessment in older patients undergoing pulmonary surgery. *J Int Med Res* 2022; **50**: 3000605221123680 [PMID: 36151758 DOI: 10.1177/03000605221123680]
- 17 **Shen Y**, Wang C, Zhong X, Wu Y, He X. Observation of Anesthetic Effect of Dexmedetomidine Combined With Intraspinal Anesthesia in Hip Arthroplasty and its Effect on Postoperative Delirium and Stress Response. *Front Surg* 2022; **9**: 928922 [PMID: 35846975 DOI: 10.3389/fsurg.2022.928922]
- 18 **Pan H**, Liu C, Ma X, Xu Y, Zhang M, Wang Y. Perioperative dexmedetomidine reduces delirium in elderly patients after non-cardiac surgery: a systematic review and meta-analysis of randomized-controlled trials. *Can J Anaesth* 2019; **66**: 1489-1500 [PMID: 31264195 DOI: 10.1007/s12630-019-01440-6]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

