

World Journal of *Clinical Cases*

World J Clin Cases 2023 July 6; 11(19): 4458-4733



Contents

Thrice Monthly Volume 11 Number 19 July 6, 2023

REVIEW

- 4458 Molecular signalling during cross talk between gut brain axis regulation and progression of irritable bowel syndrome: A comprehensive review
Singh SV, Ganguly R, Jaiswal K, Yadav AK, Kumar R, Pandey AK
- 4477 Diffusion tensor imaging in the courtroom: Distinction between scientific specificity and legally admissible evidence
van Velkinburgh JC, Herbst MD, Casper SM

MINIREVIEWS

- 4498 Inequity in the global distribution of monkeypox vaccines
Tovani-Palone MR, Doshi N, Pedersini P
- 4504 Long-term effectiveness, outcomes and complications of bariatric surgery
Gulinac M, Miteva DG, Peshevska-Sekulovska M, Novakov IP, Antovic S, Peruhova M, Snegarova V, Kabakchieva P, Assyov Y, Vasilev G, Sekulovski M, Lazova S, Tomov L, Velikova T

ORIGINAL ARTICLE

Retrospective Cohort Study

- 4513 Age, blood tests and comorbidities and AIMS65 risk scores outperform Glasgow-Blatchford and pre-endoscopic Rockall score in patients with upper gastrointestinal bleeding
Morarasu BC, Sorodoc V, Haisan A, Morarasu S, Bologa C, Haliga RE, Lionte C, Marciuc EA, Elsiddig M, Cimpoesu D, Dimofte GM, Sorodoc L

Retrospective Study

- 4531 Application of cross-migration theory in limb rehabilitation of stroke patients with hemiplegia
Lu YH, Fu Y, Shu J, Yan LY, Shen HJ
- 4544 Analysis of characteristic features in ultrasound diagnosis of fetal limb body wall complex during 11-13⁺ weeks
Ye CH, Li S, Ling L
- 4553 Network pharmacology and molecular docking-based analyses to predict the potential mechanism of Huangqin decoction in treating colorectal cancer
Li YJ, Tang DX, Yan HT, Yang B, Yang Z, Long FX
- 4567 Assessment of functional prognosis of anterior cruciate ligament reconstruction in athletes based on a body shape index
Wang YJ, Zhang JC, Zhang YZ, Liu YH

EVIDENCE-BASED MEDICINE

- 4579** Network pharmacology and molecular docking to explore Polygoni Cuspidati Rhizoma et Radix treatment for acute lung injury

Zheng JL, Wang X, Song Z, Zhou P, Zhang GJ, Diao JJ, Han CE, Jia GY, Zhou X, Zhang BQ

ORIGINAL ARTICLE**Randomized Controlled Trial**

- 4601** Ulinastatin in the treatment of severe acute pancreatitis: A single-center randomized controlled trial

Wang SQ, Jiao W, Zhang J, Zhang JF, Tao YN, Jiang Q, Yu F

- 4612** Fecal microbiota transplantation in patients with metabolic syndrome and obesity: A randomized controlled trial

da Ponte Neto AM, Clemente ACO, Rosa PW, Ribeiro IB, Funari MP, Nunes GC, Moreira L, Sparvoli LG, Cortez R, Taddei CR, Mancini MC, de Moura EGH

SYSTEMATIC REVIEWS

- 4625** Combined medial patellofemoral ligament and medial patellotibial ligament reconstruction in recurrent patellar instability: A systematic review and meta-analysis

Abbaszadeh A, Saeedi M, Hoveidaei AH, Dadgostar H, Razi S, Razi M

CASE REPORT

- 4635** Unique Roberts syndrome with bilateral congenital glaucoma: A case report

Almulhim A, Almoallem B, Alsirrhay E, Osman EA

- 4640** CK5/6-positive, P63-positive lymphoepithelioma-like hepatocellular carcinoma: A case report and literature review

Tang HT, Lin W, Zhang WQ, Qian JL, Li K, He K

- 4648** Edaravone administration and its potential association with a new clinical syndrome in cerebral infarction patients: Three case reports

Yang L, Xu X, Wang L, Zeng KB, Wang XF

- 4655** CDKN1C gene mutation causing familial Silver-Russell syndrome: A case report and review of literature

Li J, Chen LN, He HL

- 4664** Hypothetical hypoxia-driven rapid disease progression in hepatocellular carcinoma post transarterial chemoembolization: A case report

Yeo KF, Ker A, Kao PE, Wang CC

- 4670** Metastatic colon cancer treated using traditional Chinese medicine combined with chemotherapy: A case report

Deng CG, Tang MY, Pan X, Liu ZH

- 4677** Rare cause of cerebral venous sinus thrombosis: Spontaneous intracranial hypotension syndrome: A case report

Huang P

- 4684** Integrated Chinese and Western medicine in the treatment of a patient with podocyte infolding glomerulopathy: A case report
Chang MY, Zhang Y, Li MX, Xuan F
- 4692** Morbihan disease misdiagnosed as senile blepharoptosis and successfully treated with short-term minocycline and ketotifen: A case report
Na J, Wu Y
- 4698** With two episodes of right retromandibular angle subcutaneous emphysema during right upper molar crown preparation: A case report
Bai YP, Sha JJ, Chai CC, Sun HP
- 4707** Poststroke rehabilitation using repetitive transcranial magnetic stimulation during pregnancy: A case report
Jo J, Kim H
- 4713** Tuberculosis-induced aplastic crisis and atypical lymphocyte expansion in advanced myelodysplastic syndrome: A case report and review of literature
Sun XY, Yang XD, Xu J, Xiu NN, Ju B, Zhao XC
- 4723** Posterior reversible encephalopathy syndrome following uneventful clipping of an unruptured intracranial aneurysm: A case report
Hwang J, Cho WH, Cha SH, Ko JK

ACADEMIC WRITING

- 4729** Revitalizing case reports: Standardized guidelines and mentorship
Jeyaraman M, Ramasubramanian S, Jeyaraman N, Nallakumarasamy A, Sharma S

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Miroslav Vujasinovic, MD, PhD, Associate Professor, Department of Upper Abdominal Diseases, Karolinska University Hospital, Stockholm 14186, Sweden. mvujas@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Si Zhao; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

July 6, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Analysis of characteristic features in ultrasound diagnosis of fetal limb body wall complex during 11-13⁺ weeks

Cai-Hong Ye, Shuo Li, Li Ling

Specialty type: Pediatrics

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Guerriero S, Italy;
Matute JD, United States

Received: April 14, 2023

Peer-review started: April 14, 2023

First decision: April 26, 2023

Revised: May 3, 2023

Accepted: May 23, 2023

Article in press: May 23, 2023

Published online: July 6, 2023



Cai-Hong Ye, Department of Ultrasound, Yijishan Hospital Wannan Medical College, Wuhu 241001, Anhui Province, China

Shuo Li, Department of Ultrasound, Wannan Medical College, Wuhu 241001, Anhui Province, China

Li Ling, Department of Obstetrics, Wannan Medical College, Wuhu 241001, Anhui Province, China

Corresponding author: Cai-Hong Ye, Doctor, MM, Associate Chief Physician, Deputy Director, Department of Ultrasound, Yijishan Hospital Wannan Medical College, No. 2 Zheshan West Road, Wuhu 241001, Anhui Province, China. yhyjsyy@163.com

Abstract

BACKGROUND

Limb body wall complex (LBWC) is a fatal malformation characterized by major defects in the fetal abdominal or thoracic wall, visceral herniation, significant scoliosis or spina bifida, limb deformities, craniofacial deformities, and umbilical cord abnormalities (short or absent umbilical cord). Early diagnosis of this condition is of great clinical significance for clinical intervention and pregnancy decision-making. With the rapid development of fetal ultrasound medicine, early pregnancy (11-13⁺ wk) standardized prenatal ultrasound examinations have been widely promoted and applied.

AIM

To explore the value of prenatal ultrasound in the diagnosis of fetal LBWC syndrome during early pregnancy.

METHODS

The ultrasonographic data and follow-up results of 18 cases of fetal LBWC diagnosed by prenatal ultrasound during early pregnancy (11-13⁺ wk) were retrospectively analyzed, and their ultrasonographic characteristics were analyzed.

RESULTS

Among the 18 fetuses with limb wall abnormalities, there were spinal dysplasia (18/18, 100%), varying degrees of thoracoschisis and gastroschisis (18/18, 100%), limb dysplasia in 6 cases (6/18, 33%), craniocerebral malformations in 4 cases

(4/18, 22%), thickening of the transparent layer of the neck in 5 cases (5/18, 28%), and umbilical cord abnormalities in 18 cases (18/18, 100%), single umbilical artery in 5 cases.

CONCLUSION

Prenatal ultrasound in early pregnancy can detect LBWC as early as possible, and correct prenatal evaluation provides important guidance value for pregnancy decision-making and early intervention.

Key Words: Early pregnancy; Ultrasonography; Limb body wall complex; Fetus

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Limbs-body wall complex (LBWC) is a fatal malformation characterized by severe defects in the abdominal or chest wall of the fetus, resulting in protruding viscera, severe scoliosis or spina bifida, limb deformities, craniofacial malformations, and abnormal umbilical cord (short or absent). Our study focuses on the significance of using prenatal ultrasound in detecting LBWC during early pregnancy, where we also provide a retrospective analysis of ultrasound data and post-abortion consequences of 18 confirmed cases of LBWC. We aim to investigate further and identify the ultrasound characteristics of LBWC in early pregnancy to enhance our knowledge of this condition.

Citation: Ye CH, Li S, Ling L. Analysis of characteristic features in ultrasound diagnosis of fetal limb body wall complex during 11-13⁺⁶ weeks. *World J Clin Cases* 2023; 11(19): 4544-4552

URL: <https://www.wjgnet.com/2307-8960/full/v11/i19/4544.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i19.4544>

INTRODUCTION

Limb body wall complex (LBWC) is a fatal malformation characterized by major defects in the fetal abdominal or thoracic wall, visceral herniation, significant scoliosis or spina bifida, limb deformities, craniofacial deformities, and umbilical cord abnormalities (short or absent umbilical cord)[1,2]. Early diagnosis of this condition is of great clinical significance for intervention and pregnancy decision-making. With the rapid development of fetal ultrasound medicine[3,4], early pregnancy (11-13⁺⁶ wk) standardized prenatal ultrasound examinations have been widely promoted and applied. However, there are relatively few reports on early pregnancy ultrasound screening for LBWC. In this study, we retrospectively analyzed the sonographic data and post-abortion results of 18 cases diagnosed with LBWC in early pregnancy to explore the ultrasound features of prenatal diagnosis of LBWC in early pregnancy and enhance the understanding of LBWC.

MATERIALS AND METHODS

Study subjects

From March 2015 to October 2022, a total of 35486 pregnant women underwent early pregnancy (11-13⁺⁶ wk) fetal nuchal translucency (NT) ultrasound examination at our hospital. Among them, 18 cases of LBWC were detected and confirmed after induced abortion. The pregnant women's ages ranged from 21 years to 39 years, with an average age of 28 years. The gestational age of the fetuses was calculated based on the menstrual cycle, and all were singleton pregnancies.

Study methods

GE Voluson E8 and Philips EPIQ7 color Doppler ultrasound diagnostic instruments were used in our study, with a 2-8 MHz two-dimensional convex array probe. In accordance with the requirements of the Chinese "Prenatal Ultrasound Screening Guidelines", pregnant women were informed of the appropriate gestational age, content, purpose, importance, and limitations of early pregnancy prenatal ultrasound screening. In this study, all patients were placed in a supine position, and during the examination, in addition to routine checks for the number of fetuses, fetal heartbeat, placenta, and amniotic fluid, the main growth parameters of the fetus and the spectrum of the venous duct were measured. In cases with multiple pregnancies, chorionicity and amnionicity were determined. The focus was on obtaining the fetal mid-sagittal plane and measuring the crown-rump length and NT thickness on this plane, taking the average of three measurements. The presence or absence of the fetal nasal bone



DOI: 10.12998/wjcc.v11.i19.4544 Copyright ©The Author(s) 2023.

Figure 1 Ultrasound sonographic analysis of limb body wall complex. A: The fetus with the absence of the thoracic and abdominal walls and the contents seen floating within the amniotic fluid; B: The baby postdelivery confirming the evisceration of the heart, liver, stomach, large and small bowel loops; C: Fetal umbilical artery blood flow spectrum, no venous catheter blood flow detected; D: Kyphosis of the thoracolumbar spine.

was also observed.

In addition, the following ultrasound screening planes were included: starting from the fetal head and sequentially examining the skull and brain. The lateral ventricle transverse plane primarily observed the lateral ventricle level cranial halo, cerebral falx, and bilateral choroid plexus in the lateral ventricles. With the abdomen, the umbilical cord abdominal wall entrance transverse plane primarily observed the structure at the umbilical cord abdominal wall entrance. All planes were saved in the workstation. If a defect in the fetal abdominal wall or thoracoabdominal wall was found, it was necessary to carefully observe whether there were abnormal echo masses and the size, shape, and structural echoes of the masses, whether the physiological curvature of the spine had changed, the development of the limbs, and whether there were any abnormalities in the umbilical cord.

RESULTS

General information of pregnant women

As shown in Table 1, the ages of pregnant women ranged from 21 years to 39 years old, with an average of 27-years-old. Gestational age ranged from 11-13⁺⁶ wk, with an average gestational age of 12⁺⁶ wk. Among the 18 cases, 11 cases of the 18 cases were screened for Down's syndrome in the first trimester and all were found to be low risk.

Fetal NT measurement

As shown in Table 2, all 18 cases of LBWC fetuses were single live fetuses. Among the 11 cases, 5 had increased NT thickness, 2 had cystic hygromas, and 7 were not measurable.

Ultrasound sonographic analysis of LBWC

As shown in Table 3, all fetuses had severe thoracoschisis or gastroschisis (18/18, 100%), either spinal scoliosis and torsion (18/18, 100%). Among them, 6 cases (6/18, 33%) had limb developmental abnormalities, 4 cases (4/18, 22%) had cranial malformations, and 1 case had spina bifida. Umbilical cord abnormalities, such as being too short or absent, were observed in 11 cases (11/18, 61%). Color Doppler examination revealed only one umbilical artery in 5 fetuses. None of the fetuses had detectable facial abnormalities (Figure 1). All 18 pregnant women chose to terminate their pregnancies, which was

Table 1 General information of pregnant women

No.	Age	Gestational age in wk	Pregnancy history, G/P/A/L	Embryo transplantation	Down syndrome screening status
1	29	13	G1P0	No	Unchecked
2	28	13 ⁺⁶	G2P0A1	Yes	Unchecked
3	25	13 ⁺⁵	G1P0	No	Low-risk
4	21	12 ⁺³	G1P0	No	Low-risk
5	23	12 ⁺⁴	G1P0	No	Low-risk
6	26	12	G2P0A1	No	Low-risk
7	30	11 ⁺²	G2P0A1	No	Low-risk
8	29	13	G1P0	No	Low-risk
9	38	11 ⁺³	G1P0	Yes	Low-risk
10	25	13 ⁺²	G1P0	No	Low-risk
11	28	12 ⁺⁶	G2P0A1	No	Low-risk
12	24	11 ⁺⁶	G1P0	No	Low-risk
13	32	12 ⁺³	G2P1L1	No	Low-risk
14	26	12 ⁺⁴	G1P0	No	Unchecked
15	39	13 ⁺⁶	G1P0	Yes	Unchecked
16	31	13 ⁺⁵	G2P0A1	Yes	Unchecked
17	21	13 ⁺²	G1P0	No	Low-risk
18	26	13	G1P0	No	Low-risk

G: Gestation; P: Production; A: Abortion; L: Live.

confirmed after induced abortions.

DISCUSSION

Embryonic development mechanism of fetal LBWC

LBWC is an uncommon, intricate, and fatal multi-malformation syndrome characterized by encephalocele and/or craniofacial clefts, thoracoschisis or gastroschisis, and limb deformities. Other features may include spinal scoliosis, genitourinary abnormalities, and short umbilical cords. The initial diagnostic criteria for LBWC were established by Van Allen *et al*[5] in 1987, and a diagnosis can be confirmed by the presence of two of the following three malformations: any type of limb defect, encephalocele or anencephaly with facial cleft, thoracoschisis or gastroschisis with visceral herniation.

The etiology of LBWC remains uncertain, there are three pathogenic mechanisms that have been discussed by foreign scholars[6-10]. Firstly, abnormal folding of the trilaminar embryonic disc during the first 4 wk of embryonic development. Secondly, early amniotic rupture with amniotic band syndrome. Finally, early embryonic vascular rupture leading to widespread blood flow impairment.

(1) Abnormal folding of the trilaminar embryonic disc during the first 4 wk of embryonic development, which represents the complete failure of folding along the three axes (cranial, caudal, and lateral) of the body. Abnormal longitudinal folding results in thoracoschisis or gastroschisis, while abnormal transverse folding prompts abdominal contents to protrude into an expansive amniotic sac, which inserts peripherally into the placental chorionic plate, resulting in absent or short umbilical cords. As a result of the compression of abdominal viscera, the spine and thoracic cavity cannot develop symmetrically, leading to severe spinal scoliosis and trunk abnormalities. Poor spinal rotation and incomplete pelvic closure may lead to congenital limb deformities[7].

(2) Early amniotic rupture. Early amniotic rupture caused by mechanical pressure or amniotic bands can lead to severe deformities such as spinal malformations, gastroschisis, and extremely short umbilical cords if the rupture occurs at the caudal end, with the lower half of the embryo entering the extraembryonic cavity through the rupture[8]. This mechanism was questioned by Paul *et al*[11], who reported a case of intact LBWC at 10 wk of gestation without rupture of the amniotic cavity.

Table 2 Fetal nuchal translucency measurement result

No.	NT	Nasal bone visible	Ductus venosus
1	2.0	Yes	Visible, no reverse a-wave
2	Unmeasurable	No	Invisible
3	Unmeasurable	No	Visible, no reverse a-wave
4	Unmeasurable	Yes	Invisible
5	Unmeasurable	No	Invisible
6	2.3	Yes	Visible, no reverse a-wave
7	3.8	Yes	Invisible
8	Unmeasurable	No	Visible, no reverse a-wave
9	Unmeasurable	Yes	Invisible
10	1.3	Yes	Visible, no reverse a-wave
11	1.8	Yes	Invisible
12	3.4	Yes	Visible, no reverse a-wave
13	1.1	Yes	Visible, no reverse a-wave
14	4.7	Yes	Visible, no reverse a-wave
15	2.8	Yes	Visible, no reverse a-wave
16	Unmeasurable	No	Invisible
17	3.6	Yes	Visible, no reverse a-wave
18	2.4	Yes	Visible, no reverse a-wave

NT: Nuchal translucency.

And (3) early embryonic vascular rupture or developmental abnormality. Some studies[10] suggest that vascular rupture during early pregnancy (at 4-6 wk of gestation) leads to the interruption of normal blood supply to the developing embryo, resulting in ischemia, hypoxia, hemorrhagic necrosis, and abnormal development of the embryonic disc, which in turn affects the closure of the abdominal wall and extraembryonic cavity.

Arshad believes that the pathogenesis of LBWC is more inclined towards the theory of early embryonic abnormal folding[12], with the malformations associated with LBWC depending on the degree of abnormal folding of the embryonic disc. On the other hand, Sahinoglu *et al*[13] and others believe that due to the complexity of LBWC malformations, all of the above pathogenic mechanisms are possible, leading to different subgroups.

LBWC classification

There are many literature reports and controversies about the classification of LBWC, and most studies currently divide it into two types based on the characteristic abnormalities of LBWC[13-16]. Type I is based on craniofacial defects, while Type II is based on thoracoschisis or gastroschisis. However, Sahinoglu *et al*[13] proposed a new phenotypic classification, dividing it into three types, Type I still refers to craniofacial defects, while Type II refers to gastroschisis located above the umbilicus and large chest-abdominal wall defects. Type III pertains to defects located below the umbilicus, while the chest wall remains intact. This article primarily concentrates on thoracoschisis and gastroschisis, which aligns with the research conducted by Akinmoladun *et al*[17].

Ultrasonographic features of LBWC in early pregnancy

At present, early pregnancy ultrasound examination is considered the safest and most reliable imaging technique for screening severe fetal structural malformations. Owing to the relatively large extraembryonic coelom in early pregnancy, it becomes straightforward to distinctly display the extent of thoracoschisis or gastroschisis as well as the protrusion of organs in the thoracoabdominal cavity into the extraembryonic coelom, making the diagnosis relatively easy. With the widespread popularity of early pregnancy (11-13⁺6 wk) prenatal ultrasound examinations, many severe fetal structural malformations have been detected. Some scholars believe that the most characteristic ultrasonographic clue for LBWC in early pregnancy is spinal curvature[18]. The basis for ultrasound diagnosis of LBWC may be when the fetus has a large abdominal or chest-abdominal wall defect, herniation of the viscera, short or

Table 3 Ultrasound sonographic analysis of limb body wall complex

Ultrasonic anomaly						
No.	Body wall defect and bulge	Spine	Limbs	Brain	Umbilical cord	Others
1	Gastroschisis (bulge is liver)	Torsion	Both feet varus	-	Short	
2	Gastroschisis (bulge is liver)	Torsion	-	-	Undetectable	Cloacal malformation
3	Thoracoschisis and Gastroschisis (bulge are heart, liver, intestinal tube)	Twist into angles	-	Brain herniation	Short	
4	Gastroschisis (bulge are liver, gastric vesicles and intestinal tubes)	Twisted arrangement irregularity	-	Encephalocele	Undetectable	Fetal cystic hygroma
5	Gastroschisis (bulge are liver, gastric vesicles and intestinal tubes)	Torsion	Miss left upper limb	Meningoencephalocele	Undetectable	
6	Gastroschisis (bulge are liver, gastric vesicles and intestinal tubes)	Curved, middle and lower segments not shown	-		Short	
7	Gastroschisis (bulge are liver, gastric vesicles and intestinal tubes)	Twist into angles	Miss one upper limb		Undetectable	
8	Gastroschisis (bulge are liver, gastric vesicles and intestinal tubes)	Abnormal bending, abnormal alignment			Short	Fetal cystic hygroma
9	Gastroschisis (bulge is liver)	Torsion		Meningoencephalocele	Undetectable	
10	Gastroschisis (bulge are liver, gastric vesicle and intestinal tube)	Lordotic bending		-	Short	Single umbilical artery
11	Thoracoschisis and gastroschisis (bulge are heart, liver, intestinal tube and gastric vesicle)	Lordotic bending		-	Undetectable	
12	Gastroschisis (bulge are liver, intestinal tube)	Lordotic bending		-	Short	
13	Gastroschisis (bulge are liver, intestinal tube)	Angle the side		-	Short	
14	Gastroschisis and malformation (bulge are liver and intestine)	Lordotic bending	Miss left upper arm	-	Short	
15	Gastroschisis and malformation (bulge are liver, intestinal tube and gastric vesicle)	Side bending and twisting		-	Short	Single umbilical artery
16	Gastroschisis (bulge are liver, gastric vesicles and intestinal tubes)	Lordotic bending	Lower limb fusion	-	Undetectable	
17	Gastroschisis (bulge are liver, gastric vesicles and intestinal tubes)	Torsion	Miss right lower limb		Short	
18	Acromphalus (bulge is liver)	Torsion	-		Short	Megacystis

absent umbilical cord combined with spinal abnormalities[19,20].

The sonographic characteristics of the 18 LBWC fetuses in our investigation can be summarized as follows: (1) The most common and severe malformation in LBWC is the thoracoschisis or gastroschisis [21,22]. Due to the absence of abdominal wall development, the defect area is large and may be accompanied by chest wall defects, resulting in the opening of the thoracoabdominal viscera to the amniotic cavity. Prenatal ultrasound shows a large defect at the insertion of the fetal umbilical cord on the abdominal wall, with abnormal echo masses at the defect site, and the fetal liver, stomach bubble, and intestine protruding into the extraembryonic coelom. The umbilical cord is difficult to display or appears too short. Daskalakis *et al*[23] and others believe that abdominal wall defects can be roughly divided into three types, gastroschisis, umbilical protrusion, and body stalk-like malformation. LBWC can be considered when accompanied by spinal curvature or torsion, anencephaly or encephalocele, facial abnormalities, and limb development abnormalities; (2) All 18 fetuses had spinal developmental abnormalities, manifested as spinal kyphosis, torsion or scoliosis, causing the fetus to be in a flexed or hyperextended state, making it difficult to obtain the mid-sagittal section required for measuring crown-rump length in prenatal ultrasound screening guidelines, and thus unable to accurately determine the gestational age of the fetus by ultrasound. In this study, all cases were calculated by menstrual age. In this group, the earliest large abdominal wall defect was seen at 11⁺² wk. In this group, all cases had varying degrees of thoracoschisis or gastroschisis and abnormal spinal curvature; (3) Increased fetal NT,

similar to the findings of Daniilidis *et al*[24]. In our study, 4 cases out of 18 LBWC cases were normal, 5 cases had increased NT, and 9 cases could not be measured due to spinal torsion or concomitant cranial abnormalities. Although LBWC fetuses often have abnormal NT thickening or fetal edema, 8 cases in this group underwent early pregnancy Down syndrome screening and all were classified as low risk. Previous studies[13,25] have shown that fetuses with LBWC have a normal chromosomal karyotype and the recurrence rate in subsequent pregnancies is very low; (4) Cranial abnormalities. In our 18 cases, 3 had encephalocele, and 1 had exencephaly; (5) Limb abnormalities. In our study, limb abnormalities were manifested as clubfoot, limb agenesis, or fusion; and (6) Short or absent umbilical cord. Other associated malformations include megacystis, fetal cystic hygroma, and abnormal development of the cloacal cavity.

Differential diagnosis of LBWC

Owing to the numerous resemblances in ultrasound manifestations between LBWC and Cantrell's pentalogy[25], the latter primarily encompasses omphalocele, ectopia cordis, lower sternal defects, anterior diaphragmatic defects, and pericardial defects. Its hallmark features are omphalocele and ectopia cordis. Nevertheless, Cantrell's pentalogy does not involve spinal and limb developmental abnormalities, which serve as crucial distinguishing factors from LBWC[26].

CONCLUSION

As LBWC is a lethal disease, enhancing early detection is of utmost importance. Prenatal ultrasound should be well-acquainted with the sonographic features of LBWC, such as thoracoschisis and gastroschisis, severe spinal scoliosis, neural tube defects, limb abnormalities, and absence or shortening of the umbilical cord. Furthermore, when spinal scoliosis and omphalocele are observed, the likelihood of LBWC should be strongly considered. In conclusion, early pregnancy ultrasound can effectively diagnose LBWC, and precise prenatal evaluation holds significant value for informing parental decision-making.

ARTICLE HIGHLIGHTS

Research background

There are relatively few reports on early pregnancy ultrasound screening for Limb body wall complex (LBWC). In this study, we retrospectively analyzed the sonographic data and post-abortion results of 18 cases diagnosed with LBWC in early pregnancy to explore the ultrasound features of prenatal diagnosis of LBWC in early pregnancy and enhance the understanding of LBWC.

Research motivation

In this study, we retrospectively analyzed the sonographic data and post-abortion results of 18 cases diagnosed with LBWC in early pregnancy to explore the ultrasound features of prenatal diagnosis of LBWC in early pregnancy and enhance the understanding of LBWC.

Research objectives

To explore the value of prenatal ultrasound in the diagnosis of fetal LBWC syndrome during early pregnancy.

Research methods

The ultrasonographic data and follow-up results of 18 cases of fetal LBWC diagnosed by prenatal ultrasound during early pregnancy (11-13⁺ wk) were retrospectively analyzed, and their ultrasonographic characteristics were analyzed.

Research results

Among the 18 fetuses with limb wall abnormalities, there were spinal dysplasia (18/18, 100%), varying degrees of thoracoschisis and gastroschisis (18/18, 100%), limb dysplasia in 6 cases (6/18, 33%), cranio-cerebral malformations in 4 cases (4/18, 22%), thickening of the transparent layer of the neck in 5 cases (5/18, 28%), and umbilical cord abnormalities in 18 cases (18/18, 100%), single umbilical artery in 5 cases.

Research conclusions

Prenatal ultrasound in early pregnancy can detect LBWC as early as possible, and correct prenatal evaluation provides important guidance value for pregnancy decision-making and early intervention.

Research perspectives

As LBWC is a lethal disease, enhancing early detection is of utmost importance. Prenatal ultrasound should be well-acquainted with the sonographic features of LBWC, such as thoracoschisis and gastroschisis, severe spinal scoliosis, neural tube defects, limb abnormalities, and absence or shortening of the umbilical cord. Furthermore, when spinal scoliosis and omphalocele are observed, the likelihood of LBWC should be strongly considered. In conclusion, early pregnancy ultrasound can effectively diagnose LBWC, and precise prenatal evaluation holds significant value for informing parental decision-making.

FOOTNOTES

Author contributions: Ye CH contributed to the conception and design, administrative support, and manuscript writing; Ye CH and Ling L contributed to the provision of study materials or patients; all authors contributed to the data collection and assembly; Li S contributed to the data analysis and interpretation; All authors contributed to the final approval of manuscript.

Institutional review board statement: The study was approved by Institutional Review Board of Yijishan Hospital, Wannan Medical College.

Informed consent statement: All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

Conflict-of-interest statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Data sharing statement: The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Cai-Hong Ye 0009-0009-5808-8897.

S-Editor: Yan JP

L-Editor: Filipodia

P-Editor: Yan JP

REFERENCES

- Mandrekar SR**, Amoncar S, Banauliker S, Sawant V, Pinto RG. Omphalocele, exstrophy of cloaca, imperforate anus and spinal defect (OEIS Complex) with overlapping features of body stalk anomaly (limb body wall complex). *Indian J Hum Genet* 2014; **20**: 195-198 [PMID: 25400352 DOI: 10.4103/0971-6866.142906]
- Baruah P**, Ray Choudhury P. Limb body wall complex with sacrococcygeal mass and agenesis of external genitalia. *Case Rep Med* 2013; **2013**: 218626 [PMID: 23970900 DOI: 10.1155/2013/218626]
- Wang J**, Chen L, Zhou C, Wang L, Xie H, Xiao Y, Yin D, Zeng Y, Tang F, Yang Y, Zhu H, Chen X, Zhu Q, Liu Z, Liu H. Identification of copy number variations among fetuses with ultrasound soft markers using next-generation sequencing. *Sci Rep* 2018; **8**: 8134 [PMID: 29802277 DOI: 10.1038/s41598-018-26555-6]
- Yang T**, Li R, Liang N, Li J, Yang Y, Huang Q, Li Y, Cao W, Wang Q, Zhang H. The application of key feature extraction algorithm based on Gabor wavelet transformation in the diagnosis of lumbar intervertebral disc degenerative changes. *PLoS One* 2020; **15**: e0227894 [PMID: 32101549 DOI: 10.1371/journal.pone.0227894]
- Van Allen MI**, Curry C, Walden CE, Gallagher L, Patten RM. Limb-body wall complex: II. Limb and spine defects. *Am J Med Genet* 1987; **28**: 549-565 [PMID: 2962494 DOI: 10.1002/ajmg.1320280303]
- Daskalakis G**, Sebire NJ, Jurkovic D, Snijders RJ, Nicolaides KH. Body stalk anomaly at 10-14 weeks of gestation. *Ultrasound Obstet Gynecol* 1997; **10**: 416-418 [PMID: 9476328 DOI: 10.1046/j.1469-0705.1997.10060416.x]
- Hartwig NG**, Vermeij-Keers C, De Vries HE, Kagie M, Kragt H. Limb body wall malformation complex: an embryologic etiology? *Hum Pathol* 1989; **20**: 1071-1077 [PMID: 2530150 DOI: 10.1016/0046-8177(89)90225-6]
- Torpin R**. Amniochorionic Mesoblastic Fibrous Strings And Amnionic Bands: Associated Constricting Fetal Malformations Or Fetal Death. *Am J Obstet Gynecol* 1965; **91**: 65-75 [PMID: 14245093 DOI: 10.1016/0002-9378(65)90588-0]

- 9 **Russo R**, Vecchione R. Limb body wall complex: craniofacial defects as a distinctive factor. *Birth Defects Orig Artic Ser* 1996; **30**: 157-164 [PMID: 9125324]
- 10 **Van Allen MI**, Curry C, Gallagher L. Limb body wall complex: I. Pathogenesis. *Am J Med Genet* 1987; **28**: 529-548 [PMID: 2962493 DOI: 10.1002/ajmg.1320280302]
- 11 **Paul C**, Zosmer N, Jurkovic D, Nicolaides K. A case of body stalk anomaly at 10 weeks of gestation. *Ultrasound Obstet Gynecol* 2001; **17**: 157-159 [PMID: 11251927 DOI: 10.1046/j.1469-0705.2001.00248.x]
- 12 **Bhat A**, Ilyas M, Dev G. Prenatal sonographic diagnosis of limb-body wall complex: case series of a rare congenital anomaly. *Radiol Case Rep* 2016; **11**: 116-120 [PMID: 27257465 DOI: 10.1016/j.radcr.2016.02.004]
- 13 **Sahinoglu Z**, Uludogan M, Arik H, Aydin A, Kucukbas M, Bilgic R, Toksoy G. Prenatal ultrasonographical features of limb body wall complex: a review of etiopathogenesis and a new classification. *Fetal Pediatr Pathol* 2007; **26**: 135-151 [PMID: 17886024 DOI: 10.1080/15513810701563728]
- 14 **Russo R**, D'Armiento M, Angrisani P, Vecchione R. Limb body wall complex: a critical review and a nosological proposal. *Am J Med Genet* 1993; **47**: 893-900 [PMID: 8279488 DOI: 10.1002/ajmg.1320470617]
- 15 **Pumberger W**, Schaller A, Bernaschek G. Limb-body wall complex: a compound anomaly pattern in body-wall defects. *Pediatr Surg Int* 2001; **17**: 486-490 [PMID: 11527200 DOI: 10.1007/s003830000497]
- 16 **Colpaert C**, Bogers J, Hertveldt K, Loquet P, Dumon J, Willems P. Limb-body wall complex: 4 new cases illustrating the importance of examining placenta and umbilical cord. *Pathol Res Pract* 2000; **196**: 783-790 [PMID: 11186176 DOI: 10.1016/S0344-0338(00)80114-4]
- 17 **Akinmoladun J**, Bello O. Prenatal sonographic diagnosis of limb body wall complex: A rare lethal fetal anomaly. *Sahel Med J* 2019; **22**: 226 [DOI: 10.4103/smj.smj_51_18]
- 18 **Becker R**, Runkel S, Entezami M. Prenatal diagnosis of body stalk anomaly at 9 weeks of gestation. Case report. *Fetal Diagn Ther* 2000; **15**: 301-303 [PMID: 10971084 DOI: 10.1159/000021025]
- 19 **Mulita F**, Parchas N, Solou K, Tchabashvili L, Gatamati F, Iliopoulos F, Maroulis I. Postoperative Pain Scores After Open Inguinal Hernia Repair: Comparison of Three Postoperative Analgesic Regimens. *Med Arch* 2020; **74**: 355-358 [PMID: 33424089 DOI: 10.5455/medarch.2020.74.355-358]
- 20 **Jayaram PR**, Pereira FD, Barrett JA. Evaluation of dynamic ultrasound scanning in the diagnosis of equivocal ventral hernias with surgical comparison. *Br J Radiol* 2018; **91** [PMID: 29745748 DOI: 10.1259/bjr.20180056]
- 21 **Daemen JHT**, Loonen TGJ, Lozekoot PWJ, Maessen JG, Maal TJJ, Hulsewé KWE, Vissers YLJ, de Loos ER. Optical imaging versus CT and plain radiography to quantify pectus severity: a systematic review and meta-analysis. *J Thorac Dis* 2020; **12**: 1475-1487 [PMID: 32395285 DOI: 10.21037/jtd.2020.02.31]
- 22 **Lain A**, Garcia L, Gine C, Tiffet O, Lopez M. New Methods for Imaging Evaluation of Chest Wall Deformities. *Front Pediatr* 2017; **5**: 257 [PMID: 29255700 DOI: 10.3389/fped.2017.00257]
- 23 **Daskalakis G**, Pilalis A, Papadopoulos D, Antsaklis A. Body stalk anomaly diagnosed in the 2nd trimester. *Fetal Diagn Ther* 2003; **18**: 342-344 [PMID: 12913346 DOI: 10.1159/000071978]
- 24 **Daniilidis A**, Balaouras D, Chitzios D, Balaouras G, Capilna M, Asimakopoulos E. Variation of ultrasound findings in the first trimester examination of recurrent cases with trisomy 21. *J Clin Med Res* 2015; **7**: 495-498 [PMID: 25883716 DOI: 10.14740/jocmr2138w]
- 25 **Chen CP**, Lin CJ, Chang TY, Hsu CY, Tzen CY, Wang W. Second-trimester diagnosis of limb-body wall complex with literature review of pathogenesis. *Genet Couns* 2007; **18**: 105-112 [PMID: 17515306]
- 26 **Stein W**, Haller F, Hawighorst T, Emons G. Pentalogy of Cantrell vs. limb body wall complex: differential diagnosis of a severe malformation in early pregnancy. *Ultraschall Med* 2009; **30**: 598-601 [PMID: 19998210 DOI: 10.1055/s-0028-1109743]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

