

World Journal of *Clinical Cases*

World J Clin Cases 2023 July 6; 11(19): 4458-4733



Contents

Thrice Monthly Volume 11 Number 19 July 6, 2023

REVIEW

- 4458 Molecular signalling during cross talk between gut brain axis regulation and progression of irritable bowel syndrome: A comprehensive review
Singh SV, Ganguly R, Jaiswal K, Yadav AK, Kumar R, Pandey AK
- 4477 Diffusion tensor imaging in the courtroom: Distinction between scientific specificity and legally admissible evidence
van Velkinburgh JC, Herbst MD, Casper SM

MINIREVIEWS

- 4498 Inequity in the global distribution of monkeypox vaccines
Tovani-Palone MR, Doshi N, Pedersini P
- 4504 Long-term effectiveness, outcomes and complications of bariatric surgery
Gulinac M, Miteva DG, Peshevska-Sekulovska M, Novakov IP, Antovic S, Peruhova M, Snegarova V, Kabakchieva P, Assyov Y, Vasilev G, Sekulovski M, Lazova S, Tomov L, Velikova T

ORIGINAL ARTICLE

Retrospective Cohort Study

- 4513 Age, blood tests and comorbidities and AIMS65 risk scores outperform Glasgow-Blatchford and pre-endoscopic Rockall score in patients with upper gastrointestinal bleeding
Morarasu BC, Sorodoc V, Haisan A, Morarasu S, Bologa C, Haliga RE, Lionte C, Marciuc EA, Elsiddig M, Cimpoesu D, Dimofte GM, Sorodoc L

Retrospective Study

- 4531 Application of cross-migration theory in limb rehabilitation of stroke patients with hemiplegia
Lu YH, Fu Y, Shu J, Yan LY, Shen HJ
- 4544 Analysis of characteristic features in ultrasound diagnosis of fetal limb body wall complex during 11-13⁺ weeks
Ye CH, Li S, Ling L
- 4553 Network pharmacology and molecular docking-based analyses to predict the potential mechanism of Huangqin decoction in treating colorectal cancer
Li YJ, Tang DX, Yan HT, Yang B, Yang Z, Long FX
- 4567 Assessment of functional prognosis of anterior cruciate ligament reconstruction in athletes based on a body shape index
Wang YJ, Zhang JC, Zhang YZ, Liu YH

EVIDENCE-BASED MEDICINE

- 4579** Network pharmacology and molecular docking to explore Polygoni Cuspidati Rhizoma et Radix treatment for acute lung injury

Zheng JL, Wang X, Song Z, Zhou P, Zhang GJ, Diao JJ, Han CE, Jia GY, Zhou X, Zhang BQ

ORIGINAL ARTICLE**Randomized Controlled Trial**

- 4601** Ulinastatin in the treatment of severe acute pancreatitis: A single-center randomized controlled trial

Wang SQ, Jiao W, Zhang J, Zhang JF, Tao YN, Jiang Q, Yu F

- 4612** Fecal microbiota transplantation in patients with metabolic syndrome and obesity: A randomized controlled trial

da Ponte Neto AM, Clemente ACO, Rosa PW, Ribeiro IB, Funari MP, Nunes GC, Moreira L, Sparvoli LG, Cortez R, Taddei CR, Mancini MC, de Moura EGH

SYSTEMATIC REVIEWS

- 4625** Combined medial patellofemoral ligament and medial patellotibial ligament reconstruction in recurrent patellar instability: A systematic review and meta-analysis

Abbaszadeh A, Saeedi M, Hoveidaei AH, Dadgostar H, Razi S, Razi M

CASE REPORT

- 4635** Unique Roberts syndrome with bilateral congenital glaucoma: A case report

Almulhim A, Almoallem B, Alsirrhay E, Osman EA

- 4640** CK5/6-positive, P63-positive lymphoepithelioma-like hepatocellular carcinoma: A case report and literature review

Tang HT, Lin W, Zhang WQ, Qian JL, Li K, He K

- 4648** Edaravone administration and its potential association with a new clinical syndrome in cerebral infarction patients: Three case reports

Yang L, Xu X, Wang L, Zeng KB, Wang XF

- 4655** CDKN1C gene mutation causing familial Silver-Russell syndrome: A case report and review of literature

Li J, Chen LN, He HL

- 4664** Hypothetical hypoxia-driven rapid disease progression in hepatocellular carcinoma post transarterial chemoembolization: A case report

Yeo KF, Ker A, Kao PE, Wang CC

- 4670** Metastatic colon cancer treated using traditional Chinese medicine combined with chemotherapy: A case report

Deng CG, Tang MY, Pan X, Liu ZH

- 4677** Rare cause of cerebral venous sinus thrombosis: Spontaneous intracranial hypotension syndrome: A case report

Huang P

- 4684** Integrated Chinese and Western medicine in the treatment of a patient with podocyte infolding glomerulopathy: A case report
Chang MY, Zhang Y, Li MX, Xuan F
- 4692** Morbihan disease misdiagnosed as senile blepharoptosis and successfully treated with short-term minocycline and ketotifen: A case report
Na J, Wu Y
- 4698** With two episodes of right retromandibular angle subcutaneous emphysema during right upper molar crown preparation: A case report
Bai YP, Sha JJ, Chai CC, Sun HP
- 4707** Poststroke rehabilitation using repetitive transcranial magnetic stimulation during pregnancy: A case report
Jo J, Kim H
- 4713** Tuberculosis-induced aplastic crisis and atypical lymphocyte expansion in advanced myelodysplastic syndrome: A case report and review of literature
Sun XY, Yang XD, Xu J, Xiu NN, Ju B, Zhao XC
- 4723** Posterior reversible encephalopathy syndrome following uneventful clipping of an unruptured intracranial aneurysm: A case report
Hwang J, Cho WH, Cha SH, Ko JK

ACADEMIC WRITING

- 4729** Revitalizing case reports: Standardized guidelines and mentorship
Jeyaraman M, Ramasubramanian S, Jeyaraman N, Nallakumarasamy A, Sharma S

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Miroslav Vujasinovic, MD, PhD, Associate Professor, Department of Upper Abdominal Diseases, Karolinska University Hospital, Stockholm 14186, Sweden. mvujas@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Si Zhao; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

July 6, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Assessment of functional prognosis of anterior cruciate ligament reconstruction in athletes based on a body shape index

Yun-Jun Wang, Jun-Chang Zhang, Yu-Ze Zhang, Ying-Hai Liu

Specialty type: Orthopedics

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Baird PN, Australia;
Mahmud N, United States

Received: May 4, 2023

Peer-review started: May 4, 2023

First decision: May 12, 2023

Revised: May 21, 2023

Accepted: May 24, 2023

Article in press: May 24, 2023

Published online: July 6, 2023



Yun-Jun Wang, Yu-Ze Zhang, Ying-Hai Liu, Institutes of Biomedical Sciences, Shanxi University, Taiyuan 030006, Shanxi Province, China

Jun-Chang Zhang, Department of Orthopedics, First Hospital of Shanxi Medical University, Taiyuan 030001, Shanxi Province, China

Corresponding author: Ying-Hai Liu, PhD, Professor, Institutes of Biomedical Sciences, Shanxi University, No. 92 Wucheng Road, Taiyuan 030006, Shanxi Province, China.
liuyinghai3699@163.com

Abstract

BACKGROUND

A healthy body shape is essential to maintain athletes' sports level. At present, little is known about the effect of athletes' body shape on anterior cruciate ligament reconstruction (ACLR). Moreover, the relationship between body shape and variables such as knee joint function after operation and return to the field has not been well studied.

AIM

To verify the relationship between a body shape index (ABSI) and the functional prognosis of the knee after ACLR in athletes with ACL injuries.

METHODS

We reviewed 76 athletes with unilateral ACL ruptures who underwent ACLR surgery in the First Hospital of Shanxi Medical University between 2017 and 2020, with a follow-up period of more than 24 mo. First, all populations were divided into a High-ABSI group ($ABSI > 0.835$, $n = 38$) and a Low-ABSI group ($ABSI < 0.835$, $n = 38$) based on the arithmetic median (0.835) of ABSI values. The primary exposure factor was ABSI, and the outcome indicators were knee function scores as well as postoperative complications. The correlation between ABSI and postoperative knee function scores and postoperative complications after ACLR were analyzed using multifactorial logistic regression.

RESULTS

The preoperative knee function scores of the two groups were similar. The surgery and postoperative rehabilitation exercises, range of motion (ROM) compliance rate, Lysholm score, and Knee Injury and Osteoarthritis Outcome Score of the two groups gradually increased, whereas the quadriceps atrophy

index gradually decreased. The knee function scores were higher in the Low-ABSI group than in the High-ABSI group at the 24-mo postoperative follow-up ($P < 0.05$). In multifactorial logistic regression, ABSI was a risk factor of low knee joint function score after surgery, specifically low ROM scores (odds ratio [OR] = 1.31, 95% confidence interval [CI] [1.10-1.44]; $P < 0.001$), low quadriceps atrophy index (OR = 1.11, 95% CI [0.97-1.29]; $P < 0.05$), low Lysholm scores (OR = 2.34, 95% CI [1.78-2.94]; $P < 0.001$), low symptoms (OR = 1.14, 95% CI [1.02-1.34]; $P < 0.05$), low activity of daily living (OR = 1.34, 95% CI [1.18-1.65]; $P < 0.05$), low sports (OR = 2.47, 95% CI [1.78-2.84]; $P < 0.001$), and low quality of life (OR = 3.34, 95% CI [2.88-3.94]; $P < 0.001$). ABSI was also a risk factor for deep vein thrombosis of the lower limb (OR = 2.14, 95% CI [1.88-2.36], $P < 0.05$) and ACL recurrent rupture (OR = 1.24, 95% CI [0.98-1.44], $P < 0.05$) after ACLR.

CONCLUSION

ABSI is a risk factor for the poor prognosis of knee function in ACL athletes after ACLR, and the risk of poor knee function after ACLR, deep vein thrombosis of lower limb, and ACL recurrent rupture gradually increases with the rise of ABSI.

Key Words: A body shape index; Anterior cruciate ligament reconstruction; Athletes; Multifactorial logistic regression; Obesity

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: A body shape index (ABSI), an excellent body mass index, was used to substitute for the traditional body mass index to more objectively assess the degree of body size/obesity in athletes. Subsequently, multivariate logistic regression analysis was used to determine whether ABSI is a risk factor for poor knee function, lower extremity deep vein thrombosis, and fractures after anterior cruciate ligament surgery. The findings of this study have the potential to narrow the gap in previous research.

Citation: Wang YJ, Zhang JC, Zhang YZ, Liu YH. Assessment of functional prognosis of anterior cruciate ligament reconstruction in athletes based on a body shape index. *World J Clin Cases* 2023; 11(19): 4567-4578

URL: <https://www.wjgnet.com/2307-8960/full/v11/i19/4567.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i19.4567>

INTRODUCTION

Athletes are at high risk of anterior cruciate ligament (ACL) ruptures, often combined with meniscal and cartilage injuries, which can lead to knee instability and osteoarthritis (OA) if not treated promptly [1]. Due to the high demand for knee function in athletes, ACL reconstruction (ACLR) is generally recommended as early as possible to obtain knee stability after ACL rupture in athletes [2]. Given that good postoperative knee function is the foundation of an athlete's performance on the field [3], it is important to identify the factors that influence an athlete's post-ACLR knee function.

A proper body shape plays a central role in athletic performance, and the increasing obesity has a negative impact on athletes' physical performance, such as increased lower limb loading and slower movement speed [4]. As a result, it is important to regularly monitor the obesity of athletes to ensure that the mandatory physical obesity level is kept at a low level [5]. Obesity and overweight have previously been identified as high-risk variables for poor function following ACLR, and a higher body mass index (BMI) tends to predict higher knee loading and OA risk in ACLR patients, especially when the BMI reaches 25 kg/m² [6,7]. Furthermore, a high BMI can also alter post-ACLR gait biomechanics, and increase knee compression and shear forces, as well as patellofemoral pressure and knee extensor moment, which can lead to knee instability [8,9]. Therefore, an increasing number of athletes of all sizes or weights are in need of ACLR, and it is of great importance to identify risk factors that influence the outcomes and complications of ACLR in athletes.

BMI is a measurement of a person's weight and height; it is used to determine a person's fat proportion and is also a common clinical indicator for assessing overweight and obesity [10]. BMI does not directly measure body fat composition, but is defined as a positive correlation between weight and obesity at a specific height [11]. Currently, statistics on obesity are mostly based on BMI, which ranges from 18.5 to 24.9 in a healthy person, 25 to 29.9 in an overweight person, and 30+ in an obese person. However, the median distribution of "normal" BMI (approximately 22) differs considerably from the mean BMI of the current population. In addition, the correlation between BMI and the risk of various diseases is not linear [12]. Put another way, a higher BMI does not always reflect an increase in fat, as

being overweight may be due to an increase in muscle tissue, especially in athletes with higher levels of musculoskeletal development[13]. ABSI was first proposed by researchers at the City University of New York in 2012[14]. ABSI is calculated from height, weight, and waist circumference (WC), and it standardizes WC with body shape (weight and height), similar to body BMI[15]. ABSI corrects the effect of height and/or body mass to better assesses abdominal fat[16].

There is a paucity of research regarding the association between ABSI and postoperative knee function and complications in athletes with ACLR. Therefore, it is of great significance to identify risk factors associated with poorer postoperative knee function or complications in athletes with ACLR and to emphasize early intervention and guided rehabilitation to allow athletes to return to competition as soon as possible.

MATERIALS AND METHODS

Research subjects

Patients included in this retrospective cross-sectional study were professional athletes with ACL injuries who underwent arthroscopic ACLR between December 2017 and December 2020 at the Department of Orthopedics in the First Hospital of Shanxi Medical University (Shanxi, China), and all patients agreed to participate in this study. According to the Helsinki Declaration, the study was approved by the Ethics Committee of the First Hospital of Shanxi Medical University, and written informed consent was obtained from all patients. Because of the retrospective nature of our study, no additional clinical trial registration was required.

Inclusion criteria: (1) Active professional athletes aged 18-years-old to 40-years-old; (2) first-time knee arthroscopy, within 6 mo after ACL injury; (3) intraoperative definite diagnosis of unilateral ACL rupture (may be combined with meniscal or articular cartilage injury); and (4) patients who had no contraindications to surgery and agreed to undergo arthroscopic ACLR surgery and participate in this study.

Exclusion criteria: (1) Presence of lower extremity fractures or combined injuries such as medial or lateral collateral ligament or posterior cruciate ligament; (2) previous knee meniscal injury or osteoarthritis; and (3) history of previous knee surgery.

Ultimately, a total of 76 ACLR athletes were included in this study through screening.

General case information

Demographic information was collected through the medical record system and questionnaires including sex, age, type of exercise, smoking and alcohol history, preoperative physical examination (height, weight, BMI, and abdominal circumference), comorbidities (*e.g.*, hypertension, diabetes, cancer, meniscal injury), ACL injury, site of injury, and surgical status (duration of surgery, duration of tourniquet, femoral and tibial fixation).

ABSI calculation

The exposure variable in this study was ABSI, and it was calculated as follows: $BMI = \text{weight}/\text{height}^2$, and $ABSI = WC / (BMI^{2/3} \times \text{height}^{1/2})$ [17].

Grouping

First, the ABSI of all ACLR patients was calculated, and then the population was divided into a High-ABSI group ($ABSI > 0.835$, $n = 38$) and Low-ABSI group ($ABSI < 0.835$, $n = 38$) according to their arithmetic median (0.835). All patients were followed up for more than 24 mo.

ACLR surgical technique

As previously described[18], all ACLR procedures were performed on ACL patients by the same orthopedic professor under general or combined lumbar and rigid anesthesia using autologous hamstring tendons (semitendinosus and thin femoral muscles). All patients had a thigh tourniquet. Also, all patients had a standard anteromedial femoral and tibial tunnel drilled, with a final fixation of the proximal tendon bundled to the femur with an Endobutton (Smith & Nephew Inc., Memphis, TN, United States) or rounded cannulated interference (RCI) screw (Smith & Nephew, United Kingdom) and the proximal tendon bundled to the tibia with an RCI screw (Smith & Nephew, United Kingdom) or washer (Smith & Nephew UK Ltd., Watford, England, United Kingdom). Postoperatively, all patients followed a standard rehabilitation program.

Standard rehabilitation training

All patients received standard lower extremity nerve-muscle function training, including quadriceps (anterior thigh muscle group) isometric exercise. They also practiced thigh muscle tensing and relaxation, without significant pain, more than 500 times/d, and had posterior thigh muscle group

isometric exercise more than 500 times/d. Furthermore, they had pump exercises such as forceful, slow, full-range flexion and extension of the ankle joints, more than 500 times/d. Patients were clinically reviewed by the surgeon at 3 mo and 12 and 24 mo postoperatively.

Functional assessment of the knee joint

All patients completed a 24-mo postoperative follow-up to assess the functional recovery of the knees after surgeries.

Range of motion compliance rate

Range of motion (ROM) is the arc of motion through which the joint moves, and is the most basic method for assessing the motor function of the extremities[19]. Normal knee mobility should be between 140° and 150°, and active knee flexion of 120° or more after surgery is defined as compliance.

Quadriceps atrophy index

The quadriceps muscle is not only important for the overall health of the knee joint but also plays a key role in ensuring the knee joint's continued stability[20]. The quadriceps atrophy index is calculated as: atrophy index = (healthy thigh circumference - affected thigh circumference)/healthy thigh circumference × 100%. The smaller the value, the smaller the reduction in quadriceps strength and the better the recovery of muscle strength.

Knee Lysholm scale

The Lysholm Scale is a condition-specific scale for evaluating ligamentous injuries of the knee, and is also a subjective rating system in the form of a percentage questionnaire[21]. The scale consists of eight items: claudication (5 points), support (5 points), strangulation (15 points), instability (25 points), pain (25 points), swelling (10 points), stair climbing and descending (10 points), and squatting (5 points). A higher score indicates better function.

Knee Injury and Osteoarthritis Outcome Score

Knee Injury and Osteoarthritis Outcome Score (KOOS) is an assessment of the short- and long-term outcomes of treatment after a knee injury[22]. It consists of five patient-related components: symptoms, pain, activity of daily living (ADL), sports, and knee-related quality of life (QOL). A score of 0 means very poor function of that part of the joint, and a score of 100 means perfectly normal function of that part of the joint.

Return to sport

According to previous studies[18], "return to sport (RTS)" is defined as an exercise that is considered to return to the same or higher level of return to pre-injury during the follow-up period.

Postoperative complications

During the follow-up period, patients in both groups were counted for complications, including knee infection, deep vein thrombosis, internal fixation failure, and ACL re-rupture. The patients were analyzed for specific complications, and given symptomatic treatment.

Statistical analyses

IBM SPSS 27.10 software and GraphPad Prism 9 software were used for statistical analyses. The measurement data are expressed as the mean ± standard deviation, and one-way analysis of variance was used for the comparison between groups. The statistical data are expressed as frequencies (percentages), and the χ^2 test was used for the comparison between groups. The median method was used to divide all patients with ACLR into High-ASBI and Low-ASBI groups. The relationship between ASBI and postoperative knee function scores and complications after ACLR was evaluated by multifactorial logistic regression analysis. $P < 0.05$ was considered statistically significant.

RESULTS

General information

First, survey data were collected from a total of 93 athletes, and 76 ACLR athletes were included in the final analysis after excluding those patients who lacked outcomes, exposure, or did not complete follow-up (Figure 1).

As shown in Table 1, there was no significant difference in age between the High-ASBI group (ASBI > 0.835) and Low-ASBI group (ASBI < 0.835) in the preliminary analyses (25.9 ± 6.3 vs 26.4 ± 7.1 ; $P = 0.7463$). There were 57 male athletes and 19 female athletes, and there was no significant difference in sex composition between the two groups (males: 71.05% vs 78.95%; $P = 0.5970$), with both groups

Table 1 Characteristics of participants enrolled in the study, *n* (%)

Characteristics	High-ABSI group, <i>n</i> = 38	Low-ABSI group, <i>n</i> = 38	<i>P</i> value
Age in yr	25.9 ± 6.3	26.4 ± 7.1	0.7463
Age range in yr	18-38	18-39	
Sex			0.5970
Female, <i>n</i> = 19	11 (28.94)	8 (21.05)	
Male, <i>n</i> = 57	27 (71.05)	30 (78.95)	
Athletes			0.8731
Football	18	19	
Basketball	16	14	
Others	4	5	
Body mass index as kg/m ²	24.3 ± 2.2	23.9 ± 2.1	0.4201
Side			0.4838
Right	21	24	
Left	17	14	
Smoker	4 (10.53)	3 (7.895)	0.4446
Alcohol user	8 (21.05)	7 (18.42)	0.8186
Mean follow-up time in mo	26.9 ± 5.9	27.5 ± 6.3	0.6695
Operation time in min	88.3 ± 15.7	93.4 ± 16.2	0.1676
Length of stay in d	6.9 ± 1.9	6.5 ± 1.6	0.3241
Meniscus injury	14	16	0.6388
Duration of tourniquet use in min	45.9 ± 5.9	43.8 ± 6.2	0.1347
Femur fixation method			0.5558
EndoButton	36	37	
RCI screw	2	1	
Tibial fixation method			0.6422
Screw and washer	21	23	
RCI screw	17	15	
Comorbidity			0.5488
Diabetes	2	3	
Hypertension	2	2	
Cancers	1	0	

Measurement data are expressed as the mean ± standard deviation. ABSI: A body shape index; RCI: Rounded cannulated interference.

having a high percentage of males. In terms of types of athletes, soccer players had the most with 37, followed by basketball players with 28 and other athletes with 9. There were no significant differences between the High-ABSI and Low-ABS groups in terms of BMI (24.3 ± 2.2 *vs* 23.9 ± 2.1), percentage of smoking (10.53% *vs* 7.895%), and percentage of alcohol consumption (21.05% *vs* 18.42%) for individuals (all $P > 0.05$). In terms of surgery, although the High-ABSI group had a lower operative time than the Low-ABSI group, there was no significant difference between them ($[88.3 \pm 15.7]$ *vs* $[93.4 \pm 16.2]$; $P = 0.1676$).

There was no significant difference in the duration of the tourniquet ($P = 0.1347$) or the number of days in the hospital ($P = 0.3241$) between the two groups. Also, there were no significant differences in the femur ($P = 0.5558$) or tibia ($P = 0.6422$) in terms of fixation method.

Relationship between ABSI levels and prognosis of knee function in athletes after ACLR surgery

Next, the effects of ABSI level and postoperative knee function were explored. **Figure 2** shows the




Figure 1 Flow diagram of the screening and selection process in this study. ABSI: A body shape index; PCL: Posterior cruciate ligament; ACLR: Anterior cruciate ligament reconstruction.




Figure 2 Comparison of knee joint function between High-a body shape index group and Low-a body shape index group after anterior cruciate ligament reconstruction. A: Range of motion (ROM) scores; B: Quadriceps atrophy index; C: Lysholm scores; D: Return to sport (RTS) completion rate. Data are presented as the mean ± standard deviation; ^a*P* < 0.05, ^b*P* < 0.01 vs Low-a body shape index (ABSI) group.

functional recovery of the knee joint after surgery in the different ABSI groups. The results showed that the ROM scores of the two groups were similar before surgery (0 mo) and gradually improved with ongoing surgery and postoperative rehabilitation exercises (Figure 2A). Besides, the ROM scores in the

Low-ABSI group were higher than those in the High-ABSI group ($P < 0.05$; **Figure 2A**), indicating that high ABSI predicted poor knee mobility. The quadriceps muscle also influenced the knee flexion and extension function. The results showed that preoperatively (0 mo), both groups had some short-term atrophy of the quadriceps muscle due to a sudden decrease in motion caused by the ACL rupture (**Figure 2B**). At 3 mo, 12 mo, and 24 mo postoperatively, the quadriceps atrophy index was higher in the High-ABSI group than in the Low-ABSI group (**Figure 2B**). This indicates that the higher the ABSI, the lower the muscle strength of the quadriceps. In addition, the Lysholm score is an excellent indicator of knee function after knee ligament injury surgery. As shown in **Figure 2C**, before surgery, the scores of both groups were low ($P > 0.05$). With ongoing surgery and rehabilitation exercises, the Lysholm scores of both groups gradually increased, and the Lysholm scores of the Low-ABSI group were consistently higher than those of the High-ABSI group ($P < 0.05$, **Figure 2C**).

The KOOS score is a comprehensive self-rating measure of postoperative knee surgery outcomes that effectively reflects patients' perceptions of their knee health, symptoms, and function. The KOOS score consists of five subscales: symptoms, pain, ADL, sports, and QOL. As shown in **Figure 3**, the KOOS scores gradually increased with the ongoing surgery and rehabilitation in each of the five self-scales. At 3 mo postoperatively, the scores in both groups improved rapidly, but there was no significant difference in symptoms, pain, or ADL between the two. This might be attributed to the rapid relief of knee symptoms and pain with the implementation of surgery ($P > 0.05$, **Figure 3**). At the last follow-up (24 mo after surgery), all KOOS scores reached their highest values in both groups, and the Low-ABSI group had higher scores than the High-ABSI group ($P < 0.05$; **Figure 3**).

High levels of ASBI are a risk factor for poorer functional recovery of the knee after ACLR surgery in athletes

To determine whether a high level of ASBI is a risk factor for poorer postoperative knee function recovery in ACLR athletes, we performed multifactorial logistic regression analysis with ABSI as the independent variable and each knee function score at the last follow-up as the dependent variable. We turned these continuous variables into dichotomous variables. The median scores of ROM scores, quadriceps atrophy index, Lysholm scores, and KOOS (symptoms, pain, ADL, sports, and QOL) were taken for all groups, and scores greater than the median value were defined as low knee function, specifically low ROM scores (≤ 82), low quadriceps atrophy index (≥ 1.2), low Lysholm scores (≤ 83), low symptoms (≤ 88), low pain (≤ 83), low ADL (≤ 87), low sports (≤ 80), and low QOL (≤ 73).

As shown in **Table 2**, the results revealed that high ABSI was a risk factor for patients with low postoperative knee function scores, specifically low ROM scores (odds ratio [OR] = 1.31, 95% confidence interval [CI] [1.10-1.44]; $P < 0.001$), low quadriceps atrophy index (OR = 1.11, 95% CI [0.97- 1.29]; $P < 0.05$), low Lysholm scores (OR = 2.34, 95% CI [1.78-2.94]; $P < 0.001$), low symptoms (OR = 1.14, 95% CI [1.02-1.34]; $P < 0.05$), low ADL (OR = 1.34, 95% CI [1.18-1.65]; $P < 0.05$), low sports (OR = 2.47, 95% CI [1.78-2.84]; $P < 0.001$), and low QOL (OR = 3.34, 95% CI [2.88-3.94]; $P < 0.001$). Interestingly, while pain scores (higher scores were associated with less pain) were higher in the High-ABSI group than in the Low-ABSI group ($P < 0.05$; **Figure 2C**), multifactorial logistic regression showed that high ABSI was not a risk factor for pain (OR = 1.04, 95% CI [0.78-1.44]; $P = 0.06$) (**Table 2**). Ultimately, at the final follow-up, the percentage of RTS was significantly higher in the Low-ABSI group (78.95% vs 60.53%; $P < 0.05$) than in the High-ABSI group (**Figure 2D**).

Association between ASBI levels and postoperative complications in ACLR athletes

As shown in **Table 3**, with the exception that the incidence of venous thrombosis of lower limbs (21.04% vs 2.631%; $P = 0.0284$) in the High-ABSI group was higher than that in the Low-ABSI group, there was no significant difference in the incidence of knee joint infection (5.262% vs 5.262%; $P > 0.9999$), internal fixation failure (2.631% vs 0.000%; $P > 0.9999$), and ACL current failure (7.893% vs 2.631%; $P = 0.6148$) between the two groups. Further multifactorial logistic regression analysis also showed that ABSI was a risk factor for deep vein thrombosis of the lower limb (OR = 2.14, 95% CI [1.88-2.36]; $P < 0.05$) as well as ACL recurrent rupture (OR = 1.24, 95% CI [0.98- 1.44]; $P < 0.05$) (**Table 4**).

DISCUSSION

The current standardized treatment for ACL injury is ACLR, which aims to restore the function and stability of the knee joint, thus promoting RTS[23]. Although most ACL reconstructions restore the mechanical stability of the injured knee joint, the incidence of RTS is different. In 85%-90% of ACLR patients, the prognosis of knee joint function returns to normal or close to normal within 6 mo after surgery[24]. There are also some research reports that only 63% of the people restore their exercise level before the injury at the last follow-up[25].

Our study also showed that the knee joint function scores of most patients with ACLR remained stable from 1 year to 2 years after surgery. In the last follow-up 24 mo after the operation, athletes in the High-ABSI and Low-ABSI groups had 60.53% (23/38) and 78.95% (30/38) RTS completion rates, respectively. This confirms that ACLR is undoubtedly the standard operation to reconstruct the

Table 2 Multifactor logistic regression of low knee joint function score in anterior cruciate ligament reconstruction patients with low a body shape index on last follow-up

Items	OR (95%CI)	P value
Low ROM scores	1.31 (1.10-1.44) ^b	< 0.001
Low quadriceps atrophy index	1.11 (0.97-1.29) ^a	< 0.05
Low Lysholm scores	2.34 (1.78-2.94) ^b	< 0.001
Low KOOS scores		
Low symptoms	1.14 (1.02-1.34) ^a	< 0.05
Low pain	1.04 (0.78-1.44)	0.06
Low ADL	1.34 (1.18-1.65) ^a	< 0.05
Low sports	2.47 (1.78-2.84) ^b	< 0.001
Low QOL	3.34 (2.88-3.94) ^b	< 0.001

^a*P* < 0.05.^b*P* < 0.001.

Compared with the High-a body shape index group.

ABSI: A body shape index; ADL: Activity of daily living; CI: Confidence interval; KOOS: Knee Injury and Osteoarthritis Outcome Score; OR: Odds ratio; QOL: Quality of life; ROM: Range of motion.

Table 3 Postoperative complications in the High-a body shape index group and Low-a body shape index group after anterior cruciate ligament reconstruction, *n* (%)

Characteristic	High-ABSI group, <i>n</i> = 38	Low-ABSI group, <i>n</i> = 38	P value
Knee joint infection	2 (5.262)	2 (5.262)	> 0.9999
Internal fixation failure	1 (2.631)	0 (0.000)	> 0.9999
Deep vein thrombosis of lower limb	8 (21.04)	1 (2.631) ^a	0.0284
ACL recurrent rupture	3 (7.893)	1 (2.631)	0.6148
Total	14 (36.84)	4 (10.53)	0.0571

^a*P* < 0.05.

Compared with the Low-a body shape index group.

ABSI: A body shape index; ACL: Anterior cruciate ligament.

Table 4 Multifactor logistic regression of complications in anterior cruciate ligament reconstruction patients with high a body shape index on last follow-up

Items	OR (95%CI)	P value
Knee joint infection	1.00 (1.10-1.44)	> 0.05
Internal fixation failure	1.01 (0.93-1.15)	> 0.05
Deep vein thrombosis of lower limb	2.14 (1.88-2.36) ^a	< 0.05
ACL recurrent rupture	1.24 (0.98-1.44) ^a	< 0.05


^a*P* < 0.05.

Compared with the Low-a body shape index group.

ACL: Anterior cruciate ligament; CI: Confidence interval; OR: Odds ratio.

integrity of the anterior cruciate ligament and restore the function of the knee joint.

Obesity is also a risk factor for poor function after various knee surgeries[26]. Li *et al*[27] found that obesity and medial cartilage injury are strong risk factors for osteoarthritis after the first single-beam ACLR. Another prospective study involving 92 patients with ACLR also pointed out that the incidence of postoperative complications in patients with high BMI (BMI > 25 kg/m²) was slightly higher than



DOI: 10.12998/wjcc.v11.i19.4567 Copyright ©The Author(s) 2023.

Figure 3 Comparison of Knee Injury and Osteoarthritis Outcome Score between the High-a body shape index group and Low-a body shape index group after anterior cruciate ligament reconstruction. A: Symptoms; B: Pain; C: Activity of daily living (ADL); D: Sports; E: Quality of life (QOL). Data are presented as the mean \pm standard deviation; $^aP < 0.05$, $^bP < 0.01$ vs Low-a body shape index group (ABSI).

that in patients with normal BMI[21]. For obese patients with ACLR with BMI higher than 25 kg/m², the score of the International Knee Documentation Committee was lower after the operation[28]. However, there were also some different results. In addition, another study also confirmed that the preoperative BMI of patients had no significant adverse impact on the KOOS and Lysholm scores, and that there was no significant difference in the postoperative clinical results of these patients[21]. Therefore, the relationship between obesity based on BMI content and the postoperative function of ACLR is complex, because BMI may not be able to effectively measure the content of individual fat and identify muscle and obesity. ABSI has a positive correlation with visceral fat and is not affected by muscle. Thus, it can better display the content of fat than the traditional BMI.

The study also confirmed that the knee joint function scores of the High-ABSI population were lower than those of the low-ABSI population. Multivariate logistic analysis also demonstrated that high ABSI was the influencing factor of poor function after ACLR. Similar to a previous study[24], there was no significant difference between the two groups in various functional scores (ROM compliance rate, quadriceps femoris atrophy index, Lysholm score, *etc*) before surgery ($P > 0.05$). With the progress of postoperative rehabilitation, the ROM scores, quadriceps femoris atrophy index, and Lysholm score of the two groups were significantly improved compared with those before surgery ($P < 0.05$). Subsequently, it was gradually revealed that the knee joint function of the High-ABSI group was still lower than that of the Low-ABSI group. At the 6th mo and 12th mo after operation, the knee joint function score of the Low-ABSI group was significantly lower than that of the Low-ABSI group. These results also confirmed the conclusion of previous research[4] that obese people would experience increased load on the lower limbs, thus affecting their recovery of knee joints. Besides, the risk of lower limb deep venous thrombosis after operation in the High-ABSI population was also higher than that in the Low-ABSI population, since obesity is a risk factor for deep venous thrombosis (OR = 1.67, 95%CI: 1.16-2.40; $P = 0.006$), and Mendel randomization verified the causal relationship between BMI and deep vein thrombosis[29].

Fat plays an important role in the formation of deep venous thrombosis of lower limbs. Studies have shown that lipid metabolism in adipocytes participates in the process of thrombosis, and adipocytes can promote thrombosis by releasing a variety of thrombogenic factors, such as coagulation factors, platelet activating factors, platelet-derived active substances, *etc*[30]. Therefore, in the process of athletes' rehabilitation, it is necessary to take measures to reduce obesity, thus reducing the incidence rate of deep vein thrombosis.

Our investigation had some limitations. This was a retrospective study with a limited number of participants and results that only covered a brief period of time. In addition, this study is part of a larger correlation investigation. To investigate the possible causal connection between ABSI and the postoperative function of ACLR in the future, additional randomized controlled trials or Mendelian randomization will be required.

CONCLUSION

ABSI is a risk factor for poor prognosis of knee function in ACL athletes after ACLR, and the risk of poor knee function after ACLR, deep vein thrombosis of the lower limb, and ACL recurrent rupture gradually increases with the rise of ABSI.

ARTICLE HIGHLIGHTS

Research background

At present, the indicators of individual obesity only rely on body mass index (BMI) index, but BMI is often not linearly related to body fat content, which limits the research on the association between obesity and other diseases.

Research motivation

This study introduced a body shape index (ABSI), which is a body type indicator to replace traditional BMI to objectively evaluate the association between athletes' body size/obesity and anterior cruciate ligament reconstruction (ACLR).

Research objectives

Explore the relationship between knee joint function in athletes with ABSI and anterior cruciate ligament injuries after ACLR.

Research methods

Multiple logistic regression analysis was used to investigate the relationship between knee joint function scores and postoperative complications after ABSI and ACLR surgery.

Research results

The knee joint function score of the Low-ABSI group was higher than that of the High-ABSI group ($P < 0.05$). High ABSI is a risk factor for low score of knee joint function after operation, and also a risk factor for deep vein thrombosis of lower limbs.

Research conclusions

ABSI is closely related to the prognosis of knee joint function after ACLR. The rise of ABSI is likely to lead to poor knee function after ACLR and deep vein thrombosis of lower limbs.

Research perspectives

In the future, randomized controlled trials or Mendelian randomization are needed to verify the possible causal relationship between ABSI and postoperative function of ACLR.

FOOTNOTES

Author contributions: Wang YJ carried out the acquisition and interpretation of data and was the major contributor to drafting the manuscript; Wang YJ and Zhang JC carried out the clinical partial data collection and analyses; Wang YJ and Zhang YZ participated in drawing tables and diagrams; Zhang YZ was responsible for correcting the language and grammar; Zhang JC was responsible for reviewing and revising some drawings and tables; Zhang JC and Liu YH were responsible for guiding clinical discussions; Liu YH contributed to the ideas of the article and reviewed the manuscript; All authors provided final approval for publishing the manuscript.

Institutional review board statement: According to the Helsinki Declaration, the study was approved by the Ethics Committee of First Hospital of Shanxi Medical University.

Informed consent statement: The data used in this study were not involved in the patients' privacy information, so the informed consent was waived by the Ethics Committee of First Hospital of Shanxi Medical University. All patient data obtained, recorded, and managed only used for this study, and all patient information are strictly confidential, without any harm to the patient.

Conflict-of-interest statement: The authors have no conflicts of interest to declare.

Data sharing statement: The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by

external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Ying-Hai Liu [0009-0004-3105-1393](https://orcid.org/0009-0004-3105-1393).

S-Editor: Liu JH

L-Editor: Filipodia

P-Editor: Liu JH

REFERENCES

- 1 **van Yperen DT**, Reijman M, van Es EM, Bierma-Zeinstra SMA, Meuffels DE. Twenty-Year Follow-up Study Comparing Operative Versus Nonoperative Treatment of Anterior Cruciate Ligament Ruptures in High-Level Athletes. *Am J Sports Med* 2018; **46**: 1129-1136 [PMID: [29438635](#) DOI: [10.1177/0363546517751683](#)]
- 2 **Wiggins AJ**, Grandhi RK, Schneider DK, Stanfield D, Webster KE, Myer GD. Risk of Secondary Injury in Younger Athletes After Anterior Cruciate Ligament Reconstruction: A Systematic Review and Meta-analysis. *Am J Sports Med* 2016; **44**: 1861-1876 [PMID: [26772611](#) DOI: [10.1177/0363546515621554](#)]
- 3 **Wilk KE**, Arrigo CA. Rehabilitation Principles of the Anterior Cruciate Ligament Reconstructed Knee: Twelve Steps for Successful Progression and Return to Play. *Clin Sports Med* 2017; **36**: 189-232 [PMID: [27871658](#) DOI: [10.1016/j.csm.2016.08.012](#)]
- 4 **Meyer NL**, Sundgot-Borgen J, Lohman TG, Ackland TR, Stewart AD, Maughan RJ, Smith S, Müller W. Body composition for health and performance: a survey of body composition assessment practice carried out by the Ad Hoc Research Working Group on Body Composition, Health and Performance under the auspices of the IOC Medical Commission. *Br J Sports Med* 2013; **47**: 1044-1053 [PMID: [24065075](#) DOI: [10.1136/bjsports-2013-092561](#)]
- 5 **Ackland TR**, Lohman TG, Sundgot-Borgen J, Maughan RJ, Meyer NL, Stewart AD, Müller W. Current status of body composition assessment in sport: review and position statement on behalf of the ad hoc research working group on body composition health and performance, under the auspices of the I.O.C. Medical Commission. *Sports Med* 2012; **42**: 227-249 [PMID: [22303996](#) DOI: [10.2165/11597140-000000000-00000](#)]
- 6 **Luc B**, Gribble PA, Pietrosimone BG. Osteoarthritis prevalence following anterior cruciate ligament reconstruction: a systematic review and numbers-needed-to-treat analysis. *J Athl Train* 2014; **49**: 806-819 [PMID: [25232663](#) DOI: [10.4085/1062-6050-49.3.35](#)]
- 7 **Barenius B**, Ponzer S, Shalabi A, Bujak R, Norlén L, Eriksson K. Increased risk of osteoarthritis after anterior cruciate ligament reconstruction: a 14-year follow-up study of a randomized controlled trial. *Am J Sports Med* 2014; **42**: 1049-1057 [PMID: [24644301](#) DOI: [10.1177/0363546514526139](#)]
- 8 **Messier SP**, Beavers DP, Loeser RF, Carr JJ, Khajanchi S, Legault C, Nicklas BJ, Hunter DJ, Devita P. Knee joint loading in knee osteoarthritis: influence of abdominal and thigh fat. *Med Sci Sports Exerc* 2014; **46**: 1677-1683 [PMID: [25133996](#) DOI: [10.1249/MSS.0000000000000293](#)]
- 9 **Brambilla L**, Pulici L, Carimati G, Quaglia A, Prospero E, Bait C, Morengi E, Portinaro N, Denti M, Volpi P. Prevalence of Associated Lesions in Anterior Cruciate Ligament Reconstruction: Correlation With Surgical Timing and With Patient Age, Sex, and Body Mass Index. *Am J Sports Med* 2015; **43**: 2966-2973 [PMID: [26473010](#) DOI: [10.1177/0363546515608483](#)]
- 10 **Caballero B**. Humans against Obesity: Who Will Win? *Adv Nutr* 2019; **10**: S4-S9 [PMID: [30721956](#) DOI: [10.1093/advances/nmy055](#)]
- 11 **Roche AF**, Sievogel RM, Chumlea WC, Webb P. Grading body fatness from limited anthropometric data. *Am J Clin Nutr* 1981; **34**: 2831-2838 [PMID: [7315784](#) DOI: [10.1093/ajcn/34.12.2831](#)]
- 12 **Chen J**, Guan Z, Wang L, Song G, Ma B, Wang Y. Meta-analysis: overweight, obesity, and Parkinson's disease. *Int J Endocrinol* 2014; **2014**: 203930 [PMID: [24672544](#) DOI: [10.1155/2014/203930](#)]
- 13 **Prentice AM**, Jebb SA. Beyond body mass index. *Obes Rev* 2001; **2**: 141-147 [PMID: [12120099](#) DOI: [10.1046/j.1467-789x.2001.00031.x](#)]
- 14 **Krakauer NY**, Krakauer JC. A new body shape index predicts mortality hazard independently of body mass index. *PLoS One* 2012; **7**: e39504 [PMID: [22815707](#) DOI: [10.1371/journal.pone.0039504](#)]
- 15 **Christakoudi S**, Tsilidis KK, Muller DC, Freisling H, Weiderpass E, Overvad K, Söderberg S, Häggström C, Pischon T, Dahm CC, Zhang J, Tjønneland A, Halkjaer J, MacDonald C, Boutron-Ruault MC, Mancini FR, Kühn T, Kaaks R, Schulze MB, Trichopoulos A, Karakatsani A, Peppas E, Masala G, Pala V, Panico S, Tumino R, Sacerdote C, Quirós JR, Agudo A, Sánchez MJ, Cirera L, Barricarte-Gurrea A, Amiano P, Memarian E, Sonestedt E, Bueno-de-Mesquita B, May AM, Khaw KT, Wareham NJ, Tong TYN, Huybrechts I, Noh H, Aglago EK, Ellingjord-Dale M, Ward HA, Aune D, Riboli E. A Body Shape Index (ABSI) achieves better mortality risk stratification than alternative indices of abdominal obesity: results from a large European cohort. *Sci Rep* 2020; **10**: 14541 [PMID: [32883969](#) DOI: [10.1038/s41598-020-71302-5](#)]
- 16 **Gažárová M**, Gašneiderová M, Mečiarová L. Obesity diagnosis and mortality risk based on a body shape index (ABSI) and other indices and anthropometric parameters in university students. *Rocz Pansw Zakl Hig* 2019; **70**: 267-275 [PMID: [31515986](#) DOI: [10.32394/rpzh.2019.0077](#)]
- 17 **Christakoudi S**, Tsilidis KK, Evangelou E, Riboli E. A Body Shape Index (ABSI), hip index, and risk of cancer in the UK

- Biobank cohort. *Cancer Med* 2021; **10**: 5614-5628 [PMID: [34196490](#) DOI: [10.1002/cam4.4097](#)]
- 18 **Patel NK**, Sabharwal S, Hadley C, Blanchard E, Church S. Factors affecting return to sport following hamstrings anterior cruciate ligament reconstruction in non-elite athletes. *Eur J Orthop Surg Traumatol* 2019; **29**: 1771-1779 [PMID: [31321592](#) DOI: [10.1007/s00590-019-02494-4](#)]
- 19 **Mizuno T**. Effects of Dynamic Stretching Velocity on Joint Range of Motion, Muscle Strength, and Subjective Fatigue. *J Strength Cond Res* 2022; **36**: 2440-2447 [PMID: [33009348](#) DOI: [10.1519/JSC.0000000000003842](#)]
- 20 **Peck BD**, Brightwell CR, Johnson DL, Ireland ML, Noehren B, Fry CS. Anterior Cruciate Ligament Tear Promotes Skeletal Muscle Myostatin Expression, Fibrogenic Cell Expansion, and a Decline in Muscle Quality. *Am J Sports Med* 2019; **47**: 1385-1395 [PMID: [30995070](#) DOI: [10.1177/0363546519832864](#)]
- 21 **Ballal MS**, Khan Y, Hastie G, Hatcher A, Coogan S, McNicholas MJ. Functional outcome of primary hamstring anterior cruciate ligament reconstruction in patients with different body mass index classes. *Arthroscopy* 2013; **29**: 1314-1321 [PMID: [23830220](#) DOI: [10.1016/j.arthro.2013.05.005](#)]
- 22 **Collins NJ**, Prinsen CA, Christensen R, Bartels EM, Terwee CB, Roos EM. Knee Injury and Osteoarthritis Outcome Score (KOOS): systematic review and meta-analysis of measurement properties. *Osteoarthritis Cartilage* 2016; **24**: 1317-1329 [PMID: [27012756](#) DOI: [10.1016/j.joca.2016.03.010](#)]
- 23 **Czuppon S**, Racette BA, Klein SE, Harris-Hayes M. Variables associated with return to sport following anterior cruciate ligament reconstruction: a systematic review. *Br J Sports Med* 2014; **48**: 356-364 [PMID: [24124040](#) DOI: [10.1136/bjsports-2012-091786](#)]
- 24 **Zaffagnini S**, Grassi A, Marcheggiani Muccioli GM, Tsapralis K, Ricci M, Braganzoni L, Della Villa S, Marcacci M. Return to sport after anterior cruciate ligament reconstruction in professional soccer players. *Knee* 2014; **21**: 731-735 [PMID: [24593869](#) DOI: [10.1016/j.knee.2014.02.005](#)]
- 25 **Ardern CL**, Webster KE, Taylor NF, Feller JA. Return to sport following anterior cruciate ligament reconstruction surgery: a systematic review and meta-analysis of the state of play. *Br J Sports Med* 2011; **45**: 596-606 [PMID: [21398310](#) DOI: [10.1136/bjsm.2010.076364](#)]
- 26 **Mellion KM**, Grover BT. Obesity, Bariatric Surgery, and Hip/Knee Arthroplasty Outcomes. *Surg Clin North Am* 2021; **101**: 295-305 [PMID: [33743970](#) DOI: [10.1016/j.suc.2020.12.011](#)]
- 27 **Li RT**, Lorenz S, Xu Y, Harner CD, Fu FH, Irrgang JJ. Predictors of radiographic knee osteoarthritis after anterior cruciate ligament reconstruction. *Am J Sports Med* 2011; **39**: 2595-2603 [PMID: [22021585](#) DOI: [10.1177/0363546511424720](#)]
- 28 **DiSilvestro KJ**, Jauregui JJ, Glazier E, Cherkalin D, Bennett CH, Packer JD, Henn RF 3rd. Outcomes of Anterior Cruciate Ligament Reconstruction in Obese and Overweight Patients: A Systematic Review. *Clin J Sport Med* 2019; **29**: 257-261 [PMID: [31241525](#) DOI: [10.1097/JSM.0000000000000521](#)]
- 29 **Tan JS**, Liu NN, Guo TT, Hu S, Hua L. Genetically predicted obesity and risk of deep vein thrombosis. *Thromb Res* 2021; **207**: 16-24 [PMID: [34507265](#) DOI: [10.1016/j.thromres.2021.08.026](#)]
- 30 **Rothberg DL**, Makarewich CA. Fat Embolism and Fat Embolism Syndrome. *J Am Acad Orthop Surg* 2019; **27**: e346-e355 [PMID: [30958807](#) DOI: [10.5435/JAAOS-D-17-00571](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

