

World Journal of *Clinical Cases*

World J Clin Cases 2023 July 26; 11(21): 4966-5192



Contents

Thrice Monthly Volume 11 Number 21 July 26, 2023

OPINION REVIEW

- 4966 Flexner has fallen: Transitions in medical education system across time, a gradual return to pre-Flexnerian state (de-Flexnerization)
Alzerwi NAN

REVIEW

- 4975 Manipulation of the intestinal microbiome-a slow journey to primetime
Kruger-Sharabi O, Malnick SDH, Fisher D

MINIREVIEWS

- 4989 Menstrual cycle abnormalities in women with inflammatory bowel disease and effects of biological therapy on gynecological pathology
Malinauskiene V, Zuzo A, Liakina V, Kazenaite E, Stundiene I
- 4996 Exhaled volatile organic compounds for diagnosis and monitoring of asthma
Savito L, Scarlata S, Bikov A, Carratù P, Carpagnano GE, Dragonieri S

ORIGINAL ARTICLE

Retrospective Cohort Study

- 5014 Cohort analysis of pediatric intussusception score to diagnose intussusception
Rukwong P, Wangviwat N, Phewplung T, Sintusek P

Retrospective Study

- 5023 Identification of functional genes regulating gastric cancer progression using integrated bioinformatics analysis
Yu K, Zhang D, Yao Q, Pan X, Wang G, Qian HY, Xiao Y, Chen Q, Mei K
- 5035 Clinical outcomes and complications of single anastomosis duodenal-ileal bypass with sleeve gastrectomy: A 2-year follow-up study in Bogotá, Colombia
Ospina Jaramillo A, Riscanevo Bobadilla AC, Espinosa MO, Valencia A, Jiménez H, Montilla Velásquez MDP, Bastidas M
- 5047 Image-based visualization of stents in mechanical thrombectomy for acute ischemic stroke: Preliminary findings from a series of cases
Yao QY, Fu ML, Zhao Q, Zheng XM, Tang K, Cao LM
- 5056 Clinical effects of a traditional Chinese medicine nursing programme to intervene in gastric pain of the spleen and stomach with Qi deficiency
Yu YH, Yu YS, Cheng XH

- 5063** Behavior analysis and formative assessments in online oral medicine education during the COVID-19 pandemic

Ye JJ, Zhao YK, Teng ZS, Ye HW, Yuan Q, Nie X

- 5073** Clinical study of acupuncture combined with surface anesthesia using proparacaine in geriatric cataract phacoemulsification

Dong P, Wang LF, Zhang LX, Li F, Yin HS, Dou ZX, Huang XJ, Xu R, Zhang WL

Clinical Trials Study

- 5083** Safety evaluation of human umbilical cord-mesenchymal stem cells in type 2 diabetes mellitus treatment: A phase 2 clinical trial

Lian XF, Lu DH, Liu HL, Liu YJ, Yang Y, Lin Y, Xie F, Huang CH, Wu HM, Long AM, Hui CJ, Shi Y, Chen Y, Gao YF, Zhang F

Observational Study

- 5097** Time usage analysis and satisfaction comparison by occupational area according to the sex of single-parent families

Jang WH, Jang JS, Bang JH

- 5108** Evaluation of bronchoscopic direct vision glottis anesthesia method in bronchoscopy

Lang J, Guo ZZ, Xing SS, Sun J, Qiu B, Shu Y, Wang ZQ, Liu GX

CASE REPORT

- 5115** Type I Mirizzi syndrome treated by electrohydraulic lithotripsy under the direct view of SpyGlass: A case report

Liang SN, Jia GF, Wu LY, Wang JZ, Fang Z, Wang SH

- 5122** Primary ovarian angiosarcoma: Two case reports and review of literature

Zhou Y, Sun YW, Liu XY, Shen DH

- 5129** Acute myelomonocytic leukemia and T-lymphoblastic lymphoma as simultaneous bilineage hematologic malignancy treated with decitabine: A case report

Jeon SY, Lee NR, Cha S, Yhim HY, Kwak JY, Jang KY, Kim N, Cho YG, Lee CH

- 5136** Systematic review of fibroblastic rheumatism: A case report

Guo H, Liang Q, Dong C, Zhang Q, Gu ZF

- 5147** Improved super-elastic Ti-Ni alloy wire for treating adult skeletal class III with facial asymmetry: A case report

Huang CY, Chen YH, Lin CC, Yu JH

- 5160** Neuroendocrine carcinoma of the endometrium concomitant with Lynch syndrome: A case report

Siu WYS, Hong MK, Ding DC

- 5167** Myeloperoxidase-antineutrophil cytoplasmic antibody-associated vasculitis with headache and kidney involvement at presentation and with arthralgia at relapse: A case report

Zhang X, Zhao GB, Li LK, Wang WD, Lin HL, Yang N

- 5173** Rare combination of traumatic subarachnoid-pleural fistula and intracranial subdural hygromas: A case report
Chen PH, Li CR, Gan CW, Yang TH, Chang CS, Chan FH
- 5179** ADAMTS3 and FLT4 gene mutations result in congenital lymphangiectasia in newborns: A case report
Liang ZW, Gao WL
- 5187** Brucellosis presenting with pancytopenia and hearing loss: A case report
Chen X, Yang FB, Liang JY

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Baharudin Ibrahim, BPharm, PhD, Associate Professor, Pharmacist, Department of Clinical Pharmacy and Pharmacy Practice, Faculty of Pharmacy, Universiti Malaya, Kuala Lumpur 50603, Malaysia. baharudin.ibrahim@um.edu.my

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

July 26, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Clinical Trials Study

Safety evaluation of human umbilical cord-mesenchymal stem cells in type 2 diabetes mellitus treatment: A phase 2 clinical trial

Xiao-Fen Lian, Dong-Hui Lu, Hong-Li Liu, Yan-Jing Liu, Yang Yang, Yuan Lin, Feng Xie, Cai-Hao Huang, Hong-Mei Wu, Ai-Mei Long, Chen-Jun Hui, Yu Shi, Yun Chen, Yun-Feng Gao, Fan Zhang

Specialty type: Medicine, research and experimental

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C
Grade D (Fair): D
Grade E (Poor): 0

P-Reviewer: Popovic DS, Serbia; Triolo F, United States

Received: March 17, 2023

Peer-review started: March 17, 2023

First decision: April 11, 2023

Revised: May 23, 2023

Accepted: June 16, 2023

Article in press: June 16, 2023

Published online: July 26, 2023



Xiao-Fen Lian, Dong-Hui Lu, Hong-Li Liu, Yan-Jing Liu, Yuan Lin, Feng Xie, Cai-Hao Huang, Chen-Jun Hui, Yu Shi, Yun Chen, Yun-Feng Gao, Fan Zhang, Department of Endocrinology, Peking University Shenzhen Hospital, Shenzhen 518000, Guangdong Province, China

Yang Yang, Department of Endocrinology, Huizhou Central People's Hospital, Huizhou 516000, Guangdong Province, China

Hong-Mei Wu, Ai-Mei Long, Department of Endocrinology, Longgang District Central Hospital of Shenzhen, Shenzhen 518000, Guangdong Province, China

Corresponding author: Fan Zhang, MD, Doctor, Department of Endocrinology, Peking University Shenzhen Hospital, LianHua Road, Shenzhen 518000, Guangdong Province, China. bjdxszyynfm@163.com

Abstract

BACKGROUND

Progressive pancreatic β cell dysfunction is a fundamental aspect of the pathology underlying type 2 diabetes mellitus (T2DM). Recently, mesenchymal stem cell (MSC) transplantation has emerged as a new therapeutic method due to its ability to promote the regeneration of pancreatic β cells. However, current studies have focused on its efficacy, and there are few clinical studies on its safety.

AIM

To evaluate the safety of human umbilical cord (hUC)-MSC infusion in T2DM treatment.

METHODS

An open-label and randomized phase 2 clinical trial was designed to evaluate the safety of hUC-MSC transplantation in T2DM in a Class A hospital. Ten patients in the placebo group received acellular saline intravenously once per week for 3 wk. Twenty-four patients in the hUC-MSC group received hUC-MSCs (1×10^6 cells/kg) intravenously once per week for 3 wk. Diabetic clinical symptoms and signs, laboratory findings, and imaging findings were evaluated weekly for the 1st mo and then at weeks 12 and 24 post-treatment.

RESULTS

No serious adverse events were observed during the 24-wk follow-up. Four

patients (16.7%) in the hUC-MSC group experienced transient fever, which occurred within 24 h after the second or third infusion; this did not occur in any patients in the placebo group. One patient from the hUC-MSC group experienced hypoglycemic attacks within 1 mo after transplantation. Significantly lower lymphocyte levels (weeks 2 and 3) and thrombin coagulation time (week 2) were observed in the hUC-MSC group compared to those in the placebo group (all $P < 0.05$). Significantly higher platelet levels (week 3), immunoglobulin levels (weeks 1, 2, 3, and 4), fibrinogen levels (weeks 2 and 3), D-dimer levels (weeks 1, 2, 3, 4, 12, and 24), and neutrophil-to-lymphocyte ratios (weeks 2 and 3) were observed in the hUC-MSC group compared to those in the placebo group (all $P < 0.05$). There were no significant differences between the two groups for tumor markers (alpha-fetoprotein, carcinoembryonic antigen, and carbohydrate antigen 199) or blood fat. No liver damage or other side effects were observed on chest X-ray.

CONCLUSION

Our study suggested that hUC-MSC transplantation has good tolerance and high safety in the treatment of T2DM. It can improve human immunity and inhibit lymphocytes. Coagulation function should be monitored vigilantly for abnormalities.

Key Words: Type 2 diabetes mellitus; Cell transplantation; Human umbilical cord-mesenchymal stem cells; Safety; Lymphocytes; Immunity

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Diabetes mellitus is a major public health problem worldwide. Type 2 diabetes mellitus is regarded as a chronic progressive disease that arises from an impairment in the insulin-sensing mechanisms culminating in insulin resistance. Our article focused on the safety of human umbilical cord mesenchymal stem cell infusion for treating type 2 diabetes mellitus. The results suggested that human umbilical cord mesenchymal stem cell treatment can impact human immunity and inhibit lymphocytes. We should pay attention to its influence on coagulation.

Citation: Lian XF, Lu DH, Liu HL, Liu YJ, Yang Y, Lin Y, Xie F, Huang CH, Wu HM, Long AM, Hui CJ, Shi Y, Chen Y, Gao YF, Zhang F. Safety evaluation of human umbilical cord-mesenchymal stem cells in type 2 diabetes mellitus treatment: A phase 2 clinical trial. *World J Clin Cases* 2023; 11(21): 5083-5096

URL: <https://www.wjgnet.com/2307-8960/full/v11/i21/5083.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i21.5083>

INTRODUCTION

Diabetes mellitus (DM) is a major public health problem worldwide. Type 2 DM (T2DM) is the most common type of diabetes, with adults accounting for 90% of diagnoses[1]. T2DM is regarded as a chronic progressive disease that arises from an impairment in the insulin-sensing mechanisms culminating in insulin resistance. Long-term chronic hyperglycemia can cause multisystem complications, including cardiovascular and cerebrovascular diseases, retinopathy, nephropathy, diabetic foot, *etc.* Although novel medications and diet therapies continue to be developed, none have provided full protection against deterioration of β cell function[2,3]. Meanwhile, many side effects like hypoglycemia, gastrointestinal adverse reactions, heart failure, and atypical fracture have increased with drug treatment[4].

In recent years, mesenchymal stem cell (MSC) therapy has been studied extensively as a novel therapeutic option for diabetes[5,6]. Among the different types of MSCs, those from the human umbilical cord (hUC) have been widely applied in the treatment of different diseases[7]. The hUC-MSCs are a group of more primitive cells derived from neonates and express original stem cell-specific surface markers such as embryonic stem cell stage-specific surface antigen 4 and tumor rejection antigen 1-60. Compared with MSCs derived from other tissues such as bone marrow and fat, the hUC-MSCs have a more abundant content, stronger proliferation ability, and lower immunogenicity[8]. Moreover, hUC-MSCs can be sampled conveniently without damage to the health of the donor, and they do not present any ethical challenges. As such, they are attractive and preferred for clinical applications.

Studies have suggested that hUC-MSCs can promote pancreas regeneration by improving the microenvironment and restoring β cells[9,10]. With the development of hUC-MSC treatment for diabetes and its complications[11-13], the safety of hUC-MSCs is an important concern for clinicians. In practice, the clinical application of hUC-MSCs is well-tolerated (*i.e.*, safe)[14,15]. Participants reportedly suffered from mild symptoms such as fever, dizziness, and vomiting, but no cases of tumor development or death were reported. However, the effect of hUC-MSCs on tumor development remains controversial. While some studies have shown that MSCs can promote tumor progression and metastasis[16], others have suggested that MSCs can suppress tumor proliferation and apoptosis[17]. Moreover, the safety data of hUC-MSCs in diabetes treatment are insufficient. To evaluate the safety and feasibility of hUC-MSC infusion in T2DM, we designed a phase 2 clinical trial. Importantly, this was the first clinical trial of hUC-MSC infusion for T2DM treatment approved by

the China Medical Biotech Association.

MATERIALS AND METHODS

Patients

The enrolled participants were patients admitted to Peking University Shenzhen Hospital (Shenzhen, China) for T2DM, and all provided signed informed consent. The study was conducted according to the Declaration of Helsinki and approved by the Institutional Review Board of Peking University Shenzhen Hospital [IRB Approval No. (2018) 29th]. The inclusion criteria and exclusion criteria, as previously reported[18], were applied thoroughly.

hUC-MSC preparation

The hUC-MSCs were provided by Beike Biotechnology (Shenzhen, China)[19]. The isolation process involved Wharton's jelly, a gelatinous tissue around umbilical vessels, from donated hUCs. First, the primary cells were obtained by tissue block adherent culture method, followed by inoculation with 5000 cells/cm² and harvesting when the fusion degree reached 85%-90%. After continuous expansion, the fourth passage of hUC-MSCs was suspended in a 10% DMSO cryopreservation solution and stored in liquid nitrogen (-196 °C)[20] (Figure 1). In their future use as a cell stock material, the samples were thawed at 37 °C, washed to remove the DMSO cryopreservation solution, and resuspended in a compound electrolyte preservation solution (containing 5% albumin) before testing of the "final frozen product" for clinical applicability and safety (Table 1). The final frozen product that passed all tests was then placed at 2-8 °C and within 1 h was transported by the specialist to the quality control department in the hospital. The quality control department then conducted a second quality test within 3 h after receiving the sample. We followed the standard that any cell fluid applied in clinical use should be tested twice.

Once the samples passed the tests they were sent to the clinical department for intravenous infusion to the patient, which occurred within 2 h after sample receipt. The total time from the recovery of cell fluid to clinical application should be less than 12 h. The contents of the cell fluid quality test include the following: Integrity of the cell fluid package; appearance of the cell fluid (including coloration and whether there are floccules or any other precipitates or foreign matter); number of cells in the cell fluid; cell viability; endotoxin presence; and Gram stain for pathogenic bacteria.

Study design

Treatment was given for a period of 16 wk, as previously reported[18], after which all patients were reassessed. The total of 34 patients who met the inclusion and exclusion criteria were randomized into two groups by random allocation software. The hUC-MSC group received an intravenous dosage of hUC-MSCs (1×10^6 cells/kg) once per week for 3 wk. The control group was given placebo, which consisted of an acellular injection of the compound electrolyte preservation solution (containing 5% albumin but lacking hUC-MSCs).

Follow-up

The follow-up visits were conducted at weeks 1, 2, 3, 4, 12, and 24 after the first infusion (Figure 2). Experience of fever, chest tightness, chest pain, dizziness, and any other clinical symptoms experienced during the treatment were recorded, along with any adverse reactions such as cardiocerebrovascular events and tumor occurrence.

All study participants were monitored by laboratory tests for routine blood parameters, liver function, renal function, blood lipids (*e.g.*, total cholesterol and triglycerides), and coagulation indexes at weeks 1, 2, and 3 after the first infusion. We also conducted tests for diabetes antibody, specific infection indexes [hepatitis B surface antigen, antibodies against hepatitis C virus, combined detection of antigen and antibody of human immunodeficiency virus (HIV), and specific antibody against *Treponema pallidum*], and serum tumor markers [alpha-fetoprotein (AFP), carcinoembryonic antigen (CEA), and carbohydrate antigen 199 (CA199)] as well as electrocardiogram, chest X-ray, and liver ultrasound at baseline and weeks 4, 12, and 24 after the first infusion.

Statistical analysis

All statistical analyses were carried out with SPSS® 25.0 software (IBM Corp, Armonk, NY, United States). Quantitative variables were summarized as median, and categorical variables were summarized numerically. Independent sample Wilcoxon test or χ^2 test was used to assess between-group differences. Differences in proportions were analyzed by two-tailed test. *P* values < 0.05 were regarded as statistically significant.

RESULTS

Participant baseline characteristics and laboratory test findings

A total of 34 patients were included from September 2019 to September 2022 (Figure 3). After randomization, 24 of the patients were included in the treatment ("hUC-MSC") group and 10 were included in the control ("placebo") group. The clinical characteristics and baseline laboratory test findings are shown in Tables 2 and 3, respectively. No significant differences were observed between the two groups.

Table 1 Quality control standards of human umbilical cord-mesenchymal stem cells

Test	Final frozen product	Final infusion product
Visual inspection	NA	Absence of visible, particle
Morphology	Fibroblastic	NA
Viability	≥ 90%	≥ 85%
Cell count	$(4.5-6.0) \times 10^7$	According to clinical needs
Pathogen tests		
Sterility	Negative	Negative
Mycoplasma	Negative	Negative
Endotoxin	< 0.5 EU/mL	< 0.5 EU/mL
Cell surface markers		
CD73	≥ 95%	-
CD90	≥ 95%	-
CD105	≥ 95%	-
CD29	≥ 95%	-
CD34	≤ 2%	-
CD45	≤ 2%	-
CD79a	≤ 2%	-
CD14	≤ 2%	-
HLA-DR	≤ 2%	-

Not available.

Table 2 Baseline characteristics of participants

Variable	All patients <i>n</i> = 34	Placebo group <i>n</i> = 10	hUC-MSC group <i>n</i> = 24	<i>Z</i>	<i>P</i> value
Sex					0.45
Male	28 (82.4)	8 (80.0)	20 (83.3)		
Female	6 (17.6)	2 (20.0)	4 (16.7)		
Age in yr	52 (45.5, 56.0)	49.0 (44.0, 58.5)	52.0 (46.0, 56.0)	-0.19	0.85
Duration of T2DM in yr	10.00 (4.00, 14.00)	7.00 (3.75, 11.75)	10.00 (4.50, 14.00)	-0.97	0.33
BMI in kg/m ²	24.21 (23.34, 26.25)	23.72 (23.00, 26.25)	24.61 (23.25, 26.50)	-0.42	0.68
FPG in mmol/L	8.95 (8.13, 10.14)	8.82 (7.81, 9.97)	9.13 (8.18, 11.91)	-0.42	0.68
HbA1c as %	7.95 (7.40, 8.50)	8.00 (7.28, 8.35)	7.95 (7.45, 8.50)	-0.49	0.70

Data are presented as median (minimum, maximum) or number of cases (percentage). BMI: Body mass index; FPG: Fasting plasma glucose; HbA1c: Glycosylated hemoglobin; hUC-MSC: Human umbilical cord-mesenchymal stem cell; T2DM: Type 2 diabetes mellitus.

Primary safety outcome

Safety was evaluated through occurrence of adverse events (AEs) observed within 24 h after each infusion and each visit, including *via* clinical examinations and measurement of vital signs. No serious complications associated with the hUC-MSC infusion were observed.

Four patients (16.67%) from the hUC-MSC group and none from the placebo group experienced transient fever, which primarily occurred within 24 h after the second and third infusions. There was no statistically significant difference between the two groups (*P* = 0.169) (Figure 4).

Table 3 Baseline laboratory tests of participants

Variable	Placebo group	hUC-MSC group	Z	P value
	n = 10	n = 24		
Routine blood				
WBC	6.48 (5.71, 8.17)	6.87 (5.52, 8.13)	-0.227	0.821
PLT	232.0 (199.5, 253.8)	214.0 (163.0, 258.3)	-0.718	0.473
RBC	5.15 (4.58, 5.46)	4.94 (4.79, 5.45)	-0.076	0.94
NEU	3.80 (2.95, 4.89)	4.20 (2.90, 5.15)	-0.189	0.850
L	2.21 (1.69, 2.44)	2.23 (1.96, 2.61)	-0.454	0.65
MONO	0.45 (0.34, 0.53)	0.36 (0.31, 0.44)	-1.192	0.233
NLR	1.89 (1.41, 2.36)	1.64 (1.43, 2.13)	-0.227	0.821
Liver function				
ALT	21.00 (18.00, 31.25)	25.50 (16.50, 48.25)	-0.606	0.545
AST	20.50 (17.00, 23.00)	21.50 (18.00, 31.75)	-0.872	0.383
γ-GT	27.00 (20.75, 31.00)	34.50 (23.75, 52.00)	-1.703	0.089
TP	73.70 (68.88, 76.93)	71.70 (68.13, 76.78)	-0.529	0.597
GLB	29.25 (24.63, 30.98)	25.75 (23.93, 28.33)	-1.966	0.049
Renal function				
Urea	6.51 (4.68, 7.23)	5.87 (5.38, 6.61)	-0.227	0.821
Cr	65.00 (58.00, 73.00)	67.00 (56.25, 81.00)	-0.227	0.82
eGFR	106.59 (97.44, 111.71)	104.52 (84.42, 111.99)	-0.454	0.65
Blood fat				
TC	4.61 (4.06, 5.38)	3.95 (3.55, 5.40)	-1.247	0.212
TG	1.67 (1.20, 3.04)	1.87 (1.13, 2.79)	-0.094	0.925
LDL-C	2.91 (2.53, 3.29)	2.48 (2.12, 3.72)	-1.077	0.281
HDL-C	1.14 (0.91, 1.33)	1.12 (0.88, 1.46)	-0.019	0.985
Coagulation function				
PT	12.25 (12.00, 12.78)	12.15 (12.00, 12.60)	-0.494	0.621
APTT	34.05 (32.68, 37.60)	34.10 (32.58, 35.45)	-0.284	0.777
TT	17.95 (17.45, 19.00)	18.00 (17.65, 18.90)	-0.265	0.791
FIB	2.95 (2.61, 3.53)	3.07 (2.29, 3.22)	-0.302	0.762
DD	0.22 (0.22, 0.26)	0.22 (0.22, 0.30)	-0.36	0.719
Infection screening				
HBsAg	0.00 (0.00, 0.00)	0.00 (0.00, 20.31)	-1.081	0.28
HCV-Ab	0.05 (0.04, 0.06)	0.09 (0.04, 0.14)	-1.679	0.093
HIV-Ag/ Ab	0.07 (0.06, 0.08)	0.06 (0.05, 0.08)	-0.731	0.465
TP-Ab	0.04 (0.02, 0.06)	0.04 (0.04, 0.05)	-0.776	0.438
Tumor marker				
AFP	2.80 (2.20, 3.85)	1.95 (1.65, 3.10)	-1.494	0.135
CEA	1.85 (1.40, 3.50)	2.15 (1.58, 3.08)	-0.416	0.677
CA199	8.45 (5.20, 17.33)	5.55 (3.35, 15.90)	-0.643	0.52

γ-GT: Gamma-glutamyl transferase; AFP: Alpha-fetoprotein; ALT: Alanine transaminase; APTT: Activated partial thrombokinase time; AST: Aspartate

aminotransferase; CA199: Carbohydrate antigen 199; CEA: Carcinoembryonic antigen; Cr: Serum creatinine; DD: D-dimer; eGFR: Estimated glomerular filtration rate; FIB: Fibrinogen; GLB: Immunoglobulin; HBsAg: Hepatitis B surface antigen; HCV-Ab: Antibodies against hepatitis C virus; HDL-C: High-density lipoprotein cholesterol; HIV-Ag/ Ab: Combined detection of antigen and antibody of human immunodeficiency virus; hUC-MSC: Human umbilical cord mesenchymal stem cell; L: Lymphocyte absolute value; LDL-C: Low-density lipoprotein cholesterol; MONO: Monocyte absolute value; NEU: Neutrophil absolute value; NLR: Neutrophil-to-lymphocyte ratio; PLT: Platelets; PT: Prothrombin time; RBC: Red blood cell (erythrocyte); TC: Total cholesterol; TG: Triglycerides; TP: Total protein; TP-Ab: Specific antibody against *Treponema pallidum*; TT: Thrombin coagulation time; Urea: Urea nitrogen; WBC: White blood cell.




Figure 1 Stem cell manufacturing and quality control processes. FFP: Final frozen product; FIP: Final infusion product; hUC: Human umbilical cord; MSCs: Mesenchymal stem cells; P2: Second passage.




Figure 2 Flow chart of the study procedure. The patients enrolled in the present study received three infusions on days 7, 14, and 21. They participated in four visits, occurring on days 0, 28, 84, and 180. ECG: Electrocardiogram; Hu-MSCs: Human umbilical cord-mesenchymal stem cells.




Figure 3 Flow diagram for patient recruitment. Thirty-four patients were enrolled and completed the entire study procedure. No enrolled patient was excluded from the safety analysis. hUC-MSCs: Human umbilical cord-mesenchymal stem cells.





Figure 4 Fever occurrence within 24 h after infusion. A: Temperatures of the patients within 24 h after the first infusion; B: Temperatures of the patients within 24 h after the second infusion; C: Temperatures of the patients within 24 h after the third infusion. Group I: Placebo group; Group II: Human umbilical cord-mesenchymal stem cell infusion group.



DOI: 10.12998/wjcc.v11.i21.5083 Copyright ©The Author(s) 2023.

Figure 5 Influence of the human umbilical cord-mesenchymal stem cell treatment on routine blood measurements. Measurements for all indicators were taken at baseline and weeks 1, 2, 3, 4, 12, and 24 after the first infusion. A: White blood cell count; B: Neutrophil absolute value; C: Lymphocyte absolute value; D: Monocyte absolute value; E: Platelet level; F: Neutrophil-to-lymphocyte ratio. ^a $P < 0.05$. Group I: Placebo group; Group II: Human umbilical cord-mesenchymal stem cell infusion group; WBC: White blood cell; PLT: Platelet; NLR: Neutrophil-to-lymphocyte ratio.

One patient (4.17%) from the hUC-MSC group and none from the placebo group experienced nocturnal hypoglycemia. The lowest blood glucose recorded for that single patient was 59.4 mg/dL, and the level returned to normal without intervention or food intake. During the follow-up period, that patient reduced their insulin dose and did not experience hypoglycemia again.

Four patients (16.67%) in the hUC-MSC group and none from the placebo group experienced fatigue within 3 d after the first infusion; in each case, it did not affect daily activities or work. In all, the AE was relieved gradually without any intervention.

hUC-MSC infusion decreased lymphocyte levels and increased neutrophil-to-lymphocyte ratio


From the second week, patients in the hUC-MSC group showed a decrease in lymphocyte levels, with a return to the normal range after week 4. The lymphocyte levels in the hUC-MSC group were significantly lower than those in the placebo group at week 2 [1.26 (0.97, 1.87) *vs* 2.26 (2.20, 2.76); $P < 0.05$] and week 3 [1.70 (1.36, 2.13) *vs* 2.40 (2.02, 2.76); $P < 0.05$]. The neutrophil-to-lymphocyte ratio was significantly higher in the hUC-MSC group than in the placebo group at weeks 2 and 3 ($P < 0.05$ for both). The platelet levels were also significantly higher in the hUC-MSC group than in the placebo group at week 3 [243.00 (224.00, 275.25) *vs* 212.00 (178.25, 235.25); $P < 0.05$]. At the follow-up visits at weeks 4, 12, and 24, there were no statistically different findings between the two groups for the aforementioned indicators. There were also no significant differences between the two groups for white blood cell, neutrophil, and monocyte counts (Figure 5).

hUC-MSC infusion affected coagulation markers

D-dimer values were significantly higher in the hUC-MSC group than in the placebo group at all timepoints (weeks 1, 2, 3, 4, 12, and 24; $P < 0.05$ for all). The fibrinogen levels were also significantly higher in the hUC-MSC group than in the placebo group at week 2 [3.18 (2.75, 3.69) *vs* 2.78 (2.46, 2.99); $P < 0.05$] and week 3 [3.37 (3.20, 3.79) *vs* 2.87 (2.61, 2.99); $P < 0.05$]. The thrombin coagulation time was significantly lower in the hUC-MSC group than in the placebo group at week 2 [17.40 (17.00, 18.35) *vs* 18.60 (18.23, 19.10); $P < 0.01$], decreasing and stabilizing to baseline at week 3. The prothrombin time (PT) and activated partial thrombokinase time (APTT) were also monitored during treatment, but there were no significant differences found between the two groups ($P > 0.05$) (Figure 6).

hUC-MSC infusion increased immunoglobulin and did not affect liver function, renal function, or blood lipids

The immunoglobulin levels were significantly higher in the hUC-MSC group than in the placebo group at week 1 [29.60 (22.83, 31.60) *vs* 23.40 (21.98, 27.10); $P < 0.05$], week 2 [27.35 (24.93, 29.50) *vs* 23.75 (21.65, 25.60); $P < 0.05$], week 3 [28.00 (25.55, 29.65) *vs* 23.55 (22.03, 27.95); $P < 0.05$], and week 4 [28.70 (26.10, 32.60) *vs* 25.00 (23.70, 27.13); $P < 0.05$]. The liver function indexes (blood alanine transaminase, aspartate aminotransferase, gamma-glutamyl transferase, and total



DOI: 10.12998/wjcc.v11.i21.5083 Copyright ©The Author(s) 2023.

Figure 6 Influence of human umbilical cord-mesenchymal stem cell treatment on coagulation function. Measurements for all indicators were taken at baseline and weeks 1, 2, 3, 4, 12, and 24 after the first infusion. A: Prothrombin time; B: Activated partial thrombokinase time; C: Thrombin coagulation time; D: Fibrinogen; E: D-dimer. ^a*P* < 0.05. Group I: Placebo group; Group II: Human umbilical cord-mesenchymal stem cell infusion group; PT: Prothrombin time; APTT: Activated partial thrombokinase time; TT: Thrombin coagulation time; FIB: Fibrinogen; D-D: D-dimer.

protein), renal function indexes (serum creatinine, estimated glomerular filtration rate, and urea nitrogen), and blood lipid indexes (triglycerides, total cholesterol, high-density lipoprotein cholesterol, and low-density lipoprotein cholesterol) showed no significant differences between the two groups (*P* > 0.05) (Figure 7).

hUC-MSC infusion had no effect on tumor markers or on special infectious diseases

The AFP, CEA, and CA199 tumor markers, used to evaluate tumor trends, showed no differences between the two groups at any of the timepoints (Figure 8). At baseline we detected hepatitis B surface antigen in 2 patients in the placebo group and 1 patient in the hUC-MSC group. No patients in either group showed positivity for antibodies against hepatitis C virus, combined detection of HIV antigen and antibody, or specific antibody against *Treponema pallidum*. There were also no changes detected for these indicators during any of the follow-up visits after infusion.

Image test outcome

Chest X-ray and liver color ultrasound examinations showed no changes after the infusion at any of the timepoints. In addition, no nodules or tumors were detected in any patient.

DISCUSSION

Currently, MSC-based cellular therapies are considered effective for many diseases, including DM and its complications [11,21,22]. Our previous study showed that intravenous infusion of hUC-MSCs could improve blood glucose, glycated hemoglobin, and islet function in T2DM [18]. However, clinical research on the safety of hUC-MSCs in treating T2DM has been insufficient. The purpose of this study was to evaluate the safety of hUC-MSC intravenous infusion for T2DM patients. The results indicated good tolerability and safety of hUC-MSC infusion in T2DM patients; the potential disadvantageous responses observed were decreased lymphocyte counts, which may relate to escape from immune attack, and an effect on coagulation function, which may lead to thrombotic disease.

After three hUC-MSC infusions, no patients developed a thromboembolic event. Only mild AEs that spontaneously resolved or with minimal intervention (see fever below), were detected in a small number of patients. Generally, this indicates safety (without any acute infusion-related issues, allergic reactions, delayed hypersensitivity, or secondary infections) [23]. Furthermore, no patients experienced a fever after the first infusion, although 4 patients (16.7%) experienced transient fever after the second and third hUC-MSC infusions. The highest temperature recorded was 39.5 °C. The patients' body temperatures returned to normal naturally or gradually after administration of nonsteroidal anti-inflammatory drugs. Some patients also reported fatigue and hypoglycemia after the hUC-MSC infusions. Thus, post-




Figure 7 Influence of the human umbilical cord-mesenchymal stem cell treatment on liver function, renal function, and blood lipids.

Measurements for all indicators were taken at baseline and weeks 1, 2, 3, 4, 12, and 24 after the first infusion. A: Alanine transaminase; B: Aspartate transaminase; C: Gamma-glutamyl transferase; D: Total protein; E: Immunoglobulin; F: Urea nitrogen; G: Serum creatinine; H: Estimated glomerular filtration rate; I: Total cholesterol; J: Triglycerides; K: Low-density lipoprotein cholesterol; L: High-density lipoprotein cholesterol. * $P < 0.05$. Group I: Placebo group; Group II: Human umbilical cord-mesenchymal stem cell infusion group; ALT: Alanine transaminase; AST: Aspartate transaminase; γ -GT: Gamma-glutamyl transferase; TP: Total protein; GLB: Immunoglobulin; UREA: Urea nitrogen; Cr: Serum creatinine; eGFR: Estimated glomerular filtration rate; TC: Total cholesterol; TG: Triglycerides; LDL-C: Low-density lipoprotein cholesterol; HDL-C: High-density lipoprotein cholesterol.

intravenous hUC-MSC infusion immune responses were considered as mild symptoms. No novel acute cardiovascular and cerebrovascular events or tumors/nodules occurred during the follow-up.




Figure 8 Influence of the human umbilical cord-mesenchymal stem cell treatment on tumor markers. Measurements for all indicators were taken at baseline and weeks 4, 12, and 24 after the first infusion. A: Alpha-fetoprotein; B: Carcinoembryonic antigen; C: Carbohydrate antigen 199. Group I: Placebo group; Group II: Human umbilical cord-mesenchymal stem cell infusion group; AFP: Alpha-fetoprotein; CEA: Carcinoembryonic antigen; CA199: Carbohydrate antigen 199.

Lymphocytes are an essential component of the immune system. Peripheral blood lymphocytes primarily include T and B cells, which constitute a significant component of the immune system and can produce specific immune responses. MSCs are reported to be closely related to immune cells, like lymphocytes. Di Nicola *et al*[24] found that bone marrow MSCs can inhibit T lymphocyte proliferation and apoptosis. Thus, reactivation stimulates efficient proliferation, which may be relevant to soluble factor production after MSC infusion. Our study revealed that patients experienced a reduction in peripheral blood lymphocytes after 2 wk of continuous intravenous hUC-MSC infusion and recovered spontaneously after discontinuation. The hUC-MSC therapy did not impact the absolute values of lymphocytes during long-term follow-up, consistent with the findings by Di Nicola *et al*[24].

It has also been reported that MSCs can induce T cell apoptosis *via* the FAS/FASL pathway. Moreover, MSCs are known to regulate immune response intensity and promote regulatory T cell production through various paracrine effects and cell-cell contact. This induced immune tolerance diminishes the adverse impacts of lymphocytes on MSCs[24-27].

The application of embryonic stem cells is currently limited because of related teratoma formation[28]. Although research has shown that MSCs do not induce malignant transformation[29,30] and some studies have even explored the efficacy of MSC therapy on tumors[30], concerns about possible tumorigenic actions remain undeniable. Guan *et al*[23] conducted a 3-year follow-up observational study on the efficacy of hUC-MSC treatment of T2DM, and no MSC-related malignancies occurred. Similarly, we did not observe an elevation in tumor-associated antigens (AFP, CEA, and CA199) in patients treated with MSCs. Additionally, during the follow-up period, no nodules/tumors occurred in the lungs, liver, gallbladder, spleen, nor pancreas. However, our follow-up duration was relatively short, and we plan to extend it out to 3 years to thoroughly investigate potential transplantation-related complications.

To date, the roles of MSCs in coagulation and inflammation have been inadequately explored. It has, however, been shown that MSC production may lead to the release of procoagulant tissue factors (TFs) at different levels during the treatment[31]. TFs are significant determinants of cell product blood compatibility and are crucial regulators for the extrinsic coagulation pathway of cytokines. TFs can trigger negative systemic inflammatory responses under some conditions. MSCs themselves can trigger platelet production and promote platelet activation, posing a risk of thrombosis[32]. In particular, an MSC dose of $> 1 \times 10^6$ cells/kg has been shown to extend PT and APTT and to reduce concentrations of fibrinogen and factor VIII[33]. Small-dose MSC transfusion also affects PT but not APTT, indicating that MSCs mainly activate coagulation *via* TFs and extrinsic coagulation pathways.

Our study revealed that the D-dimer level was significantly elevated following hUC-MSC (1×10^6 cells/kg) intravenous infusion for three consecutive weeks and that elevation persisted to the 24th wk. Continuous transfusion can lead to transient fibrinogen elevation and thrombin coagulation time reduction. The levels recovered after discontinuation of the hUC-MSC treatment, without affecting PT and APTT. We, thus, considered this situation to be correlated with the hUC-MSC dosage and frequency. Previous studies have also found MSC dosage increases to be associated with acute AEs, including microvascular embolization[34,35].

For future clinical application of hUC-MSC treatment for T2DM, we will explore the interactive mechanism between hUC-MSCs and lymphocytes and the effect of cell dose, culturing and passaging techniques, and route of administration on the inflammatory mechanism; these findings may help to reduce and avoid related AEs in these patients. The sample size of the current study was small. At the same time, a small number of participants had delayed visits due to the coronavirus 2019 pandemic, which could have affected our results. We used a single-blind study design, and there may have been deviation during the follow-up. Reducing the error was a priority. Multiple patients were consulted, and follow-up registrants and examiners were single-blinded.

CONCLUSION

Our study demonstrated that hUC-MSCs were well tolerated during T2DM treatment and elicited no serious AEs. Transient fever, hypoglycemia, and fatigue may occur in the short-term, but no long-term AEs were detected.

Lymphocyte levels decreased and inflammatory factors increased after the intravenous infusion of hUC-MSCs. The results of our study are expected to provide a more solid theoretical basis for the continued pursuit of clinical application of MSCs in the treatment of T2DM.

ARTICLE HIGHLIGHTS

Research background

Cellular therapies represent a new opportunity for the treatment of type 2 diabetes mellitus (T2DM) and its complications. However, the safety of human umbilical cord-mesenchymal stem cells (hUC-MSCs) in clinical application has not been fully assessed.

Research motivation

We conducted a trial to evaluate the safety and tolerance of hUC-MSC infusion in T2DM treatment.

Research objectives

We hypothesized that hUC-MSC infusion may cause fevers or nodules, affect inflammatory mediators, and induce hypercoagulability. We conducted the present trial to treat T2DM patients with an hUC-MSC infusion and evaluated the safety of the hUC-MSC therapy.

Research methods

T2DM patients were enrolled and received hUC-MSC (1×10^6 cells/kg) once per week for 3 wk. The safety was assessed by clinical symptoms and signs, laboratory tests, and imaging tests. The laboratory tests included routine blood parameters, coagulation indexes, and liver function, renal function, and tumor markers. Imaging tests included electrocardiogram, chest X-ray, and ultrasound of the liver, bile, spleen, and pancreas.

Research results

During the 24-wk follow-up period, there were no serious adverse events observed. However, a few patients experienced fever, fatigue, and hypoglycemia. The lymphocyte levels were significantly decreased in the hUC-MSC group compared to the placebo group. The D-dimer level, neutrophil-to-lymphocyte ratio, and immunoglobulin level were significantly increased in the hUC-MSC group compared to the placebo group.

Research conclusions

Our study suggests that hUC-MSCs are safe for the treatment of T2DM, with only mild adverse events occurring. hUC-MSC treatment can affect immunity and inhibit lymphocytes. The influence of hUC-MSC on coagulation requires and warrants further research.

Research perspectives

Lymphocyte subsets and the inflammatory mediators of the patients in this study will be extensively followed for further analysis to elucidate the mechanisms of the observed decrease in lymphocyte numbers and the influence on immune function.

ACKNOWLEDGEMENTS

The authors acknowledge the financial support from all the funders. The authors also want to thank all patients and their families who consented to participate in the study.

FOOTNOTES

Author contributions: Zhang F designed the report; Lian XF, Lu DH, Liu HL, Liu YJ, Yang Y, Lin Y, Xie F, Huang CH, Wu HM, Long AM, Hui CJ, Shi Y, and Chen Y collected the patients' clinical data; Hui CJ provided cell technical support; Lian XF analyzed the data and wrote the paper; and all authors read and approved the final version of the manuscript.

Supported by Shenzhen Science and Technology Innovation Committee Projects, No. JCYJ20170816105416349; and Shenzhen High-Level Hospital Construction Fund, Shenzhen Key Medical Discipline Construction Fund, No. SZXK010.

Institutional review board statement: This study was approved by the Ethics Committee of the Ethical Committee of the Peking University Shenzhen Hospital [IRB of Peking University Shenzhen Hospital (2018) 29th].

Clinical trial registration statement: This study is registered in the Chinese Clinical Trial Registry, Registration No. ChiCTR2200057370.

Informed consent statement: The participants were recruited and enrolled from among patients admitted to Peking University Shenzhen Hospital for diabetes mellitus, and all provided written informed consent.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: There are no additional data.

CONSORT 2010 statement: The authors have read the CONSORT 2010 statement, and the manuscript was prepared and revised according to the CONSORT 2010 statement.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Xiao-Fen Lian 0000-0001-7950-8867; Dong-Hui Lu 0000-0001-9172-2989; Hong-Li Liu 0000-0002-2601-2857; Yan-Jing Liu 0000-0003-0489-3769; Yang Yang 0000-0002-1200-8072; Yuan Lin 0000-0003-4623-2096; Feng Xie 0000-0002-6482-272X; Cai-Hao Huang 0000-0002-2116-9724; Hong-Mei Wu 0000-0002-3054-0345; Ai-Mei Long 0000-0002-4173-8445; Yu Shi 0000-0002-3918-0630; Yun Chen 0000-0002-4975-5637; Yun-Feng Gao 0000-0003-3503-7359; Fan Zhang 0000-0001-5147-663X.

S-Editor: Wang JJ

L-Editor: Filipodia

P-Editor: Yu HG

REFERENCES

- 1 **International Diabetes Federation.** IDF diabetes atlas ninth. Dunia: International Diabetes Federation, 2019
- 2 **Levy J, Atkinson AB, Bell PM, McCance DR, Hadden DR.** Beta-cell deterioration determines the onset and rate of progression of secondary dietary failure in type 2 diabetes mellitus: the 10-year follow-up of the Belfast Diet Study. *Diabet Med* 1998; **15**: 290-296 [PMID: 9585393 DOI: 10.1002/(SICI)1096-9136(199804)15:4<290::AID-DIA570>3.0.CO;2-M]
- 3 **U.K. prospective diabetes study 16.** Overview of 6 years' therapy of type II diabetes: a progressive disease. U.K. Prospective Diabetes Study Group. *Diabetes* 1995; **44**: 1249-1258 [PMID: 7589820]
- 4 **DeFronzo RA, Eldor R, Abdul-Ghani M.** Pathophysiologic approach to therapy in patients with newly diagnosed type 2 diabetes. *Diabetes Care* 2013; **36** Suppl 2: S127-S138 [PMID: 23882037 DOI: 10.2337/dcS13-2011]
- 5 **Zhang Y, Chen W, Feng B, Cao H.** The Clinical Efficacy and Safety of Stem Cell Therapy for Diabetes Mellitus: A Systematic Review and Meta-Analysis. *Aging Dis* 2020; **11**: 141-153 [PMID: 32010488 DOI: 10.14336/AD.2019.0421]
- 6 **Kamal MM, Kassem DH.** Therapeutic Potential of Wharton's Jelly Mesenchymal Stem Cells for Diabetes: Achievements and Challenges. *Front Cell Dev Biol* 2020; **8**: 16 [PMID: 32064260 DOI: 10.3389/fcell.2020.00016]
- 7 **Zhang J, Lv S, Liu X, Song B, Shi L.** Umbilical Cord Mesenchymal Stem Cell Treatment for Crohn's Disease: A Randomized Controlled Clinical Trial. *Gut Liver* 2018; **12**: 73-78 [PMID: 28873511 DOI: 10.5009/gnl17035]
- 8 **Mebarki M, Abadie C, Larghero J, Cras A.** Human umbilical cord-derived mesenchymal stem/stromal cells: a promising candidate for the development of advanced therapy medicinal products. *Stem Cell Res Ther* 2021; **12**: 152 [PMID: 33637125 DOI: 10.1186/s13287-021-02222-y]
- 9 **Tsai PJ, Wang HS, Lin GJ, Chou SC, Chu TH, Chuan WT, Lu YJ, Weng ZC, Su CH, Hsieh PS, Sytwu HK, Lin CH, Chen TH, Shyu JF.** Undifferentiated Wharton's Jelly Mesenchymal Stem Cell Transplantation Induces Insulin-Producing Cell Differentiation and Suppression of T-Cell-Mediated Autoimmunity in Nonobese Diabetic Mice. *Cell Transplant* 2015; **24**: 1555-1570 [PMID: 25198179 DOI: 10.3727/096368914X683016]
- 10 **Khatir R, Mazurek S, Petry SF, Linn T.** Mesenchymal stem cells promote pancreatic β -cell regeneration through downregulation of FoxO1 pathway. *Stem Cell Res Ther* 2020; **11**: 497 [PMID: 33239104 DOI: 10.1186/s13287-020-02007-9]
- 11 **Cho J, D'Antuono M, Glicksman M, Wang J, Jonklaas J.** A review of clinical trials: mesenchymal stem cell transplant therapy in type 1 and type 2 diabetes mellitus. *Am J Stem Cells* 2018; **7**: 82-93 [PMID: 30510843]
- 12 **Fiori A, Terlizzi V, Kremer H, Gebauer J, Hammes HP, Harmsen MC, Bieback K.** Mesenchymal stromal/stem cells as potential therapy in diabetic retinopathy. *Immunobiology* 2018; **223**: 729-743 [PMID: 29402461 DOI: 10.1016/j.imbio.2018.01.001]
- 13 **Cao Y, Gang X, Sun C, Wang G.** Mesenchymal Stem Cells Improve Healing of Diabetic Foot Ulcer. *J Diabetes Res* 2017; **2017**: 9328347 [PMID: 28386568 DOI: 10.1155/2017/9328347]
- 14 **Madani S, Larijani B, Keshkar AA, Tootee A.** Safety and efficacy of hematopoietic and mesenchymal stem cell therapy for treatment of T1DM: a systematic review and meta-analysis protocol. *Syst Rev* 2018; **7**: 23 [PMID: 29373983 DOI: 10.1186/s13643-017-0662-9]
- 15 **Lalu MM, McIntyre L, Pugliese C, Fergusson D, Winston BW, Marshall JC, Granton J, Stewart DJ, Canadian Critical Care Trials Group.** Safety of cell therapy with mesenchymal stromal cells (SafeCell): a systematic review and meta-analysis of clinical trials. *PLoS One* 2012; **7**: e47559 [PMID: 23133515 DOI: 10.1371/journal.pone.0047559]
- 16 **Wang W, Zhong W, Yuan J, Yan C, Hu S, Tong Y, Mao Y, Hu T, Zhang B, Song G.** Involvement of Wnt/ β -catenin signaling in the mesenchymal stem cells promote metastatic growth and chemoresistance of cholangiocarcinoma. *Oncotarget* 2015; **6**: 42276-42289 [PMID: 26474277 DOI: 10.18632/oncotarget.5514]

- 17 **Lin HD**, Fong CY, Biswas A, Choolani M, Bongso A. Human Umbilical Cord Wharton's Jelly Stem Cell Conditioned Medium Induces Tumoricidal Effects on Lymphoma Cells Through Hydrogen Peroxide Mediation. *J Cell Biochem* 2016; **117**: 2045-2055 [PMID: [27392313](#) DOI: [10.1002/jcb.25501](#)]
- 18 **Lian XF**, Lu DH, Liu HL, Liu YJ, Han XQ, Yang Y, Lin Y, Zeng QX, Huang ZJ, Xie F, Huang CH, Wu HM, Long AM, Deng LP, Zhang F. Effectiveness and safety of human umbilical cord-mesenchymal stem cells for treating type 2 diabetes mellitus. *World J Diabetes* 2022; **13**: 877-887 [PMID: [36312002](#) DOI: [10.4239/wjd.v13.i10.877](#)]
- 19 **Gu J**, Huang L, Zhang C, Wang Y, Zhang R, Tu Z, Wang H, Zhou X, Xiao Z, Liu Z, Hu X, Ke Z, Wang D, Liu L. Therapeutic evidence of umbilical cord-derived mesenchymal stem cell transplantation for cerebral palsy: a randomized, controlled trial. *Stem Cell Res Ther* 2020; **11**: 43 [PMID: [32014055](#) DOI: [10.1186/s13287-019-1545-x](#)]
- 20 **Dominici M**, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini F, Krause D, Deans R, Keating A, Prockop Dj, Horwitz E. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy* 2006; **8**: 315-317 [PMID: [16923606](#) DOI: [10.1080/14653240600855905](#)]
- 21 **Zakrzewski W**, Dobrzyński M, Szymonowicz M, Rybak Z. Stem cells: past, present, and future. *Stem Cell Res Ther* 2019; **10**: 68 [PMID: [30808416](#) DOI: [10.1186/s13287-019-1165-5](#)]
- 22 **Nagaishi K**, Mizue Y, Chikenji T, Otani M, Nakano M, Konari N, Fujimiya M. Mesenchymal stem cell therapy ameliorates diabetic nephropathy via the paracrine effect of renal trophic factors including exosomes. *Sci Rep* 2016; **6**: 34842 [PMID: [27721418](#) DOI: [10.1038/srep34842](#)]
- 23 **Guan LX**, Guan H, Li HB, Ren CA, Liu L, Chu JJ, Dai LJ. Therapeutic efficacy of umbilical cord-derived mesenchymal stem cells in patients with type 2 diabetes. *Exp Ther Med* 2015; **9**: 1623-1630 [PMID: [26136869](#) DOI: [10.3892/etm.2015.2339](#)]
- 24 **Di Nicola M**, Carlo-Stella C, Magni M, Milanese M, Longoni PD, Matteucci P, Grisanti S, Gianni AM. Human bone marrow stromal cells suppress T-lymphocyte proliferation induced by cellular or nonspecific mitogenic stimuli. *Blood* 2002; **99**: 3838-3843 [PMID: [11986244](#) DOI: [10.1182/blood.v99.10.3838](#)]
- 25 **Wang L**, Zhao Y, Shi S. Interplay between mesenchymal stem cells and lymphocytes: implications for immunotherapy and tissue regeneration. *J Dent Res* 2012; **91**: 1003-1010 [PMID: [22988011](#) DOI: [10.1177/0022034512460404](#)]
- 26 **Jiang XX**, Zhang Y, Liu B, Zhang SX, Wu Y, Yu XD, Mao N. Human mesenchymal stem cells inhibit differentiation and function of monocyte-derived dendritic cells. *Blood* 2005; **105**: 4120-4126 [PMID: [15692068](#) DOI: [10.1182/blood-2004-02-0586](#)]
- 27 **Yu C**, Tang W, Lu R, Tao Y, Ren T, Gao Y. Human Adipose-derived mesenchymal stem cells promote lymphocyte apoptosis and alleviate atherosclerosis via miR-125b-1-3p/BCL11B signal axis. *Ann Palliat Med* 2021; **10**: 2123-2133 [PMID: [33725769](#) DOI: [10.21037/apm-21-49](#)]
- 28 **Thomson JA**, Itskovitz-Eldor J, Shapiro SS, Waknitz MA, Swiergiel JJ, Marshall VS, Jones JM. Embryonic stem cell lines derived from human blastocysts. *Science* 1998; **282**: 1145-1147 [PMID: [9804556](#) DOI: [10.1126/science.282.5391.1145](#)]
- 29 **Zang L**, Li Y, Hao H, Liu J, Cheng Y, Li B, Yin Y, Zhang Q, Gao F, Wang H, Gu S, Li J, Lin F, Zhu Y, Tian G, Chen Y, Gu W, Du J, Chen K, Guo Q, Yang G, Pei Y, Yan W, Wang X, Meng J, Zhang S, Ba J, Lyu Z, Dou J, Han W, Mu Y. Efficacy and safety of umbilical cord-derived mesenchymal stem cells in Chinese adults with type 2 diabetes: a single-center, double-blinded, randomized, placebo-controlled phase II trial. *Stem Cell Res Ther* 2022; **13**: 180 [PMID: [35505375](#) DOI: [10.1186/s13287-022-02848-6](#)]
- 30 **Lan T**, Luo M, Wei X. Mesenchymal stem/stromal cells in cancer therapy. *J Hematol Oncol* 2021; **14**: 195 [PMID: [34789315](#) DOI: [10.1186/s13045-021-01208-w](#)]
- 31 **Moll G**, Ankrum JA, Kamhieh-Milz J, Bieback K, Ringdén O, Volk HD, Geissler S, Reinke P. Intravascular Mesenchymal Stromal/Stem Cell Therapy Product Diversification: Time for New Clinical Guidelines. *Trends Mol Med* 2019; **25**: 149-163 [PMID: [30711482](#) DOI: [10.1016/j.molmed.2018.12.006](#)]
- 32 **Guillamat-Prats R**. Role of Mesenchymal Stem/Stromal Cells in Coagulation. *Int J Mol Sci* 2022; **23** [PMID: [36142297](#) DOI: [10.3390/ijms231810393](#)]
- 33 **Liao L**, Shi B, Chang H, Su X, Zhang L, Bi C, Shuai Y, Du X, Deng Z, Jin Y. Heparin improves BMSC cell therapy: Anticoagulant treatment by heparin improves the safety and therapeutic effect of bone marrow-derived mesenchymal stem cell cytotreatment. *Theranostics* 2017; **7**: 106-116 [PMID: [28042320](#) DOI: [10.7150/thno.16911](#)]
- 34 **Gleeson BM**, Martin K, Ali MT, Kumar AH, Pillai MG, Kumar SP, O'Sullivan JF, Whelan D, Stocca A, Khider W, Barry FP, O'Brien T, Caplice NM. Bone Marrow-Derived Mesenchymal Stem Cells Have Innate Procoagulant Activity and Cause Microvascular Obstruction Following Intracoronary Delivery: Amelioration by Antithrombin Therapy. *Stem Cells* 2015; **33**: 2726-2737 [PMID: [25969127](#) DOI: [10.1002/stem.2050](#)]
- 35 **Toma C**, Wagner WR, Bowry S, Schwartz A, Villanueva F. Fate of culture-expanded mesenchymal stem cells in the microvasculature: in vivo observations of cell kinetics. *Circ Res* 2009; **104**: 398-402 [PMID: [19096027](#) DOI: [10.1161/CIRCRESAHA.108.187724](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

