

# World Journal of *Clinical Cases*

*World J Clin Cases* 2023 August 6; 11(22): 5193-5415



**MINIREVIEWS**

- 5193 Research progress on reactive oxygen species production mechanisms in tumor sonodynamic therapy  
*Dong HQ, Fu XF, Wang MY, Zhu J*

**ORIGINAL ARTICLE****Retrospective Study**

- 5204 Combining the age-male-albumin-bilirubin-platelets score and shear wave elastography stratifies carcinogenic risk in hepatitis C patients after viral clearance  
*Masaoka R, Gytoku Y, Shirahashi R, Suda T, Tamano M*
- 5215 Changes in neurotransmitter levels, brain structural characteristics, and their correlation with PANSS scores in patients with first-episode schizophrenia  
*Xu XJ, Liu TL, He L, Pu B*
- 5224 Five-year outcomes of immediate implant placement for mandibular molars with and without chronic apical periodontitis: A retrospective study  
*Yang H, Luo D, Yuan MJ, Yang JJ, Wang DS*

**Observational Study**

- 5236 Standardization of apple cancellation test for neglect patients in Korea: An observational study  
*Jang WH, Jang JS*

**Prospective Study**

- 5244 Diabetic neuropathy results in vasomotor dysfunction of medium sized peripheral arteries  
*Ege F, Kazci Ö, Aydin S*

**SYSTEMATIC REVIEWS**

- 5252 COVID-19-induced gastrointestinal autonomic dysfunction: A systematic review  
*Elbeltagi R, Al-Beltagi M, Saeed NK, Bediwy AS*

**META-ANALYSIS**

- 5273 Meta-analysis of outcomes from drug-eluting stent implantation in infrapopliteal arteries  
*Li MX, Tu HX, Yin MC*

**CASE REPORT**

- 5288 Acute hepatitis of unknown etiology in an adult female: A case report  
*Dass L, Pacia AMM, Hamidi M*

- 5296** Zimberelimab plus chemotherapy as the first-line treatment of malignant peritoneal mesothelioma: A case report and review of literature  
*Peng XD, You ZY, He LX, Deng Q*
- 5303** Recurrent ventricular arrhythmia due to aconite intoxication successfully treated with landiolol: A case report  
*Matsuo C, Yamamoto K, Fukushima H, Yajima D, Inoue H*
- 5309** Anti-phospholipase A2 receptor-associated membranous nephropathy with human immunodeficiency virus infection treated with telitacicept: A case report  
*Wang JL, Sun YL, Kang Z, Zhang SK, Yu CX, Zhang W, Xie H, Lin HL*
- 5316** Rapid progression of heart failure secondary to radioactive iodine treatment of hyperthyroidism: A case report  
*Li ZH, Ni LJ, Liu YQ, Si DY*
- 5322** Pathological complete response to neoadjuvant alectinib in unresectable anaplastic lymphoma kinase positive non-small cell lung cancer: A case report  
*Wang LM, Zhao P, Sun XQ, Yan F, Guo Q*
- 5329** Hepatoid adenocarcinoma of the stomach with neuroendocrine differentiation: A case report and review of literature  
*Fei H, Li ZF, Chen YT, Zhao DB*
- 5338** Acquired haemophilia as a complicating factor in treatment of non-muscle invasive bladder cancer: A case report  
*Ryšánková K, Gumulec J, Grepl M, Krhut J*
- 5344** Persistent dysexecutive syndrome after pneumococcal meningitis complicated by recurrent ischemic strokes: A case report  
*Abbruzzese L, Martinelli G, Salti G, Basagni B, Damora A, Scarselli C, Peppoloni G, Podgorska A, Rosso G, Bacci M, Alfano AR, MANCUSO M*
- 5351** Treatment of refractory anti-melanoma differentiation-associated gene 5 antibody-positive dermatomyositis complicated by rapidly progressing interstitial pulmonary disease: Two case reports  
*Wang QH, Chen LH*
- 5358** TINAVI robot-assisted one-stage anteroposterior surgery in lateral position for severe thoracolumbar fracture dislocation: A case report  
*Ye S, Chen YZ, Zhong LJ, Yu CZ, Zhang HK, Hong Y*
- 5365** Individual with concurrent chest wall tuberculosis and triple-negative essential thrombocythemia: A case report  
*Xu XY, Yang YB, Yuan J, Zhang XX, Kang L, Ma XS, Yang J*
- 5373** Self-strangulation induced penile partial amputation: A case report  
*Maimaitiming ABLT, Mulati YLSD, Apizi ART, Li XD*
- 5382** Long-term rare giant sialolithiasis for 30 years: A case report and review of literature  
*Mao JS, Lee YC, Chi JCY, Yi WL, Tsou YA, Lin CD, Tai CJ, Shih LC*

- 5391** Kawasaki disease with peritonsillar abscess as the first symptom: A case report  
*Huo LM, Li LM, Peng HY, Wang LJ, Feng ZY*
- 5398** Treatment of a patient with severe lactic acidosis and multiple organ failure due to mitochondrial myopathy: A case report  
*Chen L, Shuai TK, Gao YW, Li M, Fang PZ, Christian W, Liu LP*
- 5407** Early esophageal carcinomas in achalasia patient after endoscopic submucosal dissection combined with peroral endoscopic myotomy: A case report  
*An BQ, Wang CX, Zhang HY, Fu JD*

**LETTER TO THE EDITOR**

- 5412** Caution in the use of sedation and endomyocardial biopsy for the management of pediatric acute heart failure caused by endocardial fibroelastosis  
*Xin XX, Se YY*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Etienne Andrade Munhoz, PhD, Associate Professor, Department of Dentistry, Health Science Centre, Federal University of Santa Catarina, Florianopolis 88040-379, Brazil. etiamfob@yahoo.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Si Zhao; Production Department Director: Xu Guo; Editorial Office Director: Jin-Lei Wang.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

August 6, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



Retrospective Study

# Five-year outcomes of immediate implant placement for mandibular molars with and without chronic apical periodontitis: A retrospective study

Hua Yang, Dan Luo, Mu-Jie Yuan, Jian-Jun Yang, Da-Shan Wang

**Specialty type:** Medicine, research and experimental

**Provenance and peer review:** Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B  
Grade C (Good): C, C  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Ardila CM, Colombia; Utharavalli USK, India

**Received:** May 29, 2023

**Peer-review started:** May 29, 2023

**First decision:** June 15, 2023

**Revised:** June 26, 2023

**Accepted:** July 10, 2023

**Article in press:** July 10, 2023

**Published online:** August 6, 2023



**Hua Yang, Dan Luo, Mu-Jie Yuan, Jian-Jun Yang,** Department of Oral and Maxillofacial Surgery, The Affiliated Hospital of Qingdao University, Qingdao 266003, Shandong Province, China

**Hua Yang, Dan Luo, Mu-Jie Yuan, Jian-Jun Yang, Da-Shan Wang,** School of Stomatology, Qingdao University, Qingdao 266003, Shandong Province, China

**Da-Shan Wang,** Department of Oral Implantology, The Affiliated Hospital of Qingdao University, Qingdao 266003, Shandong Province, China

**Corresponding author:** Da-Shan Wang, PhD, Professor, Department of Oral Implantology, The Affiliated Hospital of Qingdao University, No. 16 Jiangsu Road, Qingdao 266003, Shandong Province, China. [wdsapple@163.com](mailto:wdsapple@163.com)

## Abstract

### BACKGROUND

Most physicians consider molars with chronic apical periodontitis (CAP) lesions as contraindications for immediate implant placement. At the patient's request, we perform immediate implant placement of the mandibular molars with CAP in clinical practice.

### AIM

To retrospectively analyze and compare the 5-year outcomes of immediate implant placement of the mandibular molars with CAP and those without obvious inflammation.

### METHODS

The clinical data of patients with immediate implant placement of the mandibular molars in the Department of Oral and Maxillofacial Surgery, the Affiliated Hospital of Qingdao University, from June 2015 to June 2017 were collected. The patients were divided into CAP ( $n = 52$ ) and no-CAP ( $n = 45$ ) groups. Changes in bone mineral density and bone mass around implants were analyzed 5 years after implant restoration.

### RESULTS

At 5 years after implantation, the peri-implant bone mineral density was  $528.2 \pm 78.8$  Hounsfield unit (HU) in the CAP group and  $562.6 \pm 82.9$  HU in the no-CAP

group ( $P = 0.126$ ). Marginal bone resorption around implants did not differ significantly between the two groups, including buccal ( $P = 0.268$ ) or lingual ( $P = 0.526$ ) resorption in the vertical direction or buccal ( $P = 0.428$ ) or lingual ( $P = 0.560$ ) resorption in the horizontal direction. Changes in the peri-implant jump space did not differ significantly between the two groups, including the buccal ( $P = 0.247$ ) or lingual ( $P = 0.604$ ) space in the vertical direction or buccal ( $P = 0.527$ ) or lingual ( $P = 0.707$ ) space in the horizontal direction. The gray value of cone-beam computed tomography measured using Image J software can reflect the bone mineral density. In the CAP area, the gray values of the bone tissue immediately and 5 years after implant placement differed significantly from those of the surrounding bone tissue ( $P < 0.01$ ).

## CONCLUSION

The results of this study suggest that immediate implant placement of the mandibular molars with CAP can achieve satisfactory 5-year clinical results, without significant differences in the complications, survival rate, or bone tissue condition from the no-CAP mandibular molars.

**Key Words:** Molar; Chronic apical periodontitis; Dental implantation; Bone density; Treatment outcome; Retrospective study

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** This study was aimed at retrospectively analyzing and comparing the 5-year outcomes of immediate implant placement of the mandibular molars with chronic apical periodontitis and those without obvious inflammation based on changes in bone mineral density and bone mass around implants 5 years after implant restoration.

**Citation:** Yang H, Luo D, Yuan MJ, Yang JJ, Wang DS. Five-year outcomes of immediate implant placement for mandibular molars with and without chronic apical periodontitis: A retrospective study. *World J Clin Cases* 2023; 11(22): 5224-5235

**URL:** <https://www.wjgnet.com/2307-8960/full/v11/i22/5224.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v11.i22.5224>

## INTRODUCTION

Chronic apical periodontitis (CAP) is a globally prevalent infectious disease, characterized by the inflammatory response of periapical tissues and local alveolar bone destruction caused by intramedullary microbial infection[1,2]. Compared to other bones in the body, the alveolar bone can communicate directly with the outside environment through the dental pulp; if the pulp is necrotic and infected, no epithelial barrier exists to resist infection or inflammatory factors[3]. In the development of CAP, microbial infection and immune defense response jointly lead to local alveolar bone destruction[4-6]. Root canal therapy is the main treatment for CAP. However, after root canal treatment, CAP may persist as asymptomatic inflammation[7]. Persistent CAP after root canal treatment may be caused by failure to strictly follow the principle of asepsis, a poor approach design, residual accessory canals, improper use of instruments, incomplete debridement, and microleakage of the temporary or permanent prosthesis[8].

Due to the complexity of the main and accessory root canal systems and residual infection at the root canal branches and anastomosis, the mandibular molars may develop persistent CAP even after following the most stringent root canal treatment procedures, resulting in persistent destruction of the periapical bone[1,2,9]. When the periapical shadow area of the affected tooth is enlarged in the periapical radiograph, the root canal retreatment is difficult to perform, and the affected tooth cannot be retained, the method of restoring the affected tooth after extraction should be considered[10].

Since Branemark successfully implanted the first dental implant in 1965, the dental implant technology has been an effective method to restore the masticatory and esthetic functions of the dentition for patients with missing teeth[11,12]. The initial treatment protocol involves implant placement in the healed extraction socket, referring to the alveolar bone after healing for at least 5-6 mo after extraction of the affected tooth, a delayed dental implant. In 1989, Lazzara *et al*[13] introduced implants placed immediately after tooth extraction. In recent years, many studies have confirmed the reliability of implants placed during tooth extraction[14]. With advancements in the implant technology, immediate implant placement has become the first choice to restore missing teeth in clinical practice[15-17]. Hansson and Ekestubbe [18] and Ericsson *et al*[19] found that immediate dental implant surgery is minimally invasive, which reduces the risk of osteonecrosis and promotes the process of bone remodeling, shortening the healing period of the alveolar bone and promoting the transformation of woven bone into lamellar bone. Compared to delayed dental implant placement, immediate implant placement has advantages of shortening the treatment course, reducing the procedural pain of patients, and reducing the alveolar bone absorption in the surgical area[20]. Currently, immediate implant placement is mainly used in the anterior esthetic area in clinical practice[21].

Most molars cannot be retained after severe CAP. The complex shape of the extraction socket after extraction of the affected teeth, severe destruction of the periodontal soft tissue and alveolar bone, and presence of abundant inflammatory granulation tissue adversely affect the success rate of immediate implant placement[16]. In a prospective study, Alsaadi *et al*[22] found that immediate implant placement in teeth with CAP lesions resulted in a greater rate of implant failure



compared to delayed implant placement. In addition, the retrograde peri-implantitis that occurred in the study was thought to be caused by immediate implant placement at the CAP tooth position[23]. Most physicians consider the molars with severe CAP as a contraindication for immediate implant placement. However, in recent years, data from animal studies, case reports, and prospective studies have shown that the success rate of implant placement in the teeth with CAP is similar to that in the teeth without CAP[24,25].

The aim of this study was to retrospectively analyze and compare the 5-year outcomes of immediate implant placement of the mandibular molars with CAP and those without obvious inflammation based on quantitative peri-implant bone mass changes evaluated using Simplant and Image J software.

## MATERIALS AND METHODS

### Patient screening

Clinical cases of the molars that could not be retained or received immediate implant placement were collected from the Oral and Maxillofacial Surgery Department of the Affiliated Hospital of Qingdao University from June 2015 to June 2017. **Figure 1** shows the patients' screening process.

This study was conducted in accordance with Declaration of Helsinki guidelines and regulations, and all study methods were approved by the Ethics Committee of the Affiliated Hospital of Qingdao University (QYFYKYL-958311920). Informed consent was obtained and signed by all participants and their families in this study.

### Preoperative preparation

Inclusion criteria were: Age  $\geq 18$  years; no pregnancy or lactation; no systemic disease or the use of related drugs; good oral hygiene with no acute inflammation in the dentition; no retentive value of the affected teeth in the posterior mandibular region as confirmed by oral physicians and prosthodontists (**Figure 2A-C**); patient consent for immediate implant placement; and willingness to attend regular follow-ups. The same chief physician with 25 years of work experience completed the implant placement surgery and crown restoration, under assistance by doctors with more than 5 years of clinical experience.

### Surgical procedure

All study participants underwent routine disinfection and draping in the intraoral and maxillofacial regions. We used 1% iodophor for disinfection and asked the patient to gargle for 20 s before sterilizing the maxillofacial area, up to the palpebral fissure, down to the level of the hyoid bone, and bilaterally to the front of the tragus. The angular mucoperiosteal flap on the buccal side of the affected tooth was completely turned over. A high-speed turbine extraction handpiece was used to extract the affected teeth by root separation to protect the integrity of the apical septum and bone wall of the alveolar fossa to the greatest extent (**Figure 3A and B**). The inflammatory tissues on the edge and inside of the mucoperiosteal flap were pruned. The inflammatory tissue attached to the inner wall of the alveolar fossa was scraped with an appropriate type of scraping spoon and then polished with large, medium, and small ball drills until no fibrous tissue remained on the bone wall of the alveolar fossa (**Figure 3C and D**). The implant cavities were prepared step-wise, and the corresponding implants were selected and rotated into the implant socket using a ratchet wrench to ensure that the implants had been placed in the correct three-dimensional position, with the neck and shoulder of the implants placed 1.0-1.5 mm below the edge of the alveolar bone. The torque force after implant placement was tested using a force measuring wrench to ensure that the torque force exceeded 35 N. cm (**Figure 3E and F**). The implant stability quotient was measured with the resonance frequency analyzer (Osstell, Sweden) to confirm the initial stability of the implant and install the healing abutment with appropriate diameter and penetration height. The buccal-lingual mucoperiosteal flap was tightly sutured around the healing abutment (**Figure 3G**). Cone-beam computed tomography (CBCT) was performed after immediate implant placement to ensure that the implants were in the correct tooth and spatial positions (**Figure 2D-F**).

### Dental crown restoration


At 3 mo postoperatively, an impression was taken, and an all-porcelain crown was fabricated (**Figure 3H and I**).

### Evaluation index

All patients underwent CBCT before and after the implant placement and at the follow-up visits (**Figure 2**). CBCT was performed by the same dental radiologist with 10 years of experience. The clinical examination and CBCT image measurement and analysis were performed by three dentists with 10, 10, and 5 years of working experience, respectively. Each dentist measured and analyzed the CBCT scans of the two groups of patients, and the results of each patient's evaluation index were averaged and recorded. The intraclass correlation coefficient (ICC) was used to test the difference in the observation results of the three observers. According to the specificity of the evaluation indicators in this study, the peri-implant bone mineral density with the most complicated detection steps was selected for ICC calculation, which could reflect the differences in the observation results of the three observers in this study.

The bone tissue around the implant was analyzed 5 years after crown restoration. CBCT equipment and parameters were as follows: Brand model, Kava i-CAT 17-19; tube voltage, 120 kV; current, 5 mA; exposure time, 26.9 s; diameter, 16 cm; height, 11 cm; and resolution, 0.25 mm.





DOI: 10.12998/wjcc.v11.i22.5224 Copyright ©The Author(s) 2023.

**Figure 1 Patients' screening flowchart.** CAP: Chronic apical periodontitis.


The patient sat in an upright position. The dental chair was adjusted to the appropriate height according to the patient's height. The patient's head was rested on the head support. The dentist wore protective clothing for infection control, adjusted the front center line to the center of the patient's face, directed the patient's eyes to look forward, adjusted the orbital ear plane to be parallel to the ground, and held the patient's jaw bracket. "Preview" determined that the scanning range included the patient's complete upper and lower dentition and alveolar bone, and the CBCT scan was captured.

**Peripheral bone density:** The CBCT scans of the CAP and no-CAP groups 5 years after dental crown restoration were exported in the Digital Imaging and Communications in Medicine format and imported into Simplant software (Materialise Dental, Belgium). The alveolar bone density was measured around the virtual implant (Figure 4A).

**Vertical and horizontal marginal bone losses:** The implant long axis L0 and the implant shoulder plane L1 perpendicular to L0 were determined. In the vertical direction, the vertical distance from the crest of the buccal and lingual alveolar crests to L1 (H1) was measured. In the horizontal direction, the distance (W1) between L1 and the lateral side of the buccal and lingual bone walls and the intersection of the implant edge was measured (Figure 4B). Each group of data was measured thrice, with an average accuracy of 0.01 mm. The buccal and lingual vertical and horizontal marginal bone loss from immediately after implant placement to 5 years after crown restoration in the CAP and no-CAP groups were calculated.

**Changes in the jump gap:** The long axis L0 of the implant body and the shoulder plane L1 perpendicular to L0 were determined. The vertical distance from the highest point of implant-bone contact to L1 (H2) was measured immediately after surgery and 5 years after crown restoration, and the distance between L1 and the intersections of buccal and lingual bone walls and implant edges (W2) was measured (Figure 4B). The data in each group were measured thrice, with an average accuracy of 0.01 mm. Changes in buccal and lingual jump gaps in the CAP and no-CAP groups 5 years after crown restoration were calculated.

**Changes in bone density in the CAP-damaged areas:** The CBCT scans of the largest area of bone destruction in the horizontal plane of the CAP group were selected and exported in the jpg format. With the Image J software, the region of interest was selected in the image, avoiding areas such as the surrounding alveolar bone and root tip as much as possible. After determining the region of interest, the "Measure" option in the toolbar was selected, followed by the "Analyze" option to obtain the gray value of the inflammatory bone destruction area and surrounding bone tissue (Figure 4C). Differences in the gray value between the two were calculated and recorded. The gray values of the bone destruction area and surrounding bone tissue in the same area 5 years after crown restoration were obtained with the same method, and the difference in the gray values between the two was calculated and recorded. The obtained gray value cannot directly



DOI: 10.12998/wjcc.v11.i22.5224 Copyright ©The Author(s) 2023.

**Figure 2** Cone-beam computed tomography scans of the chronic apical periodontitis group. A-C: Cone-beam computed tomography (CBCT) scans of the chronic apical periodontitis (CAP) group before surgery; D-F: CBCT scans of the CAP group after immediate dental implantation; G-I: CBCT scans of the CAP group 5 years after restoration.

be used as the bone mineral density value, but the difference in the gray values between the area of inflammatory bone destruction and the surrounding bone tissue immediately and 5 years after surgery can represent the change in bone mineral density in this area.

### Statistical analysis

ICC calculations and statistical tests were performed using SPSS 20.0 (IBM, Chicago, IL). Patient age belonged to non-normally distributed data, and the Wilcoxon signed-rank test was used for the analysis. The data of peri-implant bone tissue changes (bone mineral density, marginal bone loss, jump gap, and gray value) belonged to normally distributed data and were analyzed with the independent-samples *t*-test. For patient sex, the chi-square test was used. A *P* value < 0.05 was considered to indicate statistical significance.

## RESULTS

We enrolled 97 teeth of 97 patients, with 52 patients in the CAP group, including 28 women and 24 men, aged  $35.6 \pm 5.28$  (68-18) years, and 45 patients in the no-CAP group, including 19 women and 26 men, aged  $36.8 \pm 4.79$  (66-20) years. Patient age (*P* = 0.385) or sex (*P* = 0.314) distribution did not differ significantly between the two groups (Tables 1 and 2). The peri-implant bone mineral density values measured by three observers in this study were tested for inter-observer differences, and the calculated ICC value was 0.816, falling between 0.75 and 0.9, indicating a good consistency of the observation results[26].

### Implant repair complications and retention

The clinical records of the two groups within 5 years were analyzed. In the CAP group, one patient developed central

**Table 1** Age range of the patients


Group	Max	Min	Mean	SD	P value
CAP group ( <i>n</i> = 52)	68	18	35.6	5.28	0.385
No-CAP group ( <i>n</i> = 45)	66	20	36.8	4.79	

CAP: Chronic apical periodontitis; SD: Standard deviation.

**Table 2** Sex of the patients

Group	Sex		Total	P value
	Female	Male		
CAP group ( <i>n</i> = 52)	28	24	52	0.314
No-CAP group ( <i>n</i> = 45)	19	26	45	
Total ( <i>n</i> = 97)	47	50	97	

CAP: Chronic apical periodontitis.




DOI: 10.12998/wjcc.v11.i22.5224 Copyright ©The Author(s) 2023.

**Figure 3** Immediate implant placement surgery and repair in the chronic apical periodontitis group. A: Patients were examined before surgery; B: The gingiva was incised, and the mucoperiosteal flap was raised and removed; C: The inflammatory tissue in the extraction fossa was removed, and the primary implant hole was prepared; D: Residual inflammatory tissue on the bone wall of the inflammatory lesion area was removed using a ball drill via a primary implant hole; E: Removed teeth and removed inflammatory tissue; F: Immediate implant placement; G: Non-embedding healing; H: Gum cuff; I: Crown restoration completed.

screw loosening, and one patient experienced restoration chipping. In the no-CAP group, one patient's prosthesis fell off, and one patient's prosthesis chipped off. After the corresponding treatment, the implant condition was good. The implants in the two groups were in position and functioned well, and the survival rate was 100%.





DOI: 10.12998/wjcc.v11.i22.5224 Copyright ©The Author(s) 2023.

**Figure 4 Method for measuring changes in bone tissue around implants.** A: The peri-implant bone mineral density was measured using Simplant software; B: Schemas of marginal bone resorption and jump gap measurements. L0 is the long axis of the implant. L1 is perpendicular to L0. H1 is the vertical distance from the crest of the alveolar bone to L1. W1 is the horizontal distance from the most lateral side of the alveolar bone wall to the edge of the implant. H2 is the vertical distance from the highest point of contact between the implant and the bone to L1. W2 is the width of the jump gap. C: Image J software was used to measure the gray value of the bone destruction area at the root apex of the affected tooth (the gray value of alveolar bone cone-beam computed tomography can reflect its bone mineral density).

### Changes in bone tissue around the implant


The CBCT data were imported into Simplant software to measure and compare changes in bone mineral density of peri-implant bone tissue after 5 years of implant denture restoration. The peri-implant bone densities were  $528.2 \pm 78.8$  Hounsfield unit (HU) and  $562.6 \pm 82.9$  HU after 5 years of implant restoration in the CAP and no-CAP groups, respectively. The independent-samples t-test showed no significant difference in the peri-implant bone density between the two groups ( $P = 0.126$ ).

Figure 5A and B shows the amount of bone resorption at the implant edge. In the vertical direction, the buccal marginal bone loss was  $0.43 \pm 0.16$  mm in the CAP group and  $0.47 \pm 0.14$  mm in the no-CAP group, showing no significant difference between the two groups ( $P = 0.268$ ). The marginal bone loss did not differ significantly between the CAP ( $0.43 \pm 0.14$  mm) and no-CAP ( $0.45 \pm 0.14$  mm) groups ( $P = 0.526$ ). In the horizontal direction, the buccal marginal bone loss was  $0.91 \pm 0.22$  mm in the CAP group and  $0.86 \pm 0.23$  mm in no-CAP group, showing no significant difference between the two groups ( $P = 0.428$ ). The marginal bone loss was  $0.67 \pm 0.15$  mm in the CAP group and  $0.64 \pm 0.22$  mm in the no-CAP group ( $P = 0.560$ ), showing no significant difference.

In the vertical direction, the incremental value of the highest point of buccal implant-bone contact was  $3.04 \pm 1.21$  mm in the CAP group and  $3.36 \pm 0.89$  mm in the no-CAP group, showing no significant difference ( $P = 0.247$ ). The implant bone increment at the highest point of lingual implant-bone contact was  $2.80 \pm 0.78$  mm in the CAP group and  $2.71 \pm 0.63$  mm in the no-CAP group, showing no significant difference ( $P = 0.604$ ; Figure 5C). In the horizontal direction, the width change in the buccal jump gap was  $3.44 \pm 0.95$  mm in the CAP group and  $3.30 \pm 0.83$  mm in the no-CAP group, showing no significant difference ( $P = 0.527$ ). The width change in the lingual jump gap was  $2.57 \pm 0.78$  mm in the CAP group and  $2.84 \pm 0.63$  mm in the no-CAP group, showing no significant difference ( $P = 0.707$ ; Figure 5D).

### Changes in the bone density of the CAP-damaged area

The inflammatory bone destruction area of the alveolar bone in the CAP group disappeared 5 years after implant denture



**Figure 5** Changes in bone tissue around implants. A: Vertical edge bone absorption of peri-implant (buccal,  $P = 0.268$ ; lingual,  $P = 0.526$ ); B: Horizontal edge bone absorption of peri-implant (buccal,  $P = 0.428$ ; lingual,  $P = 0.560$ ); C: Implant-bone contact peak increment (buccal,  $P = 0.247$ ; lingual,  $P = 0.604$ ); D: Changes in the jump gap (buccal,  $P = 0.527$ ; lingual,  $P = 0.707$ ).

restoration. The gray value difference between the CAP lesion area and the surrounding bone tissue was  $107.6 \pm 21.7$  immediately after surgery and  $32.5 \pm 15.3$  5 years after implant restoration, with significant differences between the two groups ( $P < 0.01$ ).

## DISCUSSION

Currently, immediate implant placement is the preferred modality to restore the missing anterior teeth[27]. For the mandibular molars, immediate implant placement is often contraindicated because of the large area of CAP lesions around the root apices and severe alveolar bone destruction[28]. Therefore, delayed dental implant placement is chosen, which prolongs the restoration time of the missing teeth and aggravates pain[29]. In this retrospective study, the patients who had received immediate implant placement and crown restoration with successful outcomes after 5 years were selected. The aim of the present study was to investigate the clinical effect of immediate implant placement for the missing mandibular molars.

Osseointegration is the decisive factor for the success of implant restoration[30]. The larger the contact area between the implant surface and the trabecular bone in the surrounding alveolar bone, the better is implant osseointegration[31]. The study revealed no significant differences in bone mineral density between the chronic periapical teeth with immediate restoration and the teeth with conventional immediate restoration 5 years after immediate restoration. Microbial antigens derived from root canal infections can stimulate specific and nonspecific immune responses in periapical tissues[32]. Macrophages, mast cells, T cells, and neutrophils are involved in the formation of CAP tissue, including cytokines and chemokines[32,33]. Therefore, the inflammatory tissue in the alveolar fossa should be completely removed to reduce the influence of inflammatory factors around the implant on the bone tissue.

The marginal bone level plays a crucial role in maintaining the stability and function of implants, with great significance in the long-term survival rate of implants[34]. As for the vertical marginal bone mass, Kakar *et al*[35] found that during immediate implant placement for the teeth with CAP, the risk of residual infection can be reduced by curetting the inflammatory granulation tissue of the extraction socket and thoroughly washing it, and the vertical marginal bone resorption can be minimized. As for the horizontal marginal bone mass, Hatting *et al*[36] found that the buccal horizontal bone mass had decreased by 0.89 mm while the lingual horizontal bone mass had decreased by 0.69 mm 1 year after immediate implant placement for a molar, similar to the results of this study. Some scholars also believe that the buccal bone wall thickness may affect horizontal bone resorption in immediate implant placement[37]. We believe that the inflammatory granulation tissue around the neck of the molars with CAP should not be ignored, owing to its relevance in marginal bone resorption of implants. Intraoperatively, the inflammatory granulation tissue can be completely removed by making an incision and raising a flap to prevent the destruction of the alveolar bone in the implant socket and reduce

marginal bone resorption of implants. In addition, we positioned the necked shoulder of the implant 1.0-1.5 mm below the alveolar bone margin to compensate for vertical and marginal bone resorptions and ensure that the threads of the implant were not exposed after the marginal bone remodeling had been stabilized.

The gap between the medial wall of the alveolar bone and the implant surface is called the jump gap[17,38]. The implant may be wrapped in blood clots after placement. At 1-2 wk postoperatively, osteoblasts form new woven bone on the implant surface and a bridge with the bone wall of the alveolar socket. As the woven bone continues to mineralize, it can transform into lamellar bone[38]. The present study involved no bone grafting in the jumping gap in the CAP or no-CAP group. The jumping gap healed naturally after the operation. Five years after implant placement and loading, CBCT showed that the original jumping gap had disappeared.

Bone mass loss is a clinical feature of CAP caused by microbial factors and immune defense responses[39]. As a chronic inflammatory disease, CAP causes an imbalance between bone resorption and remodeling, which leads to bone mass loss [28]. Bone resorption and formation are antagonistic and coupled processes of osteoblasts and osteoclasts, which together constitute the normal bone mass[40]. Several factors affect periapical bone remodeling, including microbes, human signaling pathways, and the immune system[41]. CAP may directly be caused by bacterial infection and trigger an immune response from the host, resulting in destruction of periapical tissues[40,42]. In the progression of pulpitis, the flora is simple in the early stage, but with the dominance of gram-negative anaerobic bacteria, such as porphyromonas, the complexity of root canal flora increases[42]. Of all bacterial species, 54.6% are strictly anaerobic, while anaerobic gram-negative bacteria dominate the root canals with periapical lesions[43]. In the periapical infected tissues, bacterial abundance and diversity are significantly reduced, and the microbial balance in the biofilm is disrupted[44]. Endotoxin in the cell wall of gram-negative bacteria can cause local tissue swelling and bone resorption and mobilize the host immune response. Further, its content is positively correlated with the degree of bone injury[45]. It can promote osteoclast differentiation and bone resorption induced by lipoic acid and participate in maintaining the survival of mature osteoclasts, thereby jointly causing inflammatory alveolar bone loss[45].

Therefore, in the immediate implant surgery in the present study, we should removed the inflammatory tissue in the alveolar fossa and the bacteria and inflammatory tissue in the periapical bone destruction area. The results showed no significant difference between the CAP and no-CAP groups in the gray value of the apical bone destruction area and the surrounding bone tissue 5 years after loading of immediate dental implant restoration. This indicated that the mandibular molars with various inflammatory periapical bone tissue lesions could be effectively removed by improving the implant socket approach during immediate implant placement to promote bone tissue reconstruction.

This study has some limitations. First, this was a retrospective study; therefore, the credibility of the findings is weaker. However, strict inclusion and exclusion criteria were set to minimize variables not relevant to the purpose of the study. Second, the study sample size was small; therefore, the data may not accurately represent the entire population. Third, all patients showed good compliance, which caused a selection bias. In the future, we will conduct a prospective study corresponding to this study to expand the sample size and follow-up duration and further explore the soft and hard tissue conditions of immediate dental implant placement in the mandibular molars with CAP.

## CONCLUSION

The results of this study suggest that immediate implant placement of the mandibular molars with CAP can achieve satisfactory 5-year clinical results, without significant differences in the complications, survival rate, or bone tissue condition from the no-CAP mandibular molars.

## ARTICLE HIGHLIGHTS

### Research background

At the patient's request, we tried to perform immediate implant placement of the mandibular molars with CAP in clinical practice.

### Research motivation

Most physicians consider the molars with CAP lesions as contraindications for immediate implant placement.

### Research objectives

Immediate implant placement of the mandibular molars with CAP can achieve satisfactory 5-year clinical results.

### Research methods

In this study, we retrospectively analyzed and compared the 5-year outcomes of immediate implant placement of the mandibular molars with CAP and those without obvious inflammation based on quantitative peri-implant bone mass changes evaluated using Simplant and Image J software.

### Research results

The peri-implant bone density was  $528.2 \pm 78.8$  Hounsfield unit (HU) in the CAP group and  $562.6 \pm 82.9$  HU in the no-



CAP group 5 years after implant placement. The peri-implant bone density did not differ significantly between the two groups. The marginal bone resorption or jump gap did not differ significantly between the two groups. In the CAP area, the gray values of the bone tissue immediately and 5 years after implant placement differed significantly from those of the surrounding bone tissue ( $P < 0.01$ ).

### Research conclusions

The results of this study suggest that immediate implant placement of the mandibular molars with CAP can achieve satisfactory 5-year clinical results, without significant differences in the complications, survival rate, or bone tissue condition from the no-CAP mandibular molars.

### Research perspectives

Immediate implant placement of the mandibular molars with chronic apical periodontitis (CAP) can achieve good clinical outcomes.

## FOOTNOTES

**Author contributions:** Yang H and Luo D wrote the main manuscript text; Yang H, Luo D and Yuan MJ prepared figures 1-5 and tables 1-2; Yang JJ and Wang DS contributed equally to this study, they reviewed and revised the manuscript.

**Supported by** The Qingdao Science and Technology Benefit People Project, No. 21-1-4-rkjk-20-nsh.

**Institutional review board statement:** This study was conducted in accordance with Declaration of Helsinki guidelines and regulations, and all study methods were approved by the Ethics Committee of the Affiliated Hospital of Qingdao University (QYFYKYL958311920).

**Conflict-of-interest statement:** The authors declare no competing interests.

**Data sharing statement:** Technical appendix, statistical code, and dataset available from the corresponding author at [wdsapple@163.com](mailto:wdsapple@163.com). Participants gave informed consent for data sharing.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Da-Shan Wang [0000-0002-2534-7639](https://orcid.org/0000-0002-2534-7639).

**S-Editor:** Lin C

**L-Editor:** A

**P-Editor:** Yuan YY

## REFERENCES

- 1 Azuma MM, Samuel RO, Gomes-Filho JE, Dezan-Junior E, Cintra LT. The role of IL-6 on apical periodontitis: a systematic review. *Int Endod J* 2014; **47**: 615-621 [PMID: [24224782](#) DOI: [10.1111/iej.12196](#)]
- 2 Graunaitė I, Lodiene G, Maciulskienė V. Pathogenesis of apical periodontitis: a literature review. *J Oral Maxillofac Res* 2012; **2**: e1 [PMID: [24421998](#) DOI: [10.5037/jomr.2011.2401](#)]
- 3 Silva TA, Garlet GP, Lara VS, Martins W Jr, Silva JS, Cunha FQ. Differential expression of chemokines and chemokine receptors in inflammatory periapical diseases. *Oral Microbiol Immunol* 2005; **20**: 310-316 [PMID: [16101967](#) DOI: [10.1111/j.1399-302X.2005.00232.x](#)]
- 4 Andrade AL, Nonaka CF, Gordón-Núñez MA, Freitas Rde A, Galvão HC. Immunoexpression of interleukin 17, transforming growth factor  $\beta$  1, and forkhead box P3 in periapical granulomas, radicular cysts, and residual radicular cysts. *J Endod* 2013; **39**: 990-994 [PMID: [23880265](#) DOI: [10.1016/j.joen.2013.04.028](#)]
- 5 Marçal JR, Samuel RO, Fernandes D, de Araujo MS, Napimoga MH, Pereira SA, Clemente-Napimoga JT, Alves PM, Mattar R, Rodrigues V Jr, Rodrigues DB. T-helper cell type 17/regulatory T-cell immunoregulatory balance in human radicular cysts and periapical granulomas. *J Endod* 2010; **36**: 995-999 [PMID: [20478453](#) DOI: [10.1016/j.joen.2010.03.020](#)]
- 6 Bracks IV, Armada L, Gonçalves LS, Pires FR. Distribution of mast cells and macrophages and expression of interleukin-6 in periapical cysts. *J Endod* 2014; **40**: 63-68 [PMID: [24331993](#) DOI: [10.1016/j.joen.2013.09.037](#)]
- 7 Nair PN. Pathogenesis of apical periodontitis and the causes of endodontic failures. *Crit Rev Oral Biol Med* 2004; **15**: 348-381 [PMID: [15574679](#) DOI: [10.1177/154411130401500604](#)]
- 8 Tabassum S, Khan FR. Failure of endodontic treatment: The usual suspects. *Eur J Dent* 2016; **10**: 144-147 [PMID: [27011754](#) DOI: [10.4103/1305-7456.175682](#)]
- 9 Cavalla F, Letra A, Silva RM, Garlet GP. Determinants of Periodontal/Periapical Lesion Stability and Progression. *J Dent Res* 2021; **100**: 29-36 [PMID: [32866421](#) DOI: [10.1177/0022034520952341](#)]

- 10 **Siqueira JF Jr**, Rôças IN, Ricucci D, Hülsmann M. Causes and management of post-treatment apical periodontitis. *Br Dent J* 2014; **216**: 305-312 [PMID: [24651336](#) DOI: [10.1038/sj.bdj.2014.200](#)]
- 11 **Kopp CD**. Brånemark osseointegration. Prognosis and treatment rationale. *Dent Clin North Am* 1989; **33**: 701-731 [PMID: [2680659](#)]
- 12 **Brånemark PI**. Osseointegration and its experimental background. *J Prosthet Dent* 1983; **50**: 399-410 [PMID: [6352924](#) DOI: [10.1016/s0022-3913\(83\)80101-2](#)]
- 13 **Lazzara RJ**, Testori T, Meltzer A, Misch C, Porter S, del Castillo R, Goenê RJ. Immediate Occlusal Loading (IOL) of dental implants: predictable results through DIEM guidelines. *Pract Proced Aesthet Dent* 2004; **16**: 3-15 [PMID: [15279236](#)]
- 14 **Bianchi AE**, Sanfilippo F. Single-tooth replacement by immediate implant and connective tissue graft: a 1-9-year clinical evaluation. *Clin Oral Implants Res* 2004; **15**: 269-277 [PMID: [15142088](#) DOI: [10.1111/j.1600-0501.2004.01020.x](#)]
- 15 **Schropp L**, Wenzel A, Kostopoulos L, Karring T. Bone healing and soft tissue contour changes following single-tooth extraction: a clinical and radiographic 12-month prospective study. *Int J Periodontics Restorative Dent* 2003; **23**: 313-323 [PMID: [12956475](#)]
- 16 **Ragucci GM**, Elnayef B, Criado-Cámara E, Del Amo FS, Hernández-Alfaro F. Immediate implant placement in molar extraction sockets: a systematic review and meta-analysis. *Int J Implant Dent* 2020; **6**: 40 [PMID: [32770283](#) DOI: [10.1186/s40729-020-00235-5](#)]
- 17 **Pitman J**, Seyssens L, Christiaens V, Cosyn J. Immediate implant placement with or without immediate provisionalization: A systematic review and meta-analysis. *J Clin Periodontol* 2022; **49**: 1012-1023 [PMID: [35734911](#) DOI: [10.1111/jcpe.13686](#)]
- 18 **Hansson S**, Ekstubb A. Area moments of inertia as a measure of the mandible stiffness of the implant patient. *Clin Oral Implants Res* 2004; **15**: 450-458 [PMID: [15248880](#) DOI: [10.1111/j.1600-0501.2004.01021.x](#)]
- 19 **Ericsson I**, Randow K, Nilner K, Peterson A. Early functional loading of Brånemark dental implants: 5-year clinical follow-up study. *Clin Implant Dent Relat Res* 2000; **2**: 70-77 [PMID: [11359266](#) DOI: [10.1111/j.1708-8208.2000.tb00108.x](#)]
- 20 **Blanco J**, Carral C, Argibay O, Liñares A. Implant placement in fresh extraction sockets. *Periodontol* 2000 2019; **79**: 151-167 [PMID: [30892772](#) DOI: [10.1111/prd.12253](#)]
- 21 **Araújo MG**, Wennström JL, Lindhe J. Modeling of the buccal and lingual bone walls of fresh extraction sites following implant installation. *Clin Oral Implants Res* 2006; **17**: 606-614 [PMID: [17092217](#) DOI: [10.1111/j.1600-0501.2006.01315.x](#)]
- 22 **Alsaadi G**, Quirynen M, Komárek A, van Steenberghe D. Impact of local and systemic factors on the incidence of oral implant failures, up to abutment connection. *J Clin Periodontol* 2007; **34**: 610-617 [PMID: [17433044](#) DOI: [10.1111/j.1600-051X.2007.01077.x](#)]
- 23 **Quirynen M**, Vogels R, Alsaadi G, Naert I, Jacobs R, van Steenberghe D. Predisposing conditions for retrograde peri-implantitis, and treatment suggestions. *Clin Oral Implants Res* 2005; **16**: 599-608 [PMID: [16164468](#) DOI: [10.1111/j.1600-0501.2005.01147.x](#)]
- 24 **Botticelli D**, Berglundh T, Lindhe J. Resolution of bone defects of varying dimension and configuration in the marginal portion of the peri-implant bone. An experimental study in the dog. *J Clin Periodontol* 2004; **31**: 309-317 [PMID: [15016260](#) DOI: [10.1111/j.1600-051X.2004.00502.x](#)]
- 25 **Wassdorp JA**, Evian CI, Mandracchia M. Immediate placement of implants into infected sites: a systematic review of the literature. *J Periodontol* 2010; **81**: 801-808 [PMID: [20192616](#) DOI: [10.1902/jop.2010.090706](#)]
- 26 **Koo TK**, Li MY. A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for Reliability Research. *J Chiropr Med* 2016; **15**: 155-163 [PMID: [27330520](#) DOI: [10.1016/j.jcm.2016.02.012](#)]
- 27 **De Rouck T**, Colls K, Cosyn J. Single-tooth replacement in the anterior maxilla by means of immediate implantation and provisionalization: a review. *Int J Oral Maxillofac Implants* 2008; **23**: 897-904 [PMID: [19014160](#)]
- 28 **Urban T**, Kostopoulos L, Wenzel A. Immediate implant placement in molar regions: risk factors for early failure. *Clin Oral Implants Res* 2012; **23**: 220-227 [PMID: [21457353](#) DOI: [10.1111/j.1600-0501.2011.02167.x](#)]
- 29 **Chen H**, Wang W, Gu X. Three-dimensional alveolar bone assessment of mandibular molars for immediate implant placement: a virtual implant placement study. *BMC Oral Health* 2021; **21**: 478 [PMID: [34579702](#) DOI: [10.1186/s12903-021-01849-w](#)]
- 30 **Albrektsson T**, Wennerberg A. On osseointegration in relation to implant surfaces. *Clin Implant Dent Relat Res* 2019; **21** Suppl 1: 4-7 [PMID: [30816639](#) DOI: [10.1111/cid.12742](#)]
- 31 **Buser D**, Sennerby L, De Bruyn H. Modern implant dentistry based on osseointegration: 50 years of progress, current trends and open questions. *Periodontol* 2000 2017; **73**: 7-21 [PMID: [28000280](#) DOI: [10.1111/prd.12185](#)]
- 32 **Chen ST**, Darby IB, Reynolds EC. A prospective clinical study of non-submerged immediate implants: clinical outcomes and esthetic results. *Clin Oral Implants Res* 2007; **18**: 552-562 [PMID: [17608739](#) DOI: [10.1111/j.1600-0501.2007.01388.x](#)]
- 33 **Zhang J**, Huang X, Lu B, Zhang C, Cai Z. Can apical periodontitis affect serum levels of CRP, IL-2, and IL-6 as well as induce pathological changes in remote organs? *Clin Oral Investig* 2016; **20**: 1617-1624 [PMID: [26556576](#) DOI: [10.1007/s00784-015-1646-6](#)]
- 34 **Taheri M**, Akbari S, Shamschiri AR, Shayesteh YS. Marginal bone loss around bone-level and tissue-level implants: A systematic review and meta-analysis. *Ann Anat* 2020; **231**: 151525 [PMID: [32380195](#) DOI: [10.1016/j.aanat.2020.151525](#)]
- 35 **Kakar A**, Kakar K, Leventis MD, Jain G. Immediate Implant Placement in Infected Sockets: A Consecutive Cohort Study. *J Lasers Med Sci* 2020; **11**: 167-173 [PMID: [32273958](#) DOI: [10.34172/jlms.2020.28](#)]
- 36 **Hattingh A**, De Bruyn H, Van Weehaeghe M, Hommez G, Vandeweghe S. Contour Changes Following Immediate Placement of Ultra-Wide Implants in Molar Extraction Sockets without Bone Grafting. *J Clin Med* 2020; **9** [PMID: [32759638](#) DOI: [10.3390/jcm9082504](#)]
- 37 **Naji BM**, Abdelsamea SS, Alqutaibi AY, Said Ahmed WM. Immediate dental implant placement with a horizontal gap more than two millimetres: a randomized clinical trial. *Int J Oral Maxillofac Surg* 2021; **50**: 683-690 [PMID: [32951965](#) DOI: [10.1016/j.ijom.2020.08.015](#)]
- 38 **Botticelli D**, Berglundh T, Buser D, Lindhe J. The jumping distance revisited: An experimental study in the dog. *Clin Oral Implants Res* 2003; **14**: 35-42 [PMID: [12562363](#) DOI: [10.1034/j.1600-0501.2003.140105.x](#)]
- 39 **Araujo-Pires AC**, Francisconi CF, Bigueti CC, Cavalla F, Aranha AM, Letra A, Trombone AP, Faveri M, Silva RM, Garlet GP. Simultaneous analysis of T helper subsets (Th1, Th2, Th9, Th17, Th22, Tfh, Tr1 and Tregs) markers expression in periapical lesions reveals multiple cytokine clusters accountable for lesions activity and inactivity status. *J Appl Oral Sci* 2014; **22**: 336-346 [PMID: [25141207](#) DOI: [10.1590/1678-775720140140](#)]
- 40 **Maeda K**, Kobayashi Y, Koide M, Uehara S, Okamoto M, Ishihara A, Kayama T, Saito M, Marumo K. The Regulation of Bone Metabolism and Disorders by Wnt Signaling. *Int J Mol Sci* 2019; **20** [PMID: [31698687](#) DOI: [10.3390/ijms20225525](#)]
- 41 **Jakovljevic A**, Knezevic A, Karalic D, Soldatovic I, Popovic B, Milasin J, Andric M. Pro-inflammatory cytokine levels in human apical periodontitis: Correlation with clinical and histological findings. *Aust Endod J* 2015; **41**: 72-77 [PMID: [25163634](#) DOI: [10.1111/aej.12072](#)]
- 42 **Larsen T**, Fiehn NE. Dental biofilm infections - an update. *APMIS* 2017; **125**: 376-384 [PMID: [28407420](#) DOI: [10.1111/apm.12688](#)]
- 43 **Mertowska P**, Mertowski S, Wojnicka J, Korona-Główniak I, Grywalska E, Błażewicz A, Załuska W. A Link between Chronic Kidney

- Disease and Gut Microbiota in Immunological and Nutritional Aspects. *Nutrients* 2021; **13** [PMID: 34684638 DOI: 10.3390/nu13103637]
- 44 **Carrillo C**, Peñarrocha M, Vera F, Peñarrocha D. Immunohistochemical study of Langerhans cells in periapical lesions: correlation with inflammatory cell infiltration and epithelial cell proliferation. *Med Oral Patol Oral Cir Bucal* 2010; **15**: e335-e339 [PMID: 20038917]
- 45 **Oseko F**, Yamamoto T, Akamatsu Y, Kanamura N, Iwakura Y, Imanishi J, Kita M. IL-17 is involved in bone resorption in mouse periapical lesions. *Microbiol Immunol* 2009; **53**: 287-294 [PMID: 19457170 DOI: 10.1111/j.1348-0421.2009.00123.x]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

