

# World Journal of *Clinical Cases*

*World J Clin Cases* 2023 September 26; 11(27): 6318-6669



**MINIREVIEWS**

- 6318 Characteristics of amino acid metabolism in colorectal cancer

*Xu F, Jiang HL, Feng WW, Fu C, Zhou JC*

**ORIGINAL ARTICLE****Clinical and Translational Research**

- 6327 Exploring the pharmacological mechanism of Wuzhuyu decoction on hepatocellular carcinoma using network pharmacology

*Ouyang JY, Lin WJ, Dong JM, Yang Y, Yang HK, Zhou ZL, Wang RQ*

- 6344 Identification of potential diagnostic and prognostic biomarkers for breast cancer based on gene expression omnibus

*Zhang X, Mi ZH*

**Retrospective Cohort Study**

- 6363 Treatment of proximal humeral fractures accompanied by medial calcar fractures using fibular autografts: A retrospective, comparative cohort study

*Liu N, Wang BG, Zhang LF*

**Retrospective Study**

- 6374 Effectiveness of out-fracture of the inferior turbinate with reduction nasal bone fracture

*Kim SY, Nam HJ, Byeon JY, Choi HJ*

- 6383 Prognostic model of hepatocellular carcinoma based on cancer grade

*Zhang GX, Ding XS, Wang YL*

- 6398 Oncologic efficacy of gonadotropin-releasing hormone agonist in hormone receptor-positive very young breast cancer patients treated with neoadjuvant chemotherapy

*Choi HJ, Lee JH, Jung CS, Ryu JM, Chae BJ, Lee SK, Yu JH, Kim SW, Nam SJ, Lee JE, Jung YJ, Kim HY*

- 6407 Correlation analysis of serum thyroglobulin, thyroid-stimulating hormone levels, and thyroid-cancer risk in thyroid nodule surgery

*Shuai JH, Leng ZF, Wang P, Ji YC*

- 6415 Closed thoracic drainage in elderly patients with chronic obstructive pulmonary disease complicated with spontaneous pneumothorax: A retrospective study

*Wang W, Zhu DN, Shao SS, Bao J*

**Observational Study**

- 6424 *Helicobacter pylori* eradication treatment for primary gastric diffuse large B-cell lymphoma: A single-center analysis

*Saito M, Mori A, Kajikawa S, Yokoyama E, Kanaya M, Izumiyama K, Morioka M, Kondo T, Tanei ZI, Shimizu A*

**Prospective Study**

- 6431** Effect of polyene phosphatidylcholine/ursodeoxycholic acid/ademetionine on pregnancy outcomes in intrahepatic cholestasis  
*Dong XR, Chen QQ, Xue ML, Wang L, Wu Q, Luo TF*

**SYSTEMATIC REVIEWS**

- 6440** Maternal diaphragmatic hernia in pregnancy: A systematic review with a treatment algorithm  
*Augustin G, Kovač D, Karadjole VS, Zajec V, Herman M, Hrbač P*

**META-ANALYSIS**

- 6455** Laparoscopic *vs* open radical resection in management of gallbladder carcinoma: A systematic review and meta-analysis  
*He S, Yu TN, Cao JS, Zhou XY, Chen ZH, Jiang WB, Cai LX, Liang X*

**CASE REPORT**

- 6476** Acute acquired concomitant esotropia with congenital paralytic strabismus: A case report  
*Zhang MD, Liu XY, Sun K, Qi SN, Xu CL*
- 6483** Tumor recurrence after pathological complete response in locally advanced gastric cancer after neoadjuvant therapy: Two case reports  
*Xing Y, Zhang ZL, Ding ZY, Song WL, Li T*
- 6491** Acute peritonitis secondary to post-traumatic appendicitis: A case report and literature review  
*Habachi G, Aziza B, Ben-Ammar S, Maherzi O, Houas Y, Kerkeni Y, Sahli S, Jouini R*
- 6498** Fournier's gangrene after insertion of thermo-expandable prostatic stent for benign prostatic hyperplasia: A case report  
*Jung HC, Kim YU*
- 6505** Methyl-CpG-Binding protein 2 duplication syndrome in a Chinese patient: A case report and review of the literature  
*Xing XH, Takam R, Bao XY, Ba-alwi NA, Ji H*
- 6515** Blood purification for treatment of non-liquefied multiple liver abscesses and improvement of T-cell function: A case report  
*Tang ZQ, Zhao DP, Dong AJ, Li HB*
- 6523** Eosinophilic granulomatosis with polyangiitis, asthma as the first symptom, and subsequent Loeffler endocarditis: A case report  
*He JL, Liu XY, Zhang Y, Niu L, Li XL, Xie XY, Kang YT, Yang LQ, Cai ZY, Long H, Ye GF, Zou JX*
- 6531** Left atrium veno-arterial extra corporeal membrane oxygenation as temporary mechanical support for cardiogenic shock: A case report  
*Lamastra R, Abbott DM, Degani A, Pellegrini C, Veronesi R, Pelenghi S, Dezza C, Gazzaniga G, Belliato M*

- 6537** Successful treatment of eyebrow intradermal nevi by shearing combined with electrocautery and curettage: Two case reports  
*Liu C, Liang JL, Yu JL, Hu Q, Li CX*
- 6543** Amniotic membrane mesenchymal stromal cell-derived secretome in the treatment of acute ischemic stroke: A case report  
*Lin FH, Yang YX, Wang YJ, Subbiah SK, Wu XY*
- 6551** Managing spindle cell sarcoma with surgery and high-intensity focused ultrasound: A case report  
*Zhu YQ, Zhao GC, Zheng CX, Yuan L, Yuan GB*
- 6558** Triplet regimen as a novel modality for advanced unresectable hepatocellular carcinoma: A case report and review of literature  
*Zhao Y, He GS, Li G*
- 6565** Acute diquat poisoning case with multiorgan failure and a literature review: A case report  
*Fan CY, Zhang CG, Zhang PS, Chen Y, He JQ, Yin H, Gong XJ*
- 6573** Fungal corneal ulcer after repair of an overhanging filtering bleb: A case report  
*Zhao J, Xu HT, Yin Y, Li YX, Zheng YJ*
- 6579** Combination therapy with toripalimab and anlotinib in advanced esophageal squamous cell carcinoma: A case report  
*Chen SC, Ma DH, Zhong JJ*
- 6587** Removal of a pulmonary artery foreign body during pulse ablation in a patient with atrial fibrillation: A case report  
*Yan R, Lei XY, Li J, Jia LL, Wang HX*
- 6592** Delayed-onset *micrococcus luteus*-induced postoperative endophthalmitis several months after cataract surgery: A case report  
*Nam KY, Lee HW*
- 6597** Anesthetic management of a pregnant patient with Eisenmenger's syndrome: A case report  
*Zhang Y, Wei TT, Chen G*
- 6603** Recurrence of unilateral angioedema of the tongue: A case report  
*Matsuhisa Y, Kenzaka T, Shimizu H, Hirose H, Gotoh T*
- 6613** Transverse mesocolic hernia with intestinal obstruction as a rare cause of acute abdomen in adults: A case report  
*Zhang C, Guo DF, Lin F, Zhan WF, Lin JY, Lv GF*
- 6618** Compound heterozygous mutations in tripeptidyl peptidase 1 cause rare autosomal recessive spinocerebellar ataxia type 7: A case report  
*Liu RH, Wang XY, Jia YY, Wang XC, Xia M, Nie Q, Guo J, Kong QX*

- 6624** Treatment of posterior interosseous nerve entrapment syndrome with ultrasound-guided hydrodissection: A case report  
*Qin LH, Cao W, Chen FT, Chen QB, Liu XX*
- 6631** Rapidly growing extensive polypoid endometriosis after gonadotropin-releasing hormone agonist discontinuation: A case report  
*Zhang DY, Peng C, Huang Y, Cao JC, Zhou YF*
- 6640** Preserving finger length in a patient with symmetric digital gangrene under local anesthesia: A case report  
*Kim KH, Ko IC, Kim H, Lim SY*
- 6646** Reconstruction of the lower back wound with delayed infection after spinal surgery: A case report  
*Kim D, Lim S, Eo S, Yoon JS*
- 6653** Solitary intraosseous neurofibroma in the mandible mimicking a cystic lesion: A case report and review of literature  
*Zhang Z, Hong X, Wang F, Ye X, Yao YD, Yin Y, Yang HY*
- 6664** Complete response of metastatic *BRAF* V600-mutant anaplastic thyroid cancer following adjuvant dabrafenib and trametinib treatment: A case report  
*Lee SJ, Song SY, Kim MK, Na HG, Bae CH, Kim YD, Choi YS*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Alexandru Corlateanu, MD, PhD, Reader (Associate Professor), Department of Respiratory Medicine, Nicolae Testemitanu State University of Medicine and Pharmacy, Chisinau 2001, Moldova. alexandru\_corlateanu@yahoo.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

September 26, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



## Anesthetic management of a pregnant patient with Eisenmenger's syndrome: A case report

Ying Zhang, Ting-Ting Wei, Gang Chen

**Specialty type:** Medicine, research and experimental

**Provenance and peer review:** Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B, B  
Grade C (Good): 0  
Grade D (Fair): D  
Grade E (Poor): 0

**P-Reviewer:** Ait Addi R, Morocco; Higa K, Japan

**Received:** June 24, 2023

**Peer-review started:** June 24, 2023

**First decision:** July 28, 2023

**Revised:** August 8, 2023

**Accepted:** August 17, 2023

**Article in press:** August 17, 2023

**Published online:** September 26, 2023



**Ying Zhang, Ting-Ting Wei, Gang Chen,** Department of Anesthesiology, Sir Run Run Shaw Hospital, School of Medicine, Zhejiang University, Hangzhou 310006, Zhejiang Province, China

**Corresponding author:** Gang Chen, PhD, Chief Doctor, Department of Anesthesiology, Sir Run Run Shaw Hospital, School of Medicine, Zhejiang University, No. 3 Qingchun Road, Hangzhou 310006, Zhejiang Province, China. [chengang120@zju.edu.cn](mailto:chengang120@zju.edu.cn)

### Abstract

#### BACKGROUND

Eisenmenger's syndrome (ES) is a rare complication of congenital heart disease that includes pulmonary artery hypertension and reversed or bidirectional shunts. The mortality rate of pregnant women with ES is 30%-70% due to pathophysiological deterioration. Successful perioperative management of a pregnant patient with ES is a challenge for anesthesiologists.

#### CASE SUMMARY

A 38-year-old pregnant woman was admitted to the cardiology department of our hospital at 22 wk of gestation with complaints of chest tightness and shortness of breath for 3 wk. Transthoracic echocardiography revealed a bidirectional shunt between the descending aorta and pulmonary artery after interventional closure of the patent ductus arteriosus and severe pulmonary hypertension. ES in pregnancy was our primary suspicion. The patient elected to terminate the pregnancy under adequate preoperative preparation, rigorous intraoperative monitoring, and perfect epidural anesthesia. She was discharged successfully on postoperative day 16.

#### CONCLUSION

Our experience in this case suggests that successful outcomes are possible in pregnant patients with ES for termination of pregnancy under epidural anesthesia and intensive monitoring.

**Key Words:** Eisenmenger's syndrome; Pulmonary artery hypertension; Pregnancy; Monitoring; Epidural anesthesia; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Eisenmenger's syndrome (ES) is a rare complication of congenital heart disease, with a high maternal mortality of 30%-70%. We present a rare case of a pregnant woman with ES after cardiac surgery. The patient elected to terminate the pregnancy under adequate preoperative preparation, rigorous intraoperative monitoring, and perfect epidural anesthesia. Our experience in this case suggests that successful outcomes are possible in pregnant patients with ES for termination of pregnancy under epidural anesthesia and intensive monitoring.

**Citation:** Zhang Y, Wei TT, Chen G. Anesthetic management of a pregnant patient with Eisenmenger's syndrome: A case report. *World J Clin Cases* 2023; 11(27): 6597-6602

**URL:** <https://www.wjgnet.com/2307-8960/full/v11/i27/6597.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v11.i27.6597>

## INTRODUCTION

Eisenmenger's syndrome (ES) is a rare complication of congenital heart disease during pregnancy[1]. It manifests as pulmonary artery hypertension and a right-to-left or bidirectional shunt secondary to septal defects or patent ductus arteriosus (PDA)[2]. As the mortality rate is as high as 30%-70%, pregnant women with ES are advised to terminate their pregnancy[3,4]. In recent years, advanced medical technology has allowed congenital heart disease patients to undergo corrective surgery, and some patients have decided to conceive. Although there have been a few successful births, it is still recommended to avoid pregnancy after cardiac correction due to the high mortality and maternal complication rates [5]. This case report describes a pregnant patient with ES after cardiac surgery and discusses the safety and efficacy of epidural anesthesia and intensive monitoring for termination of pregnancy.

## CASE PRESENTATION

### Chief complaints

A 38-year-old 62-kg pregnant woman (gravida 4, para 0) was admitted to the cardiology department of our hospital at 22 wk of gestation with complaints of chest tightness and shortness of breath for 3 wk.

### History of present illness

The patient's symptoms started 3 wk prior with chest tightness and shortness of breath. The degree of heart failure was New York Heart Association grade II.

### History of past illness

At 25 years of age, the patient had a history of transcatheter closure of a PDA in a local hospital. The patient stated that the PDA was completely closed by the transcatheter interventional procedure, and she attended irregular follow-up in the local hospital after the operation.

### Personal and family history

Between 2010 and 2020, the patient experienced three spontaneous abortions in the first trimester without any pregnancy discomfort. No similar family history was noted.

### Physical examination

After admission to our hospital, the patient's temperature was 36.2°C, heart rate was 75 bpm, respiratory rate was 19 breaths per minute, blood pressure was 92/53 mmHg, and oxygen saturation in an oxygen concentration of 33% was 97%. The clinical examination revealed mild cyanosis of the lips and a loud P2 systolic murmur in the second to fourth intercostal spaces of the left margin of the sternum.


### Laboratory examinations

Blood analysis revealed a mildly elevated hemoglobin level of 15.3 g/dL, with a normal hematocrit of 44.4% and a platelet count of  $1.22 \times 10^{11}/L$ . The prothrombin time was normal, and the partial thromboplastin time was slightly decreased, at 27.6 s. Electrocardiography (ECG) showed an abnormal Q wave in the high lateral border, slight ST-segment change in the lower border, right axis deviation, and clockwise transposition. Arterial blood gas analysis revealed pH: 7.47, PCO<sub>2</sub>: 28.2 mmHg, PO<sub>2</sub>: 85.3 mmHg, and SaO<sub>2</sub>: 97.4% at 21% oxygen concentration.

### Imaging examinations

Transthoracic echocardiography (TTE) revealed the occlusion (manifesting as a stronger echo) between the descending aorta and pulmonary artery (Figure 1A). Color Doppler echocardiography revealed a bidirectional shunt signal on both sides of the occlusion (Figure 1B). Spectral Doppler echocardiography revealed the pressure gradient (PG) and blood





DOI: 10.12998/wjcc.v11.i27.6597 Copyright ©The Author(s) 2023.

**Figure 1 Preoperative echocardiography.** A: Image showing the occlusion (manifesting as a stronger echo) between the descending aorta and pulmonary artery (orange arrow); B: Color Doppler echocardiography revealed a bidirectional shunt signal on both sides of the occlusion (orange arrow); C: Spectral Doppler echocardiography revealed the pressure gradient and blood velocity of the bidirectional shunt (orange arrow). Vel: Velocity; PG: Pressure gradient.

velocity of the bidirectional shunt (Figure 1C). The pulmonary artery systolic pressure (PASP) was estimated at 117 mmHg, and the brachial systolic pressure at 124 mmHg.

## FINAL DIAGNOSIS

The final diagnosis of this case was pregnancy complicated with ES.

## TREATMENT


On day 2 after her admission, the patient underwent surgery to terminate the pregnancy. In the operating room, the patient was supplemented with oxygen *via* a nasal tube. Ringer's lactate (500 mL) was slowly administered during the procedure. Standard monitoring, including SpO<sub>2</sub>, noninvasive blood pressure, and ECG, was established. Under ultrasonographic guidance and local anesthesia, an arterial line, central venous catheter, and Swan-Ganz catheter were placed for pressure monitoring and venous access. In the right lateral position, an 18-G epidural needle was inserted in the L1-L2 intervertebral space, and a 22-G epidural catheter was introduced 4 cm toward the cephalad direction in the epidural space. A 3 mL test dose of 2% lignocaine was administered, and the patient was placed in the supine position. Over a period of 15 min, a total of 9 mL of 0.75% ropivacaine was added gradually to achieve the anesthetic effect. After anesthesia, norepinephrine was given to achieve a systemic blood pressure equal to the pulmonary pressure. Over a period of 15 min, a block height of T4 was obtained in the patient, as assessed using cold stimulation and touch. There were no significant changes in her hemodynamics. The central venous pressure (CVP) was kept at 10-12 mmHg throughout the surgery. The pulmonary artery pressure was always kept lower than the systemic circulation pressure to avoid right-to-left shunt aggravation. The systemic artery pressure was kept at 110-130/60-70 mmHg, the pulmonary pressure at 100-110/50-55 mmHg, and SpO<sub>2</sub> at 98%-100% during the procedure. The procedure proceeded uneventfully, without pain or discomfort. The blood loss was estimated to be 300 mL. At the end of the operation, 1.5 mg morphine was administered epidurally for postoperative analgesia. Since we planned postoperative heparin anticoagulation, the epidural catheter was removed before the patient left the operating room to avoid epidural hematoma and other complications. After surgery, the patient was transferred to the intensive care unit for further monitoring. She was treated with alprostadil for pulmonary hypertension and heparin anticoagulation.

## OUTCOME AND FOLLOW-UP

The patient had stable hemodynamics and was transferred to the ward 6 d after surgery. The patient was discharged on the 16<sup>th</sup> day after surgery. At the follow-up visit one month after hospital discharge, the patient was asymptomatic, and a new TTE showed moderate pulmonary hypertension with a residual bidirectional shunt, which was mainly from left to right (Figure 2). The patient was advised to avoid pregnancy and to attend regular follow-up for her cardiac disease.

## DISCUSSION

In the progression of congenital heart disease, the pulmonary artery pressure exceeds the systemic circulation pressure due to the gradual increase in pulmonary vascular resistance (PVR), resulting in a bidirectional or reverse shunt, which is



DOI: 10.12998/wjcc.v11.i27.6597 Copyright ©The Author(s) 2023.

**Figure 2** Follow-up echocardiography one month after hospital discharge. A: Color Doppler echocardiography revealed only a minor bidirectional shunt signal on both sides of the occlusion (orange arrow); B: Spectral Doppler echocardiography revealed a residual bidirectional shunt, which mainly was from left to right (orange arrow).

defined as ES. Pregnant women with ES have a mortality rate as high as 30%-70% [3]. The main cause of death in such patients is pregnancy-induced decreased systemic resistance and increased cardiac afterload that exacerbates the right-to-left shunting, leading to pulmonary hypertension, right heart failure, and hypoxemia [6]. Therefore, experts unanimously recommend contraception for women with ES and, once pregnant, early termination of pregnancy.

Guidelines for the management of cardiovascular disease in pregnancy indicate that women with uncomplicated congenital heart disease have no risk or a slightly increased risk of maternal mortality after pregnancy. However, women with congenital heart disease combined with ES have a high mortality due to unbearable hemodynamic changes [7]. The factors related to maternal mortality are severe pulmonary hypertension, ES, and reduced right ventricular systolic function. A previous study showed significantly higher mortality in patients with severe pulmonary hypertension and ES [8]. Therefore, patients with congenital heart disease combined with ES are advised to avoid pregnancy and terminate any pregnancy quickly. In this case, the patient had a history of transcatheter closure of a PDA in a local hospital. She was advised to attend regular follow-up to determine her eligibility for pregnancy. At the last follow-up, she was not diagnosed with pulmonary hypertension or ES. Until the symptoms of chest tightness, she did not undergo cardiac echocardiography, and she did not have any uncomfortable symptoms during early pregnancy. Considering the high mortality rate of continued pregnancy, timely termination of pregnancy may be the best choice for pregnant women with ES.

The primary anesthetic goal in a patient with ES is to maintain cardiac output and systemic vascular resistance (SVR) and avoid any hemodynamic changes that might increase PVR and exacerbate the right-to-left shunt [9,10]. General anesthesia has clear risks and disadvantages. First, catecholamine release may increase the PVR during laryngoscopy and intubation [11]. Second, positive-pressure ventilation during general anesthesia reduces venous return and cardiac output. Third, general anesthesia increases the risk of reflux aspiration in pregnant women. In addition, induction drugs reduce myocardial contractility and SVR. Epidural anesthesia has been successfully used in pregnant patients with ES [11,12]. Epidural anesthesia reduces catecholamine levels and PVR by sympathetic blockade. Epidural anesthesia is preferable to spinal anesthesia because of its slow onset, which avoids abrupt changes in hemodynamics that increase the right-to-left shunting. Furthermore, the epidural catheter provides assurance for further anesthesia and postoperative analgesia. We consider epidural anesthesia while keeping the hemodynamics as close to normal as possible to be the best choice for these patients.

Rapid hemodynamic changes in pregnant women with ES can lead to a pulmonary hypertension crisis, right heart failure, hypoxemia, and even sudden death. The purpose of monitoring is to detect these changes early and provide timely intervention and treatment. However, the placement of invasive monitoring equipment increases the risk of infection and thromboembolism. Considering the risks and benefits, we placed an arterial catheter, central venous catheter, and pulmonary arterial catheter in this patient. Invasive blood pressure monitoring facilitates real-time supervision of hemodynamic changes. CVP monitoring helps with fluid guidance and medication use. We found that the right-to-left shunt was minimized when the CVP was maintained with an appropriate preload (CVP 10–12 mmHg). The pulmonary artery pressure was monitored to ensure a systemic resistance equal to the pulmonary artery pressure and to minimize the right-to-left shunt. The patient had severe pulmonary artery hypertension before surgery (PASP 117 mmHg). Keeping her SVR higher than her PVR was the key to reducing the right-to-left shunt. Transesophageal echocardiography is uncomfortable for awake patients. TTE is widely used for the diagnosis of ES because it is noninvasive and readily available. Intraoperative echocardiography aids in assessing shunts, pulmonary artery pressure, and right ventricular function [13]. TTE showed that the PASP decreased from 117 mmHg preoperatively (R-L: V: 2.02 m/s; PG: 16 mmHg) to 78 mmHg postoperatively (R-L: V: 1.18 m/s; PG: 6 mmHg). Therefore, early termination of pregnancy is a correct and effective decision for patients with ES.

Epidural anesthesia reduces the SVR by blocking the sympathetic nerves. Norepinephrine decreases the SVR during anesthesia because  $\beta$ -agonism increases myocardial contractility and cardiac output [14]. In this case, epidural anesthesia combined with norepinephrine successfully maintained the hemodynamic stability as close to normal as possible.

We chose not to administer oxytocin because it causes direct vasodilation and reduces the SVR, with a compensatory increase in the heart rate and cardiac output. Considering that the patient's preoperative coagulation function and platelets were normal, an epidural catheter was placed during the operation. Therefore, heparin was used for anticoagu-

lation after surgery.

## CONCLUSION

Here, we present a rare case of a pregnant woman with ES after cardiac surgery. ES is a rare complication of congenital heart disease with high maternal mortality. Our experience in this case suggests that successful outcomes are possible in pregnant patients with ES under epidural anesthesia and intensive monitoring.

## ACKNOWLEDGEMENTS

The authors thank Ji-Hong Zhu for assistance with figure preparation.

## FOOTNOTES

**Author contributions:** Zhang Y and Chen G reviewed the literature and contributed to manuscript drafting; Zhang Y and Wei TT analyzed and interpreted the imaging findings; Zhang Y, Wei TT, and Chen G were responsible for the revision of the manuscript for important intellectual content; all authors issued final approval for the version to be submitted.

**Informed consent statement:** Informed written consent was obtained from the patient for publication of this report and any accompanying images.

**Conflict-of-interest statement:** The authors declare that they have no conflict of interest to disclose.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Ying Zhang 0000-0001-8666-7276; Ting-Ting Wei 0000-0002-5540-8391; Gang Chen 0000-0003-4262-9491.

**S-Editor:** Lin C

**L-Editor:** Wang TQ

**P-Editor:** Cai YX

## REFERENCES

- 1 **Wander G**, van der Zande JA, Patel RR, Johnson MR, Roos-Hesselink J. Pregnancy in women with congenital heart disease: a focus on management and preventing the risk of complications. *Expert Rev Cardiovasc Ther* 2023; **21**: 587-599 [PMID: 37470417 DOI: 10.1080/14779072.2023.2237886]
- 2 **Yuan SM**. Eisenmenger Syndrome in Pregnancy. *Braz J Cardiovasc Surg* 2016; **31**: 325-329 [PMID: 27849306 DOI: 10.5935/1678-9741.20160062]
- 3 **Gurumurthy T**, Hegde R, Mohandas B. Anaesthesia for a patient with Eisenmenger's syndrome undergoing caesarean section. *Indian J Anaesth* 2012; **56**: 291-294 [PMID: 22923831 DOI: 10.4103/0019-5049.98780]
- 4 **Katsurahagi S**, Kamiya C, Yamanaka K, Neki R, Miyoshi T, Iwanaga N, Horiuchi C, Tanaka H, Yoshimatsu J, Niwa K, Takagi Y, Ogo T, Nakanishi N, Ikeda T. Maternal and fetal outcomes in pregnancy complicated with Eisenmenger syndrome. *Taiwan J Obstet Gynecol* 2019; **58**: 183-187 [PMID: 30910135 DOI: 10.1016/j.tjog.2019.01.002]
- 5 **Rudienė V**, Kaplerienė L, Jančauskaitė D, Meškėnė E, Palevičiūtė E, Laukytė-Slėnienė M, Gasiūnaitė D, Ramašauskaitė D, Jurevičienė E, Gumbienė L. Pregnancy in Congenital Heart Disease, Complicated by Pulmonary Arterial Hypertension-A Challenging Issue for the Pregnant Woman, the Foetus, and Healthcare Professionals. *Medicina (Kaunas)* 2022; **58** [PMID: 35454315 DOI: 10.3390/medicina58040476]
- 6 **Dachlan EG**, Amirah, Cininta N, Pranadyan R, Putri AY, Oktaviono YH, Akbar MIA. High Maternal Neonatal Mortality and Morbidity in Pregnancy with Eisenmenger Syndrome. *J Pregnancy* 2021; **2021**: 3248850 [PMID: 34616573 DOI: 10.1155/2021/3248850]
- 7 **Regitz-Zagrosek V**, Roos-Hesselink JW, Bauersachs J, Blomström-Lundqvist C, Cifková R, De Bonis M, Lung B, Johnson MR, Kintscher U, Kranke P, Lang IM, Morais J, Pieper PG, Presbitero P, Price S, Rosano GMC, Seeland U, Simoncini T, Swan L, Warnes CA; ESC Scientific Document Group. 2018 ESC Guidelines for the management of cardiovascular diseases during pregnancy. *Eur Heart J* 2018; **39**: 3165-3241 [PMID: 30165544 DOI: 10.1093/eurheartj/ehy340]
- 8 **Hartopo AB**, Anggrahini DW, Nurdianti DS, Emoto N, Dinarti LK. Severe pulmonary hypertension and reduced right ventricle systolic

- function associated with maternal mortality in pregnant uncorrected congenital heart diseases. *Pulm Circ* 2019; **9**: 2045894019884516 [PMID: 31798834 DOI: 10.1177/2045894019884516]
- 9 **Bennett JM**, Ehrenfeld JM, Markham L, Eagle SS. Anesthetic management and outcomes for patients with pulmonary hypertension and intracardiac shunts and Eisenmenger syndrome: a review of institutional experience. *J Clin Anesth* 2014; **26**: 286-293 [PMID: 24908593 DOI: 10.1016/j.jclinane.2013.11.022]
- 10 **Lopez BM**, Malhamé I, Davies LK, Gonzalez Velez JM, Marelli A, Rabai F. Eisenmenger Syndrome in Pregnancy: A Management Conundrum. *J Cardiothorac Vasc Anesth* 2020; **34**: 2813-2822 [PMID: 32381307 DOI: 10.1053/j.jvca.2020.02.053]
- 11 **Mishra L**, Pani N, Samantaray R, Nayak K. Eisenmenger's syndrome in pregnancy: Use of epidural anesthesia and analgesia for elective cesarean section. *J Anaesthesiol Clin Pharmacol* 2014; **30**: 425-426 [PMID: 25190960 DOI: 10.4103/0970-9185.137286]
- 12 **Gangakhedkar GR**, Chhabria RD, Gour SD, Palani Y. Emergency caesarean section of a patient with eisenmenger's syndrome: A tight-rope walk. *Ann Card Anaesth* 2020; **23**: 95-97 [PMID: 31929258 DOI: 10.4103/aca.ACA\_175\_18]
- 13 **Topyla-Putowska W**, Tomaszewski M, Wysokiński A, Tomaszewski A. Echocardiography in Pulmonary Arterial Hypertension: Comprehensive Evaluation and Technical Considerations. *J Clin Med* 2021; **10** [PMID: 34362015 DOI: 10.3390/jcm10153229]
- 14 **Fischer MU**, Priebe HJ. Anaesthetic management for hip arthroplasty in a 46-yr-old patient with uncorrected truncus arteriosus type IV. *Br J Anaesth* 2006; **97**: 329-332 [PMID: 16798771 DOI: 10.1093/bja/acl148]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

