

World Journal of *Clinical Cases*

World J Clin Cases 2023 October 16; 11(29): 6974-7260



MINIREVIEWS

- 6974 Applications of time series analysis in epidemiology: Literature review and our experience during COVID-19 pandemic

Tomov L, Chervenkov L, Miteva DG, Batselova H, Velikova T

ORIGINAL ARTICLE

Retrospective Cohort Study

- 6984 Acute cholangitis: Does malignant biliary obstruction *vs* choledocholithiasis etiology change the clinical presentation and outcomes?

Tsou YK, Su YT, Lin CH, Liu NJ

Retrospective Study

- 6995 Usefulness of analyzing endoscopic features in identifying the colorectal serrated sessile lesions with and without dysplasia

Wang RG, Ren YT, Jiang X, Wei L, Zhang XF, Liu H, Jiang B

- 7004 Roles of biochemistry data, lifestyle, and inflammation in identifying abnormal renal function in old Chinese

Chen CH, Wang CK, Wang CY, Chang CF, Chu TW

- 7017 Clinical efficacy and safety of Guipi decoction combined with escitalopram oxalate tablets in patients with depression

Yu J, Xu FQ

- 7026 Artificial intelligence technology and ultrasound-guided nerve block for analgesia in total knee arthroplasty

Tong SX, Li RS, Wang D, Xie XM, Ruan Y, Huang L

- 7034 Axenfeld-Reiger syndrome: A search for the missing links

Morya AK, Ramesh PV, Sinha S, Nishant P, Nain N, Ramavath RN, Gone C, Prasad R

Observational Study

- 7043 Self-management of osteoarthritis while waiting for total knee arthroplasty during the COVID-19 pandemic among older Malaysians

Mahdzir ANK, Mat S, Seow SR, Abdul Rani R, Che Hasan MK, Mohamad Yahaya NH

- 7053 "In situ bone flap" combined with vascular pedicled mucous flap to reconstruction of skull base defect

Qian M, Chen X, Zhang LY, Wang ZF, Zhang Y, Wang XJ

- 7061 Reference values of gait parameters in healthy Chinese university students: A cross-sectional observational study

Yu JS, Zhuang C, Guo WX, Chen JJ, Wu XK, Xie W, Zhou X, Su H, Chen YX, Wang LK, Li WK, Tian K, Zhuang RJ

- 7075** Effect of T-regulatory cells and interleukin-35, interleukin-10, and transforming growth factor-beta on diffuse large B-cell lymphoma

Wu H, Sun HC, Ouyang GF

META-ANALYSIS

- 7082** Meta-analysis on the effectiveness of parent education for children with disabilities

Jang J, Kim G, Jeong H, Lee N, Oh S

- 7091** Meta-analysis of the efficacy and safety of daratumumab in the treatment of multiple myeloma

Wang P, Jin SY

CASE REPORT

- 7101** Varicella-zoster virus meningitis with hypoglycorrhachia: A case report

Cao LJ, Zheng YM, Li F, Hao HJ, Gao F

- 7107** Unusual presentation of penile giant condyloma acuminatum with spontaneous prepuce perforation: A case report

Hsu FC, Yu DS, Pu TW, Wu MJ, Meng E

- 7113** Primary renal lymphoma presenting as renal failure: A case report and review of literature from 1989

Lee SB, Yoon YM, Hong R

- 7127** Intravascular ultrasonography assisted carotid artery stenting for treatment of carotid stenosis: Two case reports

Fu PC, Wang JY, Su Y, Liao YQ, Li SL, Xu GL, Huang YJ, Hu MH, Cao LM

- 7136** Mucoepidermoid carcinoma of the lung with hemoptysis as initial symptom: A case report

Xie WX, Liu R, Li Z, Zhou PL, Duan LN, Fu DD

- 7144** Co-infection of *Chlamydia psittaci* and *Tropheryma whippelii*: A case report

Du ZM, Chen P

- 7150** Surgical treatment of severe anterior capsular organized hard core cataract: A case report

Wang LW, Fang SF

- 7156** First platelet transfusion refractoriness in a patient with acute myelocytic leukemia: A case report

Tu SK, Fan HJ, Shi ZW, Li XL, Li M, Song K

- 7162** Rare finding of primary aortoduodenal fistula on single-photon emission computed tomography/computed tomography of gastrointestinal bleeding: A case report

Kuo CL, Chen CF, Su WK, Yang RH, Chang YH

- 7170** Rituximab combined with Bruton tyrosine kinase inhibitor to treat elderly diffuse large B-cell lymphoma patients: Two case reports

Zhang CJ, Zhao ML

- 7179** Use of Ilizarov technique for bilateral knees flexion contracture in Juvenile-onset ankylosing spondylitis: A case report
Xia LW, Xu C, Huang JH
- 7187** Case of takotsubo cardiomyopathy after surgical treatment of liver hydatid cyst: A case report
Altaş Y, Abdullayeva Ü
- 7193** Laparoscopic choledocholithotomy and transductal T-tube insertion with indocyanine green fluorescence imaging and laparoscopic ultrasound: A case report
Yoo D
- 7200** Hematopoietic stem cell transplantation of aplastic anemia by relative with mutations and normal telomere length: A case report
Yan J, Jin T, Wang L
- 7207** Emphysematous thrombophlebitis caused by a misplaced central venous catheter: A case report
Chen N, Chen HJ, Chen T, Zhang W, Fu XY, Xing ZX
- 7214** Aggressive angiomyxoma of the epididymis: A case report
Liu XJ, Su JH, Fu QZ, Liu Y
- 7221** Gastric and intestinal ectopic pancreas: Two case reports
Zhang H, Zhao HY, Zhang FH, Liang W
- 7227** Congenital leukemia: A case report and review of literature
Yang CX, Yang Y, Zhang FL, Wang DH, Bian QH, Zhou M, Zhou MX, Yang XY
- 7234** Imaging misdiagnosis and clinical analysis of significant hepatic atrophy after bilioenteric anastomosis: A case report
Liang SY, Lu JG, Wang ZD
- 7242** Surgical treatment of mixed cervical spondylosis with spontaneous cerebrospinal fluid leakage: A case report
Yu Z, Zhang HFZ, Wang YJ
- 7248** Simultaneous thyroglossal duct cyst with parathyroid cyst: A case report
Chen GY, Li T
- 7253** Submandibular solid-cystic mass as the first and sole manifestation of occult thyroid papillary carcinoma: A case report
Chen GY, Li T

LETTER TO THE EDITOR

- 7258** Artificial intelligence and machine learning in motor recovery: A rehabilitation medicine perspective
Swarnakar R, Yadav SL

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Zeid J Khitan, FACP, FASN, MBBS, MD, Academic Research, Director, Full Professor, Department of Medicine, Marshall University, Huntington, WV 25701, United States. zkhitan@marshall.edu

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Hua-Ge Yin; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

October 16, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Usefulness of analyzing endoscopic features in identifying the colorectal serrated sessile lesions with and without dysplasia

Rui-Gang Wang, Yu-Tang Ren, Xuan Jiang, Lai Wei, Xiao-Fei Zhang, Hao Liu, Bo Jiang

Specialty type: Medicine, research and experimental

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C
Grade D (Fair): D
Grade E (Poor): 0

P-Reviewer: Kawabata H, Japan

Received: July 8, 2023

Peer-review started: July 8, 2023

First decision: August 30, 2023

Revised: September 11, 2023

Accepted: September 23, 2023

Article in press: September 25, 2023

Published online: October 16, 2023



Rui-Gang Wang, Yu-Tang Ren, Xuan Jiang, Bo Jiang, Department of Gastroenterology, Beijing Tsinghua Changgung Hospital, School of Clinical Medicine, Tsinghua University, Beijing 102218, China

Lai Wei, Center for Hepatobiliary and Pancreatic Diseases, Beijing Tsinghua Changgung Hospital, School of Clinical Medicine, Tsinghua University, Beijing 102218, China

Xiao-Fei Zhang, Center for Clinical Epidemiology and Statistics, Beijing Tsinghua Changgung Hospital, School of Clinical Medicine, Tsinghua University, Beijing 102218, China

Hao Liu, Department of Pathology, Beijing Tsinghua Changgung Hospital, School of Clinical Medicine, Tsinghua University, Beijing 102218, China

Corresponding author: Bo Jiang, MD, Chief Doctor, Chief Physician, Doctor, Department of Gastroenterology, Beijing Tsinghua Changgung Hospital, School of Clinical Medicine, Tsinghua University, No. 168 Litang Road, Changping District, Beijing 102218, China.
jba00920@btch.edu.cn

Abstract

BACKGROUND

Sessile serrated lesions (SSLs) are often missed on colonoscopy, and studies have shown this to be an essential cause of interstitial colorectal cancer. The SSLs with dysplasia (SSL-D⁺), in particular, have a faster rate of carcinogenesis than conventional tubular adenomas. Therefore, there is a clinical need for some endoscopic features with independent diagnostic value for SSL-D⁺s to assist endoscopists in making immediate diagnoses, thus improving the quality of endoscopic examination and treatment.

AIM

To compare the characteristics of SSLs, including those with and without dysplasia (SSL-D⁺ and SSL-D⁻), based on white light and image-enhanced endoscopy, to achieve an immediate differential diagnosis for endoscopists.

METHODS

From January 2017 to February 2023, cases of colorectal SSLs confirmed by colonoscopy and histopathology at the Gastrointestinal Endoscopy Center of Beijing Tsinghua Changgung Hospital were collected. The general, endoscopic, and histopathological data were reviewed and analyzed to determine the diagnostic utility.

Univariate analysis was used to find potential diagnostic factors, and then multivariate regression analysis was performed to derive endoscopic features with independent diagnostic values for the SSL-D⁺.

RESULTS

A total of 228 patients with 253 lesions were collected as a result. There were 225 cases of colorectal SSL-D's and 28 cases of SSL-D⁺s. Compared to the colorectal SSL-D, the SSL-D⁺ was more common in the right colon ($P = 0.027$) with complex patterns of depression, nodule, and elevation based on cloud-like surfaces ($P = 0.003$), reddish ($P < 0.001$), microvascular varicose ($P < 0.001$), and mixed type (Pit II, II-O, IIIL, IV) of crypt opening based on Pit II-O ($P < 0.001$). Multifactorial logistic regression analysis indicated that lesions had a reddish color [odds ratio (OR) = 18.705, 95% confidence interval (CI): 3.684-94.974], microvascular varicose (OR = 6.768, 95%CI: 1.717-26.677), and mixed pattern of crypt opening (OR = 20.704, 95%CI: 2.955-145.086) as the independent predictors for SSL-D⁺s.

CONCLUSION

The endoscopic feature that has independent diagnostic value for SSL-D⁺ is a reddish color, microvascular varicose, and mixed pattern of crypt openings.

Key Words: Sessile serrated lesions; Dysplasia; Endoscopic features

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The colonoscopic features of colorectal sessile serrated lesions (SSLs) make them easy to be overlooked in screening, which is an essential reason for the emergence of interstage colorectal cancer. With the advancement of endoscopic techniques and refinement of the serrated carcinoma pathway, the SSL is gradually being recognized by endoscopists, especially for SSL with dysplasia (SSL-D⁺), which requires extra attention. In this study, we analyzed the endoscopic features of SSLs with and without dysplasia, and found characteristics that could independently diagnose SSL-D⁺ by multifactorial analysis, which is informative for immediate diagnosis by endoscopists.

Citation: Wang RG, Ren YT, Jiang X, Wei L, Zhang XF, Liu H, Jiang B. Usefulness of analyzing endoscopic features in identifying the colorectal serrated sessile lesions with and without dysplasia. *World J Clin Cases* 2023; 11(29): 6995-7003

URL: <https://www.wjgnet.com/2307-8960/full/v11/i29/6995.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i29.6995>

INTRODUCTION

At the end of the 20th century, a group of lesions with characteristics similar to those of hyperplastic polyposis identified by Torlakovic and Snover[1], which showed a broad-based growth pattern endoscopically but lacked heterogeneous hyperplasia and were dubbed "sessile serrated adenoma (SSA)". According to the 2010 World Health Organization (WHO) Classification of Tumors of the Digestive System, the SSA/polyp (SSA/P) is a subtype of the serrated polyp. However, the 5th edition of the WHO Classification of Gastrointestinal Tumors 2019 renamed SSA/Ps to sessile serrated lesions (SSLs) due to the progression of the serrated pathway of colorectal carcinoma[2]. SSLs have poorly defined borders, pale surfaces, and a mucus-covered cap, which is difficult to differentiate from the surrounding mucosa, resulting in both their neglect during colonoscopic screening and adverse events of incomplete resection during endoscopic treatment [3,4].

The unique endoscopic features of colorectal SSLs make it a significant risk factor for the development of interstitial colorectal cancer. Notably, with the continuous advancement and development of endoscopy techniques, equipment, and accessories, potent tools exist for the precise observation and prompt diagnosis of colorectal SSLs. Based on the pathologic diagnostic criteria, SSLs were categorized as SSLs with and without dysplasia (SSL-D⁺ and SSL-D⁻). In addition, there have been reports of SSL-D⁺s transforming into submucosal invasive carcinomas within a short period[5-7], that is to say, once a diagnosis of SSL-D⁺ has been made, the progression to serrated adenocarcinoma can be rapid.

MATERIALS AND METHODS

Research object

From January 2017 to February 2023, colorectal SSLs confirmed by histopathology during colonoscopy at the Gastrointestinal Endoscopy Center of Beijing Tsinghua Changgung Hospital were retrospectively collected. Inclusion criteria: Cases with clear endoscopic images revealing the microstructure of the lesion surface, the images have been retrospectively reviewed; cases with complete clinical and pathological data; cases with a histopathologically confirmed diagnosis of SSL-D⁻ or SSL-D⁺, it is highlighted that the histopathological diagnosis of all cases has been reviewed using

WHO Classification of Gastrointestinal Tumors 2019 criteria; and all cases that signed an informed consent form and underwent endoscopic resection. The exclusion criteria included cases with unclear endoscopic figures and a non-removed mucus cap; cases unable to identify the surface microstructure; cases without a complete resection; and cases combining ulcerative colitis and hereditary polyp syndrome.

Equipment and instruments

The endoscopic systems were colonoscopy series EC-590WM/EC-790ZP (Fuji Film, Japanese) and CF-H290I/CF-HQ290I (Olympus, Japanese), which utilized white light, blue laser imaging (BLI) and narrow band imaging (NBI) to observe and store images of lesions. In addition, an endoscopic procedure was performed using ERBE 200D high-frequency electro-surgery equipment, an Enrico Woodpecker knife (EK-410D), and a Boston Scientific single-use submucosal injection needle and snare device.

Analysis methods

Suspected SSLs were analyzed by colonoscopy in white light and BLI/NBI mode with the following characteristics: Site (left/right colon), size ($</\geq 10$ mm), surface mucus cap (no/yes, [Figure 1A](#)), surface morphology (cloud-like/mixed-like: Depressed, nodular, elevated), lesion color (pale/reddish) ([Figures 1B](#) and [C](#)), crypt black spots (no/yes, [Figure 1D](#)), and microvascular varicose (no/yes, [Figure 1E](#)), crypt opening morphology (Pit II-O type/mixed type base on Pit II-O, [Figure 1F](#)). Three seasoned endoscopists (with more than ten years of gastrointestinal endoscopy operation) examined all endoscopic figures. The decision was accepted if there was unanimity among the three endoscopists; otherwise, a vote was held.

Pathological diagnostic criteria

After colorectal SSLs were resected, the lesion surface was rinsed of mucus and impurities to expose the lesion morphology and contour, spread with a specimen needle, and fixed in 10% formaldehyde solution. The specimen was collected, embedded, sectioned, stained with hematoxylin-eosin, and observed under the microscope following the guidelines for gastrointestinal mucosal specimens and pathological examination. The histological diagnosis followed the criteria outlined in the 2019 WHO Classification of Tumors of the Digestive System and the relevant literature[2,8] ([Figures 2A](#) and [B](#)).

Statistical methods

Data management and statistical analysis were performed using SPSS 20. Case counts were expressed as cases (in %). The χ^2 test and Fisher's precision probability test were used to compare univariate data. Variables with $P < 0.1$ were included in the multivariate logistic regression analysis. $P < 0.05$ indicates a statistically significant difference.

RESULTS

General data

This study collected 228 patients with histopathologically confirmed SSL, of which 119 (52.19%) were male, and 109 (47.81%) were female. The mean age of the SSL-D⁻ group was 57.25 ± 12.02 years, while the SSL-D⁺ group was 60.00 ± 11.27 years. In the enrolled cases, a total of 253 lesions were identified (two lesions in 18 patients, three lesions in two patients, and four lesions in one patient), with 225 (88.93%) cases being colorectal SSL-D⁻s and 28 (11.07%) cases being colorectal SSL-D⁺s. Three patients were detected with one SSL-D⁻ and one SSL-D⁺. Regarding gender and age, there were no statistically significant differences between the SSL-D⁻ and SSL-D⁺ groups ($P > 0.05$).

Endoscopic features

Endoscopic features of colorectal SSL-D⁻s: 146 cases located in the right colon, 186 with diameter ≥ 10 mm, 48 with complex morphology (*e.g.*, depressions, nodules, and elevations based on cloud-like), 119 covered with mucus cap, 9 with a reddish color, 130 with black dots visible on the surface of lesions in BLI/NBI pattern and 15 with visible submucosal microvascular varicose under BLI/NBI pattern. As for crypt opening, the pattern mixed type (II, II-O, IIIL, IV) based on Pit II-O type had 2 cases.

Endoscopic features of colorectal SSL-D⁺s: 24 cases located in the right colon, 27 with diameter ≥ 10 mm, 13 with complex morphology (*e.g.*, depressions, nodules, and elevations based on cloud-like), 19 covered with mucus cap, 18 with a reddish color, 16 with black dots visible on the surface of lesions in BLI/NBI pattern, and 15 with visible submucosal microvascular varicose under BLI/NBI pattern. As for crypt opening, the pattern mixed type (II, II-O, IIIL, IV) based on Pit II-O type had 15 cases.

Compared to the SSL-D⁻, the SSL-D⁺ occurred more frequently in the right colon, and its surface displayed a complex morphology of depressions, nodules, and elevations on a cloud-like surface with a reddish hue. The crypt opening was predominantly of mixed type (II, II-O, IIIL, IV), and the lesion mucosa displayed microvascular varicose. The above endoscopic features of colorectal SSL-D⁺s were statistically distinct from those of SSL-D⁻s ($P < 0.05$). In contrast, the differences were not statistically significant ($P > 0.05$) for the lesion diameter exceeding 10 mm, with or without a mucosal cap, and the presence or absence of black spots in the BLI/NBI pattern. The general and endoscopic features data as shown in [Table 1](#).

Table 1 Comparison of the general and endoscopic characteristics of the colorectal sessile serrated lesions without dysplasia and sessile serrated lesions with dysplasia group

Factors	SSL-D ⁻ group (n = 217)	SSL-D ⁺ group (n = 22)	χ^2	P value
Gender			0.001	0.981
Male	120	15		
Female	105	13		
Age	57.25 ± 12.02	60.00 ± 11.27	1.148 ²	0.252
Location			4.899	0.027
Left colon	79	4		
Right colon	146	24		
Size				0.094 ¹
< 10 mm	39	1		
≥ 10 mm	186	27		
Surface shape			8.571	0.003
Cloud-like	177	15		
Mixed-like	48	13		
Mucus cap			2.250	0.134
No	106	9		
Yes	119	19		
Color			99.6.3	< 0.001 ¹
Pale	217	10		
Reddish	8	18		
Black spots			0.004	0.949
No	95	12		
Yes	130	16		
Microvascular varicose			57.791	< 0.001 ¹
No	212	13		
Yes	13	15		
Crypt opening				< 0.001 ¹
Type II-O	223	13		
Mixed type	2	15		


¹The Fisher's exact probability method with no statistic value.²The value of *t*.SSL-D⁺: Sessile serrated lesions with dysplasia; SSL-D⁻: Sessile serrated lesions without dysplasia.**Independent diagnostic factors of SSL-D⁺**

The right colon, mixed surface morphology, reddish color, mixed crypt opening pattern, and intramucosal microvascular varicose are potential factors for the independent diagnosis of SSL-D⁺s, according to the univariate analysis. Multifactorial logistic regression analysis of the above factors revealed that the location features of the right colon ($P = 0.172$) and complex pattern of surface shape (*e.g.*, depressions, nodules, and elevations on a cloud-like basis, $P = 0.817$) were not independent diagnostic factors for SSL-D⁺s. Meanwhile, lesion of reddish color [odds ratio (OR) = 18.705, 95% confidence interval (CI): 3.684-94.974], the finding of microvascular varicose within the mucosa (OR = 6.768, 95% CI: 1.717-26.677), and crypt opening showing the mixed type (II, II-O, IIIL, IV) based on Pit II-O (OR = 20.704, 95% CI: 2.955-145.086) were independent diagnostic factors for SSL-D⁺s (Table 2).

Table 2 Multifactorial analysis of endoscopic features that have independent diagnostic value for colorectal sessile serrated lesions with dysplasias

Factors	Wals	P value	OR	95%CI
Location (left/right colon)	1.865	0.172	3.390	0.588-19.546
Size (< 10 mm/≥ 10 mm)	0.028	0.867	1.206	0.135-10.782
Surface shape (cloud-like/mixed-like)	0.053	0.817	1.204	0.250-5.803
Color (pale/reddish)	12.482	< 0.001	18.705	3.684-94.974
Varicose microvascular (No/Yes)	7.467	0.006	6.768	1.717-26.677
Crypt opening (II-O/mixed)	9.306	0.002	20.704	2.955-145.086

OR: Odds ratio; 95%CI: 95% confidence interval.




DOI: 10.12998/wjcc.v11.i29.6995 Copyright ©The Author(s) 2023.

Figure 1 Colorectal sessile serrated lesions characterized by colonoscopy demonstrate. A: White light endoscopy of the sessile serrated lesion (SSL) surface covered with mucus cap; B: White light endoscopy of the SSL with pale color and cloud-like surface; C: White light endoscopy of the SSL-dysplasia (SSL-D) against a pale color background with localized reddish mucosa and a depression visible in the center of the cloud-like surface; D: In the endoscopic blue laser imaging (BLI) mode, the dilated crypt of the SSL appears as black spots (as shown by the arrow); E: In the endoscopic magnified BLI mode, microvascular varicose is visible on the surface of the SSL-D (as shown by arrows); F: In endoscopic magnified BLI mode, the dilated crypt of the SSL-D appears as Pit III type in the background of Pit II-O.

DISCUSSION

In this study, a univariate analysis was used to find endoscopic features of potential value in the diagnosis of colorectal SSL-D's, upon which a multivariate logistic regression analysis was performed to obtain endoscopic features of independent diagnostic value in predicting SSL-D's and their diagnostic validity. This finding serves as positive guidelines for colonoscopists' immediate endoscopic diagnosis and provides a basis for the use of appropriate endoscopic therapies.

In a population-based case-control study conducted in Denmark, the risk of colorectal cancer was significantly higher in cases of colorectal SSLs than in cases of conventional adenomas[9]. Studies suggest that it takes 7-15 years for colorectal SSL-D's to progress to SSL-D's, and after 5-7 years of follow-up, 3.03%-12.5% of SSL patients develop colorectal cancer [10]. SSL-D's proliferation progresses to colorectal cancer much earlier, with cases reported previously indicating that SSL-D's progress to invasive submucosal carcinoma within 1-2 years[11-13]. This finding suggests that ignoring SSLs is likely to increase the risk of colorectal cancer in the intermediate stages. Therefore, early identification and treatment of SSLs (especially SSL-D's) is crucial. Meanwhile, among colorectal SSLs, SSL-D's are uncommon, and only 28 cases of SSL-D's are identified in this study, representing 11.07% (28/253), similar to the structure of previous studies[14].



DOI: 10.12998/wjcc.v11.i29.6995 Copyright ©The Author(s) 2023.

Figure 2 Histopathological demonstration of colorectal the sessile serrated lesion and sessile serrated lesion-dysplasia (hematoxylin and eosin, objective magnifications 20 ×, 3DHISTECH). A: The crypt has at least one of the following histologic features: (1) Horizontal growth along the mucosal muscle layer (L or inverted T-shaped crypt structure); (2) Expansion of the crypt base (basal 1/3 of the crypt base); (3) Jagged swelling of the crypt base; and (4) Asymmetric hyperplasia (proliferative band extending laterally from the base); B: Complex structural abnormalities, including: (1) Crypt elongation, crowding, complex branching, sieve-like structures, and villi-like structures; and (2) Cytologic abnormalities of diverse morphology, either with cuboidal cells, eosinophilic cytoplasm, vesicular nuclei, and prominent nucleoli, or with elongated cells, eosinophilic cytoplasm, deep-stained nuclei, and pseudostratified nuclei, with common nuclear schizophrenic signs.

The size of colorectal SSLs > 10 mm positively correlates with SSL-D⁺' emergence, but in a study that included 48 cases of SSL-D⁺s, over one-third of cases had a diameter ≤ 10 mm[15,16]. Consistent with the present study's findings, the size of colorectal SSLs is not an independent diagnostic factor for SSL-D⁺s. Mucus cap has been confirmed as the primary distinction between colorectal SSLs and hyperplastic polyps (HPs); however, it has no diagnostic value for distinguishing the SSL-D⁻ and SSL-D⁺, which is consistent with the results of this study[16,17]. With image enhancement endoscopy (*e.g.*, BLI/NBI), the crypt openings of colorectal SSLs frequently exhibit small brownish-black dots with an enlarged crypt. Considered a critical histological distinction between SSLs and HPs, but this phenomenon has no diagnostic value for SSL-D⁻s and SSL-D⁺s[17,18].

In this study, the location of the lesion in the right colon was confirmed as a potential diagnostic factor for the SSL-D⁺ ($P = 0.027$), similar to previous studies[19,20]. However, in a multifactorial regression analysis screening for independent diagnostic factors of SSL-D⁺s, the location was not found to be an independent factor ($P = 0.172$). This is likely because previous studies have focused on the sensitivity and specificity of the single factor of right hemicolectomy in predicting SSL-D⁺s without excluding the possibility that other endoscopic features interfere with its independent diagnostic value [4,15,21,22].

In addition, endoscopic examination of the colorectal SSL-D⁺ reveals the following predominant morphology: (Semi)pedunculated, double elevations, central depressions, and reddishness[16,22]. In the univariate analysis of this study, SSL-D⁺s endoscopically demonstrated predominantly complex morphology of depressions, nodules, and elevations based on cloud-like surfaces ($P = 0.003$), indicating that the mixed morphology of the lesion is a potential diagnostic factor, which is consistent with the result of previous studies. However, in the multifactorial regression analysis, the morphology of colorectal SSL-D⁺s did not have an independent diagnostic value ($P = 0.817$), which may be explained by the fact that the morphology of SSLs is particularly susceptible to the influence of the examination environment in the intestine, and when factors such as intestinal peristalsis or the amount of gas in the intestinal lumen change, so does the morphology of the lesion.

In this study, the colorectal SSL-D⁺ is more likely to exhibit a reddish color under conventional white light endoscopy, and this is an independent diagnostic factor for SSL-D⁺ according to multifactorial regression analysis (OR = 18.705, 95%CI: 3.684-94.974), similar to previous findings[15,16]. It is not difficult to understand that when SSLs with dysplasia or cancer are diagnosed, there is a corresponding increase in the demand for blood supply.

It has been established that intramucosal microvascular varicose differs from the superficial mucosal glands' surrounding microvasculature. Therefore, at sites of colorectal SSL-D⁺s, there may be dilated and irregular capillaries, and in serrated adenocarcinomas with submucosal infiltration, their surface microvasculature or structure disappears[19,21,23]. In other words, this endoscopic feature is important for suggesting an immediate diagnosis. As this study found, the rate of microvascular varicose is higher in the SSL-D⁺ group than in the SSL-D⁻ group and the differences are statistically significant ($P = 0.006$).

The use of magnified endoscopy to observe SSLs revealed that the appearance of type III, IV, and V crypt open morphologies based on type II-O is a characteristic of the SSL-D⁺, which is a high-risk marker for the progression of serrated lesions to colorectal cancer[4,19]. In this study, the SSL-D⁺ crypt opening exhibited mixed (Pit II, IIIL, and IV) type based on Pit II-O (OR = 20.704, 95%CI: 2.955-145.086), this result is in line with previous studies. This study also found that the

mixed crypt opening of SSLs also the greater diagnostic validity than the reddish color and intramucosal microvascular varicose.

The following deficiencies persist in this study. First, there is a retrospective, single-center clinical study, and there may be some bias in the endoscopists' subjective evaluations. Second, all the cases included in this study were precancerous, which may lead to bias in the study results. The reason is that if cases of SSL-D's, SSL-D's, and SSL cancerous lesions are included, it would be realized to analyze the consecutive complete endoscopic features of such lesions in different stages, and that design might be more convincing. Third, the cases enrolled in the study were discovered by different endoscopists, which inevitably affects the evaluation's consistency.

CONCLUSION

Observation allows endoscopic features of colorectal SSLs to be effectively identified. SSL-D's should be strongly suspected when lesions exhibit a reddish hue, Pit III, IV, and V mixed crypt opening patterns based on II-O and microvascular varicosity. In addition, when performing an endoscopic procedure of the SSL-D's, *en-bloc* and curative resection should be strongly ensured.

ARTICLE HIGHLIGHTS

Research background

Missing diagnosis of sessile serrated lesions (SSLs), especially SSLs with dysplasia (SSL-D's), is an important cause of interstitial colorectal cancer, and in this study, we hoped to find endoscopic features that have independent diagnostic value for SSL-D's.

Research motivation

Previous studies on the endoscopic features of SSLs have focused on their differentiation from hyperplastic polyps and tubular adenomas, and comparisons of the endoscopic features of SSLs without dysplasia (SSL-D's) and SSL-D's have remained at the level of prediction of the sensitivity and specificity of individual endoscopic features. There have been several reports in the literature that SSL-D's have a risk of faster progression to adenocarcinoma compared to SSL-D's and conventional tubular adenomas. Therefore, it is important to look for endoscopic features that have independent predictive value for SSL-D's and assist the endoscopist in making immediate diagnoses.

Research objectives

This study looks for endoscopic features that have independent diagnostic value for colorectal SSL-D's. These features may help endoscopists to make immediate diagnoses during colonoscopy.

Research methods

In this study, endoscopic features potentially predictive of SSLs were first analyzed by univariate analysis, and then multivariate logistic regression analysis was performed to obtain endoscopic features with independent diagnostic value for predicting SSL-D's and the diagnostic validity of these endoscopic features.

Research results

In univariate analysis, location, size, surface shape, color, microvascular varices, and crypt opening pattern of colorectal SSLs had value in predicting SSL-D's. In multifactorial regression analysis, color, microvascular varices and crypt opening pattern had independent diagnostic value in predicting SSL-D's.

Research conclusions

Reddish color, microvascular varicose, and mixed pattern of crypt openings are independent diagnostic features for colorectal SSL-D's.

Research perspectives

This finding is expected to improve the immediate diagnostic accuracy of endoscopists in SSL-D's in order to inform them to use appropriate endoscopic treatment modalities.

FOOTNOTES

Author contributions: Wang RG completed the research design, data collection and analysis, and the writing and revision of the essay; Ren YT assisted in data analysis; Ren YT and Jiang X assisted in data collection; Jiang X assisted in the research design; Wei L provided guidance on the study design and method selection; Zhang XF assisted in the completion of statistical methods and participated in data analysis; Liu H assisted in completing the histopathological diagnosis of the enrolled cases; Jiang B provided guidance on the research methods and design, and assisted in completing the revision of the essay.

Institutional review board statement: The study was reviewed and approved by the Beijing Tsinghua Changgung Hospital Institutional Review Board (approval No. 21439-0-02).

Informed consent statement: All cases that signed an informed consent form and underwent endoscopic resection.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: Participants gave informed consent for data sharing.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Rui-Gang Wang 0000-0002-9053-5329; Yu-Tang Ren 0000-0002-8972-5846; Lai Wei 0000-0003-2326-1257; Bo Jiang 0000-0002-7768-7600.

S-Editor: Wang JJ

L-Editor: A

P-Editor: Wang JJ

REFERENCES

- 1 Torlakovic E, Snover DC. Serrated adenomatous polyposis in humans. *Gastroenterology* 1996; **110**: 748-755 [PMID: 8608884 DOI: 10.1053/gast.1996.v110.pm8608884]
- 2 Nagtegaal ID, Odze RD, Klimstra D, Paradis V, Rugge M, Schirmacher P, Washington KM, Carneiro F, Cree IA; WHO Classification of Tumours Editorial Board. The 2019 WHO classification of tumours of the digestive system. *Histopathology* 2020; **76**: 182-188 [PMID: 31433515 DOI: 10.1111/his.13975]
- 3 Chang LC, Tu CH, Lin BR, Shun CT, Hsu WF, Liang JT, Wang HP, Wu MS, Chiu HM. Adjunctive use of chromoendoscopy may improve the diagnostic performance of narrow-band imaging for small sessile serrated adenoma/polyp. *J Gastroenterol Hepatol* 2018; **33**: 466-474 [PMID: 28687028 DOI: 10.1111/jgh.13863]
- 4 Tanaka Y, Yamano HO, Yamamoto E, Matsushita HO, Aoki H, Yoshikawa K, Takagi R, Harada E, Nakaoka M, Yoshida Y, Eizuka M, Sugai T, Suzuki H, Nakase H. Endoscopic and molecular characterization of colorectal sessile serrated adenoma/polyps with cytologic dysplasia. *Gastrointest Endosc* 2017; **86**: 1131-1138.e4 [PMID: 28501592 DOI: 10.1016/j.gie.2017.05.006]
- 5 Oono Y, Fu K, Nakamura H, Iriguchi Y, Yamamura A, Tomino Y, Oda J, Mizutani M, Takayanagi S, Kishi D, Shinohara T, Yamada K, Matumoto J, Imamura K. Progression of a sessile serrated adenoma to an early invasive cancer within 8 months. *Dig Dis Sci* 2009; **54**: 906-909 [PMID: 18688718 DOI: 10.1007/s10620-008-0407-7]
- 6 East JE, Vieth M, Rex DK. Serrated lesions in colorectal cancer screening: detection, resection, pathology and surveillance. *Gut* 2015; **64**: 991-1000 [PMID: 25748647 DOI: 10.1136/gutjnl-2014-309041]
- 7 Bateman AC. The spectrum of serrated colorectal lesions-new entities and unanswered questions. *Histopathology* 2021; **78**: 780-790 [PMID: 33332664 DOI: 10.1111/his.14305]
- 8 Li J, Teng XD, Lai MD. [Colorectal non-invasive epithelial lesions: an update on the pathology of colorectal serrated lesions and polyps]. *Zhonghua Bing Li Xue Za Zhi* 2020; **49**: 1339-1344 [PMID: 33287530 DOI: 10.3760/cma.j.cn112151-20200330-00272]
- 9 Erichsen R, Baron JA, Hamilton-Dutoit SJ, Snover DC, Torlakovic EE, Pedersen L, Frøsløv T, Vyberg M, Hamilton SR, Sørensen HT. Increased Risk of Colorectal Cancer Development Among Patients With Serrated Polyps. *Gastroenterology* 2016; **150**: 895-902.e5 [PMID: 26677986 DOI: 10.1053/j.gastro.2015.11.046]
- 10 IJsspeert JE, Vermeulen L, Meijer GA, Dekker E. Serrated neoplasia-role in colorectal carcinogenesis and clinical implications. *Nat Rev Gastroenterol Hepatol* 2015; **12**: 401-409 [PMID: 25963511 DOI: 10.1038/nrgastro.2015.73]
- 11 Amemori S, Yamano HO, Tanaka Y, Yoshikawa K, Matsushita HO, Takagi R, Harada E, Yoshida Y, Tsuda K, Kato B, Tamura E, Eizuka M, Sugai T, Adachi Y, Yamamoto E, Suzuki H, Nakase H. Sessile serrated adenoma/polyp showed rapid malignant transformation in the final 13 months. *Dig Endosc* 2020; **32**: 979-983 [PMID: 31677187 DOI: 10.1111/den.13572]
- 12 Omori K, Yoshida K, Tamiya S, Daa T, Kan M. Endoscopic Observation of the Growth Process of a Right-Side Sessile Serrated Adenoma/Polyp with Cytological Dysplasia to an Invasive Submucosal Adenocarcinoma. *Case Rep Gastrointest Med* 2016; **2016**: 6576351 [PMID: 27437153 DOI: 10.1155/2016/6576351]
- 13 Kinoshita S, Nishizawa T, Uraoka T. Progression to invasive cancer from sessile serrated adenoma/polyp. *Dig Endosc* 2018; **30**: 266 [PMID: 29168233 DOI: 10.1111/den.12988]
- 14 Lash RH, Genta RM, Schuler CM. Sessile serrated adenomas: prevalence of dysplasia and carcinoma in 2139 patients. *J Clin Pathol* 2010; **63**: 681-686 [PMID: 20547691 DOI: 10.1136/jcp.2010.075507]
- 15 Murakami T, Sakamoto N, Ritsuno H, Shibuya T, Osada T, Mitomi H, Yao T, Watanabe S. Distinct endoscopic characteristics of sessile serrated adenoma/polyp with and without dysplasia/carcinoma. *Gastrointest Endosc* 2017; **85**: 590-600 [PMID: 27663716 DOI: 10.1016/j.gie.2016.09.018]
- 16 Murakami T, Sakamoto N, Nagahara A. Endoscopic diagnosis of sessile serrated adenoma/polyp with and without dysplasia/carcinoma. *World J Gastroenterol* 2018; **24**: 3250-3259 [PMID: 30090005 DOI: 10.3748/wjg.v24.i29.3250]

- 17 **Hazewinkel Y**, López-Cerón M, East JE, Rastogi A, Pellisé M, Nakajima T, van Eeden S, Tytgat KM, Fockens P, Dekker E. Endoscopic features of sessile serrated adenomas: validation by international experts using high-resolution white-light endoscopy and narrow-band imaging. *Gastrointest Endosc* 2013; **77**: 916-924 [PMID: 23433877 DOI: 10.1016/j.gie.2012.12.018]
- 18 **Yamashina T**, Takeuchi Y, Uedo N, Aoi K, Matsuura N, Nagai K, Matsui F, Ito T, Fujii M, Yamamoto S, Hanaoka N, Higashino K, Ishihara R, Tomita Y, Iishi H. Diagnostic features of sessile serrated adenoma/polyps on magnifying narrow band imaging: a prospective study of diagnostic accuracy. *J Gastroenterol Hepatol* 2015; **30**: 117-123 [PMID: 25088839 DOI: 10.1111/jgh.12688]
- 19 **Murakami T**, Sakamoto N, Fukushima H, Shibuya T, Yao T, Nagahara A. Usefulness of the Japan narrow-band imaging expert team classification system for the diagnosis of sessile serrated lesion with dysplasia/carcinoma. *Surg Endosc* 2021; **35**: 4528-4538 [PMID: 32909209 DOI: 10.1007/s00464-020-07967-w]
- 20 **Murakami T**, Sakamoto N, Nagahara A. Clinicopathological features, diagnosis, and treatment of sessile serrated adenoma/polyp with dysplasia/carcinoma. *J Gastroenterol Hepatol* 2019; **34**: 1685-1695 [PMID: 31158302 DOI: 10.1111/jgh.14752]
- 21 **Wang Y**, Ren YB, Yang XS, Huang YH, Zhang L, Li X, Bai P, Wang L, Fan X, Ding YM, Li HL, Lin XC. [Comparison of endoscopic features between colorectal sessile serrated adenoma/polyp with or without cytologic dysplasia and hyperplastic polyp]. *Zhonghua Yi Xue Za Zhi* 2019; **99**: 2214-2220 [PMID: 31434395 DOI: 10.3760/cma.j.issn.0376-2491.2019.28.012]
- 22 **Sano W**, Fujimori T, Ichikawa K, Sunakawa H, Utsumi T, Iwatate M, Hasuike N, Hattori S, Kosaka H, Sano Y. Clinical and endoscopic evaluations of sessile serrated adenoma/polyps with cytological dysplasia. *J Gastroenterol Hepatol* 2018; **33**: 1454-1460 [PMID: 29377243 DOI: 10.1111/jgh.14099]
- 23 **Saito S**, Tajiri H, Ikegami M. Serrated polyps of the colon and rectum: Endoscopic features including image enhanced endoscopy. *World J Gastrointest Endosc* 2015; **7**: 860-871 [PMID: 26240687 DOI: 10.4253/wjge.v7.i9.860]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

