

# World Journal of *Clinical Cases*

*World J Clin Cases* 2023 November 6; 11(31): 7508-7740



**REVIEW**

- 7508 Gut microbiome: New perspectives for type 2 diabetes prevention and treatment  
*Li SX, Guo Y*

**MINIREVIEWS**

- 7521 Endoscopic management of benign biliary strictures: Looking for the best stent to place  
*Colombo M, Forcignanò E, Da Rio L, Spadaccini M, Andreozzi M, Giacchetto CM, Carrara S, Maselli R, Galtieri PA, Pellegatta G, Capogreco A, Massimi D, Khalaf K, Hassan C, Anderloni A, Repici A, Fugazza A*
- 7530 Antibiotic resistance in patients with liver cirrhosis: Prevalence and current approach to tackle  
*Liakina V*

**ORIGINAL ARTICLE****Retrospective Study**

- 7543 Analysis of risk factors for postoperative deep vein thrombosis after craniotomy and nomogram model construction  
*Su ZJ, Wang HR, Liu LQ, Li N, Hong XY*
- 7553 Value of ultrasound and magnetic resonance imaging combined with tumor markers in the diagnosis of ovarian tumors  
*Yang Q, Zhang H, Ma PQ, Peng B, Yin GT, Zhang NN, Wang HB*
- 7562 Measurement of combined flap thickness for reconstruction of decubitus ulcer using computed tomography  
*Kim EC, Park JD, Wee SY, Kim SY*
- 7570 Does the advantage of transcutaneous oximetry measurements in diabetic foot ulcer apply equally to free flap reconstruction?  
*Lee DW, Hwang YS, Byeon JY, Kim JH, Choi HJ*

**Clinical Trials Study**

- 7583 Effects of ulinastatin therapy in deep vein thrombosis prevention after brain tumor surgery: A single-center randomized controlled trial  
*Tao YN, Han Q, Jiao W, Yang LK, Wang F, Xue S, Shen M, Wang YH*

**Observational Study**

- 7593 Network pharmacological and molecular docking study of the effect of Liu-Wei-Bu-Qi capsule on lung cancer  
*Yang Q, Li LY*

- 7610** Efficacy of  $\beta$ 2-adrenergic receptor agonist combined with corticosteroid in the treatment of children with cough variant asthma

*Cao JY, Wang YC, Deng XX*

#### Randomized Controlled Trial

- 7619** Protective effect of sevoflurane on lung function of elderly chronic obstructive pulmonary disease patients undergoing total hip arthroplasty

*Yao Y, Zhang MS, Li YB, Zhang MZ*

#### CASE REPORT

- 7629** Sunitinib-induced hyperammonemic encephalopathy in metastatic gastrointestinal stromal tumors: A case report

*Hayakawa T, Funakoshi S, Hamamoto Y, Hirata K, Kanai T*

- 7635** Simultaneous lateral and subxiphoid access methods for safe and accurate resection of a superior vena cava aneurysm: A case report

*Kim SP, Son J*

- 7640** Ultrafast power Doppler imaging for ischemic encephalopathy: A case report

*Huang LJ, Jiao JF, He Q, Luo JW, Guo Y*

- 7647** Intermittent spontaneous ovulation in patients with premature ovarian failure: Three case reports and review of literature

*Zhang WY, Wang HB, Deng CY*

- 7656** Sneddon's syndrome concurrent with cerebral venous sinus thrombosis: A case report

*Heng Y, Tang YF, Zhang XW, Duan JF, Shi J, Luo Q*

- 7663** Carcinosarcoma of the deep lobe of the parotid gland in the parapharyngeal region: A case report

*Tang YY, Zhu GQ, Zheng ZJ, Yao LH, Wan ZX, Liang XH, Tang YL*

- 7673** Malignant peripheral nerve sheath tumor with hemophilic syndrome and bone marrow fibrosis: A rare case report

*Li H, Wang L, Wu YH, Chen G, Li HX, Fan LF, Gu M, Jiang CH*

- 7680** Comparison of drug concentrations in blood and gastric lavage fluid before and after gastric lavage: A case report

*Zhou Y, Tong JL, Peng AH, Xu SY*

- 7684** Recurred forehead osteoma disseminated after previous osteoma excision: A case report

*Lee DY, Lim S, Yoon JS, Eo S*

- 7690** Renal pelvis sarcomatoid carcinoma with renal vein tumor thrombus: A case report and literature review

*Guan HY, Wang J, Wang JX, Chen QH, Lu J, He L*

- 7699** Ultrasonographic identification of lateral femoral cutaneous nerve anatomical variation in persistent meralgia paresthetica: A case report

*Park HW, Ji KS, Kim JH, Kim LN, Ha KW*

- 7706 Biliary hemorrhage caused by a malignant small round cell tumor in the common bile duct: A case report  
*Jin YL, Ruan YJ, Lu GR*
- 7712 Successive development of ischemic malignant strokes in a patient with multiple fusiform aneurysms: A case report  
*Shin DS, Yeo DK, Choi EJ*
- 7718 Isolated axillary tumor deposit consistent with primary breast carcinoma: A case report  
*Li T, Zhang WH, Liu J, Mao YL, Liu S*
- 7724 Multiple inflammatory pseudotumor formation after craniopharyngioma resection *via* an extended nasal endoscopic approach: A case report  
*Wu H, Ding YW, Yan ZC, Wei M, Wang XD, Zhang HZ*
- 7732 Huge Bartholin's cyst managed by primary marsupialization: A case report  
*Li HY, Ding DC*

**LETTER TO THE EDITOR**

- 7738 Do not forget diet and exercise during Ramadan  
*Ilias I, Tselebis A*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, How-Ran Guo, DSc, MD, MSc, Full Professor, Department of Environmental and Occupational Health, College of Medicine, National Cheng Kung University, Tainan 70428, Taiwan. hrguo@mail.ncku.edu.tw

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

*WJCC* mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJCC* as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Si Zhao*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogorgos, Maurizio Serati

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

November 6, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

# Simultaneous lateral and subxiphoid access methods for safe and accurate resection of a superior vena cava aneurysm: A case report

Sang Pil Kim, Joohyung Son

**Specialty type:** Surgery

**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

**P-Reviewer:** Chen MCY, Taiwan

**Received:** July 13, 2023

**Peer-review started:** July 13, 2023

**First decision:** September 25, 2023

**Revised:** October 10, 2023

**Accepted:** October 23, 2023

**Article in press:** October 23, 2023

**Published online:** November 6, 2023



**Sang Pil Kim**, Department of Thoracic and Cardiovascular Surgery, Kyunghee University Hospital, Seoul 02447, South Korea

**Joohyung Son**, Department of Thoracic and Cardiovascular Surgery, Pusan National University Yangsan Hospital, Yangsan-si 50612, South Korea

**Corresponding author:** Joohyung Son, MD, Assistant Professor, Department of Thoracic and Cardiovascular Surgery, Pusan National University Yangsan Hospital, Geumo-ro 20, Beomeori, Mulgeum-eup, Yangsan-si, Gyeongsangnam-do, Republic of Korea, Yangsan-si 50612, South Korea. [pnumed09@naver.com](mailto:pnumed09@naver.com)

## Abstract

### BACKGROUND

Superior vena cava (SVC) aneurysms are a relatively uncommon disease that has not been widely reported. The conventional surgical approach for treating SVC aneurysms includes open thoracotomy and mid-sternotomy. However, in this case, the aneurysm could be safely resected by thoracoscopic simultaneous lateral and subxiphoid access methods.

### CASE SUMMARY

A 58-year-old male presented with intermittent chest pain and persistent discomfort in the chest area. A chest computed tomography scan revealed a 6.2 cm aneurysm in the left innominate vein and SVC junction. For surgical resection, simultaneous lateral and subxiphoid access were planned to achieve optimal proximal and distal aneurysm control. The approach site was 1 cm below the xiphoid process, the fifth mid-axillary line and the seventh anterior axillary line on the right side. The aneurysm was resected using a stapler. The patient was discharged on the third day after chest tube removal on the second postoperative day with no particular issues.

### CONCLUSION

Aneurysms located within the mediastinum can be accessed through thoracoscopic approach without open surgery and safely resected using vascular staples.

**Key Words:** Aneurysm; Cardiopulmonary bypass; Superior vena cava; Minimal invasive surgery; Video assisted thoracic surgery; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Superior vena cava (SVC) aneurysms, which are relatively rare, may be symptomatic or cause thrombotic events if large in size, often necessitating surgical intervention. The conventional surgical approach for treating SVC aneurysms includes open thoracotomy and mid-sternotomy. However, in this case, the aneurysm could be safely resected by thoracoscopic simultaneous lateral and subxiphoid access methods. The approach site was 1 cm below the xiphoid process, the fifth mid-axillary line and the seventh anterior axillary line on the right side. To facilitate surgery, CO<sub>2</sub> insufflation was used. The aneurysm was safely resected with the use of a vascular staple.

**Citation:** Kim SP, Son J. Simultaneous lateral and subxiphoid access methods for safe and accurate resection of a superior vena cava aneurysm: A case report. *World J Clin Cases* 2023; 11(31): 7635-7639

**URL:** <https://www.wjgnet.com/2307-8960/full/v11/i31/7635.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v11.i31.7635>

## INTRODUCTION

Superior vena cava (SVC) aneurysms are a relatively uncommon disease that has not been widely reported. In some instances, where the aneurysm is large, they may cause discomfort or bleeding-related symptoms. In such cases, surgical intervention is often considered when symptoms are present, and the possibility of a thrombotic event exists. The conventional surgical approach primarily utilizes open surgery with the application of cardiopulmonary bypass (CPB) as necessary[1,2]. Herein, we present a case in which we adopted a minimally invasive approach to treat an SVC aneurysm. In addition, we successfully performed a safe and accurate procedure using simultaneous lateral and subxiphoid access methods to ensure safe access to the SVC.

## CASE PRESENTATION

### **Chief complaints**

A 58-year-old male presented with intermittent chest pain and persistent discomfort in the chest area.

### **History of present illness**

Symptoms started several months ago and there was no other present illness.

### **History of past illness**

The patient had no any History of past illness.

### **Personal and family history**

The patient had no any family history of venous aneurysm.

### **Physical examination**

On physical examination, no abnormal findings were observed.

### **Laboratory examinations**

On laboratory testing, no abnormal findings were observed.

### **Imaging examinations**


A chest computed tomography (CT) scan revealed a 6.2 cm aneurysm in the left innominate vein and SVC junction (Figure 1). The aneurysm had a saccular appearance with no evidence of a thrombus. In cases of venous aneurysms, the rupture risk is not high, but given the patient's symptoms and the large size of the aneurysm, which raised concerns regarding a potential thrombotic event, surgical resection was deemed necessary after obtaining patient consent.

## FINAL DIAGNOSIS

The final diagnosis was a 6.2 cm aneurysm in the left innominate vein and SVC junction.

## TREATMENT

Simultaneous lateral and subxiphoid access were planned to achieve optimal proximal and distal aneurysm control. The



DOI: 10.12998/wjcc.v11.i31.7635 Copyright ©The Author(s) 2023.

**Figure 1** A chest computed tomography scan revealed a 6.2 cm aneurysm in the left innominate vein and superior vena cava junction. A: Superior vena cava aneurysm of about 6.2 cm in size seen on chest computed tomography in the axial view; B: Three-dimensional reconstruction image.

patient was intubated and placed in the left semi-lateral position. A 3-cm longitudinal skin incision was made approximately 1 cm below the xiphoid process. After making the skin incision, a wound protector was inserted, and a 10 mm port was formed at the fifth mid-axillary line and the seventh anterior axillary line on the right side. Once the port was inserted, CO<sub>2</sub> insufflation at 10 mmHg was performed to retract the pericardium and create sufficient substernal space for surgery.

First, the camera was inserted on the right side to secure the substernal space, and mediastinal dissection was performed through the right side. Next, the camera was removed from the right side and inserted into the subxiphoid site, and the surgery continued. The thymus, innominate vein, and SVC were dissected to secure Aneurysm of distal and proximal site. After umbilical tape encircling, clipping was performed to reduce venous blood flow from the left side. After this procedure, the stalk of the SVC was confirmed and partially resected using a vascular stapler (Covidien, Mansfield, MA, United States) for safe resection of the aneurysm stock, after which its size was reduced by stapling. The remaining aneurysm was resected from the stalk portion using a stapler (see [Video 1](#)).

## OUTCOME AND FOLLOW-UP


The patient was discharged on the third day after chest tube removal on the second postoperative day with no particular issues. The patient was under follow-up, and no specific findings were observed on CT 1 year after surgery ([Figure 2](#)).

## DISCUSSION

Surgical approaches for treating SVC aneurysms include open thoracotomy and mid-sternotomy. In particular, a right thoracotomy has been attempted for aneurysm resection. However, exposing the left innominate vein behind the aneurysm, securing the clamping site, or encircling the vessel's roof site can be challenging. Moreover, bleeding control is problematic in cases of injury to the opposite innominate vein during dissection. Additionally, thoracotomy increases the chance of changing positions to mid-sternotomy to control bleeding. For these reasons, thoracotomy is not a commonly used approach today.

In contrast, mid-sternotomy allows for the exposure of both jugular veins to the SVC, making it easier to create a clamping site around the aneurysm or perform vessel roof encircling. CPB can be performed at any time during surgery. Currently, mid-sternotomy is widely used in most SVC aneurysm surgeries[1,3].

Surgical visibility is essential for safe SVC aneurysm resection using video-assisted thoracic surgery (VATS). The most significant complication during surgery is the rupture of the innominate vein, which can be lethal. Compared with the right approach, the subxiphoid approach can provide a clear view of the SVC and innominate vein, making it easy to secure the vascular clamp site and promptly address innominate vein rupture. The advantages of the subxiphoid approach have already been demonstrated in VATS thymectomy[4]. Moreover, the subxiphoid approach is essential for the success of minimally invasive approaches. In the present case, since the SVC is on the right side, we believed it would be safer to approach the right side simultaneously with the subxiphoid approach, rather than using the subxiphoid approach alone. There is another consideration for VATS resection. In surgery under CPB, aneurysm resection is performed using superior and inferior SVC and innominate vein clamping, followed by resection and simple suturing. However, this procedure is not possible with minimally invasive approaches. Therefore, we modified the procedure using a vascular stapler to enable aneurysmal resection using a minimally invasive approach. In addition, the vascular stapler is relatively safe for use in the pulmonary area, which has fragile blood vessels. Therefore, the authors concluded



DOI: 10.12998/wjcc.v11.i31.7635 Copyright ©The Author(s) 2023.

**Figure 2** The patient was under follow-up, and no specific findings were observed on computed tomography 1 year after surgery. A: Computed tomography image 1 year after surgery in the coronal view; B: Three-dimensional reconstruction image.

that a vascular stapler could be used for aneurysm resection.

---

## CONCLUSION

Aneurysms located within the mediastinum can be accessed through thoracoscopic approach without open surgery and safely resected using vascular staples.

---

## ACKNOWLEDGEMENTS

This study was supported by a 2021 research grant from Pusan National University Yangsan Hospital.

---

## FOOTNOTES

**Author contributions:** Son J contributed to manuscript writing; Kim SP contributed to manuscript and figure editing; all authors have read and approved the final manuscript.

**Informed consent statement:** Informed consent was obtained from the patient for the publication of the findings of the present case study and related images.

**Conflict-of-interest statement:** All the authors declare that they have no conflict of interest to disclose.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** South Korea

**ORCID number:** Joohyung Son 0000-0002-9144-5499.

**S-Editor:** Liu JH

**L-Editor:** A

**P-Editor:** Yuan YY

---

## REFERENCES

---

- 1 **Morales MM**, Anacleto A, Ferreira Leal JC, Greque VG, Souza AS Jr, Wolosker N. Saccular Superior Vena Cava Aneurysm: Case Report and Comprehensive Review. *Ann Vasc Surg* 2021; **72**: 666.e23-666.e32 [PMID: 33333194 DOI: 10.1016/j.avsg.2020.10.033]
- 2 **Kapoor H**, Gulati V, Pawley B, Lee JT. Massive fusiform superior vena cava aneurysm in a 47-year-old complicated by pulmonary embolism: A case report and review of literature. *Clin Imaging* 2022; **81**: 43-45 [PMID: 34598003 DOI: 10.1016/j.clinimag.2021.08.008]
- 3 **Honda K**, Yuzaki M, Fujimoto T, Nishimura Y. Reconstruction of the superior vena cava with an autologous pericardial patch for a giant superior vena cava aneurysm. *JTCVS Tech* 2020; **4**: 183-186 [PMID: 34318006 DOI: 10.1016/j.xjtc.2020.10.012]
- 4 **Abdellateef A**, Ali JM, Jiang G, Aresu G, Jiang L. Tips and tricks for success in subxiphoid video-assisted thoracic surgery. *J Thorac Dis* 2019; **11**: 292-301 [PMID: 30863607 DOI: 10.21037/jtd.2018.12.93]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

