

World Journal of *Clinical Cases*

World J Clin Cases 2023 December 26; 11(36): 8434-8605



Contents

Thrice Monthly Volume 11 Number 36 December 26, 2023

EDITORIAL

- 8434 Post-trans-arterial chemoembolization hepatic necrosis and biliary stenosis: Clinical characteristics and endoscopic approach

Cocca S, Carloni L, Marocchi M, Grande G, Bianchini M, Colecchia A, Conigliaro R, Bertani H

MINIREVIEWS

- 8440 Perioperative nursing care for hip arthroplasty patients with concomitant hypertension: A minireview

Ji CY, Yang LR

ORIGINAL ARTICLE

Retrospective Study

- 8447 Evaluation of response to gemcitabine plus cisplatin-based chemotherapy using positron emission computed tomography for metastatic bladder cancer

Öztürk H, Karapolat İ

- 8458 Functional magnetic resonance imaging study of group independent components underpinning item responses to paranoid-depressive scale

Stoyanov D, Paunova R, Dichev J, Kandilarova S, Khorev V, Kurkin S

EVIDENCE-BASED MEDICINE

- 8475 Mendelian randomization provides evidence for a causal effect of serum insulin-like growth factor family concentration on risk of atrial fibrillation

Lin S, Tang J, Li X, Wu G, Lin YF, Li YF

SYSTEMATIC REVIEWS

- 8486 Significance of fostering the mental health of patients with diabetes through critical time intervention

Eseadi C, Amedu AN, Aloh HE

META-ANALYSIS

- 8498 Impact of angiotensin-converting enzyme inhibitors or angiotensin receptor blockers on the mortality in sepsis: A meta-analysis

Yang DC, Xu J, Jian L, Yu Y

CASE REPORT

- 8507 Multiple sparganosis spinal infections mainly in the thoracic region: A case report

Wen GJ, Chen J, Zhang SF, Zhou ZS, Jiao GL

- 8512** Iatrogenic flexor tendon rupture caused by misdiagnosing sarcoidosis-related flexor tendon contracture as tenosynovitis: A case report
Yan R, Zhang Z, Wu L, Wu ZP, Yan HD
- 8519** Cholecystoenteric fistula in a patient with advanced gallbladder cancer: A case report and review of literature
Wang CY, Chiu SH, Chang WC, Ho MH, Chang PY
- 8527** Intraperitoneal hyaline vascular Castleman disease: Three case reports
Gao JW, Shi ZY, Zhu ZB, Xu XR, Chen W
- 8535** Iris metastasis from clear cell renal cell carcinoma: A case report
Wang TT, Chen XY, Min QY, Han YZ, Zhao HF
- 8542** Spinal cord infarction attributed to SARS-CoV-2, with post-acute sequelae of COVID-19: A case report
Oleson CV, Olsen AC, Shermon S
- 8551** Spontaneous gastric hematoma as a rare cause of acute abdomen: A case report
Budimir I, Žulec M, Eljuga K, Židak M, Lisek V
- 8557** LiNA OperaScope™ for microwave endometrial ablation for endometrial polyps with heavy menstrual bleeding: A case report
Kakinuma K, Kakinuma T, Ueyama K, Shinohara T, Okamoto R, Yanagida K, Takeshima N, Ohwada M
- 8563** Colonoscopy-induced acute appendicitis: A case report
Song XL, Ma JY, Zhang ZG
- 8568** Post-laparotomy heterotopic ossification of the xiphoid process: A case report
Lee SS
- 8574** Balloon displacement during caesarean section with pernicious placenta previa: A case report
Gu DF, Deng C
- 8581** Synchronous carotid endarterectomy and coronary artery bypass graft: Four case reports
AlGhamdi FK, Altoijry A, AlQahtani A, Aldossary MY, AlSheikh SO, Iqbal K, Alayadhi WA
- 8589** Intraoperative cardiogenic shock induced by refractory coronary artery spasm in a patient with myasthenia gravis: A case report
Hsu CW, Chang CC, Lin CS
- 8595** Effects of video game-based therapy in an adolescent with cerebral palsy: A case report
Mohd Iqbal HA, Ho WS, Zanudin A, Hisham H, Mohd Nordin NA

LETTER TO THE EDITOR

- 8603** Lyophilized recombinant human brain natriuretic peptide: A promising therapy in patients with chronic heart failure
Kourek C, Briasoulis A, Giamouzis G, Skoularigis J, Xanthopoulos A

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Marco Infante, MD, PhD, Adjunct Professor, UniCamillus, Saint Camillus International University of Health Sciences, Rome 00131, Italy.
marco.infante@unicamillus.org

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Zi-Hang Xu, Production Department Director: Xiang Li, Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

December 26, 2023

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Effects of video game-based therapy in an adolescent with cerebral palsy: A case report

Huda Aliah Mohd Iqbal, Wei Sheng Ho, Asfarina Zanudin, Hafifi Hisham, Nor Azlin Mohd Nordin

Specialty type: Medicine, research and experimental

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C, C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Batta A, India

Received: November 3, 2023

Peer-review started: November 3, 2023

First decision: November 13, 2023

Revised: November 25, 2023

Accepted: December 7, 2023

Article in press: December 7, 2023

Published online: December 26, 2023



Huda Aliah Mohd Iqbal, Wei Sheng Ho, Asfarina Zanudin, Hafifi Hisham, Nor Azlin Mohd Nordin, Physiotherapy Programme, Centre for Rehabilitation and Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Jalan Raja Muda Abdul Aziz, Kuala Lumpur 50300, Malaysia

Corresponding author: Asfarina Zanudin, PhD, Lecturer, Physiotherapy Programme, Centre for Rehabilitation and Special Needs Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Jalan Raja Muda Abdul Aziz, Kuala Lumpur 50300, Malaysia.

asfarina.zanudin@ukm.edu.my

Abstract

BACKGROUND

Herein, we report the case of a 13-year-old boy with spastic quadriplegia cerebral palsy (CP) at Gross Motor Function Classification System (GMFCS) level II, engaging in a 6-wk video game-based therapy (VBT) program. This study aimed to offer essential insights regarding VBT's impact on enhancing the physical function and improving the quality of life (QoL) of adolescents diagnosed with CP. This report provides a distinctive viewpoint that can inform and direct future clinical practices and research endeavors.

CASE SUMMARY

The boy presented with moderate mobility, balance, and overall well-being. He faced challenges with diminished lower limb strength, which affected his daily living and physical fitness capabilities. Our participant was diagnosed with spastic quadriplegic CP at GMFCS level II. He participated in a 6-wk program of VBT using a play station. This innovative approach incorporates warm-up exercises, interactive activities, and cool-down routines, targeting various movements, including single-leg stance, weight shifting, kicking, jumping, marching, and squatting. After VBT, the strength of the left hip extensor significantly increased from 199.3 N to 541.3 N. Distance covered as part of a 6-min walk test increased by 82 m. His Paediatric QoL Inventory score increased dramatically by 25.9%.

CONCLUSION

VBT is an innovative, individualized therapy that enhances physical function and QoL in CP, emphasizing its role in ambulatory patients.

Key Words: Video game; Physical function; Quality of life; Cerebral Palsy; Case report

Core Tip: This study examined the potential transformative effects of video game-based therapy (VBT) in adolescents diagnosed with cerebral palsy (CP). The 6-wk VBT program demonstrated notable improvements in lower limb strength, motor function, and overall quality of life. VBT is distinguished by its creative and patient-centric approach, which is customized to meet the unique needs of each individual. This study highlights the need to incorporate technology-driven rehabilitation approaches into treatment methods for ambulatory patients with CP. It emphasizes the significant implications of such an approach and sets the stage for future multicenter trials that consider a range of functional levels.

Citation: Mohd Iqbal HA, Ho WS, Zanudin A, Hisham H, Mohd Nordin NA. Effects of video game-based therapy in an adolescent with cerebral palsy: A case report. *World J Clin Cases* 2023; 11(36): 8595-8602

URL: <https://www.wjgnet.com/2307-8960/full/v11/i36/8595.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i36.8595>

INTRODUCTION

Cerebral palsy (CP) is the most common motor disability among children, with an estimated prevalence of one to nearly four per 1000 live births worldwide, as reported by the Centers for Disease Control and Prevention in 2022. Enhancing the quality of life (QoL) has consistently been a primary objective and robust evaluative criterion in therapeutic interventions for adolescents with CP[1,2]. Considerable research has indicated significant empirical evidence regarding the efficacy of modern physiotherapy approaches currently available for children and adolescents diagnosed with CP. Among the existing approaches, video game-based therapy (VBT) can incorporate all the aforementioned elements in the therapeutic process owing to its distinctive attributes of being centered on video content and recreational activities, while being adaptable based on the user's condition. Furthermore, in addition to the heightened excitement experienced during physical activity, VBT offers instantaneous biofeedback to improve users' movement patterns and experiences.

The current evidence indicates that VBT is a viable therapeutic option for enhancing arm function, postural control, and ambulation in adolescents with CP[3-6]. However, research on the impact of VBT on the physical functioning of children and adolescents with CP, specifically in terms of walking ability, gross motor skills, strength, and QoL, has been lacking. This inadequacy is particularly apparent in the absence of a comprehensive and reliable tool for assessing outcomes. Therefore, this study was conducted to provide essential empirical support for an alternative therapeutic approach for adolescents diagnosed with CP.

This study aimed to produce essential evidence that can assist healthcare professionals in assessing their patients while providing suggestions for their interventions. The present study examined the potential of VBT in enhancing physical function, including lower-limb strength, gross motor skills, walking capacity, and QoL in adolescents diagnosed with CP. The findings of this study provide essential information for healthcare professionals to evaluate their patients and act as a framework for guiding their interventions with patients. The results could potentially influence the physical health and overall well-being of adolescents diagnosed with CP and their respective families.

CASE PRESENTATION

Chief complaints

This case report is based on a boy who was 13 years and 3 months old he was diagnosed with spastic quadriplegia CP. His mother reported of his sedentary preference and inability to walk for long periods, especially outdoors and on inclined surfaces.

History of present illness

When diagnosed, the patient was classified as having Gross Motor Function Classification System (GMFCS) level II. He could walk without assistive equipment by employing a unique knee-gait pattern, and exhibited skills in traversing curbs, uneven surfaces, and barriers. Additionally, he could independently ascend and descend steps within his two-story home with one hand holding the rail.

History of past illness

The participant was born at full term with no other medical complications. He had mild jaundice, which is a common condition in many newborns. However, eventually, the participant's mother observed gross motor developmental deficits, including milestones, such as rolling, compared to his older siblings. Accordingly, the mother sought the expertise of a pediatrician at the hospital.

In 2009, he was referred to our hospital for various medical assessments. In mid-2010, he was officially diagnosed with spastic quadriplegia CP, which mainly explained his motor development impairments. He began his physiotherapy in

2009, following his mother's concerns about gross motor delay. Physiotherapy was conducted on a bi-monthly basis by a pediatric physiotherapist. Physiotherapy aims to provide motor training and home program strategies to effectively target the specific requirements and challenges of the participants.

Prior to VBT, his mother observed an increase in sedentary behavior and his son's difficulties in long-distance walking, especially during outdoor activities and on inclined surfaces. These concerns were the primary motivating factors behind the pursuit of novel strategies to enhance his physical capabilities and overall well-being.

Personal and family history

The patient had no family history of CP or other remarkable neurological conditions as claimed by the mother.

Physical examination

A comprehensive literature review was conducted before selecting assessments for the clinical decision-making process. Muscle strength with respect to body function was assessed using a handheld dynamometer. Additionally, general motor ability is evaluated using the Gross Motor Function Measure (GMFM)-88. The 6-min walk test (6 MWT) was used to assess walking capacity, whereas the Pediatric QoL (PedsQL) tool in Malay was used to determine the QoL. Physical examination was performed twice, at baseline and after 4 weeks of VBT.

Muscle strength: Muscle strength was tested using a micro Force Evaluation and Testing 2 (microFET®2) handheld dynamometer (Hogan Health Industries Inc., United States), a small portable device held by the examiner, and placed against the participant's limit during maximal isometric contraction. According to Goudriaan *et al*[7], a knee extension interclass correlation (ICC) value of 0.95 for children with CP indicated that the handheld dynamometer was highly reliable.

According to Neumann[8], a pathology that affects the strength, control, or extensibility of hip muscles can significantly disrupt the fluidity of many routine movements involving both functional and recreational activities. Thus, in our assessment, we focused on the hip abductor, hip extensor, and knee extensor muscles. The test was conducted in a maneuver with an ICC validated by Yazici *et al*[9] (ICC = 0.938-0.986), wherein the examiner was in a stable position, providing maximal ability to resist force by the participant, running at a slow count of four upon maximal muscle contraction. Testing was performed in the following manual muscle testing positions: Prone lying (hip extensor), side lying (hip abductor), and high sitting (knee extensor). Figure 1 shows muscle strength measurements using the MicroFET®2 handheld dynamometer.

Gross motor ability: According to the Center of Childhood Disability Research, the GMFM is a standardized observational instrument designed and validated to measure changes in gross motor function over time in children and adolescents with CP, with most items having specific descriptors for each score. The GMFM-88 has been used to assess the gross motor ability of persons with CP[10]. In our assessments, domains D (standing) and E (walking, running, and jumping) from the GMFM-88 were used to suit our participant category, which, in this case, GMFCS level II, possesses the ability to stand and walk. The items were administered according to the GMFM manual[10].

Walking capacity: The 6 MWT is a submaximal exercise test that involves walking for 6 min. ICC 0.98 was validated in the CP population by Thompson *et al*[11]. The 6 MWT was used to determine the participants walking capacity. The test was administered according to the standard protocol, implementing every instruction in accordance with the guidelines of the American Thoracic Society[12].

QoL: To explore the QoL, we chose a PedsQL parent's proxy questionnaire consisting of a 5-point Likert Scale ranging from 0 to 4, covering four multidimensional functions, namely, physical, emotional, social, and school. Adapting to the local culture, the PedsQL questionnaire was translated into Malay to ease the patient's career comprehension. The questionnaire has been validated with Cronbach's α with a range (0.7-0.98) and factor correlation values ranging from 0.1 to 0.57[13].

Laboratory examinations

The laboratory examinations are not applicable.

Imaging examinations

The imaging examinations are not applicable.

FINAL DIAGNOSIS

The final diagnosis of the case presented is spastic quadriplegia CP.

TREATMENT

The VBT intervention was initiated in the second session after an initial session, in which baseline data assessments were conducted. An interventional session was conducted by a researcher under the guidance of the patient's pediatric physio-



DOI: 10.12998/wjcc.v11.i36.8595 Copyright ©The Author(s) 2023.

Figure 1 Muscle strength measurement. A: Knee extensor; B: Hip extensor; C: Hip abductor.

therapist for 6 weeks, consisting of two sessions per week, with each session requiring approximately 1 hour. The principal objective of the program was to enhance engagement in physical activities. Balance, strength, and endurance training exercises were incorporated into the regimen using an X-Box 360 Kinect. The tasks included upper limb movements, marching, side-stepping, weight shifting, single-leg stance (SLS), kicking, squatting, and jumping. All interventions were adapted to be fun and engaging for young adolescents. To make the gaming session more fun and competitive, the researchers also participated in the gaming session during the participants' resting period to motivate them to perform the tasks at their maximal performance.

Each session started with a warm-up of 10–15 min involving upper limb movements and low-intensity exercises such as side-stepping and marching at a low pace. The session was followed by balance training, such as weight shifting, SLS, and dynamic balance training; strength training involving marching, kicking, squatting, and jumping; and endurance training, a combination of all movements and movement strategies with more significant repetitions. Following this, progress was made according to the participants' preferences and the researcher's observations. Throughout the intervention session, participants rested between tasks until they were ready for the next game. The session ended with a cooldown period of 10 min consisting of controlled breathing and stretching of the upper and lower limbs. The components and movements are listed in Table 1.

OUTCOME AND FOLLOW-UP

Initial outcomes

Clinical observation: The patient exhibited moderate thoracic scoliosis on the right side manifesting as an elevated right shoulder relative to the left shoulder. The participant could walk unaided while maintaining a mild jumping knee gait; however, he encountered challenges when sprinting and walking at high speeds. He could only walk in tandem for five steps and a SLS on the right foot for 2 s and 1 s on the left. He was only able to execute single-leg hopping with his right foot once using a double-hand grasp and was unable to conduct single-leg hopping with his left foot.

Outcome and follow-up

Clinical observation: He could run but still had a mild hopping knee gait. He could maintain the right foot SLS for 5.36 s, whereas he maintained the left foot SLS for 1.36 s. For running, he could now complete a distance of 4.5 min and 5.4 s instead of walking fast.

Description of outcomes: Post-intervention, there were marked increases in strength for his tested lower limb muscles, of which, the left hip extensor had gained the most improvement, from 199.3 N to 541.3 N, followed by the left hip abductor, from 201.1 N to 743.3 N.

Table 1 Components and tasks involved in the video game-based intervention

Component	Movements	Game
Warm up	Upper limb movements side	Dance central: Poker face
	Stepping	Kinect adventures: Space pop
	Marching	Motion explosion: Balance beam
Balance	Single leg stance	Kinect adventures: 20000 leaks
	Weight shifting	Kinect sports: Target kick
		Ice age 4: Slip slider
Strength	Kicking	Kinect sports: Target kick
	Jumping	Kinect adventures: River rush
	Marching	Kinect sports: Hurdles
		Kinect sports: Sprint
Endurance	Combination of marching, squatting, jumping, and upper limb movements	Ice age 4: Glacier hopper
		Kinect adventures: Rally ball
Cool down	Upper limb movements	Kinect adventures: Space pop
	Stepping	Motion explosion: Balance beam

The boy's gross motor function in terms of standing and advanced activities, such as walking, running, and jumping demonstrated no improvement nor deterioration. A decreased of one score in GMFM component D. According to his walking capacity, the distance increased from 368 m to 450 m, yielding an improvement of 82 m.

Assessing the QoL of the boy using the mother's proxy, the post-intervention PedsQL total scale score rose from 58.7 to 73.9, marking an increase of 17.06%. All the domains in PedsQL which are physical, emotion, social and schooling showed increased scores. The changes in muscle strength, gross motor ability, and QoL after the VBT intervention are shown in [Table 2](#).

Adherence, adverse events, and participant feedback: The participants in this study showed good adherence to the program with 100% attendance. He did not miss any appointments for the program. No adverse events were observed during the intervention period. According to the participant, VBT was fun, and he did not experience fatigue or muscle soreness after every session. The participant's caregiver also added that her child was more motivated and excited for VBT than the routine physiotherapy appointments.

DISCUSSION

The VBT intervention enhanced the physical function and QoL of a GMFCS level II adolescents with CP, as indicated by the achievement of positive results on almost all outcome measures. Furthermore, our participants demonstrated high adherence to and a desire to engage in all assigned tasks during the VBT sessions. Participants and their caregivers complied with the entire program and adhered to the schedule for each session. Consistent with findings from prior research[14,15], our participants also reported exceptional joy and amusement compared to routine interventions. Following these sessions, the patient exhibited enhancements in bodily structure, function, and QoL. The participant and his pediatric physiotherapist noted that his posture and balance improved after the intervention[14,15].

Our results are consistent with those of Gercek *et al*[16], which showed improved lower-extremity muscle strength from VBT. The bilateral hip extensors, hip abductors, and right knee extensors improved post-intervention, potentially because of the exercises performed during VBT training. The participants performed lower limb-intensive exercises during training, including marching, squatting, kicking, and jumping, at a specific velocity corresponding to the tempo of the events. As the required pace increases, stamina, endurance, and reaction time are required. Voluntary active range of motion, weight-bearing movements, and the functionality and velocity of the movements themselves must have contributed to the increase in muscle strength. Our hypothesis was consistent with findings by van Vulpen *et al*[17], showing an association of improvement in strength with the functionality of the velocity in strength training exercises.

Pre-intervention, the participant was able to perform most of the gross motor activities in GMFM component D (standing); however, he had difficulty in performing the SLS, with the right-foot SLS better than a left-foot SLS (right: 2 s; left: 1 s). After the intervention, SLS time in both feet improved, with right-foot SLS to a greater extent (right: 5.36 s; left: 1.56 s). With respect to component E (walking, running, and jumping), he possesses difficulty in performing tandem walking (5/10 steps), running 4.5 m (fast walking instead), and single leg hopping (right: One hop with two-hand held; left: 0/10 with two-hand held). Post-intervention, the changes were tandem walking (3/10 steps), running 4.5 m (able to run), and single-leg hopping (right: One hop without hand-held; left: 0/10). These changes were consistent with the findings of Gercek *et al*[16], who showed improvements in motor function post-intervention. These changes are likely due

Table 2 Changes after the video game-based therapy in muscle strength, gross motor, and quality of life

Outcome measure		Pre-intervention	Post-intervention	Change (%)
Muscle strength (N)	Left hip extensor	199.3	541.3	34 (> 100)
	Right hip extensor	147.2	535.4	388.2 (> 100)
	Left hip abductor	201.1	743.3	542.2 (> 100)
	Right hip abductor	212.6	752.2	539.6 (> 100)
	Left knee extensor	216.2	1017.9	801. (> 100)
	Right knee extensor	290.1	1035.6	745.5 (> 100)
GMFM-88 score	Dimension D standing (0-39)	36	35	1 (2.7)
	Dimension E walking, running, jumping (0-72)	65	65	-
Walking distance (m)	6 MWT	368	450	82 (5.0)
Quality of life	PedsQL in physical (0-100)	71.9	81.3	9.4 (13.0)
	PedsQL in emotion (0-100)	50.0	60.0	10 (20)
	PedsQL in social (0-100)	35.0	65.0	30 (85.7)
	PedsQL in schooling (0-100)	70.0	85.0	15 (21.4)
	PedsQL in total (0-100)	58.7	73.9	15.2 (25.9)

N: Newton; GMFM: Gross Motor Function Measure; m: Meter; 6 MWT: Six-min walk test; PedsQoL: Pediatric quality of life.

to the nature of virtual reality (VR), which addresses the critical factors of motor training, including intensity, repetition, task orientation, and multisensory environments. According to Brien and Sveistrup[18], enriched environments are known to promote neuronal plasticity changes documented using functional magnetic resonance imaging, and have shown that training in VR offers the potential for long-term learning from adaptive cerebral plasticity consistent with significant functional motor improvements in CP. Despite the minimal changes, we expect to see a more substantial improvement in the participants' gross motor skills with a longer intervention of a minimum of 8 weeks is being performed, especially in the gait component. According to Ghai S and Ghai I[4], a VBT of less than 4 times per week, 20–30 min, for more than 8 weeks can have a maximum effect on gait. Therefore, a trial utilizing these suggestions could be conducted to harness this impact. In addition, the improvement in muscle strength and gross motor function to different extents was consistent with the findings of Shin *et al*[19], who showed no correlation between muscle strength and gross motor function in CP.

Upon pre-intervention, the participant completed a distance of 368 m. After the intervention, an increment of 82 m was obtained. This finding is consistent with a previous study[20] showing that VBT intervention can improve walking capacity in adolescents with CP. Throughout our intervention, the VBT involved repetitive practice and movement correction. As pointed out by Chen *et al*[3] in a similar study utilizing VR, the repetitions used during VR sessions were high, and the child performed up to 150 reaching movements in 3 min without realizing them while playing the video games. Although the repetitions of movement in our study were not recorded, more significant repetitions within a specific time frame are required, which we believe have contributed to the build-up of endurance in the participants, thus resulting in a significant outcome.

QoL improvement as a result of overall improvement in the participants' physical and social functioning were reported with all aspects acquiring significant increment, and out of all, social functioning improved by the most (85.7%). An increase in score (from 4 to 2) was observed in Components 1 (relationships with other adolescents), 4 (inability to perform activities that other adolescents are capable of), and 5 (catching up with other adolescents' pace). Two of the three questions were based on the participants' physical abilities. With improvements in bodily functions shown in the physical functioning domain, which may have contributed to his school functioning domain improvement, the emotional and social aspects of the QoL increased significantly. Although the exact cause-and-effect relationship between these domains is still unknown, our findings indicate that the enhancements in gross motor function, walking ability, and overall physical functions, such as muscle strength, are indeed associated with the improvements observed in the social domain of the PedsQL, which are in line with the findings of Shelly *et al*[21], who stated that the psychological domain of QoL was significantly associated with the functioning level.

In conclusion, increments in muscle strength, particularly hip extensor, 6 MWT, GMFM-88 domains D and E, and PedsQL scores post-intervention, indicated that VBT could improve physical function and QoL of CP with GMFCS level II. Although the improvements are apparent, the results of this case study should be carefully interpreted as the outcomes of a single-subject case study have limited generalizability to a larger population, with limitations such as the short duration of the VBT program and follow-up assessment once participants finish the program. In the future, to ensure an optimal number of participants, recruitment planning should consider the barriers of participants visiting the clinic for VBT for two sessions per week. In addition, the PedsQL Adolescent Report should be employed to provide additional

Table 3 The pros and cons between standard care and video game-based therapy for adolescents with cerebral palsy

Standard care for cerebral palsy	Video game-based therapy for cerebral palsy
Functional training; Facilitation of normal patterns of movement; Passive range of motion; Stretching exercise; Strengthening exercise; Positioning; Manual technique; Functional electrical stimulation; Splinting; Maintenance training	Refer to Table 1
Pros Basic technique and well-known by physiotherapists; Most of the essential equipment available in the physiotherapy department	Active motor training with progressive challenges; Motivational and fun activities; Games were tailored to patients' ability, preference, and functional goals; Bi-weekly training with a structured program and assessments
Cons Patients tend to get bored with repetitive treatment	Physiotherapists need to be trained on how to operate the games; Not all physiotherapy department has the equipment and space for the VBT

VBT: Video game-based therapy.

validation of the QoL outcomes. Subsequent research should focus on the frequency and durations of VBT interventions, with specific emphasis on the duration, number of sessions per week, and localization of VBT programs.

What this case study adds to evidence-based practice

This is the first case report to examine the effect of VBT intervention in an adolescent with spastic quadriplegia and GMFCS level II CP in a local setting. This study demonstrated the clinical applicability of VBT intervention in elevating physical function with improvements in muscle strength, gross motor function, and walking capacity, ultimately improving the QoL of adolescents with CP. [Table 3](#) sums up the advantages and disadvantages of standard care and VBT in adolescents with CP.

CONCLUSION

In summary, the aforementioned case study involving a 13-year-old boy diagnosed with spastic quadriplegic CP, underscores the need for timely identification and intervention in the field of pediatric neurological disorders. Despite difficulties in the gross motor development, he has sought and engaged in ongoing physiotherapy since 2009, which highlights the compliance and support of both his family and healthcare professionals in addressing his distinct requirements.

The implementation of VBT as a novel intervention shows the potential for improving participants' physical functioning and QoL. The preliminary evaluation indicated his competence in walking while emphasizing his difficulties in navigating outdoor environments and ascending inclined slopes. The VBT, in conjunction with his family's assistance, seeks to mitigate these constraints and enhance holistic welfare.

This case highlights the importance of customized therapies for children diagnosed with CP as well as the potential advantages of rehabilitation methods that include technology. As the intervention progresses, and subsequent assessments are conducted, significant insights will be gained regarding the efficacy of VBT in treating children with spastic quadriplegic CP. These findings serve as valuable resources for informing future treatment approaches for comparable instances.

FOOTNOTES

Author contributions: Mohd Iqbal HA, Ho WS, Hisham H, and Zanudin A contributed to manuscript writing editing; Mohd Iqbal HA, Ho WS, and Zanudin A contributed to data collection; Ho WS, Mohd Iqbal HA, Zanudin A and Mohd Nordin NA contributed to data analysis; Zanudin A and Mohd Nordin NA contributed to conceptualization and supervision; all authors have read and approved the final manuscript.

Supported by the Ministry of Higher Education of Malaysia, No. FRGS/1/2021/SKK06/UKM/03/2.

Informed consent statement: Written informed consent was obtained from the patient for publication of this report and any accompanying images.

Conflict-of-interest statement: The authors declare that they have no conflict of interest to disclose.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers.

It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Malaysia

ORCID number: Asfarina Zanudin 0000-0001-6187-1019.

S-Editor: Lin C

L-Editor: A

P-Editor: Lin C

REFERENCES

- Zanudin A**, Mercer T, Samaan C, Jagadamma K, McKelvie G, van der Linden M. A Community-Based Exercise Program For Ambulant Adolescents And Young Adults With Cerebral Palsy, A Feasibility Study. *Eur J Adapt Phys Act* 2021; **14**: 1-15 [DOI: [10.5507/euj.2021.003](https://doi.org/10.5507/euj.2021.003)]
- Zanudin A**, Mercer TH, Jagadamma KC, van der Linden ML. Psychometric Properties Of Measures Of Gait Quality And Walking Performance In Young People With Cerebral Palsy: A Systematic Review. *Gait Posture* 2017; **58**: 30-40 [PMID: [28711651](https://pubmed.ncbi.nlm.nih.gov/28711651/)] DOI: [10.1016/j.gaitpost.2017.07.005](https://doi.org/10.1016/j.gaitpost.2017.07.005)]
- Chen Y**, Fanchiang HD, Howard A. Effectiveness of Virtual Reality in Children With Cerebral Palsy: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Phys Ther* 2018; **98**: 63-77 [PMID: [29088476](https://pubmed.ncbi.nlm.nih.gov/29088476/)] DOI: [10.1093/ptj/pzx107](https://doi.org/10.1093/ptj/pzx107)]
- Ghai S**, Ghai I. Virtual Reality Enhances Gait in Cerebral Palsy: A Training Dose-Response Meta-Analysis. *Front Neurol* 2019; **10**: 236 [PMID: [30984095](https://pubmed.ncbi.nlm.nih.gov/30984095/)] DOI: [10.3389/fneur.2019.00236](https://doi.org/10.3389/fneur.2019.00236)]
- Cho C**, Hwang W, Hwang S, Chung Y. Treadmill Training with Virtual Reality Improves Gait, Balance, and Muscle Strength in Children with Cerebral Palsy. *Tohoku J Exp Med* 2016; **238**: 213-218 [PMID: [26947315](https://pubmed.ncbi.nlm.nih.gov/26947315/)] DOI: [10.1620/tjem.238.213](https://doi.org/10.1620/tjem.238.213)]
- Yoo JW**, Lee DR, Cha YJ, You SH. Augmented effects of EMG biofeedback interfaced with virtual reality on neuromuscular control and movement coordination during reaching in children with cerebral palsy. *NeuroRehabilitation* 2017; **40**: 175-185 [PMID: [28222541](https://pubmed.ncbi.nlm.nih.gov/28222541/)] DOI: [10.3233/NRE-161402](https://doi.org/10.3233/NRE-161402)]
- Goudriaan M**, Nieuwenhuys A, Schless SH, Goemans N, Molenaers G, Desloovere K. A new strength assessment to evaluate the association between muscle weakness and gait pathology in children with cerebral palsy. *PLoS One* 2018; **13**: e0191097 [PMID: [29324873](https://pubmed.ncbi.nlm.nih.gov/29324873/)] DOI: [10.1371/journal.pone.0191097](https://doi.org/10.1371/journal.pone.0191097)]
- Neumann DA**. Kinesiology of the hip: a focus on muscular actions. *J Orthop Sports Phys Ther* 2010; **40**: 82-94 [PMID: [20118525](https://pubmed.ncbi.nlm.nih.gov/20118525/)] DOI: [10.2519/jospt.2010.3025](https://doi.org/10.2519/jospt.2010.3025)]
- Yazici G**, Varol F, Volkan M, Çitaker S. Reliability of make and break tests in assessing intrinsic foot muscle strength using A handheld dynamometer in healthy young adults. *Türk Fizyoterapi ve Rehabilitasyon Dergisi* 2020; **31**: 263-269 [DOI: [10.21653/tjpr.736578](https://doi.org/10.21653/tjpr.736578)]
- Te Velde A**, Morgan C. Gross motor function measure (GMFM-66 & GMFM-88) User's manual, book review. *Pediatric Physical Therapy* 2022; **34**: 88-89 [DOI: [10.1097/PEP.0000000000000858](https://doi.org/10.1097/PEP.0000000000000858)]
- Thompson P**, Beath T, Bell J, Jacobson G, Phair T, Salbach NM, Wright FV. Test-retest reliability of the 10-metre fast walk test and 6-minute walk test in ambulatory school-aged children with cerebral palsy. *Dev Med Child Neurol* 2008; **50**: 370-376 [PMID: [18355340](https://pubmed.ncbi.nlm.nih.gov/18355340/)] DOI: [10.1111/j.1469-8749.2008.02048.x](https://doi.org/10.1111/j.1469-8749.2008.02048.x)]
- American Thoracic Society**. . Guidelines for the six-minute walk test. *Am J Respir Crit Care Med* 2002; **166**: 111-117
- Ainuddin HA**, Loh SY, Chinna K, Low WY, Roslani AC. Psychometric properties of the self-report Malay version of the Pediatric Quality of Life (PedsQLTM) 4.0 Generic Core Scales among multiethnic Malaysian adolescents. *J Child Health Care* 2015; **19**: 229-238 [PMID: [24154841](https://pubmed.ncbi.nlm.nih.gov/24154841/)] DOI: [10.1177/1367493513504834](https://doi.org/10.1177/1367493513504834)]
- Viana RB**, de Oliveira VN, Dankel SJ, Loenneke JP, Abe T, da Silva WF, Morais NS, Vancini RL, Andrade MS, de Lira CAB. The effects of exergames on muscle strength: A systematic review and meta-analysis. *Scand J Med Sci Sports* 2021; **31**: 1592-1611 [PMID: [33797115](https://pubmed.ncbi.nlm.nih.gov/33797115/)] DOI: [10.1111/sms.13964](https://doi.org/10.1111/sms.13964)]
- Bryanton C**, Bossé J, Brien M, McLean J, McCormick A, Sveistrup H. Feasibility, motivation, and selective motor control: virtual reality compared to conventional home exercise in children with cerebral palsy. *Cyberpsychol Behav* 2006; **9**: 123-128 [PMID: [16640463](https://pubmed.ncbi.nlm.nih.gov/16640463/)] DOI: [10.1089/cpb.2006.9.123](https://doi.org/10.1089/cpb.2006.9.123)]
- Gercek N**, Tatar Y, Uzun S. Alternative exercise methods for children with cerebral palsy: effects of virtual vs. traditional golf training. *Int J Dev Disabil* 2022; **68**: 933-942 [PMID: [36568621](https://pubmed.ncbi.nlm.nih.gov/36568621/)] DOI: [10.1080/20473869.2021.1926853](https://doi.org/10.1080/20473869.2021.1926853)]
- van Vulpen LF**, de Groot S, Rameckers E, Becher JG, Dallmeijer AJ. Improved Walking Capacity and Muscle Strength After Functional Power-Training in Young Children With Cerebral Palsy. *Neurorehabil Neural Repair* 2017; **31**: 827-841 [PMID: [28786309](https://pubmed.ncbi.nlm.nih.gov/28786309/)] DOI: [10.1177/1545968317723750](https://doi.org/10.1177/1545968317723750)]
- Brien M**, Sveistrup H. An intensive virtual reality program improves functional balance and mobility of adolescents with cerebral palsy. *Pediatr Phys Ther* 2011; **23**: 258-266 [PMID: [21829120](https://pubmed.ncbi.nlm.nih.gov/21829120/)] DOI: [10.1097/PEP.0b013e318227ca0f](https://doi.org/10.1097/PEP.0b013e318227ca0f)]
- Shin HI**, Sung KH, Chung CY, Lee KM, Lee SY, Lee IH, Park MS. Relationships between Isometric Muscle Strength, Gait Parameters, and Gross Motor Function Measure in Patients with Cerebral Palsy. *Yonsei Med J* 2016; **57**: 217-224 [PMID: [26632404](https://pubmed.ncbi.nlm.nih.gov/26632404/)] DOI: [10.3349/ymj.2016.57.1.217](https://doi.org/10.3349/ymj.2016.57.1.217)]
- Arnoni JLB**, Pavao SL, Dos Santos Silva FP, Rocha NACF. Effects of virtual reality in body oscillation and motor performance of children with cerebral palsy: A preliminary randomized controlled clinical trial. *Complement Ther Clin Pract* 2019; **35**: 189-194 [PMID: [31003657](https://pubmed.ncbi.nlm.nih.gov/31003657/)] DOI: [10.1016/j.ctcp.2019.02.014](https://doi.org/10.1016/j.ctcp.2019.02.014)]
- Shelly A**, Davis E, Waters E, Mackinnon A, Reddihough D, Boyd R, Reid S, Graham HK. The relationship between quality of life and functioning for children with cerebral palsy. *Dev Med Child Neurol* 2008; **50**: 199-203 [PMID: [18215191](https://pubmed.ncbi.nlm.nih.gov/18215191/)] DOI: [10.1111/j.1469-8749.2008.02031.x](https://doi.org/10.1111/j.1469-8749.2008.02031.x)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

