

# World Journal of *Clinical Cases*

*World J Clin Cases* 2023 February 6; 11(4): 719-978



## Contents

Thrice Monthly Volume 11 Number 4 February 6, 2023

## MINIREVIEWS

- 719 Development and refinement of diagnostic and therapeutic strategies for managing patients with cardiogenic stroke: An arduous journey  
*Fan ZX, Liu RX, Liu GZ*
- 725 Portal vein aneurysm-etiology, multimodal imaging and current management  
*Kurtcehajic A, Zerem E, Alibegovic E, Kunosic S, Hujdurovic A, Fejzic JA*

## ORIGINAL ARTICLE

## Clinical and Translational Research

- 738 CD93 serves as a potential biomarker of gastric cancer and correlates with the tumor microenvironment  
*Li Z, Zhang XJ, Sun CY, Fei H, Li ZF, Zhao DB*

## Retrospective Study

- 756 Chest computed tomography findings of the Omicron variants of SARS-CoV-2 with different cycle threshold values  
*Ying WF, Chen Q, Jiang ZK, Hao DG, Zhang Y, Han Q*
- 764 Major depressive disorders in patients with inflammatory bowel disease and rheumatoid arthritis  
*Haider MB, Basida B, Kaur J*
- 780 Selective laser trabeculoplasty as adjunctive treatment for open-angle glaucoma *vs* following incisional glaucoma surgery in Chinese eyes  
*Zhu J, Guo J*
- 788 Efficacy of transvaginal ultrasound-guided local injections of absolute ethanol for ectopic pregnancies with intrauterine implantation sites  
*Kakinuma T, Kakinuma K, Matsuda Y, Yanagida K, Ohwada M, Kaijima H*

## Clinical Trials Study

- 797 Efficacy of incremental loads of cow's milk as a treatment for lactose malabsorption in Japan  
*Hasegawa M, Okada K, Nagata S, Sugihara S*

## Observational Study

- 809 Transdiagnostic considerations of mental health for the post-COVID era: Lessons from the first surge of the pandemic  
*Goldstein Ferber S, Shoval G, Rossi R, Trezza V, Di Lorenzo G, Zalsman G, Weller A, Mann JJ*
- 821 Effect of patient COVID-19 vaccine hesitancy on hospital care team perceptions  
*Caspi I, Freund O, Pines O, Elkana O, Ablin JN, Bornstein G*

**Randomized Clinical Trial**

- 830** Improvement of inflammatory response and gastrointestinal function in perioperative of cholelithiasis by Modified Xiao-Cheng-Qi decoction  
*Sun BF, Zhang F, Chen QP, Wei Q, Zhu WT, Ji HB, Zhang XY*

**CASE REPORT**

- 844** Metagenomic next-generation sequencing for pleural effusions induced by viral pleurisy: A case report  
*Liu XP, Mao CX, Wang GS, Zhang MZ*
- 852** *Clostridium perfringens* gas gangrene caused by closed abdominal injury: A case report and review of the literature  
*Li HY, Wang ZX, Wang JC, Zhang XD*
- 859** Is lymphatic invasion of microrectal neuroendocrine tumors an incidental event?: A case report  
*Ran JX, Xu LB, Chen WW, Yang HY, Weng Y, Peng YM*
- 866** *Pneumocystis jirovecii* diagnosed by next-generation sequencing of bronchoscopic alveolar lavage fluid: A case report and review of literature  
*Cheng QW, Shen HL, Dong ZH, Zhang QQ, Wang YF, Yan J, Wang YS, Zhang NG*
- 874** Identification of 1q21.1 microduplication in a family: A case report  
*Huang TT, Xu HF, Wang SY, Lin WX, Tung YH, Khan KU, Zhang HH, Guo H, Zheng G, Zhang G*
- 883** Double pigtail catheter reduction for seriously displaced intravenous infusion port catheter: A case report  
*Liu Y, Du DM*
- 888** Thyroid storm in a pregnant woman with COVID-19 infection: A case report and review of literatures  
*Kim HE, Yang J, Park JE, Baek JC, Jo HC*
- 896** Computed tomography diagnosed left ovarian venous thrombophlebitis after vaginal delivery: A case report  
*Wang JJ, Hui CC, Ji YD, Xu W*
- 903** Preoperative 3D reconstruction and fluorescent indocyanine green for laparoscopic duodenum preserving pancreatic head resection: A case report  
*Li XL, Gong LS*
- 909** Unusual presentation of systemic lupus erythematosus as hemophagocytic lymphohistiocytosis in a female patient: A case report  
*Peng LY, Liu JB, Zuo HJ, Shen GF*
- 918** Polyarteritis nodosa presenting as leg pain with resolution of positron emission tomography-images: A case report  
*Kang JH, Kim J*
- 922** Easily misdiagnosed complex Klippel-Trenaunay syndrome: A case report  
*Li LL, Xie R, Li FQ, Huang C, Tuo BG, Wu HC*

- 931** Benign lymphoepithelial cyst of parotid gland without human immunodeficiency virus infection: A case report  
*Liao Y, Li YJ, Hu XW, Wen R, Wang P*
- 938** Epithelioid trophoblastic tumor of the lower uterine segment and cervical canal: A case report  
*Yuan LQ, Hao T, Pan GY, Guo H, Li DP, Liu NF*
- 945** Treatment of portosystemic shunt-borne hepatic encephalopathy in a 97-year-old woman using balloon-occluded retrograde transvenous obliteration: A case report  
*Nishi A, Kenzaka T, Sogi M, Nakaminato S, Suzuki T*
- 952** Development of Henoch-Schoenlein purpura in a child with idiopathic hypereosinophilia syndrome with multiple thrombotic onset: A case report  
*Xu YY, Huang XB, Wang YG, Zheng LY, Li M, Dai Y, Zhao S*
- 962** Three cases of jejunal tumors detected by standard upper gastrointestinal endoscopy: A case series  
*Lee J, Kim S, Kim D, Lee S, Ryu K*
- 972** Omental infarction diagnosed by computed tomography, missed with ultrasonography: A case report  
*Hwang JK, Cho YJ, Kang BS, Min KW, Cho YS, Kim YJ, Lee KS*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Sahand Samieirad, DDS, MS, MSc, Associate Professor, Oral and Maxillofacial Surgery Department, Mashhad Dental School, Mashhad University of Medical Sciences, Mashhad 9178613111, Iran. samieerads@mums.ac.ir

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Si Zhao; Production Department Director: Xu Guo; Editorial Office Director: Jin-Lei Wang.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

February 6, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



## Benign lymphoepithelial cyst of parotid gland without human immunodeficiency virus infection: A case report

Yan Liao, Yan-Jie Li, Xian-Wen Hu, Rui Wen, Pan Wang

**Specialty type:** Medicine, research and experimental

**Provenance and peer review:** Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B  
Grade C (Good): C  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** ISHIDA T, Japan; liu H, United States

**Received:** October 1, 2022

**Peer-review started:** October 1, 2022

**First decision:** November 30, 2022

**Revised:** December 13, 2022

**Accepted:** January 10, 2023

**Article in press:** January 10, 2023

**Published online:** February 6, 2023



**Yan Liao, Xian-Wen Hu, Rui Wen, Pan Wang,** Department of Nuclear Medicine, Affiliated Hospital of Zunyi Medical University, Zunyi 563003, Guizhou Province, China

**Yan-Jie Li,** Department of Critical Care Medicine, The Second Affiliated Hospital of Zunyi Medical University, Zunyi 563003, Guizhou Province, China

**Corresponding author:** Pan Wang, MD, Chief Doctor, Director, Professor, Department of Nuclear Medicine, Affiliated Hospital of Zunyi Medical University, No. 149 Dalian Road, Huichuan District, Zunyi 563003, Guizhou Province, China. [1298178828@qq.com](mailto:1298178828@qq.com)

### Abstract

#### BACKGROUND

Benign lymphoepithelial cyst (BLEC) of the parotid gland is a rare benign embryonic-dysplastic cystic tumor in the anterolateral neck that occurs most commonly in human immunodeficiency virus (HIV)-positive adults and rarely in non-acquired immune deficiency syndrome patients. The main presentation is a slow-growing, painless mass, and secondary infection may cause acute inflammatory symptoms.

#### CASE SUMMARY

A 44-year-old Chinese male patient presented with a 1-year history of a mass in the left side of the neck. On physical examination, a mass similar in size and shape to a quail egg was found in the left parotid gland. The mass was tough, without tenderness, and easily moveable. The results of HIV tests, including antibody and nucleic acid tests and CD4<sup>+</sup> T cell examination, were negative. Imaging examination revealed a left parotid gland mass. The patient underwent surgical treatment, and BLEC was diagnosed based on postoperative pathology. After 2 years of follow-up, the patient survived well without related discomfort.

#### CONCLUSION

The detailed characteristics of a BLEC in a patient without HIV infection contribute to an improved understanding of this rare disease.

**Key Words:** Benign lymphoepithelial cyst; Parotid gland; Human immunodeficiency virus; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Benign lymphoepithelial cyst (BLEC) of the parotid gland is a rare lesion seldom found in non-acquired immune deficiency syndrome patients. BLECs are benign embryonic dysplastic cystic tumors that typically occur in the anterolateral part of the neck. We present a case of BLEC of the parotid gland in a non-human immunodeficiency virus -infected patient in order to improve clinicians' understanding of the disease.

**Citation:** Liao Y, Li YJ, Hu XW, Wen R, Wang P. Benign lymphoepithelial cyst of parotid gland without human immunodeficiency virus infection: A case report. *World J Clin Cases* 2023; 11(4): 931-937

**URL:** <https://www.wjgnet.com/2307-8960/full/v11/i4/931.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v11.i4.931>

## INTRODUCTION

Benign lymphoepithelial cyst (BLEC) of the parotid gland, which is also known as branchial cleft cyst, is a rare benign cystic neoplasm of embryonic dysplasia. It usually occurs in the anterolateral region of the neck but has been reported in the oral cavity or parotid gland in rare cases[1,2]. This disease mainly presents as a slow-growing tumor and is not associated with recurrence or metastasis. These tumors usually occur in patients with human immunodeficiency virus (HIV) infection and are rarely encountered in non-HIV-infected patients[3]. To improve clinicians' understanding of this rare disease, the present report describes the imaging, histopathological, and diagnostic characteristics of a parotid gland BLEC found in a 44-year-old, non-HIV-infected patient. The case description is followed by a review of the relevant literature.

## CASE PRESENTATION

### Chief complaints

A 44-year-old Chinese man was admitted to the hospital on September 1, 2020, due to a painless mass on the left side of the neck that had been present for 1 year.

### History of present illness

One year previously, the patient incidentally noticed a painless mass on the left side of his neck approximately 2.0 cm × 2.0 cm in size. The patient had neck swelling occasionally, and the severity of these symptoms gradually increased over the course of the disease. The patient reported no tenderness, redness, swelling, or heat pain at the mass; no earache, swelling, or pain around the ear; and no cheek pain, numbness or difficulty when opening the mouth.

### History of past illness

The patient had a history of hypertension for 1 month and was regularly taking nifedipine sustained release tablets and Irbesartan tablets as antihypertensive treatment.

### Personal and family history

The patient had no personal or family history of benign or malignant tumors.

### Physical examination

Physical examination on admission revealed a temperature of 36.5°C, resting respiratory rate of 20 breaths/min, heart rate of 80 bpm, and blood pressure of 117/70 mmHg. On physical examination, a mass similar in size and shape to a quail egg was found in the left parotid gland, measuring about 2.0 cm × 2.0 cm in size. The mass felt tough, without tenderness and was easily movable with no local redness, swelling, heat or pain. No abnormalities were observed in the contralateral parotid gland.

### Laboratory examinations

The laboratory test results showed a white blood cell count of  $6.16 \times 10^9/L$ , with 74% neutrophils, 135 g/L hemoglobin, and a platelet count of  $277 \times 10^9/L$ . The erythrocyte sedimentation rate was 2 mm/h, and all results for routine urine, stool, liver and kidney function tests as well as electrolyte and blood biochemical index levels were within normal ranges. The results of HIV tests, including antibody and nucleic acid tests as well as CD4+ T cell examination, were negative. Furthermore, all results of testing for carcinoembryonic antigen, neuron-specific enolase, cytokeratin 19 fragment, squamous cell carcinoma antigen, blood coagulation, and immune indexes were negative.



### Imaging examinations

On August 31, 2020, ultrasound of the neck recognized a 2.2 cm × 2.2 cm hypoechoic mass in the left parotid gland, with a clear boundary, coarse calcification foci, and no blood flow signal (Figure 1). On September 2, 2020, neck computed tomography (CT) showed a round cystic lesion in the left parotid gland, with a maximum cross-sectional area of approximately 2.3 cm × 2.2 cm, uneven density, small patches with slightly high-density shadows, and clear boundaries (Figure 2A). On contrast-enhanced CT, the cyst wall was slightly enhanced in the arterial phase, but no obvious enhancement was observed in the cyst (Figure 2B). The enhancement degree in the venous phase was similar to that in the arterial phase (Figure 2C). No abnormality was found in the right parotid gland.

### PATHOLOGY

In general, the lesion was a pale round nodule with a complete capsule and small volume, accompanied by a small amount of parotid tissue attachment. The size was about 2.2 cm × 2.0 cm × 1.8 cm. The surface of the tumor was smooth and soft, and the surface skin mucosa was not abnormal, and the color was not special. The section was cystic, and the cyst was soybean dreg-like material with a wall thickness of 0.1 cm. Under light microscopy, the cyst wall was laminated squamous epithelium without epithelial nail process and the surface layer was mostly incomplete keratosis. The epithelium was surrounded by a large number of lymphoid stromata with lymphoid follicular formation and a center of occurrence (Figure 3). The histological features of the tumor were consistent with a diagnose of BLEC. The clinical and pathological data of this case are presented in Table 1.

### FINAL DIAGNOSIS

The patient was diagnosed with a non-HIV-infected BLEC of the parotid gland on the basis of history, clinical features, and ancillary examination (imaging and histopathology).

### TREATMENT

Because BLECs are benign, most cases of BLEC have a good prognosis. The principles of treatment are early diagnosis, infection control, and complete resection of the lesion without facial nerve injury. Treatment options include observation, repeat aspiration, sclerotherapy, radiotherapy, and surgery.

During the treatment period, the patient in the present case did not stop Nifedipine and Irbesartan for treatment of hypertension. Blood pressure and electrocardiography were normal at admission, and surgical treatment was possible under close monitoring of blood pressure. During surgery, care must be taken to identify the internal and external carotid arteries and vagus nerve, hypoglossal nerve, glossopharyngeal nerve, and superior laryngeal nerve to avoid injury to these structures. Finally, the patient and his family members chose surgical treatment (the specific surgical method was termed left parotid gland tumor + partial superficial lobectomy + facial nerve exploration and protection + fascia flap plasty). The operation was successful, and the patient returned to the ward safely after surgery.

### OUTCOME AND FOLLOW-UP

The patient was followed up for 2 years. CT/magnetic resonance imaging (MRI) and color ultrasound were not performed during follow-up, because the patient felt no obvious symptoms.


### DISCUSSION

BLEC is a rare cystic neoplasm with benign embryonic dysplasia. The first case of BLEC was reported by Hildebrandt in 1895, but only 21 cases were published by researchers through 1981. With the emergence of the HIV epidemic, the incidence of BLEC in the parotid gland gradually increased, and researchers found that BLECs were closely related to HIV infection in most cases, with BLEC as one of the early clinical manifestations. Around 3%-6% of HIV-positive adults and 1%-10% of HIV-positive children experience BLEC symptoms. In sharp contrast, BLECs are rarely found in non-HIV-infected patients, and the exact prevalence of BLEC in this population has not been reported[1-4]. In the present case, the results of HIV tests, including antibody and nucleic acid tests as well as CD4+ T cell examination, were negative. In previous reports, the disease was also known as branchial cleft cyst and usually observed in either the mandibular angle at the bottom of the outer regions or in the anterior



Table 1 Clinical and pathological data of this case of benign lymphoepithelial cyst of the parotid gland	
Finding	
Age, sex	44 years, male
Family history	None
Personal history	None
Chief complaint	Left neck painless mass present for 1 year
Physical examination	A quail egg-shaped mass on the left parotid gland, about 2.0 cm × 2.0 cm in size, tough, without tenderness, easily moveable, no local redness, swelling, heat or pain, and no abnormalities in the contralateral parotid gland
Laboratory examinations	HIV tests, including antibody and nucleic acid tests and CD4+ T cell examination, were negative
Ultrasound of the neck	A 2.2 cm × 2.2 cm hypoechoic mass in the left parotid gland, with a clear boundary, coarse calcification foci, and no blood flow signal (Figure 1)
Neck CT	A round cystic lesion in the left parotid gland, with a maximum cross-sectional area of 2.3 cm × 2.2 cm, uneven density, small patches with slightly high-density shadows, and clear boundaries (Figure 2A). On enhanced scanning, the cyst wall was slightly enhanced in the arterial phase, but no obvious enhancement was observed in the cyst (Figure 2B). The enhancement degree in the venous phase was similar to that in the arterial phase (Figure 2C)
Pathology	The cyst wall was laminated squamous epithelium without epithelial nail process and the surface layer was mostly incomplete keratosis (Figure 3A); The epithelium was surrounded by a large number of lymphoid stromata with lymphoid follicular formation and a center of occurrence (Figure 3B)
Final diagnosis	Non-HIV-infected BLEC of parotid gland
Treatment	Surgical treatment (specific surgical method was termed left parotid gland tumor + partial superficial lobectomy + facial nerve exploration and protection + fascia flap plasty)
Follow-up	No recurrence of symptoms in 2-year clinical follow-up


HIV: Human immunodeficiency virus; CT: Computed tomography; BLEC: Benign lymphoepithelial cyst.



**Figure 1 Color ultrasound of the neck.** It reveals a 2.2 cm × 2.2 cm hypoechoic mass in the left parotid gland, with a clear boundary, coarse calcification foci, and no blood flow signal.


portion of the sternocleidomastoid. It was found to occur less frequently in the mouth or parotid gland. In addition, its distribution in men and women is uniform. Although catheter jam seems to be a cause, the source of the blockage is often not obvious. Furthermore, its pathogenesis remains unclear. Currently, the majority of researchers believe that extraglandular lymphoid infiltration and/or intra-glandular lymphoid hyperplasia with duct obstruction and/or epithelial embedding is the main cause.

BLEC is a slow-growing tumor not associated with recurrence or metastasis[5-8]. The main symptom of this disease is a slowly growing, painless swollen mass, often combined with local symptoms such as dysphagia, dysphagia, dyspnea and stridor, and secondary infection can manifest as acute inflammatory symptoms, such as redness, swelling and fever[3,8]. The diagnosis of BLEC mainly depends on history, clinical manifestations, and preoperative auxiliary examination. Preoperative auxiliary diagnostic procedures include CT, MRI, ultrasound and fine needle aspiration (FNA)[9]. To date, imaging characteristics of this disease have been rarely reported. By reviewing the relevant literature[1,



DOI: 10.12998/wjcc.v11.i4.931 Copyright ©The Author(s) 2023.

**Figure 2 Neck computed tomography.** A: Plain computed tomography scan. A round cystic lesion was seen in the left parotid gland, with a maximum cross-sectional area of about 2.3 cm × 2.2 cm, uneven density, small patches with slightly high-density shadow, and clear boundary; B: Arterial phase. On enhanced scanning, the cyst wall was slightly enhanced in the arterial phase, but no obvious enhancement was observed in the cyst; C: Venous phase. The enhancement degree in the venous phase was similar to that in the arterial phase (orange arrows).



DOI: 10.12998/wjcc.v11.i4.931 Copyright ©The Author(s) 2023.

**Figure 3 Histopathology of benign lymphoepithelial cyst of the parotid gland.** A: The cyst wall was laminated squamous epithelium without epithelial nail process and the surface layer was mostly incomplete keratosis; B: The epithelium was surrounded by a large number of lymphoid stromata with lymphoid follicular formation and a center of occurrence (hematoxylin–eosin staining, magnification × 200).

9], we found that the CT manifestations of a typical BLEC are a single, thin-walled round or oval cystic foci in the parotid gland, with clear boundaries and low density in the cyst. On enhanced scanning, only the cyst wall shows mild to moderate enhancement, and no enhancement is seen in the cyst. In the present case, CT showed that the small patches with slightly high-density in the parotid gland were caused by viscous fluid in the capsule or lesions complicated with infection. MRI mainly showed a single thin-walled cystic lesion in the parotid gland, which was round or oval in shape, with a clear boundary, and the cyst wall showed a low signal. Intracystic T1 weighted imaging showed low signal, while T2 weighted imaging showed high signal. Moreover, the cyst wall was significantly enhanced on enhanced scanning, but no enhancement was found in the cyst[2]. Ultrasonic features of the lesions included a circular shape with a complete capsule and smooth lining, and cystic wall blood flow signals were not rich. The epithelial cells in a BLEC may contain keratin, and shed inflammatory cells can form a viscous gel-like yellowish-white liquid[10,11]. The internal echo of the cyst can generally appear as one of three types, including coarse granular hyperecho and cloud weak echo mixed signal, clear cystic echo through sound, and cluster weak echo and cystic echo mixed. In this case, ultrasound showed typical mixed signals including coarse granular hyperechoic signal and cloud weak echo signal. FNA can be used as an important auxiliary method for the clinical diagnosis of lateral cervical lesions. The criteria for FNA cytological diagnosis of BLEC include thick, yellow, pustular fluid; non-nucleated cells; keratinocytes; and squamous epithelial cells of varying maturity[1]. Histologically, the walls of BLEC cysts are usually covered with squamous epithelium and/or, in some cases, columnar or cuboidal cells. Lymphoid tissue with or without germinal centers in subepithelial connective tissue is the most prominent morphological feature[12]. Histological examination of the mass case showed that the cyst wall was laminated squamous epithelium without epithelial nail process and the surface layer was mostly incomplete keratosis, surrounded by a large number of lymphoid stromata with lymphoid follicular formation and a center of occurrence. Differential diagnoses include: (1) Warthin tumor[2],

which is more common in middle-aged and elderly patients and more likely to occur in the posterior lower pole of the parotid gland, with uneven internal density; (2) Intramuscular benign hemangioma [13], which is a fast-flowing intramuscular mass with uneven density and visible fat density in children that requires pathological examination for diagnosis; (3) Lymphoma [14], which is more common in men over 50 years old and appears as irregular soft tissue masses with a large range and uniform density on CT, without obvious calcification, cystic degeneration or necrosis, with diffuse growth to the surrounding area and mostly without adjacent bone destruction, and mild to moderate enhancement on enhanced scanning; (4) Thyroglossal duct cysts [15], which present as a painless mass in the front of the neck and are usually dumbbell shaped and movable when the tongue is extended or swallowed; CT shows a low-density, usually monocular parenchyma lesion during embryonic thyroid migration, mostly located in the midline and associated with the hyoid bone; (5) Lymphocele [16], which typically manifest as a cystic density focus with uniform density, clear boundary, thin cyst wall, no obvious exudation and calcification, and after enhancement, the cyst wall can present slightly uniform enhancement, with no enhancement in the cyst; (6) Metastatic lymph nodes [17], which are accompanied by a history of primary tumor, and on CT exhibit uneven density, calcification, cystic or necrotizing changes, uneven edges, adhesion to surrounding tissues, and obvious annular or peripheral enhancement; and (7) Lymphadenitis [18], which is more common in children and present with local redness, swelling, heat and pain while appearing mostly oval with a thick wall and ring and uniform enhancement without obvious wall nodules and calcification, but with a blurred surrounding fat space. The current treatment principles for this disease include early diagnosis, infection control, and cyst removal, and the treatment options include observation, repeated aspiration, sclerotherapy, radiotherapy, and surgical treatment [4]. Observation in asymptomatic patients shows that repeated aspiration therapy is generally ineffective, and cysts will recur within weeks to months. Sclerotherapy is limited in scope and is generally only used for cases in which the cystic fluid can be aspirated, while radiotherapy is mostly used for HIV-infected BLEC [3]. For our patient, surgery was the best treatment at present, and attention was paid to identify and avoid injury to adjacent blood vessels and nerves during surgery. In this case, surgical treatment was chosen by the patient, and the specific surgical procedure was termed left parotid gland tumor + partial superficial lobectomy + facial nerve exploration and protection + fascia flap plasty. The potential complications of surgical treatment include recurrence, the formation of persistent fistula, and cranial nerve injury, which require close follow-up. In this case, the patient survived well with no related discomfort after 2 years of follow-up.

## CONCLUSION

BLEC in the absence of HIV infection is a rare benign cystic tumor of embryonic dysplasia, known as a branchial cleft cyst. The disease progresses slowly. Because the imaging, histopathology, and diagnosis and treatment characteristics of non-HIV-infected BLEC are poorly understood, the present case is reported to increase readers' awareness of the disease.

## ACKNOWLEDGEMENTS

We thank Medjaden Inc. for scientific editing of this manuscript.

## FOOTNOTES

**Author contributions:** Liao Y and Li YJ contributed to the study conception and writing; Liao Y and Hu XW contributed to the collection of pathological findings; Hu XW and Wen R contributed to the analysis; Li YJ collected data; Wang P reviewed and revised the first draft; all authors read and approved the final version of the manuscript.

**Supported by** Zunyi Medical College Research Start Fund, No. 2017CK-1130-038.

**Informed consent statement:** Informed written consent was obtained from the patient for publication of this report and any accompanying images.

**Conflict-of-interest statement:** The authors declare that they have no conflict of interest to disclose.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license

their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Yan Liao 0000-0001-9799-9954; Yan-Jie Li 0000-0003-3650-8052; Xian-Wen Hu 0000-0001-7251-2329; Rui Wen 0000-0002-8522-2053; Pan Wang 0000-0001-7083-0016.

**S-Editor:** Liu GL

**L-Editor:** A

**P-Editor:** Liu GL

## REFERENCES

- 1 **Najib Z**, Berrada O, Lahjaoui M, Oukessou Y, Rouadi S, Abada RA, Roubal M, Mahtar M. Cervical lymphoepithelial cyst: Case report and literature review. *Ann Med Surg (Lond)* 2021; **61**: 185-187 [PMID: 33489106 DOI: 10.1016/j.amsu.2020.12.041]
- 2 **Joshi J**, Shah S, Agarwal D, Khasgiwal A. Benign lymphoepithelial cyst of parotid gland: Review and case report. *J Oral Maxillofac Pathol* 2018; **22**: S91-S97 [PMID: 29491615 DOI: 10.4103/jomfp.JOMFP\_252\_17]
- 3 **Pillai S**, Agarwal AC, Mangalore AB, Ramaswamy B, Shetty S. Benign Lymphoepithelial Cyst of the Parotid in HIV Negative Patient. *J Clin Diagn Res* 2016; **10**: MD05-MD06 [PMID: 27190845 DOI: 10.7860/JCDR/2016/17915.7609]
- 4 **Dave SP**, Pernas FG, Roy S. The benign lymphoepithelial cyst and a classification system for lymphocytic parotid gland enlargement in the pediatric HIV population. *Laryngoscope* 2007; **117**: 106-113 [PMID: 17202938 DOI: 10.1097/01.mlg.0000024]
- 5 **Nahata V**. Branchial Cleft Cyst. *Indian J Dermatol* 2016; **61**: 701 [PMID: 27904209 DOI: 10.4103/0019-5154.193718]
- 6 **Zimmerman KO**, Hupp SR, Bourguet-Vincent A, Bressler EA, Raynor EM, Turner DA, Rehder KJ. Acute upper-airway obstruction by a lingual thyroglossal duct cyst and implications for advanced airway management. *Respir Care* 2014; **59**: e98-e102 [PMID: 24170914 DOI: 10.4187/respcare.02513]
- 7 **Fujibayashi T**, Itoh H. Lymphoepithelial (so-called branchial) cyst within the parotid gland. Report of a case and review of the literature. *Int J Oral Surg* 1981; **10**: 283-292 [PMID: 6809660 DOI: 10.1016/s0300-9785(81)80073-7]
- 8 **Chavan S**, Deshmukh R, Karande P, Ingale Y. Branchial cleft cyst: A case report and review of literature. *J Oral Maxillofac Pathol* 2014; **18**: 150 [PMID: 24959062 DOI: 10.4103/0973-029X.131950]
- 9 **Glosser JW**, Pires CA, Feinberg SE. Branchial cleft or cervical lymphoepithelial cysts: etiology and management. *J Am Dent Assoc* 2003; **134**: 81-86 [PMID: 12555960 DOI: 10.14219/jada.archive.2003.0020]
- 10 **da Silva KD**, Coelho LV, do Couto AM, de Aguiar MCF, Tarquinio SBC, Gomes APN, Mendonça EF, Batista AC, Nonaka CFW, de Sena LSB, Alves PM, Libório-Kimura TN, Louredo BVR, Câmara J, Caldeira PC. Clinicopathological and immunohistochemical features of the oral lymphoepithelial cyst: A multicenter study. *J Oral Pathol Med* 2020; **49**: 219-226 [PMID: 31782199 DOI: 10.1111/jop.12978]
- 11 **Reynolds JH**, Wolinski AP. Sonographic appearance of branchial cysts. *Clin Radiol* 1993; **48**: 109-110 [PMID: 8004887 DOI: 10.1016/s0009-9260(05)81082-7]
- 12 **Hirota J**, Maeda Y, Ueta E, Osaki T. Immunohistochemical and histologic study of cervical lymphoepithelial cysts. *J Oral Pathol Med* 1989; **18**: 202-205 [PMID: 2475614 DOI: 10.1111/j.1600-0714.1989.tb00763.x]
- 13 **Wee SJ**, Park MC, Chung CM, Tak SW. Intramuscular hemangioma in the zygomatic minor muscle: a case report and literature review. *Arch Craniofac Surg* 2021; **22**: 115-118 [PMID: 33957738 DOI: 10.7181/acfs.2021.00087]
- 14 **Thanarajasingam G**, Bennani-Baiti N, Thompson CA. PET-CT in Staging, Response Evaluation, and Surveillance of Lymphoma. *Curr Treat Options Oncol* 2016; **17**: 24 [PMID: 27032646 DOI: 10.1007/s11864-016-0399-z]
- 15 **Corvino A**, Pignata S, Campanino MR, Corvino F, Giurazza F, Tafuri D, Pinto F, Catalano O. Thyroglossal duct cysts and site-specific differential diagnoses: imaging findings with emphasis on ultrasound assessment. *J Ultrasound* 2020; **23**: 139-149 [PMID: 32052384 DOI: 10.1007/s40477-020-00433-2]
- 16 **Hekiert A**, Newman J, Sargent R, Weinstein G. Spontaneous cervical lymphocele. *Head Neck* 2007; **29**: 77-80 [PMID: 17103412 DOI: 10.1002/hed.20484]
- 17 **Suh CH**, Baek JH, Choi YJ, Lee JH. Performance of CT in the Preoperative Diagnosis of Cervical Lymph Node Metastasis in Patients with Papillary Thyroid Cancer: A Systematic Review and Meta-Analysis. *AJNR Am J Neuroradiol* 2017; **38**: 154-161 [PMID: 27789450 DOI: 10.3174/ajnr.A4967]
- 18 **Desai S**, Shah SS, Hall M, Richardson TE, Thomson JE; Pediatric Research in Inpatient Settings (PRIS) Network. Imaging Strategies and Outcomes in Children Hospitalized with Cervical Lymphadenitis. *J Hosp Med* 2020; **15**: 197-203 [PMID: 31891560 DOI: 10.12788/jhm.3333]





Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

