

# World Journal of *Clinical Cases*

*World J Clin Cases* 2023 February 26; 11(6): 1224-1433



## Contents

Thrice Monthly Volume 11 Number 6 February 26, 2023

## OPINION REVIEW

- 1224 Collagen matrix scaffolds: Future perspectives for the management of chronic liver diseases  
*Martinez-Castillo M, Altamirano-Mendoza I, Zielinski R, Priebe W, Piña-Barba C, Gutierrez-Reyes G*

## MINIREVIEWS

- 1236 Sex dimorphism and metabolic profiles in management of metabolic-associated fatty liver disease  
*Martin-Grau M, Monleon D*
- 1245 Epidemiology and etiology of chemical ocular injury: A brief review  
*Akgun Z, Selver OB*
- 1252 Review of the prevalence, diagnostics, and containment measures of the current mpox outbreak  
*Sanyaolu A, Marinkovic A, Okorie C, Prakash S, Haider N, Dixon Y, Izurieta R, Badaru O, Smith S*
- 1261 Clinical and pathophysiological understanding of the hepatorenal syndrome: Still wrong or still not exactly right?  
*Wilde B, Canbay A, Katsounas A*
- 1267 Flare of the silent pandemic in the era of the COVID-19 pandemic: Obstacles and opportunities  
*Rayan RA*
- 1275 Implications of metabolic dysfunction associated fatty liver disease in COVID-19  
*Chakraborty R, Sharma D, Kapoor DU, Dwivedi A, Khabiya R, Sen S*

## ORIGINAL ARTICLE

## Retrospective Study

- 1287 Hyperglycemia in COVID-19 infection without diabetes mellitus: Association with inflammatory markers  
*Geetha HS, Singh G, Sekar A, Gogtay M, Singh Y, Abraham GM, Trivedi N*

## Clinical Trials Study

- 1299 Efficacy of invisible advancement correction for mandibular retraction in adolescents based on Pancherz analysis  
*Kong L, Liu XQ*

## Observational Study

- 1310 Survey study of the etiology of non-traumatic altered consciousness in the Emergency Department at Suez Canal University Hospital in Egypt  
*Moussa BS, Abd Elatiff ZM, Kamal Eldin Elhadary GM*

- 1318** Metformin effect on internal carotid artery blood flow assessed by area under the curve of carotid artery Doppler in women with polycystic ovarian syndrome

*Akram W, Nori W, Abdul Ghani Zghair M*

- 1330** Effect of continuous nursing combined with respiratory exercise nursing on pulmonary function of postoperative patients with lung cancer

*Qiu QX, Li WJ, Ma XM, Feng XH*

### CASE REPORT

- 1341** Functioning gonadotroph adenoma with hyperestrogenemia and ovarian hyperstimulation in a reproductive-aged woman: A case report and review of literature

*He Y, Gao YT, Sun L*

- 1349** Clinical manifestations of adult hereditary spherocytosis with novel *SPTB* gene mutations and hyperjaundice: A case report

*Jiang N, Mao WY, Peng BX, Yang TY, Mao XR*

- 1356** Post-traumatic cauda equina nerve calcification: A case report

*Liu YD, Deng Q, Li JJ, Yang HY, Han XF, Zhang KD, Peng RD, Xiang QQ*

- 1365** Endometriosis-associated endometrioid adenocarcinoma of the fallopian tube synchronized with endometrial adenocarcinoma: A case report

*Feng JY, Jiang QP, He H*

- 1372** Gemcitabine-induced peripheral vascular disease and prolonged response in a patient with metastatic pancreatic adenocarcinoma: A case report

*Fabien MB, Elodie P, Anna S, Addeo P, Meher B*

- 1379** Epidemic Japanese B encephalitis combined with contactin-associated protein-like 2 antibody-positive autoimmune encephalitis: A case report

*Huang P*

- 1385** Acute pancreatitis as initial presentation of acute myeloid leukemia-M2 subtype: A case report

*Yang WX, An K, Liu GF, Zhou HY, Gao JC*

- 1393** Postoperative jaundice related to *UGT1A1* and *ABCB11* gene mutations: A case report and literature review

*Jiang JL, Liu X, Pan ZQ, Jiang XL, Shi JH, Chen Y, Yi Y, Zhong WW, Liu KY, He YH*

- 1403** Hidrotic ectodermal dysplasia in a Chinese pedigree: A case report

*Liao MY, Peng H, Li LN, Yang T, Xiong SY, Ye XY*

- 1410** Hepatitis A virus-associated acute acalculous cholecystitis in an adult-onset Still's disease patient: A case report and review of the literature

*Chang CH, Wang YY, Jiao Y*

- 1419** Transverse myelitis caused by herpes zoster following COVID-19 vaccination: A case report

*Cho SY, Jang BH, Seo JW, Kim SW, Lim KJ, Lee HY, Kim DJ*

- 1426 Primary malignant melanoma of the esophagus: A case report

*Wang QQ, Li YM, Qin G, Liu F, Xu YY*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Goran Augustin, MD, MSc, PhD, Assistant Professor, Senior Scientist, Surgeon, Department of Surgery, University Hospital Centre Zagreb, Zagreb 10000, Croatia. [augustin.goran@gmail.com](mailto:augustin.goran@gmail.com)

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

February 26, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



Observational Study

# Survey study of the etiology of non-traumatic altered consciousness in the Emergency Department at Suez Canal University Hospital in Egypt

Bassant S Moussa, Zeinab M Abd Elatiff, Ghada M Kamal Eldin Elhadary

**Specialty type:** Emergency medicine

**Provenance and peer review:** Invited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B  
Grade C (Good): C  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Nakaji K, Japan

**Received:** October 14, 2022

**Peer-review started:** October 14, 2022

**First decision:** December 13, 2022

**Revised:** January 6, 2022

**Accepted:** February 3, 2023

**Article in press:** February 3, 2023

**Published online:** February 26, 2023



**Bassant S Moussa, Zeinab M Abd Elatiff, Ghada M Kamal Eldin Elhadary**, Emergency Medicine Department, Suez Canal University, Faculty of Medicine, Ismailia 41522, Egypt

**Corresponding author:** Bassant S Moussa, PhD, Senior Lecturer, Emergency Medicine Department, Suez Canal University, Faculty of Medicine, Ring Road, Ismailia 41522, Egypt. [bassant\\_sayed@med.suez.edu.eg](mailto:bassant_sayed@med.suez.edu.eg)

## Abstract

### BACKGROUND

Disorders of consciousness including coma in non-trauma patients can be caused by a wide variety of pathologies affecting the central nervous system. They represent a frequent challenge in emergency medicine and are combined with a very high in-hospital mortality. Hence, early treatment of these patients is vital and increases the likelihood of a good outcome.

### AIM

To identify the causes of altered consciousness presentation to the Emergency Department at Suez Canal University Hospital.

### METHODS

This was a descriptive cross-sectional study conducted on 87 patients with acute non-traumatic disturbed level of consciousness (DLOC) at the Emergency Department.

### RESULTS

The mean age of the studied patients was  $60.5 \pm 13.6$  years. Among them, 60% were males and 40% were females. The most common cause of acute non-traumatic DLOC was systemic infection, such as sepsis and septic shock (25.3%), followed by respiratory causes (24.1%) and neurological causes (18.4%).

### CONCLUSION

The most common cause of acute non-traumatic DLOC was systemic infections followed by respiratory and neurological causes.

**Key Words:** Disturbed level of consciousness; Non-traumatic; Emergency department

**Core Tip:** Disorders of consciousness including coma in non-trauma patients can be caused by a wide variety of pathologies affecting the central nervous system. This includes life-threatening medical, neurological or neurosurgical emergencies where timely medical intervention is vital. The aim of this cross-sectional observational study was to identify the causes of acute non-traumatic altered consciousness in the Emergency Department at Suez Canal University Hospital. Our study concluded that the most common cause of acute non-traumatic disturbed level of consciousness was systemic infections followed by respiratory and neurological causes.

**Citation:** Moussa BS, Abd Elatiff ZM, Kamal Eldin Elhadary GM. Survey study of the etiology of non-traumatic altered consciousness in the Emergency Department at Suez Canal University Hospital in Egypt. *World J Clin Cases* 2023; 11(6): 1310-1317

**URL:** <https://www.wjgnet.com/2307-8960/full/v11/i6/1310.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v11.i6.1310>

## INTRODUCTION

Disorders of consciousness including coma in non-trauma patients can be caused by a wide variability of pathologies affecting the central nervous system (CNS). This includes life-threatening medical, neurological or neurosurgical emergencies where timely medical intervention is vital[1]. Five–nine percent of all patients in Emergency Departments (EDs) present with acute non-traumatic disorders of consciousness, and up to two percent of patients are in a coma at admission. Therefore, they represent a frequent challenge in emergency medicine[2,3]. Furthermore, they are combined with a very high in-hospital mortality that accounts for 25%-48% of patients in western populations[4,5].

In a previous study examining the etiology and outcome of non-traumatic coma in a tertiary pediatric ED in Egypt, the most frequent etiologies were metabolic (33%), CNS infection (28%) and intracranial hemorrhage (13%). In the ED, 50% of those patients died[6]. Hypothermia, hypotension, flaccidity and poor Glasgow coma scale (GCS) score at admission correlated significantly with mortality. Forty-eight hours after admission, poor pulse volume, poor GCS, abnormal respiratory pattern/apnea and seizures correlated significantly with mortality[6].

Generally, the underlying pathologies of patients with disturbed consciousness have been classified into primary or focal injury to the CNS and secondary affection of the CNS resulting in a diffuse brain dysfunction, such as in metabolic disorder or intoxication[7]. The reported prevalences of structural *vs* metabolic coma varies from 28%-64% and 37%-75%, respectively[8]. The early treatment of these patients is vital, and diagnoses need to be confirmed or excluded promptly because a good outcome significantly depends on early treatment ("time is brain"), *e.g.*, in meningoencephalitis[9] or basilar artery occlusion[10].

There are no data on the pattern of altered consciousness presentation in the EDs in Egypt, which could make proper preparation of the departments for the potential needs regarding these patients suboptimal. Therefore, in this study, we aimed to identify the causes of altered consciousness presentations to the ED at Suez Canal University Hospital (Ismailia, Egypt).

## MATERIALS AND METHODS

### Study design

This observational, prospective, cross-sectional study was conducted from January 2021 to January 2022.

### Study setting

The study was conducted on data of patients admitted to the ED at Suez Canal University Hospital, Ismailia, Egypt.

### Study population

This study comprised 87 patients who were ≥ 18-year-old with acute non-traumatic disturbed level of consciousness (DLOC) *i.e.* GCS < 15. Patients with a history of recent head trauma, or with a history of neurological insult with residual altered consciousness or on medications that caused an altered level of consciousness (ALC) were excluded from the study. All data of the patients included in this study were collected after receiving informed written consent from the patients' first-degree relatives.



A pre-organized questionnaire was prepared, which included age, sex, onset of DLOC, and chronic illnesses. Clinical assessment of the patients included vital signs and grading of level of consciousness using the GCS. The appropriate laboratory and radiological investigations were completed to determine etiology of the DLOC. The outcome of the patients was determined as survival or death, and the mortality rate was calculated.

### Statistical analysis

Statistical analysis was performed using Statistical Package for the Social Sciences for Windows version 22 (IBM Corp., Armonk, NY, United States). *P* values < 0.05 were considered statistically significant. The confidence interval was set at 95%.

## RESULTS

### Baseline characteristics of the study population

The present study showed that the mean age of the studied patients was  $60.5 \pm 13.6$  years (range: 18-93 years). The majority of patients (64.5%) were  $\geq 60$  years, and 23% of the patients were between the ages of 50-59 years. The majority of the patients (60.0%) were males, with a male to female ratio of 1.5:1. All characteristics are presented in [Table 1](#).

### Chronic illnesses of the cohort

Reported chronic illnesses among the patients is shown in [Table 2](#). Hypertension (HTN) and diabetes were the most common chronic illness (43.7% and 34.5%, respectively).

### Distribution of the patients according to GCS

The majority of the patients (42.5%) had GCS between 6 and 11, followed by 33.3% of patients with GCS between 11 and 14, and 24.2% of patients with GCS between 3 and 5 ([Table 3](#)).

### Radiological investigations of the cohort

Computed tomography (CT) was primarily used for radiological investigations. CT of the brain showed that 53.0% of the patients had normal CT findings, while 23.0% of patients had abnormal CT findings. CT was not performed in 24.0% of patients.

### Distribution of the causes of non-traumatic DLOC

The most common cause of acute non-traumatic DLOC was systemic infection such as sepsis and septic shock ( $n = 22$ , 25.3%), which included urinary tract infection, cellulitis, infected bedsores and sepsis. The next most common causes were respiratory ( $n = 21$ , 24.1%) and neurological ( $n = 16$ , 18.4%). Among the respiratory causes, coronavirus disease 2019 (COVID-19) accounted for 15.0% of patients ( $n = 13$ ) and pneumonia accounted for 3.5% of patients ( $n = 3$ ). Among the neurological causes, stroke represented 8.1% of patients ( $n = 7$ ) followed by intracranial hemorrhage (7.0%,  $n = 6$ ); meningitis and status epilepticus were also present at lower rates. Metabolic causes were also found (11.5%), consisting primarily of diabetic ketoacidosis and dehydration associated with electrolyte disturbances. Further causes are shown in [Table 4](#).

The mortality rate was 25.3% ( $n = 22$ ), of which 41.0% of the patients died due to systemic infection followed by 31.8% due to respiratory causes (primarily COVID-19) and 13.6% due to oncological causes.

## DISCUSSION

This cross-sectional study aimed to determine the causes of non-traumatic disorders of consciousness. We observed that the mean age of these patients was  $60.5 \pm 13.6$  years, with range of 18-93 years, and that more male patients were eligible for participation in this study than females. Our findings were similar to Jung *et al* [11], in which they observed a mean age of  $68.81 \pm 16.40$  years in patients with ALC in the ED. Most of those patients were in their 80 s, accounting for 27.09% of the patients with ALC in the ED; the patients in their 70 s and 80 s also accounted for 53.49% of their study population. Cherukuri and Dhanawade [12] studied patients in the ED of Christian Medical College Hospital, Vellore (India), a tertiary medical care center, from January 2013 to April 2013; moreover, their study population was exclusively > 18 years of age, presenting with acute undifferentiated altered mental state (AMS) *i.e.* GCS < 15, with onset of symptom(s) being no more than 1 wk before ED presentation. Patients with chronic AMS and traumatic brain injuries were excluded from the study. Ultimately, in that study, the mean age was  $52.3 \pm 17.84$  years and they observed a male predominance (62.3%).

Our study observed several chronic diseases, including HTN, diabetes, chronic kidney disease, chronic liver disease, coronary artery disease and cerebrovascular stroke. HTN and diabetes were the



**Table 1 Baseline characteristics of the study population, *n* (%)**

Variables	<i>n</i> = 87	Alive	Dead	<i>P</i> value
Age, yr				N/A
mean ± SD	60.5 ± 13.6	60.4 ± 14.1	62.6 ± 12.2	
Median (range)	75 (18-93)	75 (18-93)	48 (30-78)	
Age groups				0.4 <sup>b</sup>
18-29	3 (3.4)	3 (4.6)	0 (0)	
30-39	3 (3.4)	2 (3.1)	1 (4.5)	
40-49	5 (5.7)	4 (6.2)	1 (4.5)	
50-59	20 (23.0)	15 (23.1)	5 (22.7)	
≥ 60	56 (64.5)	41 (63.0)	15 (68.3)	
Male	52 (60.0)	40 (61.5)	12 (54.5)	0.3 <sup>b</sup>
Female	35 (40.0)	25 (38.5)	10 (45.5)	

<sup>b</sup>*P*: Non-significant.

N/A: Not applicable; SD: Standard deviation.

**Table 2 Chronic illnesses of the cohort, *n* (%)**

Chronic illnesses	<i>n</i> = 87	Dead, <i>n</i> = 22	Alive, <i>n</i> = 65	<i>P</i> value
Chronic obstructive pulmonary disease	10 (11.5)	0 (0)	10 (15.4)	0.04 <sup>a</sup>
Chronic kidney disease	12 (13.8)	6 (27.3)	6 (9.2)	0.03 <sup>a</sup>
Diabetes mellitus	30 (34.5)	10 (45.5)	20 (30.8)	0.1 <sup>b</sup>
Cancer	8 (9.2)	5 (22.7)	3 (4.6)	0.02 <sup>a</sup>
Chronic liver disease	14 (16.1)	8 (36.4)	6 (9.2)	0.006 <sup>a</sup>
Cerebrovascular stroke	14 (16.1)	2 (9.1)	12 (18.5)	0.2 <sup>b</sup>
Coronary artery disease	10 (11.5)	3 (13.6)	7 (10.8)	0.4 <sup>b</sup>
Hypertension	38 (43.7)	12 (54.5)	26 (40)	0.1 <sup>b</sup>

<sup>a</sup>*P* < 0.05.

<sup>b</sup>*P*: Non-significant.

**Table 3 Distribution of the patients according to the Glasgow coma scale, *n* (%)**

GCS	<i>n</i> = 87	Dead, <i>n</i> = 22	Alive, <i>n</i> = 65
3-5	21 (24.2)	6 (27.3)	15 (23.1)
6-10	37 (42.5)	10 (45.4)	27 (41.5)
11-14	29 (33.3)	6 (27.3)	23 (35.4)
mean ± SD	8.5 ± 3.7	7.9 ± 3.5	8.6 ± 3.8
Median (range)	11 (3-14)	11 (3-14)	11 (3-14)

GCS: Glasgow coma scale; SD: Standard deviation.

most common chronic illnesses (43.7% and 34.5%, respectively). Our findings agreed with Cherukuri and Dhanawade[12], in which 40% of their patients had type 2 diabetes mellitus and 36.8% had systemic HTN. History of smoking was recorded for 11%. These results were similar to the results of the study by Sarker *et al*[13], in which HTN, diabetes, ischemic heart disease, chronic kidney disease and chronic liver disease were identified as chronic illnesses among their study population. HTN and diabetes mellitus

**Table 4 Distribution of the causes of non-traumatic disturbed level of consciousness, *n* (%)**

Causes of disturbed level of consciousness	<i>n</i> = 87	Dead, <i>n</i> = 65	Alive, <i>n</i> = 22	<i>P</i> value
Neurological	16 (18.4)	15 (23.1)	1 (4.5)	0.5 <sup>b</sup>
Intracranial hemorrhage	6 (7.0)	6 (9.2)	0 (0)	
Stroke	7 (8.1)	6 (9.2)	1 (4.6)	
Meningitis	1 (1.1)	1 (1.5)	0 (0)	
Cavernous sinus thrombosis	1 (1.1)	1 (1.5)	0 (0)	
Status epileptics	1 (1.1)	1 (1.5)	0 (0)	
Metabolic	10 (11.5)	10 (15.4)	0 (0)	
Hypoglycemia	1 (1.1)	1 (1.5)	0 (0)	
Diabetic ketoacidosis	6 (7.0)	6 (9.2)	0 (0)	
Dehydration and poor oral feeding	3 (3.5)	3 (4.6)	0 (0)	
Systemic infection sepsis	22 (25.3)	13 (20.0)	9 (41.0)	
Respiratory	21 (24.1)	14 (21.5)	7 (31.8)	
COVID-19	13 (15.0)	7 (10.8)	6 (27.3)	
Pneumonia	3 (3.5)	3 (4.6)	0 (0)	
Respiratory failure	5 (5.8)	4 (6.2)	1 (4.5)	
Organ dysfunction	8 (9.2)	7 (10.8)	1 (4.5)	
Hepatic encephalopathy	4 (4.6)	3 (4.6)	1 (4.5)	
Uremic encephalopathy	1 (1.1)	1 (1.5)	0 (0)	
Hypertensive encephalopathy	3 (3.5)	3 (4.6)	0 (0)	
Cardiovascular	2 (2.3)	1 (1.5)	1 (4.5)	
Cardiogenic shock	1 (1.1)	1 (1.5)	0 (0)	
Pulmonary edema	1 (1.1)	0 (0)	1 (4.5)	
Oncological	5 (5.8)	2 (3.1)	3 (13.6)	
Drug overdose	1 (1.1)	1 (1.5)	0 (0)	
Unknown	2 (2.3)	2 (3.1)	0 (0)	

<sup>b</sup>*P*: Non-significant.

COVID-19: Coronavirus disease 2019.

were also the most common (26% and 17%, respectively) in their study.

Our study showed that the mean GCS was  $8.5 \pm 3.7$ . The majority of the patients (42.5%) had moderate DLOC, followed by mild and then severe DLOC. In a study by Sarker *et al*[13], more than half of the patients (53%) had GCS between 6 and 10 (moderate), followed by 27% with GCS between 3 and 5 (severe) and 20% with GCS between 11 and 14 (mild) at the time of presentation. They observed a mean GCS of  $7.7 \pm 3.1$ .

In our study, the most common cause of acute non-traumatic DLOC was systemic infection such as sepsis and septic shock ( $n = 22$ , 25.3%), followed by respiratory causes ( $n = 21$ , 24.1%) and neurological causes ( $n = 16$ , 18.4%). Similarly, Jung *et al*[11] found that the leading cause of ALC in the ED was systemic infection, which accounted for approximately 30% of the cases. The second most common cause was metabolic (21.07%), and the third most common cause was stroke (18.19%), which was diagnosed when the acute CNS symptoms were compatible with the brain lesions found on neuroimaging. Cherukuri and Dhanawade[12] found that neurological conditions were the most important cause for AMS, accounting for 37.1% of their patients. Other etiologies were metabolic and endocrine causes (18%) followed by infections (13%).

In a study by Schmidt *et al*[14], the main diagnoses were classified into acute primary brain lesions (39%), primary brain pathologies without acute lesions (25%) and pathologies that affected the brain secondarily (36%). In another study by Braun *et al*[9], in which 58% of the studied patients had neurological causes for coma (intracranial hemorrhage, stroke, and epilepsy), followed by intoxication (16.6%), cardiovascular cause (5.8%) and respiratory cause (5.5%), respectively. Also in a study by Idro

*et al*[15], infections of cerebral malaria were the primary cause of AMS in a study from Ethiopia. Studies of AMS by Sporer *et al*[16] found that substance abuse was the primary cause. The etiology and the characteristics of patients with DLOC varies between different countries and depends on many factors including regional or demographic backgrounds, location of the hospital, the national health care system and medical resources. Accordingly, our cohort was comprised of an elderly population with multiple comorbidities. Typically, elderly people are neglected in our society and do not receive prompt treatment of illnesses, which explains why we observed systemic infections as the most common cause of DLOC. In addition, our study was conducted during the COVID-19 pandemic, which may explain the number of patients that presented with acute DLOC due to respiratory causes (15% of the respiratory causes were due to COVID-19).

In our study, the mortality rate was 25.3% ( $n = 22$ ), of which 41% of the patients died due to systemic infection followed by respiratory causes (primarily COVID-19) and oncological causes. It was similar to a study by Forsberg *et al*[17], in which the total hospital mortality was 26.5%. Kekec *et al*[18] also reported higher mortality of 20.1% in patients with AMS. In contrast to our results, the mortality rate among AMS patients in the study by Cherukuri and Dhanawade[12] was 11.5%. Kanich *et al*[2], Leong *et al*[19] and Xiao *et al*[20] reported mortality rates ranging from 9% to 11%.

There were several limitations to this study. First, as this was a single-center study, the sample size was small as our center is a tertiary care facility and only receives patients 3 d/wk. Second, the scope of this study was limited to clinical practice in the ED with exclusion of traumatic DLOC, psychiatric causes and patients with cardiac arrest on presentation to ED. ALC has a wide variation of causes, and it can be challenging and time-consuming to achieve a definitive diagnosis, which may require additional evaluation that is not available in the ED. Moreover, two or more causes can occasionally be present concomitantly. The study was also carried out during outbreaks of COVID-19, which affected the causes of mortality. These obstacles make it challenging to study ALC in the ED, indicating that further study and analysis are needed.

## CONCLUSION

DLOC including coma in non-traumatic patients can be caused by a wide variety of pathologies affecting the CNS. They represent a frequent challenge in emergency medicine with a very high in-hospital mortality. Early treatment of these patients is vital and good outcomes depend on early treatment. This cross-sectional study revealed that systemic infection was the most common cause of DLOC, followed by respiratory causes and neurological diseases. Patients with systemic infections, COVID-19 and oncological diseases had the highest mortality among our studied patients.

## ARTICLE HIGHLIGHTS

### Research background

Disorders of consciousness including coma in non-traumatic patients can be caused by a wide variety of pathologies affecting the central nervous system including life-threatening medical, neurological or neurosurgical emergencies where timely medical intervention is vital. The early treatment of these patients is vital, and diagnoses need to be confirmed or excluded promptly.

### Research motivation

There are no data on the pattern of altered consciousness presentation in Emergency Departments (EDs) in Egypt, which could make proper preparation for the potential needs of these patients suboptimal.

### Research objectives

The primary objective of our study was to identify the causes of altered consciousness presentation to the ED at Suez Canal University Hospital (Egypt). The secondary objectives were to describe the treatment modalities for patients presenting with altered consciousness and to assess the mortality rate among patients presenting with altered consciousness.

### Research methods

This study was conducted on 87 patients, all of whom were  $\geq 18$ -year-old with acute non-traumatic disturbed level of consciousness (DLOC) *i.e.* Glasgow coma scale  $< 15$ . All data of the patients included in this study had been collected after receiving informed written consent from the patients' first-degree relatives. The outcomes of the patients were determined as survival or death, and the mortality rate was calculated.

# Research results

In our study, the most common cause of acute non-traumatic DLOC was systemic infection such as sepsis and septic shock ( $n = 22$ , 25.3%), followed by respiratory causes ( $n = 21$ , 24.1%) and neurological causes ( $n = 16$ , 18.4%). The mortality rate was 25.3% ( $n = 22$ ) of which 41.0% of the patients died due to systemic infection, followed by 31.8% due to respiratory causes [primarily coronavirus disease 2019 (COVID-19)] and 13.6% due to oncological causes.

# Research conclusions

The most common cause of acute non-traumatic DLOC was systemic infections followed by respiratory and neurological causes.

# Research perspectives

Further study and analysis are needed to overcome the challenges of a small sample size and outbreaks of COVID-19 encountered in our study.

# ACKNOWLEDGEMENTS

We thank all the participants in this study for their great help during this work.

# FOOTNOTES

**Author contributions:** Moussa BS and Abd Elatiff ZM collected the data; Moussa BS assessed the results; Kamal Eldin Elhadary GM and Abd Elatiff ZM wrote and revised the manuscript; all authors read and approved the final manuscript.

**Institutional review board statement:** Ethics approval and consent to participate were obtained from the Ethical Committee of the Faculty of Medicine of Suez Canal University.

**Conflict-of-interest statement:** All the authors declare having no conflicts of interest.

**Data sharing statement:** All data generated or analyzed during this study are included in this published article and supplementary materials section.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** Egypt

**ORCID number:** Bassant S Moussa 0000-0002-4413-0355.

**S-Editor:** Wang DM

**L-Editor:** A

**P-Editor:** Wang DM

# REFERENCES

- 1 Edlow JA, Rabinstein A, Traub SJ, Wijdicks EF. Diagnosis of reversible causes of coma. *Lancet* 2014; **384**: 2064-2076 [PMID: 24767707 DOI: 10.1016/S0140-6736(13)62184-4]
- 2 Kanich W, Brady WJ, Huff JS, Perron AD, Holstege C, Lindbeck G, Carter CT. Altered mental status: evaluation and etiology in the ED. *Am J Emerg Med* 2002; **20**: 613-617 [PMID: 12442240 DOI: 10.1053/ajem.2002.35464]
- 3 Martikainen K, Seppä K, Viita P, Rajala S, Laippala P, Keränen T. Transient loss of consciousness as reason for admission to primary health care emergency room. *Scand J Prim Health Care* 2003; **21**: 61-64 [PMID: 12718464 DOI: 10.1080/02834310000591]
- 4 Forsberg S, Höjer J, Enander C, Ludwigs U. Coma and impaired consciousness in the emergency room: characteristics of poisoning versus other causes. *Emerg Med J* 2009; **26**: 100-102 [PMID: 19164617 DOI: 10.1136/emj.2007.054536]
- 5 Weiss N, Regard L, Vidal C, Luque Y, Taldir G, Vallet H, Diehl JL, Fagon JY, Guerot E. Causes of coma and their evolution in the medical intensive care unit. *J Neurol* 2012; **259**: 1474-1477 [PMID: 22231871 DOI: 10.1007/s00415-011-6388-z]
- 6 Fouad H, Haron M, Halawa EF, Nada M. Nontraumatic coma in a tertiary pediatric emergency department in egypt:

- etiology and outcome. *J Child Neurol* 2011; **26**: 136-141 [PMID: [20606061](#) DOI: [10.1177/0883073810374358](#)]
- 7 **Posner JB**, Saper CB, Schiff N, Plum F. Plum and Posner's diagnosis of stupor and coma, Contemporary neurology series, vol. 71. 4th ed. New York: Oxford University Press, 2007
- 8 **Horsting MW**, Franken MD, Meulenbelt J, van Klei WA, de Lange DW. The etiology and outcome of non-traumatic coma in critical care: a systematic review. *BMC Anesthesiol* 2015; **15**: 65 [PMID: [25924678](#) DOI: [10.1186/s12871-015-0041-9](#)]
- 9 **Braun M**, Schmidt WU, Möckel M, Römer M, Ploner CJ, Lindner T. Coma of unknown origin in the emergency department: implementation of an in-house management routine. *Scand J Trauma Resusc Emerg Med* 2016; **24**: 61 [PMID: [27121376](#) DOI: [10.1186/s13049-016-0250-3](#)]
- 10 **Vergouwen MD**, Algra A, Pfefferkorn T, Weimar C, Rueckert CM, Thijs V, Kappelle LJ, Schonewille WJ; Basilar Artery International Cooperation Study (BASICS) Study Group. Time is brain(stem) in basilar artery occlusion. *Stroke* 2012; **43**: 3003-3006 [PMID: [22989501](#) DOI: [10.1161/STROKEAHA.112.666867](#)]
- 11 **Jung S**, Jeon JC, Jung CG, Cho YW, Kim KT. Altered level of consciousness in the ED. *J Neurocrit Care* 2020; **13**: 86-92 [DOI: [10.18700/jnc.200010](#)]
- 12 **Cherukuri SK**, Dhanawade VS. Altered mental status in the emergency department, a retrospective analysis. *Curr Med Issues* 2020; **18**: 300-304 [DOI: [10.4103/emi.emi\\_64\\_20](#)]
- 13 **Sarker PS**, Rahman MS, Biswas PK, Chowdhury MM, Karmaker M, Azad KA. Aetiology and Short-term Outcome of Altered Level among Patients in Medicine Department of a Tertiary Hospital. *J Med* 2017; **18**: 80-85 [DOI: [10.3329/jom.v18i2.33685](#)]
- 14 **Schmidt WU**, Ploner CJ, Lutz M, Möckel M, Lindner T, Braun M. Causes of brain dysfunction in acute coma: a cohort study of 1027 patients in the emergency department. *Scand J Trauma Resusc Emerg Med* 2019; **27**: 101 [PMID: [31699128](#) DOI: [10.1186/s13049-019-0669-4](#)]
- 15 **Idro R**, Marsh K, John CC, Newton CR. Cerebral malaria: mechanisms of brain injury and strategies for improved neurocognitive outcome. *Pediatr Res* 2010; **68**: 267-274 [PMID: [20606600](#) DOI: [10.1203/PDR.0b013e3181ee738](#)]
- 16 **Sporer KA**, Solares M, Durant EJ, Wang W, Wu AH, Rodriguez RM. Accuracy of the initial diagnosis among patients with an acutely altered mental status. *Emerg Med J* 2013; **30**: 243-246 [PMID: [22362650](#) DOI: [10.1136/emmermed-2011-200452](#)]
- 17 **Forsberg S**, Höjer J, Ludwigs U. Prognosis in patients presenting with non-traumatic coma. *J Emerg Med* 2012; **42**: 249-253 [PMID: [20542655](#) DOI: [10.1016/j.jemermed.2010.04.021](#)]
- 18 **Kekec Z**, Senol V, Koc F, Seydaoglu G. Analysis of altered mental status in Turkey. *Int J Neurosci* 2008; **118**: 609-617 [PMID: [18446577](#) DOI: [10.1080/00207450701849133](#)]
- 19 **Leong LB**, Jian KH, Vasu A, Seow E. Prospective study of patients with altered mental status: clinical features and outcome. *Int J Emerg Med* 2008; **1**: 179-182 [PMID: [19384512](#) DOI: [10.1007/s12245-008-0049-8](#)]
- 20 **Xiao HY**, Wang YX, Xu TD, Zhu HD, Guo SB, Wang Z, Yu XZ. Evaluation and treatment of altered mental status patients in the emergency department: Life in the fast lane. *World J Emerg Med* 2012; **3**: 270-277 [PMID: [25215076](#) DOI: [10.5847/wjem.j.issn.1920-8642.2012.04.006](#)]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

