

World Journal of *Clinical Cases*

World J Clin Cases 2023 February 26; 11(6): 1224-1433



Contents

Thrice Monthly Volume 11 Number 6 February 26, 2023

OPINION REVIEW

- 1224 Collagen matrix scaffolds: Future perspectives for the management of chronic liver diseases
Martinez-Castillo M, Altamirano-Mendoza I, Zielinski R, Priebe W, Piña-Barba C, Gutierrez-Reyes G

MINIREVIEWS

- 1236 Sex dimorphism and metabolic profiles in management of metabolic-associated fatty liver disease
Martin-Grau M, Monleon D
- 1245 Epidemiology and etiology of chemical ocular injury: A brief review
Akgun Z, Selver OB
- 1252 Review of the prevalence, diagnostics, and containment measures of the current mpox outbreak
Sanyaolu A, Marinkovic A, Okorie C, Prakash S, Haider N, Dixon Y, Izurieta R, Badaru O, Smith S
- 1261 Clinical and pathophysiological understanding of the hepatorenal syndrome: Still wrong or still not exactly right?
Wilde B, Canbay A, Katsounas A
- 1267 Flare of the silent pandemic in the era of the COVID-19 pandemic: Obstacles and opportunities
Rayan RA
- 1275 Implications of metabolic dysfunction associated fatty liver disease in COVID-19
Chakraborty R, Sharma D, Kapoor DU, Dwivedi A, Khabiya R, Sen S

ORIGINAL ARTICLE

Retrospective Study

- 1287 Hyperglycemia in COVID-19 infection without diabetes mellitus: Association with inflammatory markers
Geetha HS, Singh G, Sekar A, Gogtay M, Singh Y, Abraham GM, Trivedi N

Clinical Trials Study

- 1299 Efficacy of invisible advancement correction for mandibular retraction in adolescents based on Pancherz analysis
Kong L, Liu XQ

Observational Study

- 1310 Survey study of the etiology of non-traumatic altered consciousness in the Emergency Department at Suez Canal University Hospital in Egypt
Moussa BS, Abd Elatiff ZM, Kamal Eldin Elhadary GM

- 1318** Metformin effect on internal carotid artery blood flow assessed by area under the curve of carotid artery Doppler in women with polycystic ovarian syndrome

Akram W, Nori W, Abdul Ghani Zghair M

- 1330** Effect of continuous nursing combined with respiratory exercise nursing on pulmonary function of postoperative patients with lung cancer

Qiu QX, Li WJ, Ma XM, Feng XH

CASE REPORT

- 1341** Functioning gonadotroph adenoma with hyperestrogenemia and ovarian hyperstimulation in a reproductive-aged woman: A case report and review of literature

He Y, Gao YT, Sun L

- 1349** Clinical manifestations of adult hereditary spherocytosis with novel *SPTB* gene mutations and hyperjaundice: A case report

Jiang N, Mao WY, Peng BX, Yang TY, Mao XR

- 1356** Post-traumatic cauda equina nerve calcification: A case report

Liu YD, Deng Q, Li JJ, Yang HY, Han XF, Zhang KD, Peng RD, Xiang QQ

- 1365** Endometriosis-associated endometrioid adenocarcinoma of the fallopian tube synchronized with endometrial adenocarcinoma: A case report

Feng JY, Jiang QP, He H

- 1372** Gemcitabine-induced peripheral vascular disease and prolonged response in a patient with metastatic pancreatic adenocarcinoma: A case report

Fabien MB, Elodie P, Anna S, Addeo P, Meher B

- 1379** Epidemic Japanese B encephalitis combined with contactin-associated protein-like 2 antibody-positive autoimmune encephalitis: A case report

Huang P

- 1385** Acute pancreatitis as initial presentation of acute myeloid leukemia-M2 subtype: A case report

Yang WX, An K, Liu GF, Zhou HY, Gao JC

- 1393** Postoperative jaundice related to *UGT1A1* and *ABCB11* gene mutations: A case report and literature review

Jiang JL, Liu X, Pan ZQ, Jiang XL, Shi JH, Chen Y, Yi Y, Zhong WW, Liu KY, He YH

- 1403** Hidrotic ectodermal dysplasia in a Chinese pedigree: A case report

Liao MY, Peng H, Li LN, Yang T, Xiong SY, Ye XY

- 1410** Hepatitis A virus-associated acute acalculous cholecystitis in an adult-onset Still's disease patient: A case report and review of the literature

Chang CH, Wang YY, Jiao Y

- 1419** Transverse myelitis caused by herpes zoster following COVID-19 vaccination: A case report

Cho SY, Jang BH, Seo JW, Kim SW, Lim KJ, Lee HY, Kim DJ

- 1426 Primary malignant melanoma of the esophagus: A case report

Wang QQ, Li YM, Qin G, Liu F, Xu YY

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Goran Augustin, MD, MSc, PhD, Assistant Professor, Senior Scientist, Surgeon, Department of Surgery, University Hospital Centre Zagreb, Zagreb 10000, Croatia. augustin.goran@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

February 26, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Observational Study

Metformin effect on internal carotid artery blood flow assessed by area under the curve of carotid artery Doppler in women with polycystic ovarian syndrome

Wisam Akram, Wassan Nori, Muna Abdul Ghani Zghair

Specialty type: Medicine, research and experimental

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: He XM, China; Spinelli L, Italy; Zeng Y, China

Received: November 30, 2022

Peer-review started: November 30, 2022

First decision: January 19, 2023

Revised: January 19, 2023

Accepted: January 31, 2023

Article in press: January 31, 2023

Published online: February 26, 2023



Wisam Akram, Wassan Nori, Department of Obstetrics and Gynecology, Mustansiriyyah, Al Saydihya 10052, Baghdad, Iraq

Muna Abdul Ghani Zghair, Department of Radiology, Mustansiriyyah, Al Saydihya 10052, Baghdad, Iraq

Corresponding author: Wassan Nori, PhD, Academic Editor, Academic Research, Senior Researcher, Department of Obstetrics and Gynecology, Mustansiriyyah, No. 38, Al Amin Street, Al Saydihya 10052, Baghdad, Iraq. dr.wassan76@uomustansiriyyah.edu.iq

Abstract

BACKGROUND

Insulin resistance (IR) was reported in most polycystic ovarian syndrome (PCOS) cases. Metformin, a biguanide drug, successfully reduced IR. Homeostatic Model Assessment for IR (HOMA-IR) and Doppler parameters assessed metformin's effectiveness.

AIM

To verify whether the area under the curve of the internal carotid artery (AUC-ICA) Doppler wave can be a useful marker for assessing IR among PCOS cases who presented with menstrual irregularity and were treated with metformin over 6 mo.

METHODS

An observational, cross-sectional study recruited 54 eligible PCOS women; the anthropometrics were as follows: age, body mass index (BMI), menstrual cycle days, biochemical serum cholesterol, low and high-density lipoprotein, sex hormone-binding globulin, fasting blood glucose, and HOMA-IR, hormonal testosterone, luteinizing hormone over follicle-stimulating hormone ratio, and ultrasonic pulsatility index (PI) and resistance index (RI), carotid artery intima-media thickness (CIMT) and (AUC-ICA) parameters were initially recorded and repeated 3 mo and 6 mo later with metformin tab 500 mg; three times/day for 6 mo. In addition, AUC-ICA was assessed by taking repeated systolic and diastolic wave height measurements.

RESULTS

Metformin caused a progressive reduction in BMI, menstrual cycle days, biochemical hormonal, and Doppler parameters (CIMT, PI, RI, and AUC-ICA). AUC-ICA correlated strongly to all PCOS parameters. AUC-ICA correlated inversely with treatment time ($r = -0.98$, $P < 0.001$) and positively with HOMA-IR ($r = 0.98$, $P < 0.0001$). Via the best subset regression model, the AUC-ICA had the highest predictive value for HOMA-IR.

CONCLUSION

AUC-ICA preceded PI, RI, and CIMT with a strong, meaningful correlation to all PCOS parameters, making it a reliable marker for the assessment of IR, especially during metformin therapy. Further studies are recommended to promote the application in practice.

Key Words: Carotid artery intima-media thickness; Insulin resistance; Internal carotid artery Doppler; Metformin; Polycystic ovarian syndrome

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Women with polycystic ovarian syndrome (PCOS) suffer from insulin resistance and an atherogenic state evidenced by increased carotid artery intima-media thickness (CIMT). Fortunately, this increase in CIMT is reversible with metformin therapy, an insulin-sensitizing drug. Moreover, treated cases have restored ovulatory cycles and exhibited reduced androgen levels. First, we primarily aimed to examine whether the area under the curve of the internal carotid artery (AUC-ICA) Doppler is related to insulin resistance among PCOS women who presented with menstrual disturbances. Second, is to examine if AUC-ICA can be a useful marker in assessing changes in insulin resistance among treated cases with metformin for follow-up.

Citation: Akram W, Nori W, Abdul Ghani Zghair M. Metformin effect on internal carotid artery blood flow assessed by area under the curve of carotid artery Doppler in women with polycystic ovarian syndrome. *World J Clin Cases* 2023; 11(6): 1318-1329

URL: <https://www.wjgnet.com/2307-8960/full/v11/i6/1318.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i6.1318>

INTRODUCTION

Polycystic ovarian syndrome (PCOS) is still one of the most frequently challenging problems facing gynecologists worldwide. Despite much work conducted to understand its nature, its long-term complications from cardiovascular, infertility, and obesity-related problems are still major issues facing all affected women[1-3]. PCOS women are characterized by a state of life-long insulin resistance (IR) with permanently elevated serum insulin[4].

This hyperinsulinemia disturbs ovarian steroidogenesis (estrogens and progesterone) and the pituitary secretion of gonadotropins, represented by chronically elevated serum luteinizing hormones (LH) and serum testosterone[5].

Elevated serum testosterone mainly affects lipid metabolism with an elevation of low-density lipoprotein (LDL) and serum cholesterol and a reduction of serum high-density lipoprotein (HDL)[6]. This disturbed lipid metabolism; leads to acute atherosclerotic changes affecting virtually every artery in the body but mostly in the medium-sized artery like the internal carotid artery, which can be scanned by B-mode ultrasound device. The atherogenic state is evidenced by increased carotid artery intima-media thickness (CIMT); fortunately, this increase in CIMT is usually reversible with the use of insulin-sensitizing drugs among young women below 35 years of age, which makes this reduction one of the most important prognostic variables for IR complications[7].

According to Bernoulli's rule, the increased CIMT per se is associated with a reduction in the arterial diameter, which consequently increases the artery's blood flow speed[8].

According to this rule, the narrower the artery lumen is, the more blood speed per unit area in its section (inverse relationship). Modern Doppler devices allow arterial blood speed to be measured and graphed on paper strips directly from scanned arteries. With ultrasound waves beamed at 60 degrees, we can measure the difference in phase shift of the ultrasound wave[9].

The most commonly used Doppler indices to assess arterial blood flow are pulsatility and resistant index (PI and RI), respectively[10].

The PI is an ultrasonic blood flow parameter calculated from the highest, lowest, and mean Doppler frequency shifts during a given cardiac cycle. As for the RI, it estimates the resistance in a pulsatile

vascular tree[11].

However, a new parameter can be easily conducted with current Doppler devices: Measuring the area under the curve (AUC) of one complete heartbeat-associated blood flow graph in an artery represented by a systolic and diastolic wave.

Multiple readings of the heights of a systolic and diastolic wave of a single Doppler wave measure AUC under one Doppler wave. In physical terms, measuring AUC means measuring the amount of blood passing per unit area of the artery cross-section area[12]. For that, the speed of blood flow should increase while the amount of blood passed per unit sectional area of the internal carotid artery should reduce or remain the same according to the resistance distal to the vessel[12].

Metformin is an insulin sensitizer used for type 2 diabetes. Since many PCOS women are IR, metformin administration proved its efficacy in restoring ovulatory cycles and reducing androgen levels. This biguanide inhibits hepatic glycogenesis, increases insulin sensitivity in the periphery, increases glucose uptake, and decreases insulin secretion[1].

Earlier studies in the field have shown reduced atherogenic indicators by insulin sensitizers like metformin and inositol for 3-6 mo, including dyslipidemia, IR, and improved endothelial function and coronary flow. Furthermore, there was a reduction in CIMT among treated women[13]. The aim of the study is to verify whether the AUC of the internal carotid artery (AUC-ICA) Doppler is related to IR among women with PCOS who presented with menstrual disturbances. Second, to examine if AUC-ICA can be a useful marker for the assessment of IR among women treated with metformin for follow-up.

MATERIALS AND METHODS

The current study was a cross-sectional study conducted in AL Yarmouk Teaching hospital between April 2019 till December 2021. The ethical committee of Mustansiriyah University approved the study dated February/21/2019 (IRB No. 115). Participants were briefed about the study's aim and methodology; all gave their consent before enrollment, and the Helsinki declaration was followed.

From the outpatient clinic, unmarried women with PCOS who presented with menstrual cycle abnormality and/or hirsutism, the age range of 18-35 years, and a body mass index (BMI) range of 18 to and 30 were invited to participate in the study. During this period, all patients had been prescribed a metformin tab. (Merck Santé/France) 500 mg TID times /d.

The diagnosis of PCOS was made based on Rotterdam criteria, where 2 out of 3 criteria confirm PCOS diagnosis[14]. (1) Oligomenorrhea is defined as < than six cycles/ 12 mo, or amenorrhea is defined as a complete absence of the menstrual cycle (more than 90 d); (2) Clinical hyperandrogenism, with or without acne; and (3) Ultrasonic features of polycystic ovaries[15,16].

Exclusion criteria

The following groups were excluded from this study: (1) Women with hypertension, diabetes mellitus, and thyroid diseases; (2) Drug intake of insulin sensitizers, lipid-lowering medicines, anti-androgenic therapies, oral contraceptives, and steroids; (3) Participants with a BMI of more than 30; and (4) Those with missing data were also excluded.


Fifty-four participants satisfied our criteria and the patients in this study were all subjected to meticulous clinical examination and all had an initial pelvic ultrasound scan to confirm the appearance of the PCOS ovaries. For each participant, four sets of data were collected; demographic criteria, biochemical and hormonal data, and biophysical data. These data were collected at the initial visit, 3 mo, and then 6 mo later, and the patient was supplied with a data collection sheet for the dates of further scheduled visits. During this period, all patients had been prescribed a metformin tab (Merck Santé /France) 500 mg TID times /d. At the end of the study, a total of 54 women with complete data of biophysical, biochemical, and hormonal data were assessed at 0, 3, and 6 mo; all were described in the study flowchart **Figure 1**.

Initial patients' uptake

The initial uptake was based on the clinical criteria of a menstrual cycle (MC) abnormality (oligomenorrhea or amenorrhea). The patients' age was recorded and only BMI was calculated in the outpatient clinic according to the formula: Weight in kg/square meter of height. After exclusion, eligible cases were planned to be further investigated for hormonal, biochemical, and biophysical criteria.

Biochemical and hormonal assays

On the 2nd day of the menstrual cycle and after one night's fast, participants in the study were initially sent to the Teaching labs in Yarmouk Hospital for the following biochemical tests: HDL, LDL, serum cholesterol, homeostatic model assessment for insulin resistance (HOMA-IR), fasting blood glucose (FBS), and sex hormone-binding globulin (SHBG). In addition to the hormones: Follicle-stimulating hormone (FSH), LH, and serum testosterone. The HOMA-IR tested insulin sensitivity based on the formula: $HOMA-IR = \text{fasting insulin (micro U/L)} \times \text{fasting glucose (nmol/L)} / 22.5$ [17].



DOI: 10.12998/wjcc.v11.i6.1318 Copyright ©The Author(s) 2023.

Figure 1 Study flowchart. PCOS: Polycystic ovarian syndrome.

All patients were given subsequent appointments based on the initial visit to repeat all of the above investigations 3 mo and 6 mo later while on metformin therapy, in addition to recording the days of the MC.

Biophysical data uptake

On the same day of biochemical data uptake, patients were sent for a Doppler assessment of the internal carotid artery.

Examination technique

In the radiology unit, an ultrasound machine TOSHIBA, Logic p5 with a linear array probe, and a frequency range of 5-10 MHZ was used. To measure carotid media intima thickness (CIMT), a special software computer option was used to adjust angle measurements for CIMT with a two-dimensional 2D grayscale maneuver. The assessment was made for both the right and left internal carotid artery (ICA), and the mean of both readings was taken.

CIMT was defined as the distance between the lumen intima to the adventitia media layer line of interference on the far wall in the longitudinal axis[18]. In order to decrease intra- and inter-observer variability for CIMT measurements, all readings were made by the same radiologist in our department after a period of practice to master the technique.

As for blood flow parameters, PI, and RI, a Doppler study of the ICA was performed, and measurements were made to PI and RI.

The patient required no special preparation, and the examination was held while the patient lay in a supine position with their head turned to the other side and did not last more than ten minutes. As mentioned above, patients were provided with scheduled visits for further re-scan 3 mo and 6 mo later.

Measurement of area under curve for Doppler wave

To explain how we calculated the AUC, we will give an example. A sample of ICA Doppler is shown in Figure 2. In this picture, we measured the different heights of the Doppler wave (demarcated by dots in Figure 2A) as it went up and down with a facility supplied by the ultrasound device. Depending on the width of the wave, these measurements averaged between 8 and 10 for each wave; see the horizontal orange line in Figure 2A). The measurement of the heights represents the speed of blood in the systolic and diastolic velocities. The calculated heights were put in an Excel table to analyze later with simple, free software called GRAPH, which can be downloaded from: <https://www.padowan.dk/download/> [19].

In Figure 2B, a graphical simulation of the internal Carotid Artery Doppler wave in Graph software is shown, with the different heights (demarcated in red dots) used to measure the systolic and diastolic blood flow for the calculation of AUC.




Figure 2 A sample of internal carotid artery Doppler. A: It shows a real-time internal carotid artery Doppler; we have simulated the way of measuring the area under the curve by Paint app to the last wave. Note how different heights of the systolic and diastolic blood flow (demarcated with orange dots), which are used to measure the area under the curve (demarcated as orange longitudinal lines); B: It shows the graphical simulation of the internal carotid artery Doppler wave with the different heights (demarcated in red dots) used to measure the systolic and diastolic blood flow for the calculation of the area under the curve.

The AUC-ICA was calculated according to the following formula: $AUC = \int \text{Blood velocity measured at systolic and diastolic velocity for a single Doppler wave}$ [11].

AUC physically means the amount of blood passed per unit area of the internal carotid artery per single heartbeat. It was measured in unit 3. Following the initial measurement of AUC-ICA, a re-assessment was made 3 mo and 6 mo later. All data with associated biochemical and biophysical data were stored in an excel sheet for further data analysis at the end of the study (Figure 2).

Sample size calculation

The sample size was calculated according to the following equation for a cross-sectional study with quantitative variables[20]:

$$\text{Sample size} = (Z_{1-\alpha/2})^2 \text{SD}^2 / d^2.$$

$Z_{1-\alpha/2}$ = is standard normal variate = 1.96.

SD = standard deviation of the variable taken from already published studies.

d = absolute error or precision level as an operator decides.

$$\text{Sample size} = (1.96)^2 (0.35)^2 / (0.1)^2 = (3.84 \times 0.1225) / 0.01 = 43 \text{ patients.}$$

So the sample size is 43 patients, and our study involved 54 patients.

Statistical analysis

Continuous data were expressed as mean and standard deviation. Data normality was checked with the Shapiro-Wilk test, and the data were normally distributed. One-way ANOVA test was used to assess the statistical data differences at the initial visit, at 3 mo, and 6 mo later for all the above biochemical, hormonal, and biophysical study variables. Linear regression was used to evaluate the decline of the AUC-ICA within 6 mo of metformin treatment with the calculation of the correlation coefficient and associated *P* value. In addition, further linear regression was constructed between AUC as the main dependent variable *vs* all significant biochemical, hormonal and biophysical variables taken in this study with the calculation of the correlation coefficient and associated value to assess the effect of metformin treatment on those correlations for the 6 mo treatment. AUC-ICA was assessed with a freely downloadable software GRAPH and further checked with MedCalc and NCSS software. *P* values less than 0.05 were considered significant.

RESULTS

Fifty-four young, unmarried, PCOS women were collected with full hormonal, biochemical, and biophysical profiles at the initial visit, 3 mo, and 6 mo after starting the metformin tablet 500 mg/TID day. Regarding the anthropometric criteria, the mean age of the participants was 24.81 ± 3.49 years. The days of the MC showed a significant decrease ($P < 0.034$) from 57.6 ± 5.8 in the initial visit to 43 ± 5.6 and 31.97 ± 4.9 d in the second and 3rd visits, respectively. Likewise, BMI showed a significant reduction ($P < 0.04$) from 28.22 ± 0.75 in the initial visit to 25.88 ± 0.64 and 23.81 ± 0.74 kg/m², respectively.

The main demographic criteria of these women are given at the three sampling and scanning times expressed as mean and standard deviation and were described in Table 1, while the three columns were compared with a one-way ANOVA test. The results highlight a progressive increase in serum HDL and SHBG throughout the treatment period over 6 mo. FBG showed a trend decrease; however, it fails to have a statistical value. On the other hand, a significant reduction was found in serum cholesterol, LDL,

Table 1 The primary biochemical, hormonal, and ultrasonic criteria at the initial visit, 3 mo and 6 mo later, with their statistical comparison

Parameter	First visit value, <i>n</i> = 54	2 nd visit at 3 mo, <i>n</i> = 54	3 rd visit at 6 mo, <i>n</i> = 54	<i>P</i> value
Cholesterol in mg/dL	384.19 ± 13.64	337.25 ± 18.40	276.82 ± 12.69	< 0.030
HDL in mg/dL	35.70 ± 3.78	48.70 ± 4.31	62.52 ± 4.09	< 0.001
LDL in mg/dL	203.22 ± 8.74	167.41 ± 12.26	132.33 ± 12.62	< 0.040
Fasting blood sugar in mg/dL	92.22 ± 1.80	90.93 ± 2.42	92.07 ± 2.43	0.071
HOMA-IR	2.26 ± 0.07	2.02 ± 0.10	1.74 ± 0.07	< 0.001
SHBG in nmol/L	29.52 ± 7.11	58.29 ± 8.25	81.74 ± 7.49	< 0.001
Testosterone in ng/dL	108.30 ± 8.29	79.74 ± 9.45	51.56 ± 7.96	< 0.010
LH / FSH_Ratio	2.62 ± 0.12	2.23 ± 0.15	1.76 ± 0.13	< 0.001
PI	1.41 ± 0.08	1.13 ± 0.07	0.89 ± 0.07	< 0.034
RI	0.89 ± 0.04	0.77 ± 0.05	0.62 ± 0.04	< 0.040
CIMT in mm	0.92 ± 0.03	0.79 ± 0.03	0.66 ± 0.03	< 0.001
AUC- ICA Doppler	56.44 ± 4.49	42.11 ± 4.06	31.29 ± 3.93	< 0.001

Data are presented as means ± standard deviation. BMI: Body mass index; HDL: High-density lipoprotein; LDL: Low-density lipoprotein; SHBG: Sex hormone-binding globulin; FSH: Follicular stimulating hormone; LH: Luteinizing hormone; HOMA-IR: Homeostasis model assessment for insulin resistance; PI: Pulsatility index; RI: Resistance index; CIMT: Carotid artery intima-media thickness; AUC: Area under the curve; ICA: Internal carotid artery.

HOMA- IR, testosterone, LH/FSH ratio, PI, RI, and CIMT. As for AUC-ICA, it showed a progressive decrease with the treatment period.

In order to shed more light on the trend of AUC-ICA correlation with various study parameters, a linear regression was constructed in [Table 2](#).

All biochemical markers; cholesterol, HDL, LDL, SHBG, and HOMA_IR; the hormonal markers of testosterone, and LH/FSH_Ratio; and ultrasonic parameters, RI, and CIMT used in PCOS women evaluation were correlated strongly to AUC-ICA, $P < 0.05$.

A highly significant negative correlation exists between AUC-ICA Doppler *vs* HOMA-IR. An inverse correlation was confirmed between the 6 mo time (24 wk) of metformin therapy *vs* AUC-ICA with $r = -0.99$ and P value less than 0.001, which was shown in [Figure 3](#) and highlighted by a real-time Doppler scan in [Figure 4](#), where the wave peak decreased after treatment from 80 to 60 and accordingly; the AUC-ICA was reduced.

To assess the strength of association between HOMA-IR as the primary dependent variable *vs* the predictive biophysical profiles related to the ICA, namely PI, RI, CIMT, and AUC-ICA Doppler; the best subset regression model was constructed with a calculation of Mallows's coefficient described in [Table 3](#). The lowest values are shown on the AUC-ICA Doppler, which means it has the highest predictive value for HOMA-IR. While RI has the highest Mallows's coefficient value, which means it has the lowest predictive value. Both CIMT and PI lie in between the two.

DISCUSSION

Our results showed that metformin therapy for 6 mo caused a progressive improvement in BMI, menstrual cycle days, metabolic markers, hormonal parameters, and Doppler parameters (PI, RI, and CIMT). AUC-ICA exhibits a meaningful correlation to all PCOS parameters. Furthermore, it showed a progressive reduction throughout the treatment period. AUC-ICA correlated inversely with treatment time ($r = -0.98$, $P < 0.001$) and positively with HOMA-IR ($r = 0.98$, $P < 0.0001$). With the best subset regression model, the AUC-ICA had the highest predictive value for HOMA-IR.

Women with PCOS present with multiple anthropometric, metabolic, and hormonal abnormalities which are successfully reversed by metformin therapy; our results were in accordance with published studies[21].

The reduced BMI in our result was in line with a recently published meta-analysis study. Their result discussed that metformin as a monotherapy or in combination with other drugs can improve all anthropometric parameters (weight, waist-to-hip ratio, and BMI) among PCOS women. This was made irrespective of the dose and the duration of metformin use[22].

Table 2 Correlations of area under the curve - internal carotid artery Doppler vs study variables over 6 mo with metformin therapy for women recruited to this study

AUC- ICA vs parameter	Coefficient of correlation	P value
Cholesterol in mg/dL	0.81	0.03
HDL in mg/dL	-0.92	0.002
LDL in mg/dL	0.88	0.04
HOMA_IR	0.98	0.0001
SHBG in nmole/L	-0.90	0.01
Testosterone in ng/dL	0.86	0.001
LH/ FSH_ratio	0.91	0.007
PI	0.87	0.002
RI	0.8	0.009
CIMT in mm	0.95	0.0001
24 wk of metformin therapy	-0.99	0.001

Data are presented as means \pm standard deviation. HDL: High-density lipoprotein; LDL: Low-density lipoprotein; SHBG: Low hormone binding globulin; FSH: Follicular stimulating hormone; LH: Luteinizing hormone; HOMA-IR: Homeostasis model assessment for insulin resistance; PI: Homeostasis index; RI: Resistance index; CIMT: Carotid artery intima-media thickness.

Table 3 Best subset regression between homeostasis model assessment for insulin resistance as the main dependent variable vs carotid artery intima-media thickness, pulsatility indices, resistance indices and area under the curve of the carotid artery Doppler with calculation of associated Mallows coefficient

Variable	(Cp) Mallows coefficient
AUC-ICA	49.6
CIMT	155.5
PI	326.8
RI	637.0

AUC-ICA: Area under the curve for the internal carotid artery; CIMT: Carotid artery intima-media thickness; PI: Pulsatility indices; RI: Resistance indices.


The Abdalla meta-analysis declared a meaningful reduction of serum testosterone with metformin, $P < 0.0001$ with moderate grade evidence and our result was in good agreement with their results[23].

The role of IR in hyperandrogenemia is not well understood. Some authors suggest that hyperinsulinemia plays a dual role in triggering hyperandrogenemia, first *via* direct stimulation of ovarian androgens due to the insulin receptors on the theca cells, and second *via* indirect stimulation of LH secretion and suppression of SHBG production by the liver with a net increase of free androgen levels [24].

Metformin's beneficial effects on PCOS include; antihyperglycemic and reduction of IR by increasing the peripheral uptake of glucose in addition to its indirect effect on insulin levels. Our data showed a progressive reduction of FBS and HOMA-IR with the treatment, as other studies pointed out[20]. It has an anti-androgenic effect *via* reducing CYP17 cytochrome activity involved in androgen production, not to mention increasing SHBG, which consequently reduces free androgens[20].


The favorable effect of metformin in cardiovascular disease (CVD) protection has several benefits. First, it protects endothelial integrity, which is a key player in the triggering and progression of CVD, consequently reducing future CVD risk. Second, metformin may inhibit hepatic de novo lipogenesis paths to lower plasma cholesterol levels and improves the atherogenic milieu, although the exact mechanism is still unknown. Third, metformin has anti-inflammatory and anti-oxidative effects; both are accredited as an etiological factor in PCOS pathogenesis. Indirect action of metformin; *via* reducing IR and androgen, both were proposed to act independently and synergistically in the progression of atherogenic dyslipidemia and CVD risk among PCOS women[22-25].

Regarding the menstrual cycle, Yilmaz *et al*[26] studied menstrual irregularities and hirsutism in women treated with two insulin-sensitizing medications, Rosiglitazone and metformin. Their data showed an improvement in cycle regularity and hirsutism score; however, they advised Rosiglitazone



DOI: 10.12998/wjcc.v11.i6.1318 Copyright ©The Author(s) 2023.

Figure 3 Highlights an inverse correlation of area under the curve of the internal carotid artery vs metformin treatment time ($r = -0.99$, $P < 0.001$).



DOI: 10.12998/wjcc.v11.i6.1318 Copyright ©The Author(s) 2023.

Figure 4 A real-time Doppler scan. A: A real-time Doppler scan showing progressive reduction of the area under the curve of the internal carotid artery with treatment. Note the decrease in the peak of the wave in a subsequent visit 3 mo apart; from 80 to 60; B: A real-time Doppler scan showing the progressive reduction of the area under the curve of the internal carotid artery with treatment. Note the reduced wave peak to 60 following treatment compared to previous Figure 4A.

over metformin for greater patient acceptance and hirsutism improvement. In line with their result, our data showed significant improvement in MC days and regularity[26].

Androgen excess tends to associate with hyperinsulinemia and IR. This alliance seems to worsen menstrual cycle abnormalities. Several studies showed that many women with metformin therapy have ovulated and had a pregnancy. In contrast, patients with higher levels of serum testosterone were more likely to be infertile[27,28].

There was a debate on which contributes more to improving MC by metformin therapy, reduced IR, reduced androgen, or reduced BMI caused by metformin. The Ezech *et al*[29] study results highlighted that menstrual cycle abnormality correlated positively with IR severity and not with hyperandrogenemia with adjustment of IR confounders, including BMI. In fact, amenorrheic PCOS women had the worst IR. Our results showed an improved IR and menstrual cycle pattern with metformin use, which suggests a clinical implication for daily practice; tracking down MC changes is easy and is free of charge, and it can reflect an improvement in the IR that underlies PCOS syndrome[29]. Prior studies demonstrated that MC abnormalities among adolescents would not impact future reproductive performance[30]. However, that was not the case for older women (> 30 years) who presented with oligomenorrheic and hirsute; they had lower reproductive performance than their age-matched controls [31].

IR lies at the heart of PCOS pathogenesis and severity and is a prognostic marker for response to treatment[32]. IR is universally assessed by HOMA-IR, calculated from the fasting serum insulin levels and blood sugar. Nevertheless, the assessment of HOMA-IR is costly and time-consuming, and this has pushed many researchers to find alternatives to assess IR[33,34].

Since 2010 many papers have reported that women with PCOS have universally increased intima-media thickness IMT in all medium-sized arteries like the carotid and ovarian artery[35]. Since then, many researchers have declared an excellent correlation between CIMT and HOMA-IR, and the current study results are in good agreement with the aforementioned studies[36].

Blood flow parameters were used to monitor HOMA-IR among PCOS cases during metformin use. In this study, both PI and RI had a progressive reduction, consistent with results obtained by Kaya *et al* [37]. Their study highlighted the beneficial role of adding metformin to combined contraceptive pills among PCOS women compared to those treated with combined contraceptive pills alone. CIMT and flow-mediated dilatation (FMD) was improved with the metformin group. The author attributes this to the effect of metformin on endothelial integrity, which is responsible for nitric oxide release, reducing oxidative stress, and the correction of altered cellular signaling pathways[27,37,38].

The foundation of this study is a simple physical fact. Bernoulli's rule states that fluid velocity has an inverse correlation to the diameter of a vessel[8]. The increased ICA-IMT will cause the passing of blood to be faster. Since the area under the curve physically means the amount of blood passed per one sectional area in the vessel wall, consequently, AUC might represent a sensitive marker to HOMA-IR [11].

Scientists have discussed that measuring insulin sensitivity through surrogate markers of insulin sensitivity (HOMA-IR and QUICKI) is no longer enough and raised the necessity for more acute ways [37-40]. Although IMT and FMD were investigated in PCOS, we believe that estimating the change of blood speed mirrored by AUC is more accurate, as described in earlier work[38,39].

Indeed, the strong correlations of AUC-ICA with all biochemical, hormonal, and Doppler parameters (PI, RI, and CIMT) support our hypothesis; furthermore, the progressive reduction of AUC-ICA represents the initial positive predictive ability of AUC-ICA as a possible novel marker for predicting IR. The best subset regression added more strength to our result, which confirmed; that AUC-ICA had the highest predictive value; it preceded PI, RI, and CIMT. The parameters mentioned above are already validated in IR, and thus AUC-ICA use needs no external validation.

Study limitations

The treatment period is relatively short, so we cannot be sure of the future implications of metformin treatment in the long run. The study type is another limitation, as in any cross-sectional study, the exact cause-and-effect link cannot be determined because the IR and AUC-ICA were simultaneously measured[40]. Intra- and inter-observer variability for AUC-ICA measurements is another limitation. Finally, the COVID-19 pandemic has seriously affected the duration of the study due to the repeated lockdown conducted in Iraq at that time[40,41].

Study strengths

AUC is a novel marker proposed by the current study, calculated by serial readings of Doppler wave height using freely downloadable software from the web, namely GRAPH software. It was easily calculated, free of extra charge, as it can be integrated into pelvic scan sessions. Its strong association with HOMA-IR and the duration of the treatment added to its cost make it superior to HOMA-IR in terms of cost-benefit analysis. For that, AUC-ICA is recommended as a reliable predictive marker for IR, follow-up, and prognostic value, especially during metformin therapy. Further studies are warranted for AUC-ICA application in clinical practice.

CONCLUSION

Measurement of the area under the curve of ICA is a promising marker for assessing IR in polycystic ovarian syndrome cases during the metformin therapy period. AUC-ICA showed strong significant correlations to PCOS parameters and had a superior cost-benefit analysis over HOMA-IR. Further studies are recommended to explore future applications in practice.

ARTICLE HIGHLIGHTS

Research background

Insulin resistance (IR) is implicated in many aspects of polycystic ovarian syndrome (PCOS) pathogenesis. Metformin effectively decreased IR. Improved IR was evaluated *via* homeostatic model assessment for IR (HOMA-IR) and Doppler parameters; mainly carotid artery intima-media thickness (ICA-IMT). The area under the curve of the internal carotid artery (AUC-ICA) Doppler wave was examined as a helpful marker for determining IR among PCOS cases presented with menstrual irregularity and treated by metformin over 6 mo.

Research motivation

Much research has shown that surrogate measures of insulin sensitivity are no longer sufficient for evaluating insulin sensitivity, which has increased the need for new direct methods. Demonstrating changes in blood flow mirrored by AUC appeared to be more dependable; as indicated in earlier work, ICA-IMT was already examined in PCOS.

Research objectives

To ascertain if IR is related to the AUC-ICA Doppler in PCOS-affected women with menstrual irregularities. The second goal is to analyze the reliability of AUC-ICA as a helpful marker for monitoring IR in women who have received metformin treatment.

Research methods

The study enrolled 54 PCOS women in a cross-sectional study. *Anthropometric data* included patient age, body mass index (BMI), menstrual cycle days, *biochemical parameters*: serum cholesterol, low and high-density lipoprotein, sex hormone-binding globulin, fasting blood glucose, and HOMA-IR, *hormonal parameters*: testosterone, luteinizing hormone over follicle-stimulating hormone ratio, and *ultrasonic parameters*: (CIMT, PI, RI, and AUC-ICA). Measurements of the systolic and diastolic wave height were repeated in order to evaluate the AUC-ICA following metformin tab-500 mg; three times/d for 6 mo. Metformin caused a progressive reduction in BMI, menstrual cycle days, biochemical hormonal, and Doppler parameters (CIMT, PI, RI, and AUC-ICA). AUC-ICA correlated strongly to all PCOS parameters. AUC-ICA correlated inversely with treatment time ($r = -0.98$, $P < 0.001$) and positively with HOMA-IR ($r = 0.98$, $P < 0.0001$). *Via* the best subset regression model, the AUC-ICA had the highest predictive value for HOMA-IR.

Research results

BMI, menstrual cycle days, biochemical hormonal, and Doppler markers (CIMT, PI, RI, and AUC-ICA were all gradually reduced by metformin treatment). All PCOS indicators and AUC-ICA had significant correlations. AUC-ICA had a negative correlation ($r = -0.98$, $P < 0.001$) with treatment time and a positive correlation ($r = 0.98$, $P < 0.0001$) with HOMA-IR. The AUC-ICA demonstrated the best subset regression model's maximum predictive value for HOMA-IR.

Research conclusions

AUC-ICA was a reliable marker for the assessment of IR, especially during metformin medication. AUC-ICA preceded PI, RI, and CIMT and showed a high, meaningful correlation to other PCOS markers. For further use in practice, more research is suggested.

Research perspectives

The area under the curve of the internal carotid artery had a significant correlation with HOMA-IR and the length of metformin therapy, not to mention it has a superior cost-benefit analysis over HOMA-IR. AUC-ICA is a reliable indicator of IR, follow-up, and prognostic value, particularly while using metformin.

FOOTNOTES

Author contributions: Akram W and Nori W designed research and reviewed the data, wrote and revised the letter; Zghair MAG collected and analyzed the data; All authors have read and agreed on the final version of the manuscript.

Institutional review board statement: The ethical committee of Mustansiriyah University approved the study dated February/21/2019 (IRB No. 115).

Informed consent statement: All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

Conflict-of-interest statement: All authors declare that they have no potential or real conflicts related to this paper.

Data sharing statement: All data are available on reasonable request from the corresponding author.

STROBE statement: The authors have read the STROBE Statement – checklist of items, and the manuscript was prepared and revised according to the STROBE Statement – checklist of items.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license

their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Iraq

ORCID number: Wisam Akram 0000-0002-1968-7004; Wassan Nori 0000-0002-8749-2444; Muna Abdul Ghani Zghair 0000-0002-6219-1215.

S-Editor: Wang LL

L-Editor: Filipodia

P-Editor: Wang LL

REFERENCES

- 1 **Reyes-Muñoz E**, Sathyapalan T, Rossetti P, Shah M, Long M, Buscema M, Valenti G, La Rosa VL, Cianci S, Vitale SG. Polycystic Ovary Syndrome: Implication for Drug Metabolism on Assisted Reproductive Techniques-A Literature Review. *Adv Ther* 2018; **35**: 1805-1815 [PMID: 30311070 DOI: 10.1007/s12325-018-0810-1]
- 2 **Ali AI**, Hassan WNM, Alrawi S. A Copeptin as a Predictor Marker for Insulin Resistance Among Women with Polycystic Ovary Syndrome. *CWHR* 2022; **18**: e081221198670 [DOI: 10.2174/1573404817666211208152049]
- 3 **Zhang J**, Xu JH, Qu QQ, Zhong GQ. Risk of Cardiovascular and Cerebrovascular Events in Polycystic Ovarian Syndrome Women: A Meta-Analysis of Cohort Studies. *Front Cardiovasc Med* 2020; **7**: 552421 [PMID: 33282917 DOI: 10.3389/fcvm.2020.552421]
- 4 **Pani A**, Gironi I, Di Vieste G, Mion E, Bertuzzi F, Pintaudi B. From Prediabetes to Type 2 Diabetes Mellitus in Women with Polycystic Ovary Syndrome: Lifestyle and Pharmacological Management. *Int J Endocrinol* 2020; **2020**: 6276187 [PMID: 32587614 DOI: 10.1155/2020/6276187]
- 5 **Puttabyatappa M**, Padmanabhan V. Ovarian and Extra-Ovarian Mediators in the Development of Polycystic Ovary Syndrome. *J Mol Endocrinol* 2018; **61**: R161-R184 [PMID: 29941488 DOI: 10.1530/JME-18-0079]
- 6 **Kakoly NS**, Moran LJ, Teede HJ, Joham AE. Cardiometabolic risks in PCOS: a review of the current state of knowledge. *Expert Rev Endocrinol Metab* 2019; **14**: 23-33 [PMID: 30556433 DOI: 10.1080/17446651.2019.1556094]
- 7 **Song P**, Fang Z, Wang H, Cai Y, Rahimi K, Zhu Y, Fowkes FGR, Fowkes FJI, Rudan I. Global and regional prevalence, burden, and risk factors for carotid atherosclerosis: a systematic review, meta-analysis, and modelling study. *Lancet Glob Health* 2020; **8**: e721-e729 [PMID: 32353319 DOI: 10.1016/S2214-109X(20)30117-0]
- 8 **Pisano A**. Continuity Equation and Bernoulli's Theorem: Airplanes, Venturi Masks, and Other Interesting Things (for Anesthesiologists and Intensivists). *Physics for Anesthesiologists and Intensivists*, 2021 [DOI: 10.1007/978-3-030-72047-6_7]
- 9 **Panayides AS**, Amini A, Filipovic ND, Sharma A, Tsaftaris SA, Young A, Foran D, Do N, Golemati S, Kurc T, Huang K, Nikita KS, Veasey BP, Zervakis M, Saltz JH, Pattichis CS. AI in Medical Imaging Informatics: Current Challenges and Future Directions. *IEEE J Biomed Health Inform* 2020; **24**: 1837-1857 [PMID: 32609615 DOI: 10.1109/JBHI.2020.2991043]
- 10 **Manzoor I**, Bacha R, Gilani SA. Sonographic association of polycystic ovaries with intraovarian arterial pulsatility and resistive index. *Gynecol Endocrinol* 2019; **35**: 851-853 [PMID: 31062996 DOI: 10.1080/09513590.2019.1612357]
- 11 **Wang F**, Jin P, Feng Y, Fu J, Wang P, Liu X, Zhang Y, Ma Y, Yang Y, Yang A, Feng X. Flexible Doppler ultrasound device for the monitoring of blood flow velocity. *Sci Adv* 2021; **7**: eabi9283 [PMID: 34705515 DOI: 10.1126/sciadv.abi9283]
- 12 **Chirinos JA**, Segers P, Hughes T, Townsend R. Large-Artery Stiffness in Health and Disease: JACC State-of-the-Art Review. *J Am Coll Cardiol* 2019; **74**: 1237-1263 [PMID: 31466622 DOI: 10.1016/j.jacc.2019.07.012]
- 13 **Paul C**, Laganà AS, Maniglio P, Triolo O, Brady DM. Inositol's and other nutraceuticals' synergistic actions counteract insulin resistance in polycystic ovarian syndrome and metabolic syndrome: state-of-the-art and future perspectives. *Gynecol Endocrinol* 2016; **32**: 431-438 [PMID: 26927948 DOI: 10.3109/09513590.2016.1144741]
- 14 **Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group**. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome. *Fertil Steril* 2004; **81**: 19-25 [PMID: 14711538 DOI: 10.1016/j.fertnstert.2003.10.004]
- 15 **Verhoef SJ**, Wielink MC, Achterberg EA, Bongers MY, Goossens SMTA. Absence of menstruation in female athletes: why they do not seek help. *BMC Sports Sci Med Rehabil* 2021; **13**: 146 [PMID: 34814941 DOI: 10.1186/s13102-021-00372-3]
- 16 **Hickey M**, Doherty DA, Atkinson H, Sloboda DM, Franks S, Norman RJ, Hart R. Clinical, ultrasound and biochemical features of polycystic ovary syndrome in adolescents: implications for diagnosis. *Hum Reprod* 2011; **26**: 1469-1477 [PMID: 21478180 DOI: 10.1093/humrep/der102]
- 17 **Celik O**, Acbay O. Effects of metformin plus rosuvastatin on hyperandrogenism in polycystic ovary syndrome patients with hyperlipidemia and impaired glucose tolerance. *J Endocrinol Invest* 2012; **35**: 905-910 [PMID: 22522778 DOI: 10.3275/8371]
- 18 **Jabbour R**, Ott J, Eppel W, Frigo P. Carotid intima-media thickness in polycystic ovary syndrome and its association with hormone and lipid profiles. *PLoS One* 2020; **15**: e0232299 [PMID: 32330202 DOI: 10.1371/journal.pone.0232299]
- 19 **Meun C**, Gunning MN, Louwers YV, Peters H, Roos-Hesseling J, Roeters van Lennep J, Rueda Ochoa OL, Appelman Y, Lambalk N, Boersma E, Kavousi M, Fauser BC, Laven JS, CREW consortium. The cardiovascular risk profile of middle-aged women with polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 2020; **92**: 150-158 [PMID: 31638273 DOI: 10.1111/cen.14444]

- 10.1111/cen.14117]
- 20 **Chow S**, Shao J, Wang H, Lokhnygina Y. Sample Size Calculations in Clinical Research: Third Edition, 2017 [DOI: 10.1201/9781315183084]
 - 21 **Costello M**, Shrestha B, Eden J, Sjoblom P, Johnson N. Insulin-sensitising drugs versus the combined oral contraceptive pill for hirsutism, acne and risk of diabetes, cardiovascular disease, and endometrial cancer in polycystic ovary syndrome. *Cochrane Database Syst Rev* 2007; CD005552 [PMID: 17253562 DOI: 10.1002/14651858.CD005552.pub2]
 - 22 **Guan Y**, Wang D, Bu H, Zhao T, Wang H. The Effect of Metformin on Polycystic Ovary Syndrome in Overweight Women: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Int J Endocrinol* 2020; **2020**: 5150684 [PMID: 33014044 DOI: 10.1155/2020/5150684]
 - 23 **Abdalla MA**, Shah N, Deshmukh H, Sahebkar A, Östlundh L, Al-Rifai RH, Atkin SL, Sathyapalan T. Impact of pharmacological interventions on anthropometric indices in women with polycystic ovary syndrome: A systematic review and meta-analysis of randomized controlled trials. *Clin Endocrinol (Oxf)* 2022; **96**: 758-780 [PMID: 34918367 DOI: 10.1111/cen.14663]
 - 24 **El Hayek S**, Bitar L, Hamdar LH, Mirza FG, Daoud G. Poly Cystic Ovarian Syndrome: An Updated Overview. *Front Physiol* 2016; **7**: 124 [PMID: 27092084 DOI: 10.3389/fphys.2016.00124]
 - 25 **Nafisa A**, Gray SG, Cao Y, Wang T, Xu S, Wattoo FH, Barras M, Cohen N, Kamato D, Little PJ. Endothelial function and dysfunction: Impact of metformin. *Pharmacol Ther* 2018; **192**: 150-162 [PMID: 30056057 DOI: 10.1016/j.pharmthera.2018.07.007]
 - 26 **Yilmaz M**, Karakoç A, Törüner FB, Cakir N, Tiras B, Ayvaz G, Arslan M. The effects of rosiglitazone and metformin on menstrual cyclicity and hirsutism in polycystic ovary syndrome. *Gynecol Endocrinol* 2005; **21**: 154-160 [PMID: 16335907 DOI: 10.1080/09513590500231627]
 - 27 **Xu Y**, Qiao J. Association of Insulin Resistance and Elevated Androgen Levels with Polycystic Ovarian Syndrome (PCOS): A Review of Literature. *J Healthc Eng* 2022; **2022**: 9240569 [PMID: 35356614 DOI: 10.1155/2022/9240569]
 - 28 **Fornes R**, Simin J, Nguyen MH, Cruz G, Crisosto N, van der Schaaf M, Engstrand L, Brusselaers N. Pregnancy, perinatal and childhood outcomes in women with and without polycystic ovary syndrome and metformin during pregnancy: a nationwide population-based study. *Reprod Biol Endocrinol* 2022; **20**: 30 [PMID: 35130922 DOI: 10.1186/s12958-022-00905-6]
 - 29 **Ezeh U**, Ezeh C, Pisarska MD, Azziz R. Menstrual dysfunction in polycystic ovary syndrome: association with dynamic state insulin resistance rather than hyperandrogenism. *Fertil Steril* 2021; **115**: 1557-1568 [PMID: 33602559 DOI: 10.1016/j.fertnstert.2020.12.015]
 - 30 **Hudecova M**, Holte J, Olovsson M, Sundström Poromaa I. Long-term follow-up of patients with polycystic ovary syndrome: reproductive outcome and ovarian reserve. *Hum Reprod* 2009; **24**: 1176-1183 [PMID: 19168874 DOI: 10.1093/humrep/den482]
 - 31 **West S**, Lashen H, Bloigu A, Franks S, Puukka K, Ruokonen A, Järvelin MR, Tapanainen JS, Morin-Papunen L. Irregular menstruation and hyperandrogenaemia in adolescence are associated with polycystic ovary syndrome and infertility in later life: Northern Finland Birth Cohort 1986 study. *Hum Reprod* 2014; **29**: 2339-2351 [PMID: 25085801 DOI: 10.1093/humrep/deu200]
 - 32 **Shorakae S**, Ranasinha S, Abell S, Lambert G, Lambert E, de Courten B, Teede H. Inter-related effects of insulin resistance, hyperandrogenism, sympathetic dysfunction and chronic inflammation in PCOS. *Clin Endocrinol (Oxf)* 2018; **89**: 628-633 [PMID: 29992612 DOI: 10.1111/cen.13808]
 - 33 **Rudvik A**, Månsson M. Evaluation of surrogate measures of insulin sensitivity - correlation with gold standard is not enough. *BMC Med Res Methodol* 2018; **18**: 64 [PMID: 29940866 DOI: 10.1186/s12874-018-0521-y]
 - 34 **Pantoja-Torres B**, Toro-Huamanchumo CJ, Urrunaga-Pastor D, Guarnizo-Poma M, Lazaro-Alcantara H, Paico-Palacios S, Del Carmen Ranilla-Seguín V, Benites-Zapata VA; Insulin Resistance and Metabolic Syndrome Research Group. High triglycerides to HDL-cholesterol ratio is associated with insulin resistance in normal-weight healthy adults. *Diabetes Metab Syndr* 2019; **13**: 382-388 [PMID: 30641729 DOI: 10.1016/j.dsx.2018.10.006]
 - 35 **Kaya MG**, Gunebakmaz O, Zencir C, Yilmazsoy A, Karadag M, Topsakal R, Ergin A, Kelestimur F. An assessment of the elastic properties of the aorta in nonobese women with polycystic ovary syndrome. *Fertil Steril* 2010; **94**: 2402-2405 [PMID: 20493476 DOI: 10.1016/j.fertnstert.2010.04.002]
 - 36 **Kupreeva M**, Diane A, Lehner R, Watts R, Ghosh M, Proctor S, Vine D. Effect of metformin and flutamide on insulin, lipogenic and androgen-estrogen signaling, and cardiometabolic risk in a PCOS-prone metabolic syndrome rodent model. *Am J Physiol Endocrinol Metab* 2019; **316**: E16-E33 [PMID: 30153063 DOI: 10.1152/ajpendo.00018.2018]
 - 37 **Kaya MG**, Yildirim S, Calapkorur B, Akpek M, Unluhizarci K, Kelestimur F. Metformin improves endothelial function and carotid intima media thickness in patients with PCOS. *Gynecol Endocrinol* 2015; **31**: 401-405 [PMID: 25791462 DOI: 10.3109/09513590.2015.1006188]
 - 38 **Nori W**, Fleeh NH, Akram W. Will the area under the curve of the umbilical artery Doppler predict fetal growth restriction at 34 weeks of gestation among pre-eclamptic women? 2nd International Conference On Engineering & Science, 2021 [DOI: 10.1063/5.0069008]
 - 39 **Zhu S**, Cheng C, Wang LL, Zhao DJ, Zhao YL, Liu XZ. Prognostic values of optic nerve sheath diameter for comatose patients with acute stroke: An observational study. *World J Clin Cases* 2022; **10**: 12175-12183 [PMID: 36483822 DOI: 10.12998/wjcc.v10.i33.12175]
 - 40 **Solem RC**. Limitation of a cross-sectional study. *Am J Orthod Dentofacial Orthop* 2015; **148**: 205 [PMID: 26232823 DOI: 10.1016/j.ajodo.2015.05.006]
 - 41 **Nori W**, Akram W. Effect of gender on the reliability of COVID-19 rapid antigen test among elderly. *World J Clin Cases* 2022; **10**: 10820-10822 [PMID: 36312479 DOI: 10.12998/wjcc.v10.i29.10820]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

