

# World Journal of *Clinical Cases*

*World J Clin Cases* 2023 February 26; 11(6): 1224-1433



## Contents

Thrice Monthly Volume 11 Number 6 February 26, 2023

## OPINION REVIEW

- 1224 Collagen matrix scaffolds: Future perspectives for the management of chronic liver diseases  
*Martinez-Castillo M, Altamirano-Mendoza I, Zielinski R, Priebe W, Piña-Barba C, Gutierrez-Reyes G*

## MINIREVIEWS

- 1236 Sex dimorphism and metabolic profiles in management of metabolic-associated fatty liver disease  
*Martin-Grau M, Monleon D*
- 1245 Epidemiology and etiology of chemical ocular injury: A brief review  
*Akgun Z, Selver OB*
- 1252 Review of the prevalence, diagnostics, and containment measures of the current mpox outbreak  
*Sanyaolu A, Marinkovic A, Okorie C, Prakash S, Haider N, Dixon Y, Izurieta R, Badaru O, Smith S*
- 1261 Clinical and pathophysiological understanding of the hepatorenal syndrome: Still wrong or still not exactly right?  
*Wilde B, Canbay A, Katsounas A*
- 1267 Flare of the silent pandemic in the era of the COVID-19 pandemic: Obstacles and opportunities  
*Rayan RA*
- 1275 Implications of metabolic dysfunction associated fatty liver disease in COVID-19  
*Chakraborty R, Sharma D, Kapoor DU, Dwivedi A, Khabiya R, Sen S*

## ORIGINAL ARTICLE

## Retrospective Study

- 1287 Hyperglycemia in COVID-19 infection without diabetes mellitus: Association with inflammatory markers  
*Geetha HS, Singh G, Sekar A, Gogtay M, Singh Y, Abraham GM, Trivedi N*

## Clinical Trials Study

- 1299 Efficacy of invisible advancement correction for mandibular retraction in adolescents based on Pancherz analysis  
*Kong L, Liu XQ*

## Observational Study

- 1310 Survey study of the etiology of non-traumatic altered consciousness in the Emergency Department at Suez Canal University Hospital in Egypt  
*Moussa BS, Abd Elatiff ZM, Kamal Eldin Elhadary GM*

- 1318** Metformin effect on internal carotid artery blood flow assessed by area under the curve of carotid artery Doppler in women with polycystic ovarian syndrome  
*Akram W, Nori W, Abdul Ghani Zghair M*
- 1330** Effect of continuous nursing combined with respiratory exercise nursing on pulmonary function of postoperative patients with lung cancer  
*Qiu QX, Li WJ, Ma XM, Feng XH*

**CASE REPORT**

- 1341** Functioning gonadotroph adenoma with hyperestrogenemia and ovarian hyperstimulation in a reproductive-aged woman: A case report and review of literature  
*He Y, Gao YT, Sun L*
- 1349** Clinical manifestations of adult hereditary spherocytosis with novel *SPTB* gene mutations and hyperjaundice: A case report  
*Jiang N, Mao WY, Peng BX, Yang TY, Mao XR*
- 1356** Post-traumatic cauda equina nerve calcification: A case report  
*Liu YD, Deng Q, Li JJ, Yang HY, Han XF, Zhang KD, Peng RD, Xiang QQ*
- 1365** Endometriosis-associated endometrioid adenocarcinoma of the fallopian tube synchronized with endometrial adenocarcinoma: A case report  
*Feng JY, Jiang QP, He H*
- 1372** Gemcitabine-induced peripheral vascular disease and prolonged response in a patient with metastatic pancreatic adenocarcinoma: A case report  
*Fabien MB, Elodie P, Anna S, Addeo P, Meher B*
- 1379** Epidemic Japanese B encephalitis combined with contactin-associated protein-like 2 antibody-positive autoimmune encephalitis: A case report  
*Huang P*
- 1385** Acute pancreatitis as initial presentation of acute myeloid leukemia-M2 subtype: A case report  
*Yang WX, An K, Liu GF, Zhou HY, Gao JC*
- 1393** Postoperative jaundice related to *UGT1A1* and *ABCB11* gene mutations: A case report and literature review  
*Jiang JL, Liu X, Pan ZQ, Jiang XL, Shi JH, Chen Y, Yi Y, Zhong WW, Liu KY, He YH*
- 1403** Hidrotic ectodermal dysplasia in a Chinese pedigree: A case report  
*Liao MY, Peng H, Li LN, Yang T, Xiong SY, Ye XY*
- 1410** Hepatitis A virus-associated acute acalculous cholecystitis in an adult-onset Still's disease patient: A case report and review of the literature  
*Chang CH, Wang YY, Jiao Y*
- 1419** Transverse myelitis caused by herpes zoster following COVID-19 vaccination: A case report  
*Cho SY, Jang BH, Seo JW, Kim SW, Lim KJ, Lee HY, Kim DJ*

- 1426 Primary malignant melanoma of the esophagus: A case report

*Wang QQ, Li YM, Qin G, Liu F, Xu YY*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Goran Augustin, MD, MSc, PhD, Assistant Professor, Senior Scientist, Surgeon, Department of Surgery, University Hospital Centre Zagreb, Zagreb 10000, Croatia. [augustin.goran@gmail.com](mailto:augustin.goran@gmail.com)

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jin-Lei Wang*.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

February 26, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



## Post-traumatic cauda equina nerve calcification: A case report

Yan-Dong Liu, Qiang Deng, Jun-Jie Li, Hai-Yun Yang, Xian-Fu Han, Kai-Dong Zhang, Ran-Dong Peng, Qian-Qian Xiang

**Specialty type:** Neurosciences

**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C, C

Grade D (Fair): 0

Grade E (Poor): 0

**P-Reviewer:** Mohey NM, Egypt; Zharikov YO, Russia

**Received:** October 2, 2022

**Peer-review started:** October 2, 2022

**First decision:** January 5, 2023

**Revised:** January 18, 2023

**Accepted:** February 3, 2023

**Article in press:** February 3, 2023

**Published online:** February 26, 2023



**Yan-Dong Liu, Hai-Yun Yang, Xian-Fu Han, Qian-Qian Xiang**, Graduate School, Gansu University of Traditional Chinese Medicine, Lanzhou 730000, Gansu Province, China

**Qiang Deng, Jun-Jie Li**, Spinal Disease Diagnosis and Treatment Center, Gansu Provincial Hospital of Traditional Chinese Medicine, Lanzhou 730050, Gansu Province, China

**Kai-Dong Zhang**, First Orthopedic Department, Lanzhou Traditional Chinese Medicine Orthopedic Hospital, Lanzhou 730050, Gansu Province, China

**Ran-Dong Peng**, Department of Osteomyelitis, Gansu Provincial Hospital of Traditional Chinese Medicine, Lanzhou 730000, Gansu Province, China

**Corresponding author:** Qiang Deng, MM, Chief Physician, Spinal Disease Diagnosis and Treatment Center, Gansu Provincial Hospital of Traditional Chinese Medicine, No. 418 Guazhou Road, Jianlan Road Street, Lanzhou 730050, Gansu Province, China.

[1327171163@qq.com](mailto:1327171163@qq.com)

### Abstract

#### BACKGROUND

Post-traumatic cauda equina nerve calcification is extremely rare in clinical practice, and its etiology, pathogenesis, treatment and prognosis are unclear. There are few studies and reports on Post-traumatic cauda equina nerve calcification, and this review reports a case of Post-traumatic cauda equina nerve calcification for reference.

#### CASE SUMMARY

A 52-year-old patient presented to our hospital with a history of lumbar spinal stenosis and a lumbar vertebral fracture caused by trauma. The patient's right lower limb had weakness in hip flexion, knee extension and plantarflexion with muscle strength grade 3, right ankle dorsiflexion and thumb dorsiflexion with muscle strength grade 0. The patient's skin sensation below the right knee plane disappeared. The patient's Computed tomography (CT) data showed signs of cauda equina nerve calcification and the terminal filaments in the plane of the third to fifth lumbar vertebrae. After treatment the patient's symptoms were slightly relieved.

#### CONCLUSION

We provide an extremely rare case of Post-traumatic cauda equina nerve calcification and offer a conservative treatment plan. However, the etiology, mechanism and treatment of Post-traumatic cauda equina nerve calcification are

still unclear. This requires scholars to conduct more research and exploration in this area.

**Key Words:** Post-traumatic; Calcification; Cauda equina nerve; Spinal Cord Injury; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** We report an extremely rare case of Post-traumatic cauda equina nerve calcification and speculate on its etiology and mechanism, providing material for scholars to study the etiology, mechanism and treatment of post-traumatic cauda equina nerve calcification.

**Citation:** Liu YD, Deng Q, Li JJ, Yang HY, Han XF, Zhang KD, Peng RD, Xiang QQ. Post-traumatic cauda equina nerve calcification: A case report. *World J Clin Cases* 2023; 11(6): 1356-1364

**URL:** <https://www.wjgnet.com/2307-8960/full/v11/i6/1356.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v11.i6.1356>

## INTRODUCTION

Post-traumatic cauda equina nerve calcification refers to the deposition of calcium in ten pairs of lumbosacral nerve roots below the conus medullaris, resulting in various clinical symptoms[1]. The cauda equina nerve is the bridge between the spinal cord and the peripheral nerve, connecting the pseudounipolar neurons of the dorsal root ganglion and the spinal cord neurons. Because of its special anatomical structure (only a layer of nerve inner membrane, lack of corresponding protective tissue), it is vulnerable to mechanical injury. The nutrition of cauda equina nerve comes from cerebrospinal fluid and blood, and there is a relatively anemia area in the anastomosis between the spinal artery and the root artery supplying the cauda equina nerve, so the cauda equina nerve is more prone to decompensation due to mechanical compression[2]. Post-traumatic cauda equina nerve calcification is extremely rare clinically, and its etiology, pathogenesis, treatment, and prognosis are unclear. No studies or reports on Post-traumatic cauda equina nerve calcification have been inquired, and a case of Post-traumatic cauda equina nerve calcification is reported here for reference.

## CASE PRESENTATION

### Chief complaints

On May 13, 2022, a 53 years female community worker with 35 years of progressive lumbar kyphosis was admitted to our hospital with symptoms worsening over 7 mo, claudication and right knee retroflexion.

### History of present illness

The patient complained of lumbar spine fracture and bilateral lower limb paralysis due to injury from fall more than 35 years ago, and underwent emergency surgery in other hospitals. Two years after the surgery, the motor and sensory functions of the left lower limb gradually recovered, while the right lower limb had weakness in hip flexion, knee extension and plantarflexion, with muscle strength of about grade 3. He was admitted to the hospital 7 mo ago after a fall during a walk in the park, resulting in right knee retroflexion and right ankle sprain. In the last week, the patient's symptoms were aggravated without any obvious cause, and the right hip was accompanied by radiating pain and numbness in the right lower limb, especially in the lateral and posterior side of the right lower limb. During the pain attack, the patient took metamizole sodium tablets (0.5g po st). After taking metamizole sodium tablets, the patient's pain was slightly relieved. The patient complained of numbness and trapped pain in the lumbosacral region with radiating pain and numbness in the right lower extremity, especially the lateral and posterior sides of the right lower limb. The patient has difficulty squatting down, and the pain increases when bending down. The patient had no symptoms such as chills, high fever, spontaneous sweating, night sweating, dizziness, nausea and vomiting. The patient's sleep and urination were normal. Since the patient's stool is dry, laxatives are used to help him defecate for 3 days.

### History of past illness

The patient had a history of hypertension for more than 3 years, with blood pressure up to 150/100 mmHg. Patients regularly take nifedipine sustained-release tablets to control their blood pressure, which has been controlled at present. The patient underwent internal fixation of the lumbar spine at an



another hospital 35 years ago and had no adverse blood transfusion reactions during the operation. Had a cesarean section in another hospital more than 30 years ago.

### **Personal and family history**

The patient had no relevant personal or family history.

### **Physical examination**

The patient's Lumbar kyphosis and right knee retroflexion. The patient had limited anterior flexion and posterior extension of the lumbar region with percussion pain and pressure pain at the spinous process and paraspinal process of the third lumbar vertebra - fifth lumbar vertebra (L3-5). The patient's right lower limb was positive for lasague sign and supine abdominal sticking test. Patient's muscle strength: iliopsoas (Second lumbar vertebra / fourth lumbar vertebra) (L2/4) muscle strength, quadriceps (L2/4) muscle strength, gluteus maximus Fifth lumbar vertebra/second sacral vertebra (L5/S2) muscle strength: right: grade 3, left: grade 4+. biceps femoris (L5/S2) muscle strength, tibialis anterior L4/5 muscle strength, thumb extension (L4/S1) muscle strength: right: grade 0, left: grade 4+. gastrocnemius, hallux valgus (L5/S2) muscle strength: right: grade 3, left: grade 4+. The patient's right Achilles tendon reflex disappeared, the skin sensation below the knee joint plane of the right lower limb disappeared, and the skin sensation below the knee joint plane of the left knee decreased. The patient's pathological reflex was not elicited.

### **Laboratory examinations**

Erythrocyte Sedimentation Rate: 24 mm/h. Complete biochemistry: Apolipoprotein A1: 0.77 g/L, High density lipoprotein-C: 0.84 mmol/L, Albumin ratio (A/G): 1.18, Triglycerides (TG): 2.78 mmol/L, Homocysteine (HCY): 21 µmol/L, Glutamyl transpeptidase (γ-GT): 218 U/L, Creatinine (CREA): 34 µmol/L, Rheumatoid factor (RF):42.26, C-reactive protein (CRP): 11.12 mg/L. Bone densitometry suggested decreased bone mass.

### **Imaging examinations**

Lumbar spine Computed tomography (CT) showed: L5 plane end filament and cauda equina nerve calcification signs (Figure 1). The X-ray film of lumbar vertebrae shows that the sequence of lumbar vertebrae is abnormal, L3 vertebral body is diseased and shifted backward, L2-3 intervertebral space disappears, L2-3, L3-4, L4-5, L5-S1 intervertebral space narrows, and all vertebrae and vertebral facet joints have hyperosteoecy (Figure 2). The X-ray film of the right knee joint showed that the right knee joint space was narrow inside and wide outside, the articular surface was sclerotic, and the lower edge of the articular surface was spotted with low density. Hyperosteoecy appeared on the upper edge of patella, tibial plateau, tibial intercondylar crest and femoral condyle. The density of the suprapatellar bursa is increased and the surrounding soft tissue is swollen (Figure 3). The X-ray film of the full length of the spine shows that L3-4 vertebral body is diseased and shifted backward, the lumbar spine protrudes backward at the L2-3 plane, the intervertebral space at the L2-3 segment is narrowed, and the L3 vertebral body is wedge-shaped and flattened (Figure 4). The lumbar spine Magnetic Resonance Imaging (MRI) showed: Lumbar vertebrae protrude backward, L3 vertebrae show wedge-shaped changes, spinal canal stenosis on the same level of L3, intervertebral space narrowing on the same level of L2-3, endplate inflammation in the L3-4 intervertebral space, Schmorl node formation near T10-11 vertebral body, degeneration and bulge of L3-4 and L4-5 intervertebral discs, and many abnormal strip signals in the filum terminale (Figure 5). The right knee MRI showed: The cartilage of the medial femoral condyle and tibial plateau is worn, the cartilage is denatured, the posterior cruciate ligament of the right knee is torn, the medial collateral ligament is injured, the anterior and posterior corners of the medial and lateral meniscus of the right knee are worn, the right knee joint has effusion, and the medial head bursa of the gastrocnemius has effusion (Figure 6). Lumbar vertebral body (plain scan + 3D reconstruction) CT showed: Lumbar vertebral body (plain scan + 3D reconstruction) CT showed: L2-3 plane lumbar lordosis, L2-3 vertebral space narrowing, L3 vertebral body wedge-shaped flattening, L3, 4 vertebral body adjacent edge patchy high density. There are some bone defects in the lamina and spinous process of L3 and L4 vertebrae, and the vertebral canal at the same plane is deformed (Figure 7).


### **Neurological examination**

Motor nerve conduction velocity measurement (both lower extremities), sensory nerve conduction velocity measurement (both lower extremities), neuroelectrogram, needle electrode electromyography results showed: Neurogenic damage in both lower extremities.

## **FINAL DIAGNOSIS**


Based on this patient's imaging data and his symptoms, we diagnosed this patient with Post-traumatic cauda equina nerve calcification.





DOI: 10.12998/wjcc.v11.i6.1356 Copyright ©The Author(s) 2023.

**Figure 1 Lumbar spine computed tomography.** L5 plane end filament and cauda equina nerve calcification signs. A: L5 upper edge of vertebral body; B: L5 center of vertebral body.




DOI: 10.12998/wjcc.v11.i6.1356 Copyright ©The Author(s) 2023.

**Figure 2 The X-ray film of lumbar vertebrae.** The sequence of lumbar vertebrae is abnormal, L3 vertebral body is diseased and shifted backward, L2-3 intervertebral space disappears, L2-3, L3-4, L4-5, L5-S1 intervertebral space narrows, and all vertebrae and vertebral facet joints have hyperosteoecy. A: Orthostatic lumbar X-ray film; B: Lateral lumbar X-ray film.


## TREATMENT

We use the combination of traditional Chinese and western medicine to treat the patient's clinical symptoms symptomatically. Use mannitol injection, dexamethasone sodium phosphate injection, pregabalin capsules, etocoxib, mecobalamin tablets, sodium glutamate injection, and compound betamethasone injection according to scientific drug use rules to dehydrate and detumescence, nourish nerves, diminish inflammation and pain. In addition, we also use radio frequency electrotherapy, traditional Chinese medicine directional penetration, intracutaneous acupuncture, radioactive shock wave and other physical therapy methods to activate blood circulation and remove blood stasis, warm meridians and relieve pain.



DOI: 10.12998/wjcc.v11.i6.1356 Copyright ©The Author(s) 2023.

**Figure 3 The X-ray film of the right knee joint.** The right knee joint space was narrow inside and wide outside, the articular surface was sclerotic, and the lower edge of the articular surface was spotted with low density. Hyperosteoegeny appeared on the upper edge of patella, tibial plateau, tibial intercondylar crest and femoral condyle. The density of the suprapatellar bursa is increased and the surrounding soft tissue is swollen. A: Orthostatic Knee joint X-ray film; B: Lateral Knee joint X-ray film.



DOI: 10.12998/wjcc.v11.i6.1356 Copyright ©The Author(s) 2023.


**Figure 4 The X-ray film of the full length of the spine.** L3-4 vertebral body is diseased and shifted backward, the lumbar spine protrudes backward at the L2-3 plane, the intervertebral space at the L2-3 segment is narrowed, and the L3 vertebral body is wedge-shaped and flattened. A: The Orthostatic X-ray film of the full length of the spine; B: The Orthostatic X-ray film of lumbar spine and pelvis.

## OUTCOME AND FOLLOW-UP

We advise patients to maintain a light diet and recommend active functional exercise. Continue to take etoricoxib tablets, methylcobalamin tablets, and pregabalin capsules. The patient was also advised to carry out a variety of blood-activating and stasis-removing types of physical therapy. After completing regular follow-up (1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> and months after discharge), the patient's limb muscle strength changed to grade II and limb sensation appeared. The patient had good compliance.


## DISCUSSION

Nerve calcification usually occurs in the cranial nerve and the dental nerve. It is generally accepted that nerve calcification is divided into two main types: Physiological and pathological calcification. Physiological calcification is due to the natural formation of calcified spots on the nerve with age. Pathological calcification is caused by degenerative diseases, parasitic infections, ischemic necrosis of the surrounding tissue of the nerve, oxidative stress, and apoptosis[3]. However, pathological calcification of cauda equina nerve is extremely rare. We speculate that cauda equina nerve calcification in this patient may be secondary to acute trauma and surgical scar, thus stimulating local lesions. In addition, lumbar spinal stenosis, lumbar disc herniation, spinal tumors and spinal infection are also possible causes of cauda equina nerve calcification[4]. L2-5, S1-5 and the 10 pairs of nerve roots emanating from their caudal segments together form the cauda equine nerve. The cauda equina nerve, in the usual sense, is the nerve root below L2 that innervates the pelvis as well as the perineum. It is a very important nerve in the human body[5]. Once the cauda equina nerve undergoes pathological changes, it will lead to dysfunction of sensory nerve, motor nerve, autonomic nerve, *etc.*, thus causing



DOI: 10.12998/wjcc.v11.i6.1356 Copyright ©The Author(s) 2023.


**Figure 5 The lumbar spine magnetic resonance imaging.** Lumbar vertebrae protrude backward, L3 vertebrae show wedge-shaped changes, spinal canal stenosis on the same level of L3, intervertebral space narrowing on the same level of L2-3, endplate inflammation in the L3-4 intervertebral space, Schmorl node formation near T10-11 vertebral body, degeneration and bulge of L3-4, L4-5 and L5-S1 intervertebral discs, and many abnormal strip signals in the filum terminale. A: L5-S1 Plain magnetic resonance imaging (MRI) scan of L5-S1 intervertebral disc; B: L3 MRI plain scan.



DOI: 10.12998/wjcc.v11.i6.1356 Copyright ©The Author(s) 2023.

**Figure 6 The right knee magnetic resonance imaging.** The cartilage of the medial femoral condyle and tibial plateau is worn, the cartilage is denatured, the posterior cruciate ligament of the right knee is torn, the medial collateral ligament is injured, the anterior and posterior corners of the medial and lateral meniscus of the right knee are worn, the right knee joint has effusion, and the medial head bursa of the gastrocnemius has effusion. A: Lateral magnetic resonance imaging (MRI) of knee; B: Oblique MRI of knee.

other body dysfunction. as opposed to the cauda equina nerve calcification, Cauda equina injury is relatively common in clinical practice[6]. The clinical symptoms of cauda equina compression were first reported by Verbiest in 1949 and were named Cauda Equine Syndrome (CES). The clinical symptoms of cauda equina nerve injury mainly include Low back pain, numbness of one or both lower limbs, weakness of lower limbs, weakness or disappearance of tendon reflex, sensory abnormalities in the sellar region, sphincter dysfunction, urinary incontinence, and sexual dysfunction. The cauda equina nerve is easy to be damaged (for example, the free nucleus pulposus and epidural hematoma will cause the cauda equina nerve to be compressed), because the cauda equina nerve has no protective sheath of connective tissue and is particularly sensitive to compression. The recovery after peripheral nerve injury is very slow, which may be due to the lack of blood supply. The nutrition of cauda equina nerve also comes from cerebrospinal fluid and blood, and the recovery of cauda equina nerve after injury is slower than that of peripheral nerve. Therefore, we can boldly speculate that the mechanism of cauda equina nerve calcification may be that the surrounding tissues of cauda equina nerve suffer from mechanical compression or tissue necrosis, which affects the blood supply of cauda equina nerve. Due to long-term



DOI: 10.12998/wjcc.v11.i6.1356 Copyright ©The Author(s) 2023.

**Figure 7 Lumbar vertebral body (plain scan + 3D reconstruction) computed tomography.** Lumbar vertebral body (plain scan + 3D reconstruction) computed tomography (CT) showed: L2-3 plane lumbar lordosis, L2-3 vertebral space narrowing, L3 vertebral body wedge-shaped flattening, L3, 4 vertebral body adjacent edge patchy high density. There are some bone defects in the lamina and spinous process of L3 and L4 vertebrae, and the vertebral canal at the same plane is deformed. A: Lumbar vertebral body CT (3D reconstruction); B: Lumbar vertebral body CT (plain scan).

poor blood supply, nerve dehydration and poor calcium and phosphorus metabolism lead to calcium salt deposition, which eventually leads to cauda equina nerve calcification[7,8].

There are many reports on Idiopathic Basal Ganglia Calcification (IBGC) in the existing literature, but the specific cause is still unclear, and scholars believe that it is related to the following factors: (1) Genetic studies show that IBCG patients have autosomal recessive or dominant inheritance; (2) Vitrification during atherosclerosis can lead to calcium deposition, so some scholars believe that local calcium accumulation is related to changes in vascular osmotic pressure; (3) The abnormal level metabolism of iron and calcium phosphate, especially the decrease of serum ferritin level and iron binding capacity, will also lead to IBCG; (4) Exogenous toxic substances continuously stimulate and activate glutamate receptor, thus producing neurotoxin effect, leading to calcium ion deposition; (5) It has been reported in the literature that during brain biopsy of patients with IBCG, there are immune inflammatory cells infiltrating around calcification points and ESR is accelerated, while C-reactive protein, rheumatoid factor (RF) and antinuclear antibody (ANA) are significantly increased. So the occurrence of IBCG is also related to the immune system. It is worth noting here that the ESR, C-reactive protein, RF and ANA of the cases reported by us are higher than the normal values. This is consistent with the study of IBCG; and (6) The IBCG has also been suggested to be associated with carbonic anhydrase II deficiency. Regrettably, the epidemiological characteristics and specific treatment methods of basal ganglia calcification have not been studied[9,10,11].

The benign tumor of the cauda equina nerve may also lead to the calcification of the cauda equina nerve. We call this benign tumor the neurilemmoma of the cauda equina nerve. spinal cord neurilemmoma often occur in the spinal cord nerve roots, but only a few occur in the cauda equina nerve. Because the neurilemmoma of the cauda equina nerve has good activity and broad intradural space, it usually has no obvious pain symptoms. The report of neurilemmoma calcification is extremely rare. In 2012, Dr. Seung Jae Hyun and others from South Korea reported that a 21 year old patient with dystrophic neurilemmoma of cauda equina had calcification, and the neurilemmoma was completely removed by surgery. However, this report does not explain the relationship between trauma and calcification of cauda equina neurilemmoma[12]. Paraganglioma is a neuroendocrine tumor located outside the adrenal gland. Paragangliomas originating from filum terminale or cauda equina nerve are rare (accounting for 1% of all paragangliomas and 3%-4% of all tumors in the lower lumbar spine). Calcification of cauda equina nerve paraganglioma is very rare. It is noteworthy that Professor M Vural once reported cauda equina paraganglioma with obvious calcification characteristics, but did not clarify its cause and pathogenesis[13]. Professor J Rot é s Querol once reported a case of cauda equina syndrome caused by ankylosing spondylitis, which caused calcification of lumbosacral meninges. However, the CT images of the cases reported by Professor J Rot é s Querol showed perispinal calcification rather than central calcification[14]. Intradural lumbar disc herniation is another important cause of intraspinal calcification and even cauda equina nerve calcification. Intradural lumbar disc herniation means that the free nucleus pulposus punctures the fibrous ring, posterior longitudinal ligament and dura mater, and directly compresses the spinal cord or nerve after entering the subdural cavity, resulting in acute spinal cord or cauda equina nerve injury. Intradural disc herniation is a rare type of disc herniation, with a incidence rate of 0.26%-0.3%. The most common location of the disease was the lumbar spine, accounting for 92%[15]. Surgical exploration is the only standard for the diagnosis of intradural lumbar disc herniation. Although it is difficult to make a definite diagnosis of intradural lumbar disc herniation through imaging and symptoms, some special signal signs shown by



imaging data can well indicate the existence of free intradural nucleus pulposus. For example, the Y sign refers to the prominent nucleus pulposus separating the dura and arachnoid membranes into two lines in a "Y" shape. Over time, the free nucleus pulposus entering the dura will directly compress the cauda equina nerve and calcification will occur to some extent[16]. We speculate that the calcification of the free nucleus pulposus will probably cause the calcification of the cauda equina nerve compressed by it.

Since cauda equina nerve calcification in this patient is secondary to trauma stress, it is necessary to discuss arachnoiditis during injury. There are also some literatures that study the problems related to cauda equina nerve calcification from this perspective. In 1999, I. G. Bilgen reported a case of cauda equina syndrome caused by ankylosing spondylitis, a patient with adhesive arachnoiditis. CT data of the patient showed signs of dural calcification. His MRI data showed that the cauda equina had adhesions with the dorsal arachnoid membrane. They believe that arachnoiditis is responsible for cauda equina syndrome and curvilinear dural calcification[17]. In 2021, Brunner A *et al*[18] introduced a case of intradural calcification caused by chronic adhesive arachnoiditis and performed a dural incision exploration, and found nerve calcification structures in the dural sac. Brunner A *et al*[18] believed that the main cause of calcification was the inflammation of arachnoid membrane. Arachnoid inflammation leads to excessive proliferation of arachnoid cells, which leads to excessive production of collagen tissue and rapid formation of intradural scars. Subsequently, osteoblasts are generated, leading to progressive intradural ossification. However, the pathogenesis of ossifying arachnoiditis has not been completely clarified. El Asri AC *et al*[19] reported a rare case of post-traumatic Arachnoiditis ossificans of the cauda equina nerve. El Asri AC *et al*[19] clearly proposed that trauma, surgery, infection and subarachnoid hemorrhage are the main causes of post traumatic arachnoiditis ossificans of the cauda equina nerve.

Since cases of Post-traumatic cauda equina nerve calcification are extremely rare in clinical practice, our research on this disease has not been carried out and the literature that we have been able to study is very scarce. Therefore, it is necessary that we carry out studies on the etiology, mechanisms, treatment and prognosis of this disease.

## CONCLUSION

We provide a rare case of Post-traumatic cauda equina nerve calcification. This case reports the symptoms, imaging data and treatment of a patient with Post-traumatic cauda equina nerve calcification, and analyzes the etiology and mechanism of Post-traumatic cauda equina nerve calcification.

## FOOTNOTES

**Author contributions:** Liu YD, Deng Q, Li JJ, Zhang KD contributed to conceptualization; Liu YD, Li JJ, Peng RD contributed to data curation and investigation; Deng Q conducted the project administration; Liu YD, Han XF, Yang HY and Xiang QQ wrote the original draft, and reviewed and edited the manuscript; all authors issued final approval for the version to be submitted.

**Supported by** National Natural Science Foundation of China, No. 82060879; Natural Science Foundation of Gansu Province, No. 20JR10RA356 and No. 2022-0405-JCC-1430; Lanzhou Science and Technology Plan Project, No. 2022-3-30; Paikouen-Spine Pathological Fracture Vertebral Body Strengthening Treatment Special Fund Project, No. BK-JP2018004; Local Projects Transferred by the Central Government in 2021, No. 20210200111.

**Informed consent statement:** Informed written consent was obtained from the patient for publication of this report and any accompanying images.

**Conflict-of-interest statement:** All authors declare that they have no conflict of interest to disclose.

**CARE Checklist (2016) statement:** All authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Qiang Deng 0000-0003-1193-9307; Xian-Fu Han 0000-0002-3240-497X.

**S-Editor:** Ma YJ

**L-Editor:** A

P-Editor: Ma YJ

## REFERENCES

- 1 **Joshi A**, Chitale N, Phansopkar P. The Impact of Physical Therapy Rehabilitation on Pain and Function in a Patient With Cauda Equina Syndrome. *Cureus* 2022; **14**: e28131 [PMID: [36134093](#) DOI: [10.7759/cureus.28131](#)]
- 2 **Fukatsu S**, Ogiwara S, Imada H, Ikemune S, Tamaru JI, Saita K. Chronic spontaneous epidural hematoma in the lumbar spine with cauda equina syndrome and severe vertebral scalloping mimicking a spinal tumor: a case report. *BMC Musculoskelet Disord* 2022; **23**: 508 [PMID: [35637479](#) DOI: [10.1186/s12891-022-05463-x](#)]
- 3 **Isen DR**, Gaddamanugu S, Kline LB. Optic Nerve Sheath Calcification. *Ophthalmology* 2022 [PMID: [35773073](#) DOI: [10.1016/j.ophtha.2022.03.013](#)]
- 4 **Nasim O**, Eskander B, Rustam Z, Pantelias C, Moverley R. Expediting the Management of Suspected Cauda Equina Syndrome (CES) in the Emergency Department Through Clinical Pathway Design at a District General Hospital: A Quality Improvement Project. *Cureus* 2022; **14**: e32722 [PMID: [36570114](#) DOI: [10.7759/cureus.32722](#)]
- 5 **Ando T**, Watanabe H, Riku Y, Yoshida M, Goto Y, Ando R, Fujino M, Ito M, Koike H, Katsuno M, Iwasaki Y. Neurogenic intermittent claudication caused by vasculitis in the cauda equina: an autopsy case report. *Eur Spine J* 2022 [PMID: [36416968](#) DOI: [10.1007/s00586-022-07458-7](#)]
- 6 **Khashan M**, Ofir D, Grundshtein A, Kuzmenko B, Salame K, Nir D, Hochberg U, Lidar Z, Regev GJ. Minimally invasive discectomy vs open laminectomy and discectomy for the treatment of cauda equina syndrome: A preliminary study and case series. *Front Surg* 2022; **9**: 1031919 [PMID: [36311945](#) DOI: [10.3389/fsurg.2022.1031919](#)]
- 7 **Rascón-Ramírez FJ**. Spinal cord stimulation and cauda equina syndrome: Could it be a valid option? *Neurocirugía (Astur: Engl Ed)* 2022; **33**: 90-94 [PMID: [35248303](#) DOI: [10.1016/j.neucie.2020.12.002](#)]
- 8 **Moussa MK**, Alkefrawi P, Elkhaili JK. Large central disc herniation causing cauda equina syndrome in an adolescent. A case report. *Int J Surg Case Rep* 2021; **79**: 119-122 [PMID: [33454631](#) DOI: [10.1016/j.ijscr.2020.12.066](#)]
- 9 **Li M**, Fu Q, Xiang L, Zheng Y, Ping W, Cao Y. SLC20A2-Associated Idiopathic basal ganglia calcification (Fahr disease): a case family report. *BMC Neurol* 2022; **22**: 438 [PMID: [36397039](#) DOI: [10.1186/s12883-022-02973-y](#)]
- 10 **Durante A**, Audino N, Cristiano M, Tanga M, Martino MT, Noschese I, D'Auria D, Pinto F. Basal ganglia calcification: a Fahr's disease case report. *Radiol Case Rep* 2021; **16**: 3055-3059 [PMID: [34429801](#) DOI: [10.1016/j.radcr.2021.07.042](#)]
- 11 **Peters MEM**, de Brouwer EJM, Bartstra JW, Mali WPTM, Koek HL, Rozemuller AJM, Baas AF, de Jong PA. Mechanisms of calcification in Fahr disease and exposure of potential therapeutic targets. *Neurol Clin Pract* 2020; **10**: 449-457 [PMID: [33299674](#) DOI: [10.1212/CPJ.0000000000000782](#)]
- 12 **Hyun SJ**, Rhim SC. Giant cauda equina schwannoma with dystrophic calcifications : case report and review of the literature. *J Korean Neurosurg Soc* 2012; **51**: 105-108 [PMID: [22500204](#) DOI: [10.3340/jkns.2012.51.2.105](#)]
- 13 **Vural M**, Arslantas A, Isiksoy S, Adapinar B, Atasoy M, Soylemezoglu F. Gangliocytic paraganglioma of the cauda equina with significant calcification: first description in pediatric age. *Zentralbl Neurochir* 2008; **69**: 47-50 [PMID: [18393166](#) DOI: [10.1055/s-2007-985162](#)]
- 14 **Rotés-Querol J**, Tolosa E, Roselló R, Granados J. Progressive cauda equina syndrome and extensive calcification/ossification of the lumbosacral meninges. *Ann Rheum Dis* 1985; **44**: 277-280 [PMID: [3872639](#) DOI: [10.1136/ard.44.4.277](#)]
- 15 **Francio VT**, Wie CS, Murphy MT, Neal MT, Lyons MK, Gibbs WN, Strand NH. Multispecialty perspective on intradural disc herniation: diagnosis and management - A case report. *Anesth Pain Med (Seoul)* 2022; **17**: 221-227 [PMID: [35378571](#) DOI: [10.17085/apm.21100](#)]
- 16 **Thohar Arifin M**, Ikbar K N, Brilliantika SP, Bakhtiar Y, Bunyamin J, Muttaqin Z. Challenges in intradural disc herniation diagnosis and surgery: A case report. *Ann Med Surg (Lond)* 2020; **58**: 156-159 [PMID: [32983437](#) DOI: [10.1016/j.amsu.2020.08.022](#)]
- 17 **Bilgen IG**, Yuntun N, Ustun EE, Oksel F, Gumusdis G. Adhesive arachnoiditis causing cauda equina syndrome in ankylosing spondylitis: CT and MRI demonstration of dural calcification and a dorsal dural diverticulum. *Neuroradiology* 1999; **41**: 508-511 [PMID: [10450845](#) DOI: [10.1007/s002340050793](#)]
- 18 **Brunner A**, Leoni M, Eustacchio S, Kurschel-Lackner S. Spinal Arachnoiditis Ossificans: A Case-Based Update. *Surg J (N Y)* 2021; **7**: e174-e178 [PMID: [34307874](#) DOI: [10.1055/s-0041-1731448](#)]
- 19 **El Asri AC**, El Mostarchid B, Akhaddar A, Baallal H, Dao I, Naama O, Gazzaz M, Boucetta M. Arachnoiditis ossificans of the cauda equina. *Br J Neurosurg* 2012; **26**: 547-548 [PMID: [22239274](#) DOI: [10.3109/02688697.2011.645916](#)]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

