

World Journal of *Clinical Cases*

World J Clin Cases 2023 March 6; 11(7): 1434-1668



Contents

Thrice Monthly Volume 11 Number 7 March 6, 2023

OPINION REVIEW

- 1434 Reconstruction surgery in head and neck cancer patients amidst the COVID-19 pandemic: Current practice and lessons for the future

Lizambri D, Giacalone A, Shah PA, Tovani-Palone MR

REVIEW

- 1442 Risk factors and digital interventions for anxiety disorders in college students: Stakeholder perspectives

Liu XQ, Guo YX, Xu Y

MINIREVIEWS

- 1458 Immune-related adverse events induced by programmed death protein-1 inhibitors from the perspective of lymphoma immunotherapy

Hou YZ, Zhang Q, Bai H, Wu T, Chen YJ

ORIGINAL ARTICLE

Clinical and Translational Research

- 1467 Analysis of differentially expressed genes related to cerebral ischaemia in young rats based on the Gene Expression Omnibus database

Xia Y, Liu H, Zhu R

Retrospective Study

- 1477 Deep learning-assisted diagnosis of femoral trochlear dysplasia based on magnetic resonance imaging measurements

Xu SM, Dong D, Li W, Bai T, Zhu MZ, Gu GS

- 1488 Facial basal cell carcinoma: A retrospective study of 67 cases

Khalil AA, Enezei HH, Aldelaimi TN, Al-Ani RM

CASE REPORT

- 1498 Successful multidisciplinary therapy for a patient with liver metastasis from ascending colon adenocarcinoma: A case report and review of literature

Tan XR, Li J, Chen HW, Luo W, Jiang N, Wang ZB, Wang S

- 1506 Accessory renal arteries - a source of hypertension: A case report

Calinoiu A, Guluta EC, Rusu A, Minca A, Minca D, Tomescu L, Gheorghita V, Minca DG, Negreanu L

- 1513 Synchronous multiple primary malignant neoplasms in breast, kidney, and bilateral thyroid: A case report

Jia MM, Yang B, Ding C, Yao YR, Guo J, Yang HB

- 1521** Invasive breast carcinoma with osteoclast-like stromal giant cells: A case report
Wang YJ, Huang CP, Hong ZJ, Liao GS, Yu JC
- 1528** Retroperitoneal and abdominal bleeding in anticoagulated COVID-19 hospitalized patients: Case series and brief literature review
Evrev D, Sekulovski M, Gulina M, Dobrev H, Velikova T, Hadjidekov G
- 1549** Hyperthyroidism and severe bradycardia: Report of three cases and review of the literature
He YL, Xu WX, Fang TY, Zeng M
- 1560** Isolated cerebral mucormycosis that looks like stroke and brain abscess: A case report and review of the literature
Chen CH, Chen JN, Du HG, Guo DL
- 1569** Gastric ectopic pancreas combined with synchronous multiple early gastric cancer: A rare case report
Zhao ZY, Lai YX, Xu P
- 1576** Manifestation of the malignant progression of glioma following initial intracerebral hemorrhage: A case report
Xu EX, Lu SY, Chen B, Ma XD, Sun EY
- 1586** Four kinds of antibody positive paraneoplastic limbic encephalitis: A rare case report
Huang P, Xu M
- 1593** Spontaneous fracture of a titanium mesh cranioplasty implant in a child: A case report
Zhang R, Gao Z, Zhu YJ, Wang XF, Wang G, He JP
- 1600** Rheumatic valvular heart disease treated with traditional Chinese medicine: A case report
Chen WH, Tan Y, Wang YL, Wang X, Liu ZH
- 1607** Mucosa-associated lymphoid tissue lymphoma of the trachea treated with radiotherapy: A case report
Zhen CJ, Zhang P, Bai WW, Song YZ, Liang JL, Qiao XY, Zhou ZG
- 1615** Bow-and-arrow sign on point-of-care ultrasound for diagnosis of pacemaker lead-induced heart perforation: A case report and literature review
Chen N, Miao GX, Peng LQ, Li YH, Gu J, He Y, Chen T, Fu XY, Xing ZX
- 1627** Prostate lymphoma with renal obstruction; reflections on diagnosis and treatment: Two case reports
Chen TF, Lin WL, Liu WY, Gu CM
- 1634** Pulmonary nocardiosis with bloodstream infection diagnosed by metagenomic next-generation sequencing in a kidney transplant recipient: A case report
Deng ZF, Tang YJ, Yan CY, Qin ZQ, Yu N, Zhong XB
- 1642** Primary yolk sac tumor in the abdominal wall in a 20-year-old woman: A case report
Wang Y, Yang J

- 1650** Misdiagnosis of food-borne foreign bodies outside of the digestive tract on magnetic resonance imaging: Two case reports

Ji D, Lu JD, Zhang ZG, Mao XP

- 1656** IgG4-related kidney disease complicated with retroperitoneal fibrosis: A case report

He PH, Liu LC, Zhou XF, Xu JJ, Hong WH, Wang LC, Liu SJ, Zeng JH

LETTER TO THE EDITOR

- 1666** Commentary on a case report and literature review of acute carotid stent thrombosis

Willman M, Lucke-Wold B

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Baharudin Ibrahim, BPharm, PhD, Associate Professor, Pharmacist, Department of Clinical Pharmacy and Pharmacy Practice, Faculty of Pharmacy, Universiti Malaya, Kuala Lumpur 50603, Malaysia. baharudin.ibrahim@um.edu.my

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJCC as 1.534; IF without journal self cites: 1.491; 5-year IF: 1.599; Journal Citation Indicator: 0.28; Ranking: 135 among 172 journals in medicine, general and internal; and Quartile category: Q4. The WJCC's CiteScore for 2021 is 1.2 and Scopus CiteScore rank 2021: General Medicine is 443/826.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Si Zhao; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

March 6, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Isolated cerebral mucormycosis that looks like stroke and brain abscess: A case report and review of the literature

Cai-Hong Chen, Jing-Nan Chen, Hang-Gen Du, Dong-Liang Guo

Specialty type: Medicine, research and experimental

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Kaushik P, India; Liu Y, China

Received: November 25, 2022

Peer-review started: November 25, 2022

First decision: December 13, 2022

Revised: January 7, 2023

Accepted: February 10, 2023

Article in press: February 10, 2023

Published online: March 6, 2023



Cai-Hong Chen, Dong-Liang Guo, Department of Neurology, Hangzhou Ninth People's Hospital, Hangzhou 311225, Zhejiang Province, China

Jing-Nan Chen, Hang-Gen Du, Department of Neurosurgery, The Second Affiliated Hospital of Zhejiang University of Traditional Chinese Medicine, Hangzhou 310005, Zhejiang Province, China

Corresponding author: Jing-Nan Chen, MD, Doctor, Professor, Department of Neurosurgery, The Second Affiliated Hospital of Zhejiang University of Traditional Chinese Medicine, No. 318 Chaowang Road, Gongshu District, Hangzhou 310005, Zhejiang Province, China. 1028713842@qq.com

Abstract

BACKGROUND

Cerebral mucormycosis is an infectious disease of the brain caused by fungi of the order Mucorales. These infections are rarely encountered in clinical practice and are often misdiagnosed as cerebral infarction or brain abscess. Increased mortality due to cerebral mucormycosis is closely related to delayed diagnosis and treatment, both of which present unique challenges for clinicians.

CASE SUMMARY

Cerebral mucormycosis is generally secondary to sinus disease or other disseminated disease. However, in this retrospective study, we report and analyze a case of isolated cerebral mucormycosis.

CONCLUSION

The constellation of symptoms including headaches, fever, hemiplegia, and changes in mental status taken together with clinical findings of cerebral infarction and brain abscess should raise the possibility of a brain fungal infection. Early diagnosis and prompt initiation of antifungal therapy along with surgery can improve patient survival.

Key Words: Cerebral mucormycosis; Stroke; Brain abscess; Diagnosis; Treatment; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Cerebral mucormycosis is an infectious disease of the brain caused by fungi of the order Mucorales. These infections are rarely encountered in clinical practice and often misdiagnosed as cerebral infarction or brain abscess. Diagnosis and treatment are challenging, and as such this disease is often associated with a high mortality rate. Through this case study, we aim to help further understand the pathophysiology of cerebral mucormycosis and suggest strategies for the improvement of clinical diagnosis and treatment thereof.

Citation: Chen CH, Chen JN, Du HG, Guo DL. Isolated cerebral mucormycosis that looks like stroke and brain abscess: A case report and review of the literature. *World J Clin Cases* 2023; 11(7): 1560-1568

URL: <https://www.wjgnet.com/2307-8960/full/v11/i7/1560.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i7.1560>

INTRODUCTION

Cerebral mucormycosis is a life-threatening fungal infection[1]. Although modern technology advances and various imaging techniques are available to aid its diagnosis and treatment, this disease still carries a mortality rate as high as 62%[2-4]. Mucormycosis can affect various areas of the body, including rhino-cerebral tissue, lung, skin, and gastrointestinal tract, and can even present as a disseminated infection. Cerebral mucormycosis is the most common presentation of infection caused by Mucorales fungi and can occur *via* direct invasion of rhino-orbital structures or by hematogenous spread to the brain[5].

Cerebral mucormycosis cases can be found across the globe, with 47 countries reporting this infection. North America accounts for the majority of reported cases (130/345, 37.68%), followed by Asia (87/345, 25.22%). The incidence of cerebral mucormycosis is higher in males (60%) and the age distribution of affected patients is wide (2 mo to 89 years)[6]. Cerebral mucormycosis is generally diagnosed in patients with underlying medical conditions, with diabetes being the most common[7]. Amphotericin B deoxycholate (AmB) antifungal therapy and surgical debridement are the most commonly utilized methods for the treatment of cerebral mucormycosis.

Isolated cerebral mucormycosis is very rare. Due to the lack of specific clinical manifestations, the misdiagnosis rate is high in the early stages of the disease. Up to 90% of cases are reported to go undiagnosed and untreated[8,9]. The most common symptoms are orbital and neurological, followed by headache. Neurological symptoms are often associated with other extracerebral symptoms, with fever reported in greater than 1 in 5 cases[10,11]. Therefore, the presence of any of these symptoms in immunocompromised patients requires a high degree of suspicion for brain fungal infection. Here, we present a case of isolated cerebral mucormycosis followed by discussion and review of the current literature.

CASE PRESENTATION

Chief complaints

A 49-year-old female patient presented to the outpatient clinic with headache for 11 d, and left upper and lower extremity weakness accompanied by fever for 1 wk.

History of present illness

The patient had persistent pain mainly due to right frontal and parietal headache, which could not be relieved after rest. The patient was hospitalized at the Department of Neurology with a body temperature of 39.2 °C.

History of past illness

The patient had type 2 diabetes. A diagnostic consultation did not identify any foci of infection. She had no past history of otitis media, sinusitis, or head trauma.

Personal and family history

The patient denied having any other relevant personal history and her familial history was unremarkable.

Physical examination

The physical examination revealed intact consciousness with a Glasgow coma scale (GCS) score of 15, a soft neck, left upper and lower extremity muscular strength (1 out of 5), a shallow left nasolabial fold, and intact deep and superficial sensation.

Laboratory examinations

A complete blood count with differential revealed a white blood cell count of $13.4 \times 10^9/L$ (normal reference range: $4.0\text{--}10.0 \times 10^9/L$), 80.0% neutrophils (normal reference range: 50%–70%), 12.1% lymphocytes (normal reference range: 20%–40%), and C-reactive protein level of 50.1 mg/L (normal reference range: 0–10 mg/L). Urinalysis revealed a white blood cell count of 229/ μg (normal reference range: 0–26/ μg). Urine culture grew *Enterococcus faecalis*. Blood and sputum cultures were negative.

Imaging examinations

No obvious abnormalities were observed in a computed tomography (CT) scan of the chest. Cardiac Doppler ultrasound revealed mild mitral regurgitation. Magnetic resonance imaging (MRI) of the head with and without contrast demonstrated the following: (1) Large-scale abnormal signal lesions in the right frontal lobe and right basal ganglia with low signal on T1-weighted imaging, slightly higher signal on T2-weighted imaging, and high signal on diffusion-weighted imaging (DWI); (2) Enhancement of the gyrus around the lesion without significant space-occupying effect; and (3) Multiple long T2 signals and enhanced DWI signals in the left cerebral hemisphere (Figure 1). CT angiography of cervical arteries demonstrated occlusion of the right internal carotid artery (Figure 2A–D). Further examination of the arteries by magnetic resonance angiography showed that there was no development of the right internal carotid artery skull base and intracranial segments, and the anterior and posterior cerebral arteries were supplied by the traffic branch (Figure 2E and F).

Further diagnostic work-up


All available clinical evidence up to this point suggested cerebral infarction, with the only unexplained symptoms being persistent fever and headache. Per antibacterial susceptibility testing of the urine culture, piperacillin and tazobactam were administered intravenously, in addition to the initiation of standard treatment for cerebral infarction. In the following week, the patient's temperature fluctuated between 38 °C and 39 °C, and her headache did not improve significantly. Two consecutive cerebrospinal fluid tests revealed the following: (1) Greater than 100 nucleated cells, most of which were neutrophils; (2) Slightly elevated glucose and protein levels; (3) Normal chloride levels; and (4) Opening pressures of 170 mmH₂O and 210 mmH₂O. No bacteria or fungi were found in cultures of the cerebrospinal fluid. Considering the possibility of brain abscess, the patient was empirically treated with meropenem and vancomycin. However, as discussed above, a CT scan of the head showed a low-density shadow in the right frontal lobe and right basal ganglia. The right basal ganglia showed focal enhancement, the right lateral ventricle was slightly compressed, and there was slight midline shift to the left. In light of these findings, 100 mL intravenous mannitol, 250 mL intravenous glycerol fructose, and intravenous sodium chloride were infused every 12 h. The patient's temperature continued to fluctuate between 37 °C and 38 °C, and her headache persisted. Approximately 1 wk later, the patient's temperature suddenly rose to 39 °C. A review of the head CT revealed a large area of infarction corresponding to the right middle cerebral artery. The density of some areas was slightly higher, the midline was shifted, and cerebral palsy developed (Figure 2G and H). Accordingly, right craniotomy and drainage were performed, with turbid serosanguineous fluid and no signs of abscess observed intraoperatively. After 1 wk, the patient's level of consciousness decreased (GCS, E3V4M4). Repeat MRI of the head was still consistent with cerebral infarction (Figure 3A–F). However, after communicating with the patient's family, resection of the right brain lesion was performed and tissue was sent for gross pathologic and histologic examination. Findings reported by pathology included purulent brain tissue with visible fungal structures revealed by special staining to be consistent with mucormycosis (Figure 4). On the 10th d after the brain lesion resection, repeat head CT showed a large low-density shadow, and the midline returned to normal (Figure 3G and H).

FINAL DIAGNOSIS

A final diagnosis of cerebral mucormycosis was made following histopathological examination of the resected brain lesion.

TREATMENT

Following surgical resection of the right cerebral lesion, the patient received therapy with intravenous amphotericin B.



DOI: 10.12998/wjcc.v11.i7.1560 Copyright ©The Author(s) 2023.


Figure 1 Head magnetic resonance imaging findings. A-D: In the right frontal lobes and basal ganglia, large-scale abnormal signal lesions were found with low signal on T1-weighted imaging and slightly higher signal on T2-weighted imaging; E-H: Diffusion-weighted imaging signals were enhanced in the right frontal lobe and right basal ganglia, and multiple spotted high signals were observed in the left hemisphere; I and J: After enhancement, no ring enhancement was observed around the lesion, and there was no obvious space-occupying effect.

OUTCOME AND FOLLOW-UP

There were signs of improvement in the first 3 d following resection of the lesion. However, the patient gradually deteriorated and brain death was pronounced 1 mo later.

DISCUSSION

Cerebral mucormycosis is a life-threatening acute or chronic central nervous system disease caused by Mucorales fungi. In most cases, cerebral mucormycosis is caused by the spread of infection from the sinuses, and isolated cerebral mucormycosis is extremely rare. In a meta-analysis of 929 cases of mucormycosis, 30% described central nervous system disease, of which only 16% were confined to the central nervous system[2]. In other studies, isolated cerebral mucormycosis was found to account for up to 8% of all cases of mucormycosis[12-14]. Several risk factors are described in the literature, including intravenous drug use, diabetes mellitus, malignancy, solid and bone marrow transplantation, and iron overload status. Intravenous drug use is the most important risk factor for the development of isolated cerebral mucormycosis[15,16]. Verma *et al*[17] reported 30 cases of isolated cerebral mucormycosis, of which 17 were notable for history of intravenous drug abuse. Diabetes mellitus is the strongest risk factor for development of other mucormycosis infections, and in the meta-analysis of 929 cases mentioned above, 36% of patients with mucormycosis had a history of diabetes. It is believed that diabetic patients are susceptible to mucormycosis due to the fungal ketoreductase system allowing the



DOI: 10.12998/wjcc.v11.i7.1560 Copyright ©The Author(s) 2023.


Figure 2 Vascular examination and head computed tomography review. A-D: The right internal carotid artery was occluded, and the distal end was not shown. No obvious abnormalities were observed in the left carotid artery and vertebral artery; E and F: The right internal carotid artery skull base and intracranial segment were occluded, and the anterior and posterior cerebral arteries were supplied by the traffic branch; G and H: The right frontal lobe and basal ganglia showed lamellar low-density shadows, most of which tended to be liquid density, the right lateral ventricle was slightly stressed, and the midline was shifted to the left. RICA: Right internal carotid artery.

organisms to metabolize ketone bodies. In addition, the hyperglycemia and acidosis often found in the setting of diabetes reduce neutrophil chemotaxis and adhesion to fungal hyphae as well as impair the inhibition of Mucorales spores and mycelia by alveolar macrophages[18]. In the present case, the patient had a history of poorly controlled diabetes mellitus, and as such carried an increased susceptibility to mucormycosis infection due to these immunosuppressive mechanisms.

As mentioned previously, the most common symptoms of cerebral mucormycosis are orbital and neurological, followed by headache. Neurological symptoms are often related to other extracerebral symptoms such as fever. In our case, the symptoms demonstrated in the patient were headache, hemiplegia, and fever. In a review of 13 cases of isolated cerebral mucormycosis, altered mental status (54%) and headache (51%) were common[6]. Considering the presenting symptoms of our patient and those of patients discussed in the literature, it is apparent that these nonspecific signs and symptoms are largely responsible for the challenges faced by clinicians in diagnosing cerebral mucormycosis, as many other conditions must be considered in the differential diagnosis. To further add to these challenges, imaging findings can also vary widely. For example, Kursun *et al*[10] report cavernous sinus involvement as the most common brain imaging sign in 27 patients with cerebral mucormycosis, whereas in our case, abscess-like enhancing lesions were prominent.


In addition to the nonspecific presenting symptoms and imaging findings discussed above, many reported cerebrospinal fluid cultures are essentially negative, and culture conditions necessary for the growth of Mucorales fungi are inconsistently reported[19]. Histopathological examination of involved tissue can confirm the diagnosis; however, this requires invasive surgery. Taken together, these difficulties underscore the importance of a high clinical suspicion of cerebral mucormycosis and careful differential diagnosis.

Histopathological visualization of broad, ribbon-like, pauciseptate hyphae with wide-angle branching supports the diagnosis of mucormycosis. Acute inflammation with predominant neutrophilia is usually present, but may be difficult to identify in patients with underlying neutropenia. In addition, histopathological identification of mucormycosis does not provide genus and species information nor antifungal susceptibility data, and has limited ability to detect mixed fungal infections[20,21]. New diagnostic methods, including matrix-assisted laser desorption/ionization-time of flight (commonly known as MALDI-TOF) testing of serum and cryosections, can be used to aid diagnosis[22]. Unfortunately, molecular techniques may not be available at all institutions, and histopathological examination remains the gold standard for the diagnosis of isolated cerebral mucormycosis[23,24]. In our case, the histologic



DOI: 10.12998/wjcc.v11.i7.1560 Copyright ©The Author(s) 2023.

Figure 3 Head magnetic resonance imaging and postoperative head computed tomography review. A-D: Flaky T2 mixed signal was observed in the right frontal lobe and basal ganglia, there was a mixed signal on diffusion-weighted imaging (DWI), and the right lateral ventricle was compressed. In the left cerebral hemisphere, multiple long T2 signals were observed, some of which were high on DWI, and the midline was shifted to the left; E and F: After enhancement, partial ring enhancement was observed; G and H: On the 10th d after the brain lesion resection, repeat head computed tomography showed a large low-density shadow, and the midline returned to normal.



DOI: 10.12998/wjcc.v11.i7.1560 Copyright ©The Author(s) 2023.

Figure 4 Histopathological examination findings. A and B: Hematoxylin-eosin staining (100 and 400 × magnification) showed a large area of necrotic brain tissue, focal suppurative inflammation, and a blood vessel infiltrated by non-septate hyphae (black arrows); C: Periodic acid-silver methenamine staining (400 × magnification) showed sporangium, characteristic of Mucorales fungi (black arrows); D: Periodic acid-Schiff staining (400 × magnification) showed multiple broad hyphae.

diagnosis was confirmed by hematoxylin-eosin staining and periodic acid-silver methenamine staining of resected brain tissue.

Mucorales fungi are extremely invasive, and their hyphae can invade blood vessels. The growth of hyphae in the lumen of the blood vessels with associated damage to the endothelium can cause vascular occlusion, and formed fungal emboli can easily cause cerebral infarction. The blood vessels usually involved are the basilar artery and carotid artery, and the extravasation of fungal elements can lead to brain abscess formation[25,26]. In our case, the right internal carotid artery was completely occluded, and the MRI of the head demonstrated infarction in the right frontal lobe and right basal ganglia, likely due to impediment of blood flow by fungal emboli. However, infarction may also be a consequence of abscess formation.

The treatment of patients with mucormycosis is challenging, and even in adequately treated patients, the associated mortality rate is still high. Standard treatments include AmB and its lipid formulation and surgical debridement[27,28]. In most retrospective studies, AmB and its lipid formulation are the preferred treatment and are part of any potential combination therapy. Posaconazole in combination

with AmB is reported to offer treatment benefits when compared to antifungal monotherapy[29]. In some reports, posaconazole may be a more advantageous treatment strategy when used as the first line treatment for cerebral mucormycosis in diabetic patients with antifungal-resistant infections[30,31]. Moreover, vascular invasion by Mucorales fungi leads to vascular occlusion and necrosis, in turn significantly reducing drug distribution to target tissue. This emphasizes the importance of surgical debridement in the treatment of cerebral mucormycosis, especially in the setting of vascular occlusion [32-34]. Debridement can aid in treatment by reducing the microbiological load and altering anaerobic and microaerobic conditions conducive to fungal reproduction. However, in many case studies, death occurred despite surgical debridement and adequate antifungal treatment. These deaths are largely attributed to the rapid invasion of the brain by Mucorales fungi, delayed surgical treatment, and progressive infections[35,36]. The combination of surgery and medication is superior to either treatment alone. Roden *et al*[2] reported in a review of 929 cases that the survival rate was 61% (324 of 532) in the AmB alone treatment group, 57% (51 of 90) in the surgery alone treatment group, and 70% (328 of 470) in the antifungal and surgical combined treatment group.

A literature review suggests that the risk factors for death of patients with cerebral mucormycosis can be categorized as follows: (1) Susceptibility to infection (*e.g.*, history of diabetes and long-term use of broad-spectrum antibiotics); (2) Site of infection (*e.g.*, Chandra *et al*[37] found in a retrospective analysis of 68 cases that basal ganglia lesions carried increased mortality due to higher iron availability in that region); and (3) Delay in diagnosis and treatment (*e.g.*, studies finding that a treatment delay of longer than 6 d[38] or 1 wk[39] was associated with increased mortality).

CONCLUSION

Isolated cerebral mucormycosis is rare and diagnosis is challenging and requires a high degree of clinical suspicion. When a patient presents with headache, fever, hemiplegia, and altered mental status, his/her history and physical examination results should be closely followed by an investigation of the possibility of fungal infection. Histopathology can reveal the pathological characteristics of Mucorales fungi, and molecular methods (MALDI-TOF testing of serum and cryosections) may help to confirm the diagnosis and identification of pathogenic species. Early diagnosis and initiation of antifungal therapy and surgery can improve patient survival.

FOOTNOTES

Author contributions: Chen CH contributed to data collection and manuscript writing; Chen JN treated the patient and contributed to the conceptualization and supervision of the entire work; all authors have read and approved the final manuscript.

Informed consent statement: Written informed consent was obtained from the patient for publication of this case report.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

CARE Checklist (2016) statement: All authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Cai-Hong Chen 0000-0002-9270-144X; Jing-Nan Chen 0000-0002-1555-1601.

S-Editor: Fan JR

L-Editor: Wang TQ

P-Editor: Fan JR

REFERENCES

- Kennedy KJ, Daveson K, Slavin MA, van Hal SJ, Sorrell TC, Lee A, Marriott DJ, Chapman B, Halliday CL, Hajkowicz K, Athan E, Bak N, Cheong E, Heath CH, Morrissey CO, Kidd S, Beresford R, Blyth C, Korman TM, Robinson JO, Meyer W, Chen SC; Australia and New Zealand Mycoses Interest Group of the Australasian Society for Infectious Diseases. Mucormycosis in Australia: contemporary epidemiology and outcomes. *Clin Microbiol Infect* 2016; **22**: 775-781 [PMID: 26806139 DOI: 10.1016/j.cmi.2016.01.005]
- Roden MM, Zaoutis TE, Buchanan WL, Knudsen TA, Sarkisova TA, Schaufele RL, Sein M, Sein T, Chiou CC, Chu JH, Kontoyiannis DP, Walsh TJ. Epidemiology and outcome of zygomycosis: a review of 929 reported cases. *Clin Infect Dis* 2005; **41**: 634-653 [PMID: 16080086 DOI: 10.1086/432579]
- Kontoyiannis DP, Azie N, Franks B, Horn DL. Prospective antifungal therapy (PATH) alliance® : focus on mucormycosis. *Mycoses* 2014; **57**: 240-246 [PMID: 24147728 DOI: 10.1111/myc.12149]
- Skiada A, Pagano L, Groll A, Zimmerli S, Dupont B, Lagrou K, Lass-Flörl C, Bouza E, Klimko N, Gaustad P, Richardson M, Hamal P, Akova M, Meis JF, Rodriguez-Tudela JL, Roilides E, Mitrousia-Ziouva A, Petrikos G; European Confederation of Medical Mycology Working Group on Zygomycosis. Zygomycosis in Europe: analysis of 230 cases accrued by the registry of the European Confederation of Medical Mycology (ECMM) Working Group on Zygomycosis between 2005 and 2007. *Clin Microbiol Infect* 2011; **17**: 1859-1867 [PMID: 21199154 DOI: 10.1111/j.1469-0691.2010.03456.x]
- Ermak D, Kanekar S, Specht CS, Wojnar M, Lowden M. Looks like a stroke, acts like a stroke, but it's more than a stroke: a case of cerebral mucormycosis. *J Stroke Cerebrovasc Dis* 2014; **23**: e403-e404 [PMID: 25106832 DOI: 10.1016/j.jstrokecerebrovasdis.2014.02.024]
- Salehi M, Mahmoudi S, Reza Hosseini O, Hashemi SJ, Ahmadi K, Aala F, Khajavirad N, Alijani N, Izadi A, Getso MI, Abdollahi A, Salami A, Khatami SR, Adibimehr A, Hedayat Yaghoobi M, Sabahi M, Pazooki B, Yazdi F, Zebardast J, Saifi A, Hasan Nezhad M, Mardani M, Khodavaisy S. The Epidemiological, Clinical, Mycological, and Pathological Features of Rhino-cerebral Mucormycosis: A Systematic Review. *Iran J Pathol* 2022; **17**: 112-121 [PMID: 35463720 DOI: 10.30699/IJP.2022.538690.2721]
- Jeong W, Keighley C, Wolfe R, Lee WL, Slavin MA, Kong DCM, Chen SC. The epidemiology and clinical manifestations of mucormycosis: a systematic review and meta-analysis of case reports. *Clin Microbiol Infect* 2019; **25**: 26-34 [PMID: 30036666 DOI: 10.1016/j.cmi.2018.07.011]
- Dong N, Jordan AE, Shen X, Wu X, Guo X, Zhao H, Wang Y, Wang D, Fang Q. Rhino-Orbital Cerebral Mucormycosis in a Patient With Diabetic Ketoacidosis: A Case Report and Literature Review. *Front Neurol* 2022; **13**: 815902 [PMID: 35599740 DOI: 10.3389/fneur.2022.815902]
- Walsh TJ, Gamaletsou MN, McGinnis MR, Hayden RT, Kontoyiannis DP. Early clinical and laboratory diagnosis of invasive pulmonary, extrapulmonary, and disseminated mucormycosis (zygomycosis). *Clin Infect Dis* 2012; **54** Suppl 1: S55-S60 [PMID: 22247446 DOI: 10.1093/cid/cir868]
- Kursun E, Turunc T, Demiroglu YZ, Alişkan HE, Arslan AH. Evaluation of 28 cases of mucormycosis. *Mycoses* 2015; **58**: 82-87 [PMID: 25590855 DOI: 10.1111/myc.12278]
- Chikley A, Ben-Ami R, Kontoyiannis DP. Mucormycosis of the Central Nervous System. *J Fungi (Basel)* 2019; **5** [PMID: 31288475 DOI: 10.3390/jof5030059]
- Zhang GJ, Zhang SK, Wang Z, Zhu YX, Kong J, Huang LL, Guo YJ, Wang YJ, Zou RC, Xie CM. Fatal and Rapid Progressive Isolated Cerebral Mucormycosis Involving the Bilateral Basal Ganglia: A Case Report. *Front Neurol* 2020; **11**: 295 [PMID: 32373057 DOI: 10.3389/fneur.2020.00295]
- Meyerowitz EA, Sanchez S, Mansour MK, Triant VA, Goldberg MB. Isolated Cerebral Mucormycosis in Immunocompetent Adults who Inject Drugs: Case Reports and Systematic Review of the Literature. *Open Forum Infect Dis* 2020; **7**: ofaa552 [PMID: 33409328 DOI: 10.1093/ofid/ofaa552]
- Mamali V, Koutserimpas C, Zarkotou O, Vrioni G, Samonis G. Isolated Cerebral Mucormycosis Caused by Lichtheimia Species in a Polytrauma Patient. *Diagnostics (Basel)* 2022; **12** [PMID: 35204449 DOI: 10.3390/diagnostics12020358]
- Ha NT, Lowery M, Woo J, Mehta Y, Bhanot N. Brain lesion in a recreational drug user: Isolated cerebral mucormycosis. *IDCases* 2020; **22**: e00979 [PMID: 33083231 DOI: 10.1016/j.idcr.2020.e00979]
- Kerezoudis P, Watts CR, Bydon M, Dababneh AS, Deyo CN, Frye JM, Kelley PC, Kemp AM, Palraj BV, Pupillo GT. Diagnosis and Treatment of Isolated Cerebral Mucormycosis: Patient-Level Data Meta-Analysis and Mayo Clinic Experience. *World Neurosurg* 2019; **123**: 425-434.e5 [PMID: 30415043 DOI: 10.1016/j.wneu.2018.10.218]
- Verma A, Brozman B, Petit CK. Isolated cerebral mucormycosis: report of a case and review of the literature. *J Neurol Sci* 2006; **240**: 65-69 [PMID: 16269155 DOI: 10.1016/j.jns.2005.09.010]
- Blitzer A, Lawson W. Fungal infections of the nose and paranasal sinuses. Part I. *Otolaryngol Clin North Am* 1993; **26**: 1007-1035 [PMID: 8290280]
- Farid S, AbuSaleh O, Liesman R, Sohail MR. Isolated cerebral mucormycosis caused by Rhizomucor pusillus. *BMJ Case Rep* 2017; **2017** [PMID: 28978601 DOI: 10.1136/bcr-2017-221473]
- Sangoi AR, Rogers WM, Longacre TA, Montoya JG, Baron EJ, Banaei N. Challenges and pitfalls of morphologic identification of fungal infections in histologic and cytologic specimens: a ten-year retrospective review at a single institution. *Am J Clin Pathol* 2009; **131**: 364-375 [PMID: 19228642 DOI: 10.1309/AJCP99OOOZSNISCZ]
- Guarner J, Brandt ME. Histopathologic diagnosis of fungal infections in the 21st century. *Clin Microbiol Rev* 2011; **24**: 247-280 [PMID: 21482725 DOI: 10.1128/CMR.00053-10]
- Tissot F, Agrawal S, Pagano L, Petrikos G, Groll AH, Skiada A, Lass-Flörl C, Calandra T, Viscoli C, Herbrecht R. ECIL-6 guidelines for the treatment of invasive candidiasis, aspergillosis and mucormycosis in leukemia and hematopoietic stem cell transplant patients. *Haematologica* 2017; **102**: 433-444 [PMID: 28011902 DOI: 10.3324/haematol.2016.152900]
- Yang M, Lee JH, Kim YK, Ki CS, Huh HJ, Lee NY. Identification of mucorales from clinical specimens: a 4-year experience in a single institution. *Ann Lab Med* 2016; **36**: 60-63 [PMID: 26522761 DOI: 10.3343/alm.2016.36.1.60]

- 24 **Dadwal SS**, Kontoyiannis DP. Recent advances in the molecular diagnosis of mucormycosis. *Expert Rev Mol Diagn* 2018; **18**: 845-854 [PMID: [30203997](#) DOI: [10.1080/14737159.2018.1522250](#)]
- 25 **Al-Otaibi F**, Albloushi M, Alhindi H, Timms MS. Carotid artery occlusion by rhinoorbitocerebral mucormycosis. *Case Rep Surg* 2012; **2012**: 812420 [PMID: [23198252](#) DOI: [10.1155/2012/812420](#)]
- 26 **Fu KA**, Nguyen PL, Sanossian N. Basilar artery territory stroke secondary to invasive fungal sphenoid sinusitis: a case report and review of the literature. *Case Rep Neurol* 2015; **7**: 51-58 [PMID: [25873889](#) DOI: [10.1159/000380761](#)]
- 27 **Lanternier F**, Dannaoui E, Morizot G, Elie C, Garcia-Hermoso D, Huerre M, Bitar D, Dromer F, Lortholary O; French Mycosis Study Group. A global analysis of mucormycosis in France: the RetroZygo Study (2005-2007). *Clin Infect Dis* 2012; **54** Suppl 1: S35-S43 [PMID: [22247443](#) DOI: [10.1093/cid/cir880](#)]
- 28 **Cornely OA**, Arikian-Akdagli S, Dannaoui E, Groll AH, Lagrou K, Chakrabarti A, Lanternier F, Pagano L, Skiada A, Akova M, Arendrup MC, Boekhout T, Chowdhary A, Cuenca-Estrella M, Freiburger T, Guinea J, Guarro J, de Hoog S, Hope W, Johnson E, Kathuria S, Lackner M, Lass-Flörl C, Lortholary O, Meis JF, Meletiadis J, Muñoz P, Richardson M, Roilides E, Tortorano AM, Ullmann AJ, van Diepeningen A, Verweij P, Petrikos G; European Society of Clinical Microbiology and Infectious Diseases Fungal Infection Study Group; European Confederation of Medical Mycology. ESCMID and ECMM joint clinical guidelines for the diagnosis and management of mucormycosis 2013. *Clin Microbiol Infect* 2014; **20** Suppl 3: 5-26 [PMID: [24479848](#) DOI: [10.1111/1469-0691.12371](#)]
- 29 **Biswas D**, Kotwal A, Kakati B, Ahmad S. Amphotericin B Resistant *Apophysomyces elegans* Causing Rhino-oculo-Cerebral Mucormycosis in an Immunocompetent Host. *J Clin Diagn Res* 2015; **9**: DD01-DD02 [PMID: [26435947](#) DOI: [10.7860/JCDR/2015/13929.6272](#)]
- 30 **Tarani L**, Costantino F, Notheis G, Wintergerst U, Venditti M, Di Biasi C, Friederici D, Pasquino AM. Long-term posaconazole treatment and follow-up of rhino-orbital-cerebral mucormycosis in a diabetic girl. *Pediatr Diabetes* 2009; **10**: 289-293 [PMID: [18828793](#) DOI: [10.1111/j.1399-5448.2008.00465.x](#)]
- 31 **Anders UM**, Taylor EJ, Martel JR, Martel JB. Acute orbital apex syndrome and rhino-orbito-cerebral mucormycosis. *Int Med Case Rep J* 2015; **8**: 93-96 [PMID: [25945068](#) DOI: [10.2147/IMCRJ.S83036](#)]
- 32 **Garas G**, Choudhury N, Farrell R. Invasive fatal rhino-orbito-cerebral mucormycosis in diabetic ketoacidosis. *JRSM Short Rep* 2010; **1**: 57 [PMID: [21234129](#) DOI: [10.1258/shorts.2010.010089](#)]
- 33 **Angali RK**, Jeshtadi A, Namala VA, Gannepalli A. Fatal rhino-orbito-cerebral mucormycosis in a healthy individual. *J Oral Maxillofac Pathol* 2014; **18**: 460-463 [PMID: [25949007](#) DOI: [10.4103/0973-029X.151355](#)]
- 34 **Chagini Z**, Didehdar M, Khoshbayan A, Rajaei S, Salehi M, Shariati A. Epidemiology, clinical features, diagnosis and treatment of cerebral mucormycosis in diabetic patients: A systematic review of case reports and case series. *Mycoses* 2020; **63**: 1264-1282 [PMID: [32965744](#) DOI: [10.1111/myc.13187](#)]
- 35 **Bavikar P**, Mehta V. Rhino-Orbital-Cerebral Mucormycosis: A Fatal Complication of Uncontrolled Diabetes Mellitus. *Cureus* 2017; **9**: e1841 [PMID: [29348987](#) DOI: [10.7759/cureus.1841](#)]
- 36 **Raffa LH**. Rhino-orbito-cerebral mucormycosis following penetrating keratoplasty. *J Surg Case Rep* 2019; **2019**: rjz314 [PMID: [31832132](#) DOI: [10.1093/jscr/rjz314](#)]
- 37 **Chandra S**, Sharma S, Vats R, Pandey S. Isolated cerebral mucormycosis masquerading as a tumor in an immunocompetent patient. *Autops Case Rep* 2021; **11**: e2020233 [PMID: [33968820](#) DOI: [10.4322/acr.2020.233](#)]
- 38 **Yohai RA**, Bullock JD, Aziz AA, Markert RJ. Survival factors in rhino-orbital-cerebral mucormycosis. *Surv Ophthalmol* 1994; **39**: 3-22 [PMID: [7974189](#) DOI: [10.1016/s0039-6257\(05\)80041-4](#)]
- 39 **Cano P**, Horseman MA, Surani S. Rhinocerebral mucormycosis complicated by bacterial brain abscess. *Am J Med Sci* 2010; **340**: 507-510 [PMID: [20861715](#) DOI: [10.1097/MAJ.0b013e3181f0aad1](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

