

World Journal of *Clinical Cases*

World J Clin Cases 2024 January 26; 12(3): 466-670



Contents

Thrice Monthly Volume 12 Number 3 January 26, 2024

EDITORIAL

- 466 Is medical management useful in Moyamoya disease?
Muengtaweepongsa S, Panpattanakul V
- 474 Metabologenomics and network pharmacology to understand the molecular mechanism of cancer research
Tutar Y

ORIGINAL ARTICLE

Case Control Study

- 479 Significance of oxidative stress and antioxidant capacity tests as biomarkers of premature ovarian insufficiency: A case control study
Kakinuma K, Kakinuma T
- 488 Colorectal resections for malignancy: A pilot study comparing conventional *vs* freehand robot-assisted laparoscopic colectomy
Cawich SO, Plummer JM, Griffith S, Naraynsingh V

Retrospective Study

- 495 Ultrasound diagnosis of congenital Morgagni hernias: Ten years of experience at two Chinese centers
Shi HQ, Chen WJ, Yin Q, Zhang XH

Observational Study

- 503 Genetic investigation of the ubiquitin-protein ligase E3A gene as putative target in Angelman syndrome
Manoubi W, Mahdouani M, Hmida D, Kdissa A, Rouissi A, Turki I, Gueddiche N, Soyah N, Saad A, Bouwkamp C, Elgersma Y, Mougou-Zerelli S, Gribaa M

Prospective Study

- 517 Benefit in physical function and quality of life to nonsurgical treatment of varicose veins: Pilot study
Kim GM, Kim B, Jang M, Park JH, Bae M, Lee CW, Kim JW, Huh U

SYSTEMATIC REVIEWS

- 525 Emerging roles of microRNAs as diagnostics and potential therapeutic interest in type 2 diabetes mellitus
Shrivastav D, Singh DD

META-ANALYSIS

- 538 Impact of body mass index on adverse kidney events in diabetes mellitus patients: A systematic-review and meta-analysis
Wan JF, Chen Y, Yao TH, Wu YZ, Dai HZ

CASE REPORT

- 551 Epithelioid malignant peripheral nerve sheath tumor of the bladder and concomitant urothelial carcinoma: A case report
Ozden SB, Simsekoglu MF, Sertbudak I, Demirdag C, Gurses I
- 560 Simultaneous type III congenital esophageal atresia and patent ductus arteriosus in a low-weight patient: A case report
Ma YY, Chen JR, Yang SW, Wang SY, Cao X, Wu J
- 565 Marginal zone lymphoma with severe rashes: A case report
Bai SJ, Geng Y, Gao YN, Zhang CX, Mi Q, Zhang C, Yang JL, He SJ, Yan ZY, He JX
- 575 Inetetamab combined with pyrotinib and chemotherapy in the treatment of breast cancer brain metastasis: A case report
Dou QQ, Sun TT, Wang GQ, Tong WB
- 582 Adult rhabdomyosarcoma combined with acute myeloid leukemia: A case report
Zheng L, Zhang FJ
- 587 Special electromyographic features in a child with paramyotonia congenita: A case report and review of literature
Yi H, Liu CX, Ye SX, Liu YL
- 596 Removal of a guide-wire sliding into abdominal cavity *via* transgastric natural orifice transluminal endoscopic surgery: A case report
Chen SJ, Zhang DY, Lv YT, Bai FH
- 601 Polyneuropathy, organomegaly, endocrinopathy, M-protein, skin changes syndrome with dilated cardiomyopathy: A case report
Li JR, Feng LY, Li JW, Liao Y, Liu FQ
- 607 Ischemic colitis induced by a platelet-raising capsule: A case report
Wang CL, Si ZK, Liu GH, Chen C, Zhao H, Li L
- 616 Brain abscess from oral microbiota approached by metagenomic next-generation sequencing: A case report and review of literature
Zhu XM, Dong CX, Xie L, Liu HX, Hu HQ
- 623 Carrimycin in the treatment of acute promyelocytic leukemia combined with pulmonary tuberculosis: A case report
Yang FY, Shao L, Su J, Zhang ZM
- 630 Rare esophageal carcinoma-primary adenoid cystic carcinoma of the esophagus: A case report
Geng LD, Li J, Yuan L, Du XB
- 637 Early selective enteral feeding in treatment of acute pancreatitis: A case report
Kashintsev AA, Anisimov SV, Nadeeva A, Proutski V

- 643** Pathological diagnosis and immunohistochemical analysis of giant retrosternal goiter in the elderly: A case report
Meng YC, Wu LS, Li N, Li HW, Zhao J, Yan J, Li XQ, Li P, Wei JQ
- 650** Cerebral syphilitic gumma misdiagnosed as brain abscess: A case report
Mu LK, Cheng LF, Ye J, Zhao MY, Wang JL
- 657** Primary anaplastic lymphoma kinase-positive large B-cell lymphoma of the left bulbar conjunctiva: A case report
Guo XH, Li CB, Cao HH, Yang GY
- 665** Porocarcinoma in a palm reconstructed with a full thickness skin graft: A case report
Lim SB, Kwon KY, Kim H, Lim SY, Koh IC

ABOUT COVER

Peer Reviewer of *World Journal of Clinical Cases*, Kazuhiro Katada, MD, PhD, Assistant Professor, Molecular Gastroenterology and Hepatology, Graduate School of Medical Science, Kyoto Prefectural University of Medicine, Kyoto 6028566, Japan. katada@koto.kpu-m.ac.jp

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Zi-Hang Xu, Production Department Director: Xiang Li, Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

January 26, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Removal of a guide-wire sliding into abdominal cavity via transgastric natural orifice transluminal endoscopic surgery: A case report

Shi-Ju Chen, Da-Ya Zhang, Yan-Ting Lv, Fei-Hu Bai

Specialty type: Medicine, research and experimental

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Kumar R, India

Received: October 3, 2023

Peer-review started: October 3, 2023

First decision: December 6, 2023

Revised: December 16, 2023

Accepted: January 4, 2024

Article in press: January 4, 2024

Published online: January 26, 2024



Shi-Ju Chen, Da-Ya Zhang, Yan-Ting Lv, Graduate School, Hainan Medical University, Haikou 571199, Hainan Province, China

Fei-Hu Bai, Department of Gastroenterology, The Second Affiliated Hospital of Hainan Medical University, 570216 Haikou, Hainan Province, China

Fei-Hu Bai, Department of Gastroenterology, Clinical Medical Center of Hainan Province, Haikou 570216, Hainan Province, China

Corresponding author: Fei-Hu Bai, Doctor, Chief Physician, Department of Gastroenterology, The Second Affiliated Hospital of Hainan Medical University, Yehai Avenue, No. 368 Longhua District, 570216 Haikou, Hainan Province, China. baifeihu_hy@163.com

Abstract

BACKGROUND

Guidewire slippage into the peritoneal cavity during clinical operations is extremely rare. Therefore, this paper aims to report a successful case of guidewire removal using transgastric natural orifice transluminal endoscopic surgery (NOTES). The goal is to enhance physicians' understanding of the management plan for this unique scenario and provide a valuable reference for clinical practice.

CASE SUMMARY

A 64-year-old man presented with abdominal distension and was diagnosed with cirrhosis combined with massive ascites. To proceed with treatment, the patient underwent ultrasound-guided peritoneal puncture and underwent catheterization and drainage. Unfortunately, a 0.035-inch guidewire slipped into the abdominal cavity during the procedure. Following a comprehensive evaluation and consultation by a multidisciplinary team, the guidewire was successfully removed using NOTES.

CONCLUSION

This case highlights the potential consideration of transgastric NOTES removal when encountering a foreign body, such as a guidewire, within the abdominal cavity.

Key Words: Guidewire; Abdominal cavity; Natural orifice transluminal endoscopic

surgery; Cirrhosis; Massive ascites; Case report

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Guidewires play a crucial role in facilitating various medical procedures. Whereas previous reports primarily focused on guidewire breakage and subsequent entrapment within peripheral blood vessels during interventional procedures, this case highlights a unique incident where a guidewire slipped into the abdominal cavity. After extensive multidisciplinary consultations, which took into account the patient's age, frail physical condition, and fair coagulation function, the abdominal guidewire was successfully removed using natural orifice transluminal endoscopic surgery in accordance with individual wishes of the patient. Consequently, the patient experienced a smooth recovery without fever or abdominal pain. This comprehensive article meticulously delineates the management of this exceptional case, which presents an inevitable scenario in clinical practice, ultimately offering valuable insights for healthcare practitioners.

Citation: Chen SJ, Zhang DY, Lv YT, Bai FH. Removal of a guide-wire sliding into abdominal cavity *via* transgastric natural orifice transluminal endoscopic surgery: A case report. *World J Clin Cases* 2024; 12(3): 596-600

URL: <https://www.wjgnet.com/2307-8960/full/v12/i3/596.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v12.i3.596>

INTRODUCTION

Guidewires are pivotal in various clinical procedures, encompassing guidance, support, and exchange. While the retention of guidewires within the body is typically deemed uncommon, recent estimates suggest that, on average, two instances of guidewire retention during central venous catheter insertion arise each month[1]. Nevertheless, only one case involving the retention of guidewires within the abdominal cavity has been previously documented[2]. This article presents the pioneering retrieval of abdominal guidewires utilizing a gastrointestinal natural-tunnel endoscopic procedure (natural orifice transluminal endoscopic surgery, NOTES), thus providing clinicians with a valuable point of reference.

CASE PRESENTATION

Chief complaints

A 64-year-old male patient reported that the guide wire had slipped into the abdominal cavity for 2 d.

History of present illness

The patient, a 64-year-old male, was admitted to the hospital due to abdominal distension. A diagnosis of cirrhosis with extensive ascites was established. To proceed with treatment, the patient underwent ultrasound-guided peritoneal puncture and underwent catheterization and drainage. Unfortunately, a 0.035-inch guidewire inadvertently entered the patient's abdominal cavity during the procedure. Notably, the patient did not experience significant discomfort at the time, and following effective communication, he exhibited understanding and willingness to undergo surgery for the removal of the foreign object.

History of past illness

The patient had been diagnosed with viral hepatitis B for three years and had been consistently receiving entecavir antiviral therapy. Additionally, the patient had been diagnosed with diabetes mellitus for the same duration; however, there had been no regular adherence to medication or monitoring of blood glucose levels, thus rendering the control of blood glucose unknown.

Personal and family history

The patient's personal and familial medical history did not reveal any notable or significant findings.

Physical examination

The patient exhibited normal vital signs, while the abdominal examination revealed distension and a slight sensation of pressure in the upper right quadrant without any indication of rebound pain.

Laboratory examinations

The blood tests revealed the following results: white blood cell count of 9.12×10^9 , platelet count of 91×10^9 , and hemoglobin level of 84 g/L. In terms of liver function, alanine aminotransferase measured 46 U/L, aspartate aminotrans-



DOI: 10.12998/wjcc.v12.i3.596 Copyright ©The Author(s) 2024.

Figure 1 The abdominal X-ray plain film revealed the presence of a distinctive, well-defined elongated shadow in the right middle and lower abdomen, extending to the L2 vertebrae superiorly, as well as the suprapubic branch inferiorly. Additionally, the film showed the presence of visible tubes within the pelvic region.

ferase measured 43 U/L, total bilirubin measured 67.3 $\mu\text{mol/L}$, and prothrombin time registered at 15.6 s. Renal function and electrolyte tests demonstrated normal results.

Imaging examinations

The abdominal X-ray examination revealed the presence of elongated dense shadows, prominently visible in the middle and lower right abdomen, extending into the pelvic region ([Figure 1](#)).

FINAL DIAGNOSIS

The object found within the abdominal cavity was identified as a guidewire, which was presumed to have inadvertently entered during peritoneal puncture placement.

TREATMENT


Considering the patient's advanced age and deteriorating health condition, the decision was made to remove the guidewire through transgastric NOTES. The procedure was conducted under anesthesia and involved the following steps ([Figure 2](#) and [Video](#)): (1) An incision was created in the anterior wall of the gastric antrum using a DualKnife and IT-knife; (2) Exploration of the abdominal cavity allowed for the identification and extraction of the guidewire with the assistance of foreign body forceps through the gastric wall incision; and (3) The wound was treated using thermocoagulation forceps, and the gastric wall incision was subsequently closed with endoclips.

OUTCOME AND FOLLOW-UP

Following the procedure, the patient did not experience any episodes of abdominal pain, fever, or other discomfort. A subsequent abdomen X-ray displayed no dense shadows within the abdominal cavity ([Figure 3](#)). The patient was discharged after one-week post-procedure, and during the one-year follow-up period, the patient remained in good health with no complaints of recurrent abdominal discomfort.


DISCUSSION

The concept of endoscopic laparotomy utilizing the natural lumen was initially introduced by Kalloo *et al*[3] in the United States in 1998. Subsequently, in 2004, successful transgastric endoscopic cholecystectomy procedures were performed on pigs. In 2005, the American Society for Gastrointestinal Endoscopy and the Society of American Gastrointestinal



DOI: 10.12998/wjcc.v12.i3.596 Copyright ©The Author(s) 2024.

Figure 2 The sequential steps of the endoscopic treatment procedures. A: Incision of the gastric wall; B: The abdominal cavity was explored; C: Discovery of the guidewire; D: Removal of the guidewire.



DOI: 10.12998/wjcc.v12.i3.596 Copyright ©The Author(s) 2024.

Figure 3 The abdominal stereotaxic plain film revealed a prominent, irregularly shaped hyperdense shadow within the abdominal cavity, accompanied by minimal intestinal gas.

Endoscopic Surgeons collaborated to propose the notion of NOTES, which has gained widespread adoption since its inception[4]. NOTES represents an evolving minimally invasive technique aimed at mitigating abdominal trauma and its associated complications[5]. When compared to conventional and laparoscopic surgeries, NOTES offers several advantages, including minimal scarring, reduced trauma, diminished pain, shorter surgical pathways, and accelerated healing[4,6].

With the advancement of various interventional therapies, adverse events caused by auxiliary devices like guide wires, guide catheters, and PEG tubes have become increasingly evident. The safe and swift removal of foreign bodies assumes particular significance. Previous reports of guidewire retention predominantly arise from instances where a broken guidewire inadvertently slipped into the bloodstream during procedures such as coronary angiography[1,7]. Unlike those reported cases, the guidewire in the present study was inadvertently left within the abdominal cavity, with only one similar case reported in the literature[2]. In that previous case, the guidewire was retrieved using ultrasound-guided gastroscopic foreign body forceps *via* a transcutaneous route, which proved to be relatively challenging and unsuitable for patients with extensive ascites or complex guidewire locations. Given the patient's advanced age and weakened state, traditional approaches involving laparoscopy or open surgery were deemed inappropriate. Taking into account the patient's favorable coagulation function and platelet count, the multidisciplinary team ultimately decided on the transgastric NOTES approach. This well-established surgical method may serve as a valuable reference for clinical practice.

CONCLUSION

Guidewire slippage into the abdominal cavity is a rare occurrence in clinical practice. When faced with such circumstances, the safe and efficient extraction of the guidewire becomes a pertinent consideration. This study advocates for the utilization of the transgastric NOTES technique as an advantageous approach for guidewire removal from the abdominal cavity.

FOOTNOTES

Author contributions: Chen SJ, Zhang DY and Lv YT participated in the research design; Chen SJ, Zhang DY, Lv YT and Bai FH participated in the performance of the research and in the data analysis; Zhang DY and Lv YT participated in the writing of the article; Chen SJ, Zhang DY and Lv YT contributed equally to this work.

Supported by the Specific Research Fund of The Innovation Platform for Academicians of Hainan Province, No. YSPTZX202313; Hainan Province Clinical Medical Center, No. 2021818; Hainan Provincial Health Industry Research Project, No. 22A200078; and Hainan Provincial Postgraduate Innovation Research Project, No. Qhyb2022-133.

Informed consent statement: All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

Conflict-of-interest statement: All authors disclosed no financial relationships.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Da-Ya Zhang 0000-0001-6133-8919; Fei-Hu Bai 0000-0002-1560-6131.

S-Editor: Gao CC

L-Editor: A

P-Editor: Chen YX

REFERENCES

- 1 **Mariyaselvam MZA**, Patel V, Young HE, Blunt MC, Young PJ. Central Venous Catheter Guidewire Retention: Lessons From England's Never Event Database. *J Patient Saf* 2022; **18**: e387-e392 [PMID: 33512864 DOI: 10.1097/PTS.0000000000000826]
- 2 **Zhu Z**, Zhu J, Qiao Z. Ultrasound guidance for the removal of guide-wire sliding into abdominal cavity. *Asian J Surg* 2022; **45**: 1484-1486 [PMID: 35283026 DOI: 10.1016/j.asjsur.2022.02.059]
- 3 **Kalloor AN**, Singh VK, Jagannath SB, Niiyama H, Hill SL, Vaughn CA, Magee CA, Kantsevov SV. Flexible transgastric peritoneoscopy: a novel approach to diagnostic and therapeutic interventions in the peritoneal cavity. *Gastrointest Endosc* 2004; **60**: 114-117 [PMID: 15229442 DOI: 10.1016/s0016-5107(04)01309-4]
- 4 **Rattner D**, Kalloor A; ASGE/SAGES Working Group. ASGE/SAGES Working Group on Natural Orifice Transluminal Endoscopic Surgery. October 2005. *Surg Endosc* 2006; **20**: 329-333 [PMID: 16402290 DOI: 10.1007/s00464-005-3006-0]
- 5 **Liu BR**, Cui GX, Zhang XY, Kaur K, Song JT, Si Y, Kong XC. Pure transgastric NOTES ovarian cystectomy: the first human procedure. *Endoscopy* 2014; **46** Suppl 1 UCTN: E199-E200 [PMID: 24756294 DOI: 10.1055/s-0034-1365290]
- 6 **Wang S**, Guan F, Wang F, Wen P. Transgastric natural orifice transluminal endoscopic surgery (NOTES) to remove a foreign body from the abdominal cavity of a 12-year old boy. *Endoscopy* 2022; **54**: E110-E112 [PMID: 33784757 DOI: 10.1055/a-1407-9775]
- 7 **Xue Y**, Dai L, Jiang W, Zhang H. Off-pump coronary artery bypass concomitant with retrieval of broken guide wire stuck in the brachial artery: a case report. *BMC Cardiovasc Disord* 2021; **21**: 46 [PMID: 33482724 DOI: 10.1186/s12872-021-01867-0]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

