

World Journal of *Clinical Cases*

World J Clin Cases 2024 January 26; 12(3): 466-670



Contents

Thrice Monthly Volume 12 Number 3 January 26, 2024

EDITORIAL

- 466 Is medical management useful in Moyamoya disease?
Muengtaweepongsa S, Panpattanakul V
- 474 Metabologenomics and network pharmacology to understand the molecular mechanism of cancer research
Tutar Y

ORIGINAL ARTICLE

Case Control Study

- 479 Significance of oxidative stress and antioxidant capacity tests as biomarkers of premature ovarian insufficiency: A case control study
Kakinuma K, Kakinuma T
- 488 Colorectal resections for malignancy: A pilot study comparing conventional *vs* freehand robot-assisted laparoscopic colectomy
Cawich SO, Plummer JM, Griffith S, Naraynsingh V

Retrospective Study

- 495 Ultrasound diagnosis of congenital Morgagni hernias: Ten years of experience at two Chinese centers
Shi HQ, Chen WJ, Yin Q, Zhang XH

Observational Study

- 503 Genetic investigation of the ubiquitin-protein ligase E3A gene as putative target in Angelman syndrome
Manoubi W, Mahdouani M, Hmida D, Kdissa A, Rouissi A, Turki I, Gueddiche N, Soyah N, Saad A, Bouwkamp C, Elgersma Y, Mougou-Zerelli S, Gribaa M

Prospective Study

- 517 Benefit in physical function and quality of life to nonsurgical treatment of varicose veins: Pilot study
Kim GM, Kim B, Jang M, Park JH, Bae M, Lee CW, Kim JW, Huh U

SYSTEMATIC REVIEWS

- 525 Emerging roles of microRNAs as diagnostics and potential therapeutic interest in type 2 diabetes mellitus
Shrivastav D, Singh DD

META-ANALYSIS

- 538 Impact of body mass index on adverse kidney events in diabetes mellitus patients: A systematic-review and meta-analysis
Wan JF, Chen Y, Yao TH, Wu YZ, Dai HZ

CASE REPORT

- 551 Epithelioid malignant peripheral nerve sheath tumor of the bladder and concomitant urothelial carcinoma: A case report
Ozden SB, Simsekoglu MF, Sertbudak I, Demirdag C, Gurses I
- 560 Simultaneous type III congenital esophageal atresia and patent ductus arteriosus in a low-weight patient: A case report
Ma YY, Chen JR, Yang SW, Wang SY, Cao X, Wu J
- 565 Marginal zone lymphoma with severe rashes: A case report
Bai SJ, Geng Y, Gao YN, Zhang CX, Mi Q, Zhang C, Yang JL, He SJ, Yan ZY, He JX
- 575 Inetetamab combined with pyrotinib and chemotherapy in the treatment of breast cancer brain metastasis: A case report
Dou QQ, Sun TT, Wang GQ, Tong WB
- 582 Adult rhabdomyosarcoma combined with acute myeloid leukemia: A case report
Zheng L, Zhang FJ
- 587 Special electromyographic features in a child with paramyotonia congenita: A case report and review of literature
Yi H, Liu CX, Ye SX, Liu YL
- 596 Removal of a guide-wire sliding into abdominal cavity *via* transgastric natural orifice transluminal endoscopic surgery: A case report
Chen SJ, Zhang DY, Lv YT, Bai FH
- 601 Polyneuropathy, organomegaly, endocrinopathy, M-protein, skin changes syndrome with dilated cardiomyopathy: A case report
Li JR, Feng LY, Li JW, Liao Y, Liu FQ
- 607 Ischemic colitis induced by a platelet-raising capsule: A case report
Wang CL, Si ZK, Liu GH, Chen C, Zhao H, Li L
- 616 Brain abscess from oral microbiota approached by metagenomic next-generation sequencing: A case report and review of literature
Zhu XM, Dong CX, Xie L, Liu HX, Hu HQ
- 623 Carrimycin in the treatment of acute promyelocytic leukemia combined with pulmonary tuberculosis: A case report
Yang FY, Shao L, Su J, Zhang ZM
- 630 Rare esophageal carcinoma-primary adenoid cystic carcinoma of the esophagus: A case report
Geng LD, Li J, Yuan L, Du XB
- 637 Early selective enteral feeding in treatment of acute pancreatitis: A case report
Kashintsev AA, Anisimov SV, Nadeeva A, Proutski V

- 643** Pathological diagnosis and immunohistochemical analysis of giant retrosternal goiter in the elderly: A case report
Meng YC, Wu LS, Li N, Li HW, Zhao J, Yan J, Li XQ, Li P, Wei JQ
- 650** Cerebral syphilitic gumma misdiagnosed as brain abscess: A case report
Mu LK, Cheng LF, Ye J, Zhao MY, Wang JL
- 657** Primary anaplastic lymphoma kinase-positive large B-cell lymphoma of the left bulbar conjunctiva: A case report
Guo XH, Li CB, Cao HH, Yang GY
- 665** Porocarcinoma in a palm reconstructed with a full thickness skin graft: A case report
Lim SB, Kwon KY, Kim H, Lim SY, Koh IC

ABOUT COVER

Peer Reviewer of *World Journal of Clinical Cases*, Kazuhiro Katada, MD, PhD, Assistant Professor, Molecular Gastroenterology and Hepatology, Graduate School of Medical Science, Kyoto Prefectural University of Medicine, Kyoto 6028566, Japan. katada@koto.kpu-m.ac.jp

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Zi-Hang Xu, Production Department Director: Xiang Li, Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

January 26, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Cerebral syphilitic gumma misdiagnosed as brain abscess: A case report

Li-Kun Mu, Li-Feng Cheng, Jing Ye, Meng-Yan Zhao, Jin-Long Wang

Specialty type: Clinical neurology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C, C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Kaur M, United States

Received: November 15, 2023

Peer-review started: November 15, 2023

First decision: December 15, 2023

Revised: December 22, 2023

Accepted: January 2, 2024

Article in press: January 2, 2024

Published online: January 26, 2024



Li-Kun Mu, Li-Feng Cheng, Jing Ye, Meng-Yan Zhao, Jin-Long Wang, Department of Neurosurgery, Shengli Oilfield Central Hospital, Dongying 257099, Shandong Province, China

Corresponding author: Li-Kun Mu, MMed, Neurosurgeon, Department of Neurosurgery, Shengli Oilfield Central Hospital, No. 31 Jinan Road, Dongying 257099, Shandong Province, China. sjwmlk@163.com

Abstract

BACKGROUND

Cerebral syphilitic gumma is a relatively rare clinical disease. Its clinical manifestations are non-specific, and the imaging manifestations are similar to other intracranial occupying lesions, often misdiagnosed as tumors or abscesses. There are few reports on this disease in the relevant literature. To our knowledge, we have reported the first case of cerebral syphilitic gumma misdiagnosed as a brain abscess. We report this case and provide useful information for clinical doctors on neurosyphilis diseases.

CASE SUMMARY

We report the case to explore the diagnostic essentials of cerebral syphilitic gumma and attempt to mitigate the rates of misdiagnosis and missed diagnosis by equipping physicians with knowledge of neurosyphilis characteristics. The clinical diagnosis and treatment of a patient with cerebral syphilitic gumma were reported. Clinical manifestations, classifications, and diagnostic points were retrospectively analyzed. The patient was admitted to the hospital with fever and limb weakness. Brain magnetic resonance imaging showed multiple space-occupying lesions and a positive serum *Treponema pallidum* gelatin agglutination test. The patient was misdiagnosed as having a brain abscess and underwent a craniotomy. A postoperative pathological diagnosis of syphilis gumma was made. The patient improved and was discharged after penicillin anti-syphilis treatment. Follow-up recovery was satisfactory.

CONCLUSION

Cerebral syphilitic gumma is rare in clinical practice, and it is often misdiagnosed and missed. Clinical diagnosis should be considered in combination with multiple examinations.

Key Words: Neurosyphilis; Cerebral syphilitic gumma; Brain abscess; Misdiagnosis; Treatment; Case report

Core Tip: Cerebral syphilitic gumma is a relatively rare clinical disease. Its clinical manifestations are non-specific, and the imaging manifestations are similar to other intracranial occupying lesions, often misdiagnosed as tumors or abscesses. There are few reports on this disease in the relevant literature. To our knowledge, we have reported the first case of cerebral syphilitic gumma misdiagnosed as a brain abscess. We report this case and provide useful information for clinical doctors on neurosyphilis diseases.

Citation: Mu LK, Cheng LF, Ye J, Zhao MY, Wang JL. Cerebral syphilitic gumma misdiagnosed as brain abscess: A case report. *World J Clin Cases* 2024; 12(3): 650-656

URL: <https://www.wjgnet.com/2307-8960/full/v12/i3/650.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v12.i3.650>

INTRODUCTION

Cerebral syphilitic gumma is a late form of syphilis caused by *Treponema pallidum* (*T. pallidum*), which invades the central nervous system and often develops 3-20 years after syphilis infection. Cerebral syphilis gummata is a rare type of neurosyphilis with an incidence of only 2%, and the affected population is mainly middle-aged and elderly men who are mostly newly diagnosed with intracranial masses[1]. With an insidious onset, nonspecific clinical manifestations, and the imaging findings resembling other intracranial space-occupying ring enhancements, the condition poses challenges in differential diagnosis, making it susceptible to misdiagnosis and mistreatment. Surgical treatment, often mistaken for a tumor or abscess, necessitates pathological examination for accurate confirmation of the diagnosis[2]. The clinical data of a patient with cerebral syphilis gummata admitted to the Department of Neurosurgery at Shengli Oilfield Central Hospital are reported below to further deepen our understanding of this disease.

CASE PRESENTATION

Chief complaints

A 55-year-old female patient was admitted to the hospital with “intermittent fever with weakness of right limbs for half a month” on June 28, 2019.

History of present illness

Half a month ago, the patient experienced right limb weakness without obvious cause, coinciding with two episodes of low-grade fever with diarrhea. After oral administration of cephalosporins at a local clinic (details unknown), the patient exhibited marked improvement in diarrhea and return to normal body temperature. However, the right-sided weakness persisted, and there the patient tended to lean towards the right rear while standing. Subsequently, she was referred to our neurology clinic, and brain magnetic resonance imaging (MRI) showed multiple intracranial space-occupying lesions with surrounding edema. Considering the possibility of a brain abscess, metastases were not excluded (Figure 1). On the same day, the patient was admitted to our department with a diagnosis of brain abscess from the neurology clinic. Since the onset of disease, the patient had paroxysmal headache, dizziness, and blurred vision, without limb twitching or numbness.

History of past illness

The patient had a history of “liver abscess” for half a year and underwent puncture drainage of liver abscess and anti-infection treatment in the Department of Hepatobiliary Surgery of our hospital.

Personal and family history


The pus culture from the puncture revealed *Klebsiella pneumoniae*, and the remaining medical history was unremarkable.

Physical examination

Physical examination on admission revealed a temperature of 36.9 °C; heart rate, 86 beats/min; respiration, 18 breaths/min; and blood pressure, 131/75 mmHg, and physical examination of the heart, lungs, and abdomen showed no significant abnormalities.

Laboratory examinations

Laboratory tests indicated a white blood cell (WBC) count of $10.9 \times 10^9/L$ ($3.5-9.5 \times 10^9/L$), neutrophil percentage of



DOI: 10.12998/wjcc.v12.i3.650 Copyright ©The Author(s) 2024.

Figure 1 Magnetic resonance imaging images captured on June 28, 2019. A: Enhanced magnetic resonance imaging image; B: T1 image; C: T2 image; D: FLAIR image; E: Diffusion-weighted imaging image.


82.4% (40%-75%), and direct neutrophil count or $9 \times 10^9/L$ ($1.8-6.3 \times 10^9/L$). Other blood biochemical parameters showed no significant abnormalities.

Imaging examinations

Neurological examination revealed clear consciousness, clear and fluent speech, normal orientation, intact calculation, and memory functions. Pupils were equal and round bilaterally, responding well to light sensitivity. Fundus examination identified bilateral papilledema. Eye movements were well-coordinated, and there were no abnormalities in the examination of other cranial nerves. Muscle strength was graded as V in the left limb and IV in the right limb, with normal muscle tone. Tendon reflexes were brisk (++), while pathologic reflexes were absent (-). Sensory and synkinesis examinations showed no abnormalities, and there was a normal sense of resistance in the neck.


MULTIDISCIPLINARY EXPERT CONSULTATION

Preliminary diagnosis included brain abscess and post-puncture drainage of the liver abscess. Abdominal ultrasound after admission showed a 14 mm × 9 mm heterogeneous echogenic area in the right liver; however, positron emission tomography and computed tomography revealed no evidence of malignancy. Considering the patient's medical history, a hematogenous brain abscess was suspected. The initial approach involved dehydration to reduce intracranial pressure, administration of meropenem for anti-infection, and fluid replacement. Lumbar puncture (July 1, 2019) revealed a cerebrospinal fluid pressure of 280 mm H₂O (1 mm H₂O = 0.0098 kPa). Routine examination indicated a WBC count of $49 \times 10^9/L$, while biochemical and cytological examinations showed no significant abnormality. After 1 wk of treatment, the symptoms of headache and blurred vision were slightly improved, but no significant improvement in the weakness of the right limb was noted. A repeated brain MRI (July 8, 2019) revealed multiple intracranial space-occupying lesions, with surrounding edema showing only slight reduction than in the previous examination (Figure 2).



DOI: 10.12998/wjcc.v12.i3.650 Copyright ©The Author(s) 2024.

Figure 2 Intraoperative image of left occipital lobe lesion.



DOI: 10.12998/wjcc.v12.i3.650 Copyright ©The Author(s) 2024.


Figure 3 Pathological section.

FINAL DIAGNOSIS

The patient did not support the diagnosis of brain abscess during treatment; subsequently, a thorough inquiry into the medical history was conducted again. The patient reported a history of unprotected sexual activity following a divorce. A physical examination conducted 1 year prior had revealed a positive *T. pallidum* antibody in the blood, with no subsequent specialized treatment administered. Four preoperative examinations were performed in the hospital, revealing a *T. pallidum* antibody level of 15.77 S/CO and a positive *T. pallidum* particle agglutination test. Syphilis confirmatory tests indicated weak positivity for anti-P47 and anti-P15 antibodies and positivity for anti-P45, anti-P17, and anti-*T. pallidum* antibody IgG. According to the relevant data and the imaging characteristics of the patient, the possibility of cerebral syphilitic gumma could not be ruled out, and pathological biopsy was recommended for some lesions. After obtaining consent from the patient and her family, a complete left occipital craniotomy brain lesion resection was performed on July 11, 2019 (Figure 3). The postoperative pathology was indicative of cerebral syphilitic gumma (Figure 4A).

TREATMENT

On July 12, a dermatology consultation was requested, and the recommended treatment plan included: (1) Conducting serological examinations for syphilis in sexual partners and close contacts; and (2) administering intravenous penicillin at 4 million U ivdrip every 6 h $\times$ 14 d, subsequently switched to intramuscular benzathine penicillin at 2.4 million U every week $\times$ 3 times.



DOI: 10.12998/wjcc.v12.i3.650 Copyright ©The Author(s) 2024.

Figure 4 Magnetic resonance images. A: Enhanced magnetic resonance imaging (MRI) image captured on July 8, 2019; B: Enhanced MRI image captured on July 25, 2019; C: Enhanced MRI image captured on March 1, 2020.

OUTCOME AND FOLLOW-UP

The patient's symptoms resolved after treatment. Repeated brain MRI (July 25, 2019) showed that the multiple intracranial lesions and surrounding edema were significantly reduced compared to the previous imaging (Figure 4B). Brain MRI (March 1, 2020) indicated disappearance of multiple intracranial lesions (Figure 4C).

DISCUSSION

Cerebral syphilitic gumma occurs when *T. pallidum*, originating from a primary infection, spreads throughout the whole body, penetrates the blood-brain barrier, and induces chronic infectious diseases in the central nervous system. Its incidence has increased in recent years[3,4]. Cerebral syphilitic gumma is divided into: (1) Asymptomatic; (2) interstitial syphilis (meningovascular syphilis); (3) parenchymal syphilis (paralytic dementia, spinal phthisis); and (4) gummatous syphilis. Gumma is a late-onset neurosyphilis with a long incubation period that occurs 8-12 years after syphilis infection [5]. Gummatous lesions may appear anywhere in the brain tissue, such as the posterior fossa, pons, midbrain, cerebello-pontine angle, corpus callosum, or cerebral convexity[6]. Imaging examinations showed lesions that were either single or multiple. Computed tomography scans exhibited low or isodense areas, with possible ring enhancement. On MR T1WI, the lesions appeared as low or isointensity, while on T2WI, they showed high signal intensity. Post-enhancement, nodular or ring enhancement was observed. However, these imaging findings were similar to other intracranial space-occupying lesions, such as malignant meningioma, glioma, metastases, tuberculoma, cryptococcal tumor, sarcoma, and brain abscess, which were not characteristic of syphilis gumma, making differentiation challenging[7]. Hence, surgical resection or craniotomy biopsy has been easily misdiagnosed as neoplastic lesions[6]. This patient had multiple intracranial syphilis gumma located in the frontal lobe, occipital lobe, and paraventricular region, showing central necrosis, meningeal tail sign, significant edema around the lesion, and a mild mass effect. These findings are similar to those of brain abscesses and can easily lead to misdiagnosis.

Upon entering the central nervous system, *T. pallidum* adheres to hyaluronidase on the cell membrane of the vessel's inner wall, using its own mucopolysaccharide as a receptor. This interaction leads to the decomposition of the mucopolysaccharide of the vessel, resulting in vascular collapse and the development of obliterative arteritis or periarteritis. Because the body's inflammatory cells infiltrate and develop edema, granulomatous changes occur locally in the damaged area, with caseous necrosis in the center, mostly surrounded by a large area of the edema zone, and the adjacent meninges are often thickened owing to inflammatory reactions. Patients often present with symptoms of increased intracranial pressure, such as headache, vomiting, or blurred vision, which are similar to other lesions of the nervous system and are not characteristic of syphilis gumma; however, a history of unprotected sexual behavior significantly contributes to the diagnostic considerations for this disease and warrants careful attention.

The specimens examined included serum and cerebrospinal fluid, and *T. pallidum* nonspecific antibody tests rapid plasma reagin test (RPR) and toluidine red unheated serum test are commonly used for screening. Specific antibody tests such as the *T. pallidum* particle agglutination assay (TPPA), *T. pallidum* hemagglutination assay, and *T. pallidum* antibody absorption test can be used to confirm the diagnosis of *T. pallidum* infection. Positive specific antibodies indicate syphilis infection but cannot distinguish whether the infection is short-term or long-term, whereas nonspecific antibodies can reflect the activity of syphilis infection, and the combination of the two can further confirm the diagnosis. Patients often exhibit abnormal changes in WBC counts and protein levels in the cerebrospinal fluid. Laboratory tests are crucial for the diagnosis and treatment of this disease; however, these results may be false-positive or false-negative. Therefore, based

on a full understanding of the clinical history, a comprehensive analysis of imaging examinations and laboratory findings can facilitate a preoperative diagnosis of this disease. Nonetheless, postoperative pathological biopsy remains the gold standard for confirming the diagnosis.

After the diagnosis is confirmed, the patient must be actively treated with antibiotics against syphilis. Remission is often associated with the normalization of cerebrospinal fluid cell counts and reduction of protein levels, such as the reduction and conversion of cerebrospinal fluid RPR and TPPA, indicating good treatment results[8]. Postoperative follow-up showed that the main clinical symptoms of this patient significantly improved and disappeared, and re-examination of the brain MRI showed that the number of lesions was significantly reduced, indicating that the therapeutic effect was satisfactory. Currently, penicillin is still the recommended drug of choice for the treatment of neurosyphilis, as it can significantly improve the clinical symptoms of patients and reduce intracranial lesions. According to early, adequate, regular, and whole-course treatment principles, alternative drugs such as ceftriaxone, doxycycline, or erythromycin should be selected for patients allergic to penicillin.

Causes of misdiagnosis may include: (1) The patient's admission with symptoms resembling "intermittent fever", and a history of "liver abscess", often leading to misdiagnosis as bacterial infectious diseases; (2) cerebral syphilitic gumma's rarity in clinical practice, where patients often present to dermatology departments, and neurosurgeons may lack sufficient awareness of this disease; (3) inadequate attention to the patient's sexual history by medical staff and oversight of the syphilis serological examination results upon admission, contributing to misdiagnosis; and (4) the lack of clear specificity in imaging findings for cerebral syphilitic gumma, further complicating the diagnostic process.

CONCLUSION

The clinical manifestations of neurosyphilis are complex and similar to those of many neurological and other systemic diseases. Because patients themselves and doctors often misdiagnose and miss the diagnosis owing to the abovementioned reasons, clinicians should enhance their understanding of neurosyphilis, heighten disease awareness during clinical processes, and consider neurosyphilis when facing unexplained central nervous system infections, cerebrovascular issues, parenchymal damage, or cases with inadequate therapeutic response, especially if there is uncertainty about the underlying cause. Routine syphilis serology and cerebrospinal fluid examinations are crucial in such scenarios to exclude the possibility of neurosyphilis. This proactive approach will aid in avoiding missed diagnoses and misdiagnoses, ensuring prompt identification and treatment of this manageable disease.

ACKNOWLEDGEMENTS

The authors would like to thank the patient for permitting us to use her clinical information in this study.

FOOTNOTES

Author contributions: Mu LK and Cheng LF conceived and designed the study; Mu LK, Cheng LF, and Ye J provided the study materials and collected the cases; Mu LK and Zhao MY wrote the manuscript; Mu LK and Wang JL confirmed the authenticity of the raw data.

Informed consent statement: Informed written consent was obtained from the patients for the publication of this report and any accompanying images.

Conflict-of-interest statement: The authors report having no relevant conflicts of interest for this article.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Li-Kun Mu 0009-0008-9965-9433.

S-Editor: Chen YL

L-Editor: Filipodia

P-Editor: Xu ZH

REFERENCES

- 1 **Li JC**, Mahta A, Kim RY, Saria M, Kesari S. Cerebral syphilitic gumma: a case report and review of the literature. *Neurol Sci* 2012; **33**: 1179-1181 [PMID: [22167653](#) DOI: [10.1007/s10072-011-0878-8](#)]
- 2 **Liu LL**, Zheng WH, Tong ML, Liu GL, Zhang HL, Fu ZG, Lin LR, Yang TC. Ischemic stroke as a primary symptom of neurosyphilis among HIV-negative emergency patients. *J Neurol Sci* 2012; **317**: 35-39 [PMID: [22482824](#) DOI: [10.1016/j.jns.2012.03.003](#)]
- 3 **Karsan N**, Barker R, O'Dwyer JP. Clinical reasoning: the "great imitator". *Neurology* 2014; **83**: e188-e196 [PMID: [25422404](#) DOI: [10.1212/WNL.0000000000001033](#)]
- 4 **Timmermans M**, Carr J. Neurosyphilis in the modern era. *J Neurol Neurosurg Psychiatry* 2004; **75**: 1727-1730 [PMID: [15548491](#) DOI: [10.1136/jnnp.2004.031922](#)]
- 5 **Singh AE**, Romanowski B. Syphilis: review with emphasis on clinical, epidemiologic, and some biologic features. *Clin Microbiol Rev* 1999; **12**: 187-209 [PMID: [10194456](#) DOI: [10.1128/CMR.12.2.187](#)]
- 6 **Darwish BS**, Fowler A, Ong M, Swaminathan A, Abraszko R. Intracranial syphilitic gumma resembling malignant brain tumour. *J Clin Neurosci* 2008; **15**: 308-310 [PMID: [18187326](#) DOI: [10.1016/j.jocn.2006.07.023](#)]
- 7 **Agrons GA**, Han SS, Husson MA, Simeone F. MR imaging of cerebral gumma. *AJNR Am J Neuroradiol* 1991; **12**: 80-81 [PMID: [1899524](#)]
- 8 **Casolla B**, Ristori G, Romano A, Bozzao A, Orzi F. Subacute multicranial neuropathy revealing an early case of meningeal syphilis. *Neurol Sci* 2015; **36**: 1033-1034 [PMID: [25739944](#) DOI: [10.1007/s10072-015-2125-1](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

