

# World Journal of *Clinical Cases*

*World J Clin Cases* 2024 February 16; 12(5): 872-1038



## Contents

Thrice Monthly Volume 12 Number 5 February 16, 2024

## EDITORIAL

- 872 Is it possible to anchor a tooth with photobiomodulation?  
*Dominguez A*
- 875 Strengthening pharmacotherapy research for COVID-19-induced pulmonary fibrosis  
*Liu YM, Zhang J, Wu JJ, Guo WW, Tang FS*

## ORIGINAL ARTICLE

## Clinical and Translational Research

- 880 Causal associations between gastroesophageal reflux disease and essential hypertension: A bidirectional Mendelian randomization study  
*Wei N, Liu MH, Song YH*
- 891 Serum urate is associated with an increased risk of inflammatory bowel disease: A bidirectional Mendelian randomization study  
*Zhang S, Fang X, Kang L, Sui XY, Liu M, Luo YJ, Fu S, Li ZS, Zhao SB, Bai Y*

## Retrospective Cohort Study

- 903 Effect of health education based on information-motivation-behavioral skills model on patients with unilateral vestibular dysfunction  
*Shi Q, Wu RJ, Liu J*

## Retrospective Study

- 913 Removal of intrahepatic bile duct stone could reduce the risk of cholangiocarcinoma: A single-center retrospective study in South Korea  
*Kim TI, Han SY, Lee J, Kim DU*
- 922 Effect of nursing on postoperative respiratory function and mental health of lung cancer patients  
*Yang X, Yin D, Chen SQ*
- 931 Value of glucose transport protein 1 expression in detecting lymph node metastasis in patients with colorectal cancer  
*Kim H, Choi SY, Heo TY, Kim KR, Lee J, Yoo MY, Lee TG, Han JH*

## Observational Study

- 942 Clinical efficacy and mechanism study of mid-frequency anti-snoring device in treating moderate obstructive sleep apnea-hypopnea syndrome  
*Qian B, Chen ZJ, Wang YS, Hu XY, Hu XB, Zheng YH*
- 951 Urinary metabolic profiles during *Helicobacter pylori* eradication in chronic gastritis  
*An WT, Hao YX, Li HX, Wu XK*

- 966 Clinical significance of platelet mononuclear cell aggregates in patients with sepsis and acute respiratory distress syndrome

*Huang CM, Li JJ, Wei WK*

### CASE REPORT

- 973 Left ventricular thrombosis caused cerebral embolism during venoarterial extracorporeal membrane oxygenation support: A case report

*Bai YB, Zhao F, Wu ZH, Shi GN, Jiang N*

- 980 Abnormal uterine bleeding successfully treated *via* ultrasound-guided microwave ablation of uterine myoma lesions: Three case reports

*Kakinuma T, Kakinuma K, Okamoto R, Yanagida K, Ohwada M, Takeshima N*

- 988 Omental fibroma combined with right indirect inguinal hernia masquerades as a scrotal tumor: A case report

*Zhou P, Jin CH, Shi Y, Ma GQ, Wu WH, Wang Y, Cai K, Fan WF, Wang TB*

- 995 Imaging, pathology, and diagnosis of solitary fibrous tumor of the pancreas: A case report and review of literature

*Wang WW, Zhou SP, Wu X, Wang LL, Ruan Y, Lu J, Li HL, Ni XL, Qiu LL, Zhou XH*

- 1004 Neuroimaging features in a patient with non-ketotic hyperglycaemic seizures: A case report

*Wu J, Feng H, Zhao Y, Li J, Li T, Li K*

- 1010 Novel approach of ultrasound-guided lateral recess block for a patient with lateral recess stenosis: A case report

*Yang J, Li XL, Li QB*

- 1018 Ankylosing spondylitis coexisting with *Clonorchis sinensis* infection: A case report

*Yi TX, Liu W, Leng WF, Wang XC, Luo L*

- 1025 Hematuria after nocturnal exercise of a man: A case report

*Bai MJ, Yang ST, Liu XK*

### LETTER TO THE EDITOR

- 1029 Response letter to "Acute cholangitis: Does malignant biliary obstruction *vs* choledocholithiasis etiology change the outcomes?" with imaging aspects

*Aydin S, Irgul B*

- 1033 Exploring multifaceted factors in chronic kidney disease risk: A comprehensive analysis of biochemistry, lifestyle, and inflammation in elderly Chinese individuals

*Cardona F*

- 1036 Transcranial direct current stimulation efficacy in trigeminal neuralgia

*Fasilis T, Gatzonis S, Patrikelis P, Korfiatis S, Alexoudi A*

**ABOUT COVER**

Peer Reviewer of *World Journal of Clinical Cases*, Abdulkarim Hasan, MD, MSc, Consultant Physician-Scientist, Doctor, Lecturer, Department of Pathology, Al-Azhar University, Faculty of Medicine, Cairo 11884, Egypt. doctorabdulkarim7@gmail.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Hua-Ge Yin; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

February 16, 2024

**COPYRIGHT**

© 2024 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Observational Study

## Clinical significance of platelet mononuclear cell aggregates in patients with sepsis and acute respiratory distress syndrome

Chong-Min Huang, Juan-Juan Li, Wei-Ke Wei

**Specialty type:** Medicine, research and experimental**Provenance and peer review:** Unsolicited article; Externally peer reviewed.**Peer-review model:** Single blind**Peer-review report's scientific quality classification**Grade A (Excellent): 0  
Grade B (Very good): B, B  
Grade C (Good): 0  
Grade D (Fair): 0  
Grade E (Poor): 0**P-Reviewer:** Ghimire R, Nepal**Received:** November 16, 2023**Peer-review started:** November 16, 2023**First decision:** December 26, 2023**Revised:** December 30, 2023**Accepted:** January 24, 2024**Article in press:** January 24, 2024**Published online:** February 16, 2024**Chong-Min Huang**, Department of Intensive Medicine, The Second People's Hospital of Haining City, Haining 314419, Zhejiang Province, China**Juan-Juan Li, Wei-Ke Wei**, Department of Integrated Traditional Chinese and Western Medicine, Qingyang Hospital, Hangzhou 310023, Zhejiang Province, China**Corresponding author:** Chong-Min Huang, MD, Doctor, Department of Intensive Medicine, The Second People's Hospital of Haining City, No. 85 Guoque Road, Maqiao Street, Haining 314419, Zhejiang Province, China. [15325307291@163.com](mailto:15325307291@163.com)

## Abstract

## BACKGROUND

The diagnosis of sepsis combined with acute respiratory distress syndrome (ARDS) has increased owing to the enhanced awareness among medical professionals and the continuous development of modern medical technologies, while early diagnosis of ARDS still lacks specific biomarkers. One of the main pathogenic mechanisms of sepsis-associated ARDS involves the actions of various pathological injuries and inflammatory factors, such as platelet and white blood cells activation, leading to an increase of surface adhesion molecules. These adhesion molecules further form platelet-white blood cell aggregates, including platelet-mononuclear cell aggregates (PMAs). PMAs has been identified as one of the markers of platelet activation, here we hypothesize that PMAs might play a potential biomarker for the early diagnosis of this complication.

## AIM

To investigate the expression of PMAs in the serum of patients with sepsis complicated by ARDS and its clinical significance.

## METHODS

We selected 72 hospitalized patients diagnosed with sepsis as the study population between March 2019 and March 2022. Among them, 30 patients with sepsis and ARDS formed the study group, while 42 sepsis patients without ARDS comprised the control group. After diagnosis, venous blood samples were immediately collected from all patients. Flow cytometry was employed to analyze the expression of PMAs, platelet neutrophil aggregates (PNAs), and platelet aggregates (PLyAs) in the serum. Additionally, the Acute Physiology and Chronic Health Evaluation (APACHE) II score was calculated for each patient, and receiver operating characteristic curves were generated to assess diagnostic value.

## RESULTS

The study found that the levels of PNAs and PLyAs in the serum of the study group were higher than those in the control group, but the difference was not statistically significant ( $P > 0.05$ ). However, the expression of PMAs in the serum of the study group was significantly upregulated ( $P < 0.05$ ) and positively correlated with the APACHE II score ( $r = 0.671$ ,  $P < 0.05$ ). When using PMAs as a diagnostic indicator, the area under the curve value was 0.957, indicating a high diagnostic value ( $P < 0.05$ ). Furthermore, the optimal cutoff value was 8.418%, with a diagnostic sensitivity of 0.819 and specificity of 0.947.

## CONCLUSION

In summary, the serum levels of PMAs significantly increase in patients with sepsis and ARDS. Therefore, serum PMAs have the potential to become a new biomarker for clinically diagnosing sepsis complicated by ARDS.

**Key Words:** Sepsis; Acute respiratory distress syndrome; Platelet leukocyte aggregates; Platelet mononuclear cell aggregates, Biomarker

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Our research aimed to investigate the expression of platelet mononuclear cell aggregates (PMAs) in the serum of patients with sepsis complicated by acute respiratory distress syndrome (ARDS) and its clinical significance. The results indicate that the serum levels of PMAs significantly increase in patients with sepsis and ARDS. Therefore, serum PMAs have the potential to become a new biomarker for clinically diagnosing sepsis complicated by ARDS.

**Citation:** Huang CM, Li JJ, Wei WK. Clinical significance of platelet mononuclear cell aggregates in patients with sepsis and acute respiratory distress syndrome. *World J Clin Cases* 2024; 12(5): 966-972

**URL:** <https://www.wjgnet.com/2307-8960/full/v12/i5/966.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v12.i5.966>

## INTRODUCTION

Sepsis is a common and critical clinical condition, and in recent years, the number of cases diagnosed as sepsis with concomitant acute respiratory distress syndrome (ARDS) has increased, owing to the enhanced awareness among medical professionals and the continuous development of modern medical diagnostic technologies[1]. Due to the high mortality rate associated with sepsis, it has attracted widespread attention in the clinical community. Despite various diagnostic tools available, early diagnosis of ARDS still lacks specific biomarkers. Recent studies have confirmed that one of the main pathogenic mechanisms of sepsis-associated ARDS involves the actions of various pathological injuries and inflammatory factors[2,3].

In this study, platelets and white blood cells in the patient's body are activated, leading to an increase in surface adhesion molecules. These adhesion molecules further form platelet-white blood cell aggregates, including platelet-neutrophil aggregates (PNAs), platelet aggregates (PLyAs), and platelet-mononuclear cell aggregates (PMAs)[4]. Activated platelets bind to monocytes and neutrophils, with the noteworthy observation that the binding of platelets to monocytes precedes that to neutrophils. PMAs are considered one of the markers of platelet activation[5]. The objective of this study is to investigate whether PMAs in the serum of sepsis patients with ARDS can serve as effective biomarkers for the early diagnosis of this complication.

## MATERIALS AND METHODS

### Study subjects and diagnostic criteria

This study included 72 adult sepsis patients admitted to our hospital between March 2019 and March 2022. The diagnosis was in accordance with the "International Guidelines for Management of Sepsis and Septic Shock: 2016[6]." Following the 2012 Berlin definition[7], patients were categorized into the study group (sepsis with ARDS,  $n = 30$ ) and the control group (sepsis alone,  $n = 42$ ). Exclusion criteria comprised pregnancy with blood system diseases, pure blood system diseases, HIV infection, ongoing chemotherapy, use of immunosuppressive agents or antiplatelet drugs, pulmonary interstitial fibrosis, and acute exacerbation of chronic obstructive pulmonary disease. The study was approved by our hospital's medical ethics committee, and written informed consent was obtained from all patients or their authorized representatives.



### Sample collection

Three milliliters of peripheral venous blood were collected from all confirmed sepsis patients immediately upon admission, placed in anticoagulant tubes, and preserved and transported to the Shanghai Lanwei Medical Laboratory for further testing *via* ice pack refrigeration. To ensure accuracy, the entire blood collection process strictly adhered to standardized procedures to prevent platelet activation-induced errors.

### Sample testing

Patients underwent blood gas analysis, and the oxygenation index  $\text{PaO}_2/\text{FiO}_2$  was calculated. Flow cytometry was employed to classify platelet-mononuclear cell aggregates in the peripheral blood of both sepsis patient groups, including PLyAs, PMAs, and PNAs. The site of infection was recorded, and the nature of the pathogenic bacteria was determined through blood culture.

### Acute physiology and chronic health evaluation II Score

Within 24 h of admission, all confirmed sepsis patients had their physiological indicators meticulously recorded by the attending physician, who then calculated the Acute Physiology and Chronic Health Evaluation (APACHE) II score, noting the worst values.

### Statistical analysis

Statistical analysis was conducted using SPSS 20.0 software. Descriptive data are presented as mean  $\pm$  SD, and *t*-tests were used for comparisons. Receiver operating characteristic (ROC) curves were generated to determine the optimal cutoff value, sensitivity, and specificity of serum PMAs in diagnosing sepsis with ARDS. The significance level was set at  $P < 0.05$ .

## RESULTS

### General data comparison between the two groups

According to the data in Table 1, there were no significant differences ( $P > 0.05$ ) observed in general information, such as age, gender, infection site, pathogen, and oxygenation index  $\text{PaO}_2/\text{FiO}_2$ , between sepsis patients with ARDS and those with sepsis alone.

### Comparison of PNAs, PLyAs, and PMAs between the two groups

As shown in Table 2, the serum levels of PNAs and PLyAs in sepsis patients with ARDS were slightly higher than those in the sepsis-alone group; however, these differences were not statistically significant ( $P > 0.05$ ). Nevertheless, the serum levels of PMAs in sepsis patients with ARDS were significantly higher than those in the sepsis-alone group, and this difference was statistically significant ( $P < 0.05$ ).

### Comparison of APACHE II scores between the two groups

According to the data in Table 3, the APACHE II scores of sepsis patients with ARDS were significantly higher than those of sepsis patients without ARDS, and this difference was statistically significant ( $P < 0.05$ ).

### Correlation analysis between PMAs and APACHE II scores

As illustrated in Figure 1A, the PMAs levels in sepsis patients with ARDS were significantly higher than those in sepsis patients without ARDS. Further linear correlation analysis revealed a positive correlation between PMAs and APACHE II scores in patients ( $r = 0.671$ ,  $P < 0.05$ ).

### Diagnostic value of various indicators for ARDS

Using PMAs and APACHE II scores as test variables and ARDS as the state variable, ROC curves were fitted. When using PMAs as the test variable, the area under the curve (AUC) was 0.957, indicating a significant diagnostic value ( $P < 0.05$ ). The optimal cutoff value for PMAs was 8.418%, with a diagnostic sensitivity of 0.819 and specificity of 0.947. When using APACHE II scores as the test variable, the AUC was 0.940, indicating a significant diagnostic value ( $P < 0.05$ ). The optimal cutoff value for APACHE II scores was 17.115, with a diagnostic sensitivity of 0.837 and specificity of 0.844. Refer to Tables 4 and 5, and Figure 1B for detailed results.

## DISCUSSION

Sepsis, as a common complication of severe infections, trauma, acute abdomen, and major surgeries in clinical practice, spans multiple disciplines such as internal medicine, surgery, and gynecology. It leads to multi-organ dysfunction, poor prognosis, and a high mortality rate[8]. The lungs are particularly susceptible to the effects of sepsis, causing pathological damage that is closely related to patient prognosis. Sepsis-induced multi-organ pathology includes the aggregation of white blood cells and platelets at the site of infection, disseminated intravascular coagulation, endothelial damage, resulting in the loss of surfactant on the alveolar surface, and activation of oxidative stress responses. These mechanisms

**Table 1 Comparison of general data between the two groups of patients (mean  $\pm$  SD), *n* (%)**

	Study group ( <i>n</i> = 30)	Control group ( <i>n</i> = 42)	$\chi^2/t$	<i>P</i> value
Male/female	16/14	25/17	0.475	0.523
Age	51.6 $\pm$ 11.4	48.6 $\pm$ 14.7	1.234	0.224
Infection			1.587	0.904
Urinary tract infection	4 (13.3)	5 (11.9)		
Hematogenous infection	3 (10)	4 (9.5)		
Abdominal infection	6 (20)	8 (19.1)		
Pulmonary infection	15 (50)	22 (52.4)		
Others	2 (6.7)	3 (7.1)		
Microbiology			2.88	0.518
Fungus	3 (10)	5 (11.9)		
G-	12 (40)	17 (40.5)		
G+	5 (16.7)	8 (19)		
Mixed infection	6 (20)	7 (16.7)		
Unknown cause	4 (13.3)	5 (11.9)		
PaO <sub>2</sub> /FiO <sub>2</sub>	141.85 $\pm$ 29.44	145.35 $\pm$ 30.28	11.27	0.912

**Table 2 Comparison of platelet neutrophil aggregates, platelet aggregates, and platelet-mononuclear cell aggregates between the two groups (mean  $\pm$  SD)**

Groups	<i>n</i>	PNAs (%)	PLyAs (%)	PMAs (%)
Study group	30	14.15 $\pm$ 8.93	15.42 $\pm$ 6.97	27.18 $\pm$ 6.14 <sup>1</sup>
Control group	42	13.87 $\pm$ 9.24	14.78 $\pm$ 3.24	17.29 $\pm$ 2.05

<sup>1</sup>Compared to the Control groups, the serum levels of platelet-mononuclear cell aggregates increased in sepsis patients with acute respiratory distress syndrome (*P* < 0.05).

APACHE II: Acute Physiology and Chronic Health Evaluation II Scores; PMAs: Platelet-mononuclear cell aggregates; PNAs: platelet neutrophil aggregates; PLyAs: Platelet aggregates.

**Table 3 Comparison of Acute Physiology and Chronic Health Evaluation II Scores between the two groups (mean  $\pm$  SD)**

Groups	<i>n</i>	APACHE II
Study group	30	35.17 $\pm$ 5.44 <sup>1</sup>
Control group	42	23.39 $\pm$ 4.24

<sup>1</sup>Compared to the Control groups, the Acute Physiology and Chronic Health Evaluation II scores increased in sepsis patients with acute respiratory distress syndrome (*P* < 0.05).

APACHE II: Acute Physiology and Chronic Health Evaluation II Scores.

collectively contribute to the development of severe lung injury[9]. Due to the activation of inflammatory reactions and coagulation mechanisms in sepsis patients, they are prone to developing ARDS, with pathological manifestations in the lungs characterized by increased permeability of the alveolar-capillary barrier, pulmonary tissue edema, and severe hypoxemia. After the onset of typical injury symptoms, some sepsis patients may rapidly deteriorate within a short period, progressing to ARDS, thus affecting their prognosis[10]. Venous blood samples are the most readily available and suitable for laboratory testing. Among various specimens, patient serum is primarily used as a biological specimen for accurately and rapidly assessing the severity of sepsis.

In sepsis, damage to endothelial cells leads to activating inflammatory cells and platelets. Activated inflammatory cells release a large number of inflammatory and cellular factors through a cascade reaction, promoting endothelial cell apoptosis and monocyte release of chemokines. Platelets and white blood cells interact in the microcirculation of damaged tissues, forming platelet-white blood cell aggregates. This process further accelerates the release of inflam-



Table 4 Area under the curve for various parameters


Parameter	AUC	SE	P value	95%CI	
				Upper limit	Lower limit
PMAs	0.957	0.022	< 0.05	0.914	0.974
APACHE II	0.93	0.021	< 0.05	0.872	0.981

APACHE II: Acute Physiology and Chronic Health Evaluation II Scores; PMAs: Platelet-mononuclear cell aggregates; AUC: Area under the curve.

Table 5 Diagnostic values for various parameters

Parameter	Cutoff value	Sensitivity	Specificity	Positive predictive value	Negative predictive value
PMAs	8.418	0.819	0.947	0.956	0.819
APACHE II	17.115	0.837	0.844	0.829	0.877

APACHE II: Acute Physiology and Chronic Health Evaluation II Scores; PMAs: Platelet-mononuclear cell aggregates.



DOI: 10.12998/wjcc.v12.i5.966 Copyright ©The Author(s) 2024.

Figure 1 Platelet-mononuclear cell aggregates and Acute Physiology and Chronic Health Evaluation II scores. A: Linear correlation between platelet-mononuclear cell aggregates and Acute Physiology and Chronic Health Evaluation (APACHE) II scores; B: Diagnostic Value of platelet-mononuclear cell aggregates and APACHE II Scores for acute respiratory distress syndrome. APACHE II: Acute Physiology and Chronic Health Evaluation II Scores; PMA: Platelet-mononuclear cell aggregate.

matory factors such as interleukin and tumor necrosis factor- $\alpha$ [11]. The worsening of the inflammatory response leads to endothelial cell swelling, necrosis and shedding, further worsening the patient's condition. Therefore, platelet-white blood cell aggregates in the serum play a crucial intermediary role between platelet activation and inflammatory response.

The results of this study indicate that the serum levels of PMAs in sepsis patients with ARDS were significantly higher than those in sepsis patients without ARDS ( $P < 0.05$ ), confirming the utility of PMAs as a beneficial indicator for diagnosing sepsis with ARDS. The APACHE II scoring system is commonly used to assess the severity and prognosis of critically ill patients[12], have confirmed the application of the APACHE II score in predicting mortality in sepsis patients. The current study demonstrates that the APACHE II scores of sepsis patients with ARDS were significantly higher than those of sepsis patients without ARDS and were positively correlated with serum PMAs levels ( $P < 0.05$ ). This further confirms the clinical importance of serum PMAs in the early diagnosis of sepsis with ARDS.

Study limitation

One limitation of this study is the relatively small sample size, with a total of 72 hospitalized patients included in the analysis. The limited sample size may affect the generalizability of the findings to a broader population. Additionally, the study focused on patients from a single hospital, which may introduce institutional biases and limit the external validity of the results. Including a more diverse and larger sample from multiple medical centers could enhance the robustness and applicability of the study findings. Furthermore, the retrospective nature of the study poses inherent limitations. The reliance on historical data collected from medical records may lead to incomplete or missing information. The retrospective design also prevents the researchers from controlling the data collection process, potentially introducing biases in the selection of patients or in the measurement of variables. A prospective study with a carefully designed protocol and standardized data collection procedures would provide stronger evidence and allow for better control of confounding variables. The study primarily focused on the expression of PMAs in the serum as a potential biomarker for

sepsis complicated by ARDS. While the findings suggest a significant association, the study does not explore the underlying mechanisms or causality between elevated PMAs levels and the development of ARDS in sepsis patients. Further mechanistic studies are needed to elucidate the pathways through which PMAs may contribute to the pathogenesis of ARDS in sepsis. Finally, the study does not address the specificity of PMAs as a biomarker, and its utility in distinguishing sepsis with ARDS from other conditions that may present with similar clinical manifestations. Future research should explore the specificity and sensitivity of PMAs in differentiating various respiratory and systemic disorders to better understand its diagnostic value in a broader clinical context.

## CONCLUSION

In conclusion, the elevation of serum PMAs levels is closely associated with the release of inflammatory factors. Although the exact mechanism of PMAs still requires further research, current studies suggest that its changes have important clinical significance in early diagnosing sepsis with ARDS. Therefore, PMAs may serve as one of the biomarkers for early diagnosing sepsis with ARDS.

## ARTICLE HIGHLIGHTS

### Research background

The diagnosis of sepsis combined with acute respiratory distress syndrome (ARDS) has increased owing to the enhanced awareness among medical professionals and the continuous development of modern medical technologies, while early diagnosis of ARDS still lacks specific biomarkers. One of the main pathogenic mechanisms of sepsis-associated ARDS involves the actions of various pathological injuries and inflammatory factors, such as platelet and white blood cells activation, leading to an increase of surface adhesion molecules. These adhesion molecules further form platelet-white blood cell aggregates, including platelet-mononuclear cell aggregates (PMAs). PMAs has been identified as one of the markers of platelet activation, here we hypothesize that PMAs might play a potential biomarker for the early diagnosis of this complication.

### Research motivation

To investigate whether PMAs could be a potential biomarker for the early diagnosis of sepsis combined with ARDS.

### Research objectives

To investigate the clinical significance of PMAs in patients with sepsis complicated by ARDS.

### Research methods

72 patients diagnosed with sepsis were enrolled in the study between March 2019 and March 2022. Among them, 30 patients with sepsis and ARDS formed the study group, while 42 sepsis patients without ARDS comprised the control group. After diagnosis, venous blood samples were immediately collected from all patients. Flow cytometry was employed to analyze the expression of PMAs, platelet neutrophil aggregates (PNAs), and Platelet Aggregates (PLyAs) in the serum. Additionally, the Acute Physiology and Chronic Health Evaluation (APACHE) II score was calculated for each patient, and Receiver operating characteristic curves were generated to assess diagnostic value.

### Research results

The levels of PNAs and PLyAs in the serum of the study group were higher than those in the control group, but the difference was not statistically significant ( $P > 0.05$ ). However, the expression of PMAs in the serum of the study group was significantly upregulated ( $P < 0.05$ ) and positively correlated with the APACHE II score ( $r=0.671$ ,  $P < 0.05$ ). When using PMAs as a diagnostic indicator, the area under the curve value was 0.957, indicating a high diagnostic value ( $P < 0.05$ ). Furthermore, the optimal cutoff value was 8.418%, with a diagnostic sensitivity of 0.819 and specificity of 0.947.

### Research conclusions

The serum levels of PMAs significantly increase in patients with sepsis and ARDS, which might have the potential to become a new biomarker for clinically diagnosing sepsis complicated by ARDS.

### Research perspectives

Our study provides a new method for the early diagnosis of sepsis combined with ARDS, which is the detection of serum PMAs. More samples should be enrolled to confirm this method in the future study.

## FOOTNOTES

**Author contributions:** Huang CM performed the study; Li JJ analyzed the data; Wei WK designed the research and wrote the manuscript;

All authors have read and approve the final manuscript.

**Institutional review board statement:** The study was reviewed and approved by the Institutional Review Board of the Second People's Hospital of Haining City.

**Informed consent statement:** All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

**Conflict-of-interest statement:** All the authors report no relevant conflicts of interest for this article.

**Data sharing statement:** No additional data are available.

**STROBE statement:** The authors have read the STROBE Statement – checklist of items, and the manuscript was prepared and revised according to the STROBE Statement – checklist of items.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/Licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Chong-Min Huang [0009-0001-7015-3577](https://orcid.org/0009-0001-7015-3577).

**S-Editor:** Li L

**L-Editor:** A

**P-Editor:** Li L

## REFERENCES

- 1 **Hu Q**, Hao C, Tang S. From sepsis to acute respiratory distress syndrome (ARDS): emerging preventive strategies based on molecular and genetic researches. *Biosci Rep* 2020; **40** [PMID: [32319516](https://pubmed.ncbi.nlm.nih.gov/32319516/) DOI: [10.1042/BSR20200830](https://doi.org/10.1042/BSR20200830)]
- 2 **Li X**, Jamal M, Guo P, Jin Z, Zheng F, Song X, Zhan J, Wu H. Irisin alleviates pulmonary epithelial barrier dysfunction in sepsis-induced acute lung injury via activation of AMPK/SIRT1 pathways. *Biomed Pharmacother* 2019; **118**: 109363 [PMID: [31545277](https://pubmed.ncbi.nlm.nih.gov/31545277/) DOI: [10.1016/j.biopha.2019.109363](https://doi.org/10.1016/j.biopha.2019.109363)]
- 3 **Chen Y**, Qiu C, Cai W. Identification of key immune genes for sepsis-induced ARDS based on bioinformatics analysis. *Bioengineered* 2022; **13**: 697-708 [PMID: [34898369](https://pubmed.ncbi.nlm.nih.gov/34898369/) DOI: [10.1080/21655979.2021.2012621](https://doi.org/10.1080/21655979.2021.2012621)]
- 4 **Le Joncour A**, Biard L, Vautier M, Bugaut H, Mekinian A, Maalouf G, Vieira M, Marcelin AG, Rosenzweig M, Klatzmann D, Corvol JC, Paccoud O, Carillion A, Salem JE, Cacoub P, Boulaftali Y, Saadoun D. Neutrophil-Platelet and Monocyte-Platelet Aggregates in COVID-19 Patients. *Thromb Haemost* 2020; **120**: 1733-1735 [PMID: [33124027](https://pubmed.ncbi.nlm.nih.gov/33124027/) DOI: [10.1055/s-0040-1718732](https://doi.org/10.1055/s-0040-1718732)]
- 5 **Hottz ED**, Azevedo-Quintanilha IG, Palhinha L, Teixeira L, Barreto EA, Pão CRR, Righy C, Franco S, Souza TML, Kurtz P, Bozza FA, Bozza PT. Platelet activation and platelet-monocyte aggregate formation trigger tissue factor expression in patients with severe COVID-19. *Blood* 2020; **136**: 1330-1341 [PMID: [32678428](https://pubmed.ncbi.nlm.nih.gov/32678428/) DOI: [10.1182/blood.2020007252](https://doi.org/10.1182/blood.2020007252)]
- 6 **Rhodes A**, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, Kumar A, Sevransky JE, Sprung CL, Nunnally ME, Rochwerg B, Rubenfeld GD, Angus DC, Annane D, Beale RJ, Bellinhan GJ, Bernard GR, Chiche JD, Coopersmith C, De Backer DP, French CJ, Fujishima S, Gerlach H, Hidalgo JL, Hollenberg SM, Jones AE, Karnad DR, Kleinpell RM, Koh Y, Lisboa TC, Machado FR, Marini JJ, Marshall JC, Mazuski JE, McIntyre LA, McLean AS, Mehta S, Moreno RP, Myburgh J, Navalesi P, Nishida O, Osborn TM, Perner A, Plunkett CM, Ranieri M, Schorr CA, Seckel MA, Seymour CW, Shieh L, Shukri KA, Simpson SQ, Singer M, Thompson BT, Townsend SR, Van der Poll T, Vincent JL, Wiersinga WJ, Zimmerman JL, Dellinger RP. Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock: 2016. *Intensive Care Med* 2017; **43**: 304-377 [PMID: [28101605](https://pubmed.ncbi.nlm.nih.gov/28101605/) DOI: [10.1007/s00134-017-4683-6](https://doi.org/10.1007/s00134-017-4683-6)]
- 7 **ARDS Definition Task Force**, Ranieri VM, Rubenfeld GD, Thompson BT, Ferguson ND, Caldwell E, Fan E, Camporota L, Slutsky AS. Acute respiratory distress syndrome: the Berlin Definition. *JAMA* 2012; **307**: 2526-2533 [PMID: [22797452](https://pubmed.ncbi.nlm.nih.gov/22797452/) DOI: [10.1001/jama.2012.5669](https://doi.org/10.1001/jama.2012.5669)]
- 8 **Chiu C**, Legrand M. Epidemiology of sepsis and septic shock. *Curr Opin Anaesthesiol* 2021; **34**: 71-76 [PMID: [33492864](https://pubmed.ncbi.nlm.nih.gov/33492864/) DOI: [10.1097/ACO.0000000000000958](https://doi.org/10.1097/ACO.0000000000000958)]
- 9 **Huang M**, Cai S, Su J. The Pathogenesis of Sepsis and Potential Therapeutic Targets. *Int J Mol Sci* 2019; **20** [PMID: [31671729](https://pubmed.ncbi.nlm.nih.gov/31671729/) DOI: [10.3390/ijms2015376](https://doi.org/10.3390/ijms2015376)]
- 10 **Li W**, Li D, Chen Y, Abudou H, Wang H, Cai J, Wang Y, Liu Z, Liu Y, Fan H. Classic Signaling Pathways in Alveolar Injury and Repair Involved in Sepsis-Induced ALI/ARDS: New Research Progress and Prospect. *Dis Markers* 2022; **2022**: 6362344 [PMID: [35726235](https://pubmed.ncbi.nlm.nih.gov/35726235/) DOI: [10.1155/2022/6362344](https://doi.org/10.1155/2022/6362344)]
- 11 **Liu D**, Huang SY, Sun JH, Zhang HC, Cai QL, Gao C, Li L, Cao J, Xu F, Zhou Y, Guan CX, Jin SW, Deng J, Fang XM, Jiang JX, Zeng L. Sepsis-induced immunosuppression: mechanisms, diagnosis and current treatment options. *Mil Med Res* 2022; **9**: 56 [PMID: [36209190](https://pubmed.ncbi.nlm.nih.gov/36209190/) DOI: [10.1186/s40779-022-00422-y](https://doi.org/10.1186/s40779-022-00422-y)]
- 12 **Akinosoglou K**, Kapsokosta G, Mouktaroudi M, Rovina N, Kaldis V, Stefanos A, Kontogiorgi M, Giamarellos-Bourboulis E, Gogos C; Hellenic Sepsis Study Group. Diabetes on sepsis outcomes in non-ICU patients: A cohort study and review of the literature. *J Diabetes Complications* 2021; **35**: 107765 [PMID: [33187869](https://pubmed.ncbi.nlm.nih.gov/33187869/) DOI: [10.1016/j.jdiacomp.2020.107765](https://doi.org/10.1016/j.jdiacomp.2020.107765)]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [office@baishideng.com](mailto:office@baishideng.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

