World Journal of *Clinical Cases*

World J Clin Cases 2024 February 26; 12(6): 1039-1195





Published by Baishideng Publishing Group Inc

W J C C World Journal of Clinical Cases

Contents

Thrice Monthly Volume 12 Number 6 February 26, 2024

EDITORIAL

- 1039 Lateral clavicle fracture-plating options and considerations Muthu S, Annamalai S, Kandasamy V
- 1045 Tumor deposits in axillary adipose tissue in patients with breast cancer: Do they matter? Mubarak M, Rashid R, Shakeel S

MINIREVIEWS

1050 New strategies in the diagnosis and treatment of immune-checkpoint inhibitor-mediated colitis Velikova T. Krastev B. Gulinac M. Zashev M. Graklanov V. Peruhova M

ORIGINAL ARTICLE

Retrospective Cohort Study

1063 Correlative factors of poor prognosis and abnormal cellular immune function in patients with Alzheimer's disease

Bai H, Zeng HM, Zhang QF, Hu YZ, Deng FF

1076 Bipolar hip arthroplasty using conjoined tendon preserving posterior lateral approach in treatment of displaced femoral neck fractures

Yan TX, Dong SJ, Ning B, Zhao YC

Retrospective Study

Association of preschool children behavior and emotional problems with the parenting behavior of both 1084 parents

Wang SM, Yan SQ, Xie FF, Cai ZL, Gao GP, Weng TT, Tao FB

- 1094 Assessment of the triglyceride glucose index in adult patients with chronic diarrhea and constipation Zhu JY, Liu MY, Sun C
- 1104 Acute pancreatitis as a complication of acute COVID-19 in kidney transplant recipients Basic-Jukic N, Juric I, Katalinic L, Furic-Cunko V, Sesa V, Mrzljak A

Observational Study

1111 Clinical analysis of 12 cases of ovarian neuroendocrine carcinoma Xing XY, Zhang W, Liu LY, Han LP

META-ANALYSIS

1120 Efficacy and safety of remimazolam in bronchoscopic sedation: A meta-analysis Zhou Y, Zhao C, Tang YX, Liu JT



Contents

Thrice Monthly Volume 12 Number 6 February 26, 2024

CASE REPORT

- 1130 Simple bone cysts of the proximal humerus presented with limb length discrepancy: A case report Lin CS, Lin SM, Rwei SP, Chen CW, Lan TY
- 1138 Postoperative abdominal herpes zoster complicated by intestinal obstruction: A case report Dong ZY, Shi RX, Song XB, Du MY, Wang JJ
- 1144 Clinical evolution of antisynthetase syndrome-associated interstitial lung disease after COVID-19 in a man with Klinefelter syndrome: A case report

Wu XX, Cui J, Wang SY, Zhao TT, Yuan YF, Yang L, Zuo W, Liao WJ

- 1150 Giant bile duct dilatation in newborn: A case report Quan DW, Li PG, Xu XH, Liu SQ
- Left atrial appendage occluder detachment treated with transthoracic ultrasound combined with digital 1157 subtraction angiography guided catcher: A case report

Yu K, Mei YH

- 1163 Adult sigmoid intussusception resembling rectal prolapse: A case report Tsai TJ, Liu YS
- 1169 Gigantic occipital epidermal cyst in a 56-year-old female: A case report Wei Y, Chen P, Wu H
- 1174 Autoimmune hepatitis-primary biliary cholangitis overlap syndrome complicated by various autoimmune diseases: A case report

Qin YJ, Gao T, Zhou XN, Cheng ML, Li H

- 1182 Parotid metastasis of rare lung adenocarcinoma: A case report Yan RX, Dou LB, Wang ZJ, Qiao X, Ji HH, Zhang YC
- 1190 Management of retroperitoneal high-grade serous carcinoma of unknown origin: A case report Hsieh WL, Ding DC



Contents

Thrice Monthly Volume 12 Number 6 February 26, 2024

ABOUT COVER

Peer Reviewer of World Journal of Clinical Cases, Madhan Jeyaraman, MS, PhD, Assistant Professor, Sri Lalithambigai Medical College and Hospital, Dr MGR Educational and Research Institute University, Chennai 600095, India. madhanjeyaraman@gmail.com

AIMS AND SCOPE

The primary aim of World Journal of Clinical Cases (WJCC, World J Clin Cases) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports[®] cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Zi-Hang Xu; Production Department Director: Xu Gue; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL World Journal of Clinical Cases	INSTRUCTIONS TO AUTHORS https://www.wjgnet.com/bpg/gerinfo/204	
ISSN	GUIDELINES FOR ETHICS DOCUMENTS	
ISSN 2307-8960 (online)	https://www.wjgnet.com/bpg/GerInfo/287	
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH	
April 16, 2013	https://www.wjgnet.com/bpg/gerinfo/240	
FREQUENCY	PUBLICATION ETHICS	
Thrice Monthly	https://www.wjgnet.com/bpg/GerInfo/288	
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT	
Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati	https://www.wjgnet.com/bpg/gerinfo/208	
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE	
https://www.wjgnet.com/2307-8960/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242	
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS	
February 26, 2024	https://www.wjgnet.com/bpg/GerInfo/239	
COPYRIGHT	ONLINE SUBMISSION	
© 2024 Baishideng Publishing Group Inc	https://www.f6publishing.com	

© 2024 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: office@baishideng.com https://www.wjgnet.com



W J C C World Journal of Clinical Cases

Submit a Manuscript: https://www.f6publishing.com

World J Clin Cases 2024 February 26; 12(6): 1130-1137

DOI: 10.12998/wjcc.v12.i6.1130

ISSN 2307-8960 (online)

CASE REPORT

Simple bone cysts of the proximal humerus presented with limb length discrepancy: A case report

Cing Syue Lin, Shang Ming Lin, Syang-Peng Rwei, Chin-Wen Chen, Tsung-Yu Lan

Specialty type: Orthopedics

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): 0 Grade C (Good): C Grade D (Fair): 0 Grade E (Poor): 0

P-Reviewer: Liu TF, China

Received: October 1, 2023 Peer-review started: October 1, 2023

First decision: January 2, 2024 Revised: January 10, 2024 Accepted: January 30, 2024 Article in press: January 30, 2024 Published online: February 26, 2024



Cing Syue Lin, Tsung-Yu Lan, Department of Orthopedic Surgery, Far Eastern Memorial Hospital, New Taipei City 220, Taiwan

Shang Ming Lin, Department of Materials and Textiles, Asia Eastern University of Science and Technology, New Taipei City 220, Taiwan

Syang-Peng Rwei, Chin-Wen Chen, Institute of Organic and Polymeric Materials, National Taipei University of Technology, Taipei City 10608, Taiwan

Corresponding author: Tsung-Yu Lan, Doctor, MD, Chief Doctor, Surgeon, Department of Orthopedic Surgery, Far Eastern Memorial Hospital, No. 21 Sec. 2, Nanya S. Road, Banciao District, New Taipei City 220, Taiwan. gbwsh0130@gmail.com

Abstract

BACKGROUND

Simple bone cysts (SBC) are benign tumor-like bone lesions typically identified in children. While SBC may lead to growth disturbances or growth arrest, such cases are uncommon. The mechanisms behind these observations remain unclear. Additionally, research on the etiology of SBC remains inconclusive, and there has been no consensus on the appropriate timing and methodology for treatment.

CASE SUMMARY

Here, we present our experience in the successful surgical management of a 10year-old girl with SBC, who presented with a pathological fracture complicated by malunion of the displaced fracture, varus deformity, and limb length discrepancy. We hypothesized two possible etiologies for the patient's growth arrest and subsequent humerus varus deformity: (1) Direct disruption of the physis by fluid from the cyst itself; and (2) damage to the epiphysis due to repetitive pathological fractures associated with SBC. In addressing this case, surgical intervention was undertaken to correct the proximal humerus varus deformity. This approach offered the advantages of simultaneously correcting angular abnormalities, achieving mild limb lengthening, providing definitive SBC treatment, and reducing the overall treatment duration.

CONCLUSION

As per current literature, acute correction of acute angular deformity in proximal humeral SBC is not well comprehended. However, in this specific case, acute correction was considered an optimal solution.

WJCC | https://www.wjgnet.com

Key Words: Bone cyst; Solitary cysts; Benign; Humerus; Growth arrest; Limb length discrepancy; Case report

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: We successfully operated on a 10-year-old girl with simple bone cysts (SBC), who presented with varus deformity, limb length discrepancy, and malunion resulting from a neglected pathological fracture. We proposed two potential causes for the humeral growth arrest and varus deformity in this patient: (1) Direct disruption of the physis by fluid from the cyst itself; and (2) damage to the epiphysis due to recurrent pathological fractures associated with SBC. Presently, the literature lacks a comprehensive understanding of the acute correction of angular deformity in proximal humeral SBC. However, in this specific case, acute correction was considered an ideal option.

Citation: Lin CS, Lin SM, Rwei SP, Chen CW, Lan TY. Simple bone cysts of the proximal humerus presented with limb length discrepancy: A case report. World J Clin Cases 2024; 12(6): 1130-1137 URL: https://www.wjgnet.com/2307-8960/full/v12/i6/1130.htm DOI: https://dx.doi.org/10.12998/wjcc.v12.i6.1130

INTRODUCTION

Simple bone cysts (SBC), also known as solitary bone cysts are benign, fluid-filled tumor-like bone lesions commonly found in the proximal humerus (70%) or femur (25%) of children and adolescents[1,2]. While SBC may occasionally result in growth disturbances or growth arrest, such cases are uncommon. The underlying mechanisms behind this remain unclear[3]. This article presents a case involving a 10-year-old girl who initially complained of left arm pain attributed to an angulated old fracture, accompanied by limb length discrepancy. Additionally, a review of the current literature on SBC is included in this discussion.

CASE PRESENTATION

Chief complaints

A 10-year-old Chinese girl presented at our orthopedics clinic with a history of restricted left shoulder elevation and left upper limb length discrepancy spanning six months.

History of present illness

The patient experienced difficulty lifting her left arm during undressing, leading her mother to observe a noticeable length difference in the left arm. Concerned about these issues, the mother brought her to our clinic for evaluation.

History of past illness

The patient had no pre-existing medical conditions. No history of trauma, fever, local infection, or other precipitating factors in the left shoulder were noted.

Personal and family history

The patient's mother denied any family history of tumors.

Physical examination

During physical examination, it was observed that the patient's left arm was approximately 4-5 cm shorter than the right. The active range of motion (ROM) for the left shoulder was limited due to pain (Table 1). However, the distal circulation was found to be intact, and no focal neurological signs or symptoms were observed.

Laboratory examinations

Serum C-reactive protein levels were within the normal range, and routine blood and urine analyses showed no abnormalities. A tumor marker workup, which included alpha-fetoprotein, cancer antigen 125, carbohydrate antigen 19-9, prostate-specific antigen, and carcinoembryonic antigen, all revealed values within normal limits.

Imaging examinations

Initial imaging studies indicated an angulated old fracture at the proximal left humeral neck (Figure 1). Subsequent examination with magnetic resonance imaging (MRI) revealed marked deformity of the left humeral head and possible malunion of the displaced fracture in the left humeral neck. No evidence of cortical discontinuity, intraosseous patho-logies, or periosteal reaction was noted (Figure 2).



Table 1 The preoperative and postoperative active range of motion of left shoulder			
	Preoperative active ROM	Postoperative-1 month active ROM	Postoperative-1 year active ROM
Flexion	70	176	180
Abduction	80	175	180
External rotation	45	90	95
Internal rotation	10	65	75

ROM: Range of motion.







Figure 2 Preoperative magnetic resonance images and pathological results of the patient. Preoperative T2-weighted magnetic resonance images of the left humerus showed a marked deformity of the humeral head and a potential malunion of the displaced fracture in the left humeral neck.

FINAL DIAGNOSIS

Considering the patient's medical history, the final diagnosis indicated a case of left humerus SBC with varus malunion or suspected postinfection sequelae of osteomyelitis.



Baisbideng® WJCC | https://www.wjgnet.com



Figure 3 Surgical procedure for acute correction of the proximal humerus for the varus deformity. The red area indicates the location of the simple bone cyst (SBC). A: Osteotomy was performed in the most angulated area, removing the deformity site of the SBC; B: The gap was filled with allograft (donor femoral head); C: The proximal humerus was fixed with an anatomical locking plate (Depuy-Synthesis®, Raynham, MA, United States) in a valgus position.

TREATMENT

Surgical intervention was recommended for this case of left humerus SBC with varus malunion or suspected postinfection sequela of osteomyelitis. The procedure involved placing the patient under general anesthesia in a semi-sitting position. The anterior approach was utilized to expose the proximal metaphysis and diaphysis of the left humerus. Osteotomy was performed at the most angulated area, and curettage over the lesion site was performed, with the specimens sent for pathological study. The proximal humerus was then fixed with an anatomical locking plate (Depuy-Synthes, Raynham, MA, United States) in a valgus position and the gap was filled with an allograft (donor femoral head) (Figure 3). The patient has not received any additional medication therapy beyond the surgical intervention.

OUTCOME AND FOLLOW-UP

Postoperatively, the patient was immobilized in a sling for 2 wk. The left shoulder length discrepancy was reduced by 2 cm. After 1 month, she exhibited full active ROM in the left shoulder, with no reported wound complications or infection (Table 1 and Figure 4). Follow-up radiographs indicated no progression of the lesion; the osteotomy site showed solid union at 1 month and 1 year postoperatively (Figure 1). The histological report revealed degenerated cortical bone and cartilage tissue, with fibrous cystic wall with fibrin-like materials, cholesterol clefts and surrounding some new bone formation (Figure 5), leading to the diagnosis of SBC. While no pain or discomfort was reported, the implant was removed one year later based on the parents' preference. Subsequent radiographs showed that the left proximal humerus had healed with union and was well-aligned (Figure 1). One year following implant removal surgery, radiographic assessment revealed favorable remodeling and alignment of the left proximal humerus. Physical examination showed full active ROM and normal muscle power in the left shoulder (Table 1 and Figure 1). She exhibits the ability to execute motor milestone tasks in accordance with her chronological age. Furthermore, the length discrepancy between both hands was maintained at a 2 cm difference.

DISCUSSION

SBC are benign metaphyseal lytic lesions predominantly occurring in males^[3]. Despite research, the etiology of SBC remains inconclusive [4,5]. Most cases of SBC are asymptomatic and are discovered incidentally. Spontaneous resolution of SBC occurs in only about 5% to 10% of cases[6,7]. Symptomatic cases are often associated with pathological fractures (63%-87%)[1,8] and less than 15% of cysts resolve spontaneously after a fracture occurs[5,9]. Presently, there is no consensus or established guidelines on the optimal timing and method of treating SBC[2,6,7,10]. To the best of our knowledge, the acute correction of angular deformity in proximal humeral SBC is not yet fully understood.

The primary differential diagnoses for pathological fractures in children aged 5-10 years include SBC, aneurysmal bone cysts (ABC), non-ossifying fibroma, and osteomyelitis[11]. In our patient, the lesion was located in the proximal humerus, consistent with the predisposing location of SBC. As our patient exhibited no signs of inflammation, bruises, or swelling, and had not received antibiotic treatment, chronic osteomyelitis was ruled out. Given the similar clinical symptoms of SBC, ABC, and non-ossifying fibroma, imaging and pathology played a crucial role in differentiating these





Figure 4 Two-week postoperative photos of the patient. A: Full active range of motion of the left shoulder; B: About 2-3 cm length discrepancy on the affected side.



Figure 5 Pathological results of the patient. A: Histological results showed fibrous cyst wall with fibrin-like material and calcification; B: Histological results showed cystic wall with cholesterol clefts and surrounding some new bone formation, all of which may be indicative of a diagnosis of a simple bone cyst.

diseases.

Radiographic findings in our case revealed an old, angulated fracture at the proximal left humeral neck (Figure 1). Although the characteristic cystic lesions of SBC were not visibly evident on radiography and MRI, the possibility of SBC could not be ruled out. Furthermore, MRI showed a smoother sclerotic junction with the absence of fat stranding or inflammation in the surrounding soft tissue (Figure 2), suggesting fluid reabsorption and healing. These findings, combined with the histological report, were indicative of a benign cause.

A pathological fracture occurred in association with the benign lesion, leading to the subsequent displacement of the fracture and resulting in a malunion at the left humeral neck. The histological report showed fibrous cystic wall with fibrin-like materials, cholesterol clefts and surrounding some new bone formation, consistent with SBC. Considering the patient's history, the results suggest fibrosis with chronic inflammation, supporting a diagnosis of SBC. Pathologic fractures can often remain asymptomatic until the moment of fracture or may occur after lesions that cause prodromal pain. However, they typically do not present with accompanying symptoms.

As such, we posit that this patient experienced an initially asymptomatic pathologic fracture due to low-energy trauma, resulting in a delayed diagnosis and treatment. The lack of a proper diagnosis and timely intervention contributed to the development of malunion in the old, displaced pathological fracture at the left humeral neck, along with subsequent issues of limb length discrepancy and angular deformity.

On rare occasions, SBC may lead to growth disturbances, and, the underlying mechanism remains a subject of debate. Some evidence suggests that the cyst fluid itself may directly disrupt the physis, causing growth disturbances[12-14]. Other potential processes include: (1) Fractures through the cyst, damaging the physis and resulting in growth arrest; (2) direct extension of the cyst through the physis; and (3) iatrogenic damage to a developing physis due to surgical removal

Baishidena® WJCC | https://www.wjgnet.com





of a cyst in proximity [6,12-18]. Regardless of the etiology, growth disturbances ultimately result in angular deformity and/or limb length discrepancy in the affected limb. In the case of our patient, limb length discrepancy was evident before surgical treatment, making it less likely that her injury was related to iatrogenic damage.

We proposed two hypotheses to explain the etiology of our patient's condition. First, we considered the possibility of direct extension of the SBC through the epiphysis on the medial side, or the direct disruption of the physis by the cyst fluid itself, leading to growth arrest and subsequent varus deformity of the humerus. Given that the proximal physis contributes to approximately 80% of the humerus' development, any disturbance in this region can result in significant upper limb length discrepancy[6]. The additional varus angulation of the humeral head occurred as a sequela of the proximal humerus fracture (Figure 6).

The second hypothesis proposed was that the epiphysis was damaged by repetitive pathological fractures due to SBC. The bone affected by SBC is weakened and prone to pathological fractures, which can, in turn, damage the epiphysis. As a consequence of these pathological fractures, varus humeral head angulation developed. The muscles pulling on the proximal humerus allowed pathological fractures to heal with an increasing varus deformity, giving rise to the typical complaint of severe shoulder ROM limitations in abduction and flexion. Furthermore, due to a defective medial epi-



Raishideng® WJCC | https://www.wjgnet.com

physis, the humerus developed further varus deformity and a length discrepancy (Figure 6).

We presented two surgical intervention options for the patient. The first option involved acute correction of the proximal humerus varus deformity. The alternative was distraction osteogenesis with external fixation, such as Ilizarov or Taylor spatial frame, combined with internal lengthening nails for limb lengthening and angle correction. Following careful consideration, the parents opted for the first treatment due to its shorter course, while recognizing the inherent limitation in correcting limb length discrepancy. Considering the nearly closed proximal humeral physis in the patient, the likelihood of developing subsequent length disparities was deemed low. Minor discrepancies in arm length, up to 5 or 6 cm, are generally well-tolerated without causing significant functional constraints, given the primary role of the upper extremity, which is non-weight-bearing[18,19].

Thus, the surgical treatment of acute correction for the proximal humerus varus deformity was performed. This approach offers the advantage of simultaneously correcting angular abnormalities, providing mild limb lengthening, definitive cyst therapy, and a shortened course of treatment. Consequently, we believe that the acute correction of the varus deformity was the optimal option in this case.

CONCLUSION

In conclusion, we presented a case of SBC presenting with a pathological fracture, complicated with malunion of the displaced fracture, varus deformity, and limb length discrepancy. The discussion included differential diagnoses of SBC, alongside a comprehensive review of the current literature. In our case, the patient's limb length discrepancy and varus deformity emerged as complications of the SBC and pathological fracture. It underscores the importance of close followup in patients with SBC to monitor the potential occurence of growth arrest.

FOOTNOTES

Author contributions: Lan TY, Lin SM, Rwei SP, and Chen CW contributed to conceptualization and supervision; Lin CS contributed to data collection, software, paper review, manuscript writing and editing; Lan TY contributed to clinical examination, surgery, original draft preparation and editing; All authors have read and agreed to the published version of the final manuscript.

Informed consent statement: Informed consent was obtained from the subject involved in the study. Written informed consent has been obtained from the patient to publish this paper.

Conflict-of-interest statement: All authors declare that they have no conflict of interest to disclose.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: Taiwan

ORCID number: Cing Syue Lin 0000-0002-3508-3104; Shang Ming Lin 0000-0002-6077-2657; Syang-Peng Rwei 0000-0001-6169-5852; Chin-Wen Chen 0000-0001-6966-7573; Tsung-Yu Lan 0000-0003-0222-422X.

S-Editor: Liu IH L-Editor: A P-Editor: Xu ZH

REFERENCES

- Zhang K, Wang Z, Zhang Z. Comparison of curettage and bone grafting combined with elastic intramedullary nailing vs curettage and bone 1 grafting in the treatment of long bone cysts in children. Medicine (Baltimore) 2019; 98: e16152 [PMID: 31232970 DOI: 10.1097/MD.000000000016152]
- 2 Donaldson S, Chundamala J, Yandow S, Wright JG. Treatment for unicameral bone cysts in long bones: an evidence based review. Orthop Rev (Pavia) 2010; 2: e13 [PMID: 21808696 DOI: 10.4081/or.2010.e13]
- Deventer N, Deventer N, Gosheger G, de Vaal M, Vogt B, Budny T. Current strategies for the treatment of solitary and aneurysmal bone cysts: 3 A review of the literature. J Bone Oncol 2021; 30: 100384 [PMID: 34367902 DOI: 10.1016/j.jbo.2021.100384]
- Zhang P, Zhu N, Du L, Zheng J, Hu S, Xu B. Treatment of simple bone cysts of the humerus by intramedullary nailing and steroid injection. 4 BMC Musculoskelet Disord 2020; 21: 70 [PMID: 32019514 DOI: 10.1186/s12891-020-3054-6]
- Sung AD, Anderson ME, Zurakowski D, Hornicek FJ, Gebhardt MC. Unicameral bone cyst: a retrospective study of three surgical treatments. 5



Clin Orthop Relat Res 2008; 466: 2519-2526 [PMID: 18679761 DOI: 10.1007/s11999-008-0407-0]

- Ahn JI, Park JS. Pathological fractures secondary to unicameral bone cysts. Int Orthop 1994; 18: 20-22 [PMID: 8021063 DOI: 6 10.1007/BF001801731
- Chuo CY, Fu YC, Chien SH, Lin GT, Wang GJ. Management strategy for unicameral bone cyst. Kaohsiung J Med Sci 2003; 19: 289-295 7 [PMID: 12873037 DOI: 10.1016/S1607-551X(09)70475-9]
- Erol B, Onay T, Topkar OM, Tokyay A, Aydemir AN, Okay E. A comparative study for the treatment of simple bone cysts of the humerus: 8 open curettage and bone grafting either without instrumentation or with intramedullary nailing. J Pediatr Orthop B 2017; 26: 5-13 [PMID: 27341120 DOI: 10.1097/BPB.000000000000353]
- 9 Neer CS 2nd, Francis KC, Marcove RC, Terz J, Carbonara PN. Treatment of unicameral bone cyst. A follow-up study of one hundred seventyfive cases. J Bone Joint Surg Am 1966; 48: 731-745 [PMID: 15580740]
- 10 Hou HY, Wu K, Wang CT, Chang SM, Lin WH, Yang RS. Treatment of unicameral bone cyst: a comparative study of selected techniques. J Bone Joint Surg Am 2010; 92: 855-862 [PMID: 20360508 DOI: 10.2106/JBJS.I.00607]
- 11 Canavese F, Samba A, Rousset M. Pathological fractures in children: Diagnosis and treatment options. Orthop Traumatol Surg Res 2016; 102: S149-S159 [PMID: 26774903 DOI: 10.1016/j.otsr.2015.05.010]
- 12 Stanton RP, Abdel-Mota'al MM. Growth arrest resulting from unicameral bone cyst. J Pediatr Orthop 1998; 18: 198-201 [PMID: 9531401]
- Gupta AK, Crawford AH. Solitary bone cyst with epiphyseal involvement: confirmation with magnetic resonance imaging. A case report and 13 review of the literature. J Bone Joint Surg Am 1996; 78: 911-915 [PMID: 8666611]
- Ovadia D, Ezra E, Segev E, Hayek S, Keret D, Wientroub S, Lokiec F. Epiphyseal involvement of simple bone cysts. J Pediatr Orthop 2003; 14 23: 222-229 [PMID: 12604955]
- 15 Cohen J. Unicameral bone cysts. a current synthesis of reported cases. Orthop Clin North Am 1977; 8: 715-736 [PMID: 335313]
- Violas P, Salmeron F, Chapuis M, Sales de Gauzy J, Bracq H, Cahuzac JP. Simple bone cysts of the proximal humerus complicated with 16 growth arrest. Acta Orthop Belg 2004; 70: 166-170 [PMID: 15165020]
- Haims AH, Desai P, Present D, Beltran J. Epiphyseal extension of a unicameral bone cyst. Skeletal Radiol 1997; 26: 51-54 [PMID: 9040144 17 DOI: 10.1007/s002560050191]
- Reif TJ, Matthias J, Fragomen AT, Rozbruch SR. Limb Length Discrepancy and Angular Deformity due to Benign Bone Tumors and Tumor-18 like Lesions. J Am Acad Orthop Surg Glob Res Rev 2021; 5 [PMID: 33720060 DOI: 10.5435/JAAOSGlobal-D-20-00214]
- 19 Lee FY, Schoeb JS, Yu J, Christiansen BD, Dick HM. Operative lengthening of the humerus: indications, benefits, and complications. J Pediatr Orthop 2005; 25: 613-616 [PMID: 16199941 DOI: 10.1097/01.bpo.0000164868.97060.bb]



WJCC | https://www.wjgnet.com



Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

