# World Journal of *Clinical Cases*

World J Clin Cases 2024 March 6; 12(7): 1196-1381





Published by Baishideng Publishing Group Inc

W J C C World Journal of Clinical Cases

# Contents

# Thrice Monthly Volume 12 Number 7 March 6, 2024

# **EDITORIAL**

1196 Relevance of sleep for wellness: New trends in using artificial intelligence and machine learning Nag DS, Swain A, Sahu S, Chatterjee A, Swain BP

# **MINIREVIEWS**

1200 Expect the unexpected: Brown tumor of the mandible as the first manifestation of primary hyperparathyroidism

Majic Tengg A, Cigrovski Berkovic M, Zajc I, Salaric I, Müller D, Markota I

1205 Research progress in spasmodic torticollis rehabilitation treatment

Zhang S, Zeng N, Wu S, Wu HH, Kong MW

# **ORIGINAL ARTICLE**

# **Clinical and Translational Research**

1215 Investigating the causal associations between five anthropometric indicators and nonalcoholic fatty liver disease: Mendelian randomization study

Xiao XP, Dai YJ, Zhang Y, Yang M, Xie J, Chen G, Yang ZJ

1227 Causal role of immune cells in obstructive sleep apnea hypopnea syndrome: Mendelian randomization study

Zhao HH. Ma Z. Guan DS

# **Case Control Study**

1235 Significant risk factors for intensive care unit-acquired weakness: A processing strategy based on repeated machine learning

Wang L, Long DY

# **Retrospective Cohort Study**

1243 Perioperative and long-term results of ultrasonography-guided single- and multiple-tract percutaneous nephrolithotomy for staghorn calculi

Cheng RX, Dai N, Wang YM, Qi P, Chen F

# **Retrospective Study**

Clinical characteristics of testicular torsion and factors influencing testicular salvage in children: A 12-year 1251 study in tertiary center

Gang XH, Duan YY, Zhang B, Jiang ZG, Zhang R, Chen J, Teng XY, Zhang DB



Contents

# Thrice Monthly Volume 12 Number 7 March 6, 2024

# **META-ANALYSIS**

1260 Effectiveness of sensory integration therapy in children, focusing on Korean children: A systematic review and meta-analysis

Oh S, Jang JS, Jeon AR, Kim G, Kwon M, Cho B, Lee N

1272 Safety and efficacy comparison of remimazolam and propofol for intravenous anesthesia during gastroenteroscopic surgery of older patients: A meta-analysis

Li FZ, Zhao C, Tang YX, Liu JT

# **CASE REPORT**

- 1284 Sporadic gastrinoma with refractory benign esophageal stricture: A case report Chen QN, Bai BQ, Xu Y, Mei Q, Liu XC
- 1290 Efficacy of borneol-gypsum in skin regeneration and pain control in toxic epidermal necrolysis: A case report

Yang LW, Zhang LJ, Zhou BB, Lin XY, Chen YT, Qin XY, Tian HY, Ma LL, Sun Y, Jiang LD

1296 Extended survival with metastatic pancreatic cancer under fruquintinib treatment after failed chemotherapy: Two case reports

Wu D, Wang Q, Yan S, Sun X, Qin Y, Yuan M, Wang NY, Huang XT

- 1305 Reconstruction of cervical necrotizing fasciitis defect with the modified keystone flap technique: Two case reports Cho W, Jang EA, Kim KN
- 1313 Reversal of complete atrioventricular block in dialysis patients following parathyroidectomy: A case report Xu SS, Hao LH, Guan YM
- 1320 Treatment of bilateral developmental dysplasia of the hip joint with an improved technique: A case report Yu XX, Chen JY, Zhan HS, Liu MD, Li YF, Jia YY
- 1326 Misdiagnosis of synovial sarcoma - cellular myofibroma with SRF-RELA gene fusion: A case report Zhou Y. Sun YW. Liu XY. Shen DH
- 1333 Heterochronous multiple primary prostate cancer and lymphoma: A case report Liang JL, Bu YQ, Peng LL, Zhang HZ
- 1339 Cardiac remodeling in patients with atrial fibrillation reversing bradycardia-induced cardiomyopathy: A case report Gao DK, Ye XL, Duan Z, Zhang HY, Xiong T, Li ZH, Pei HF

1346 Microsurgical management of radicular cyst using guided tissue regeneration technique: A case report Gómez Mireles JC, Martínez Carrillo EK, Alcalá Barbosa K, Gutiérrez Cortés E, González Ramos J, González Gómez LA, Bayardo González RA, Lomelí Martínez SM

1356 Delayed neurological dysfunction following posterior laminectomy with lateral mass screw fixation: A case report and review of literature

Yan RZ, Chen C, Lin CR, Wei YH, Guo ZJ, Li YK, Zhang Q, Shen HY, Sun HL



Contor	World Journal of Clinical Cases
Conter	Thrice Monthly Volume 12 Number 7 March 6, 2024
1365	Translocation of a fish spike from the pharynx to the thyroid gland: A case report
	Li D, Zeng WT, Jiang JG, Chen JC
1371	Double plasma molecular adsorption system for Stevens-Johnson syndrome/toxic epidermal necrolysis: A case report
	Tan YW, Liu LP, Zhang K

# **LETTER TO THE EDITOR**

Enhancing competency of clinical research nurses: A comprehensive training and evaluation framework 1378 Liu YX, Xu Y



# Contents

Thrice Monthly Volume 12 Number 7 March 6, 2024

# **ABOUT COVER**

Peer Reviewer of World Journal of Clinical Cases, Narendra Pamidi, PhD, Assistant Professor, Department of Anatomy, Melaka Manipal Medical College, Karnataka 576104, India. narendra.pamidi@gmail.com

# **AIMS AND SCOPE**

The primary aim of World Journal of Clinical Cases (WJCC, World J Clin Cases) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

# **INDEXING/ABSTRACTING**

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

# **RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Si Zhao; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang,

NAME OF JOURNAL World Journal of Clinical Cases	INSTRUCTIONS TO AUTHORS https://www.wjgnet.com/bpg/gerinfo/204
ISSN	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 2307-8960 (online)	https://www.wjgnet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
April 16, 2013	https://www.wjgnet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Thrice Monthly	https://www.wjgnet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT
Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati	https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
https://www.wjgnet.com/2307-8960/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS
March 6, 2024	https://www.wjgnet.com/bpg/GerInfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2024 Baishideng Publishing Group Inc	https://www.f6publishing.com

© 2024 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: office@baishideng.com https://www.wjgnet.com



W J C C World Journal of Clinical Cases

Submit a Manuscript: https://www.f6publishing.com

World J Clin Cases 2024 March 6; 12(7): 1205-1214

DOI: 10.12998/wjcc.v12.i7.1205

ISSN 2307-8960 (online)

MINIREVIEWS

# Research progress in spasmodic torticollis rehabilitation treatment

Shuang Zhang, Ni Zeng, Shuang Wu, Hui-Hui Wu, Mo-Wei Kong

Specialty type: Rehabilitation

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

# Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): 0 Grade C (Good): 0 Grade D (Fair): D Grade E (Poor): 0

P-Reviewer: Bernardes A, Portugal

Received: October 21, 2023 Peer-review started: October 21, 2023 First decision: December 28, 2023 Revised: December 31, 2023 Accepted: February 6, 2024 Article in press: February 6, 2024 Published online: March 6, 2024



Shuang Zhang, Ni Zeng, Shuang Wu, Hui-Hui Wu, Department of Rehabilitation, The Affiliated Hospital of Guizhou Medical University, Guiyang 550018, Guizhou Province, China

Mo-Wei Kong, Department of Cardiology, Guiqian International General Hospital, Guiyang 550018, Guizhou Province, China

Corresponding author: Mo-Wei Kong, MD, Doctor, Department of Cardiology, Guiqian International General Hospital, No. 1 Dongfeng Avenue, Guiyang 550018, Guizhou Province, China. 1600181272@qq.com

# Abstract

Spasmodic torticollis (ST) is a focal dystonia that affects adults, causing limited muscle control and impacting daily activities and quality of life. The etiology and curative methods for ST remain unclear. Botulinum toxin is widely used as a firstline treatment, but long-term usage can result in reduced tolerance and adverse effects. Rehabilitation therapy, with its minimal side effects and low potential for harm, holds significant clinical value. This article explores the effectiveness of adjunctive therapies, including exercise therapy, transcranial magnetic stimulation, shockwave therapy, neuromuscular electrical stimulation, vibration therapy, electromyographic biofeedback, and acupuncture, in the treatment of ST. The aim is to provide clinicians with additional treatment options and to discuss the efficacy of rehabilitation therapy for ST.

Key Words: Spasmodic torticollis; Rehabilitation therapy; Botulinum toxin; Exercise therapy; Adjunctive therapy; Effectiveness

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Spasmodic torticollis (ST) is a focal dystonia characterized by involuntary contractions of the neck muscles, significantly impacting daily activities and quality of life. Botulinum toxin is a widely used first-line treatment for ST, but long-term use can lead to reduced efficacy and potential side effects. Surgical interventions may have associated complications. Rehabilitation therapy, including exercise therapy, has potential clinical value as a low-risk treatment option. It can be used as an adjunctive therapy for ST, showing efficacy in improving clinical outcomes and reducing tolerance to botulinum toxin.



Citation: Zhang S, Zeng N, Wu S, Wu HH, Kong MW. Research progress in spasmodic torticollis rehabilitation treatment. *World J Clin Cases* 2024; 12(7): 1205-1214 URL: https://www.wjgnet.com/2307-8960/full/v12/i7/1205.htm DOI: https://dx.doi.org/10.12998/wjcc.v12.i7.1205

# INTRODUCTION

Spasmodic torticollis (ST), a focal dystonia affecting adults, is typified by involuntary muscle contractions in the neck. Profoundly impacting daily activities and quality of life, the etiology of ST remains as yet undefined, and there is no affirmed cure[1-4]. Recurrent botulinum toxin injections are currently the most pervasive first-line therapeutic approach for ST, but long-term usage can foster lowered patient tolerance, thus causing the alleviating effects on muscle spasms to gradually diminish and potentially result in side effects. Surgical treatments are opted for when the efficacy of botulinum toxin and oral medications fall short, but these treatments may lead to associated complications[1-8]. Consequently, rehabilitation therapy, with its minor side effects and low risk, exhibits substantial clinical value. Rehabilitation approaches described in this article, including exercise therapy, transcranial magnetic stimulation, shockwave therapy, neuromuscular electrical stimulation, vibration therapy, electromyographic biofeedback, and acupuncture, are potentially advantageous as adjunctive therapies for ST. However, their efficacies are currently underexplored. The goal of this article is to elucidate the effectiveness of these specific rehabilitation treatment regimens for ST, thereby equipping clinicians with a broader spectrum of therapeutic options.

# **EXERCISE THERAPY**

Exercise therapy, a significant component of physical therapy, employs specific movements or professional equipment for active and/or passive exercise training to regain normal bodily functions[9]. The management of ST may involve techniques for joint mobility, soft tissue stretching, strength exercises, aerobic training, and balance training. Exercise therapy is less frequently used in isolation for alleviating muscle spasms in ST patients, but it is often combined with botulinum toxin treatment. As an adjunctive therapy for ST, it has demonstrated beneficial efficacy and could offer a therapeutic direction for clinical practitioners[10-15], as shown in Table 1. Further studies have revealed that exercise therapy can reduce the dosage of subsequent botulinum toxin injections and maintain its therapeutic effect for a longer period[13,16]. In light of the established efficacy of botulinum toxin in treating ST patients, clinicians might also consider integrating exercise therapy to improve the clinical outcome and mitigate the problem of reduced tolerance resulting from long-term botulinum toxin injections.

# TRANSCRANIAL MAGNETIC STIMULATION THERAPY

Transcranial magnetic stimulation (TMS) is a non-invasive, painless brain stimulation technique, capable not only of investigating the functional roles of specific brain regions by interfering with their neurons[17] but also of relieving muscle tone disorders by modulating the excitability and plasticity of pathogenic brain areas [18]. Increasing studies have pointed to a link between abnormal muscle tone and the cerebellum[19-21]. TMS can activate Purkinje cells in the posterior cerebellum (the VIII lobule and crus II). Such activation inhibits the dentate nucleus, which delivers excitatory responses through synaptic relays in the ventrolateral thalamus to the contralateral primary motor cortex (M1), resulting in suppression of the contralateral M1 due to a decrease in dentato-thalamo-cortical excitation - this form of cortical inhibition is referred to as cerebellar inhibition[22]. TMS can reduce muscle tone and alleviate spasms by reducing the excitability of primary motor cortex through stimulating the cerebellum[23]. Many studies have shown the efficacy of TMS in treating different spasms, as shown in Table 2[24-29]. An experiment comparing TMS and botulinum toxin therapy for the treatment of ST found the latter superior; however, it is worth noting that post-botulinum toxin injection, half of the patients experienced reversible neck weakness and swallowing difficulties, while no adverse reactions were apparent after TMS treatment, indicating good safety[30]. Yang et al[31] established that the use of TMS in patients with ST results in a shortened resting period for the orbicularis oris, orbicularis oculi, and sternocleidomastoid muscles compared to the normal control group. In addition, patients with torticollis exhibit a delayed onset of the resting period for the sternocleidomastoid muscle, this substantiates that in patients with localized muscular tone disorders like ST, there is a decline in the inhibitory functions of the motor cortex and the presence of generalization, thereby affirming the efficacy of TMS in treating ST. If a patient with ST responds poorly to botulinum toxin therapy or refuses this treatment, given the non-invasive nature of TMS, we should consider its broader application to ensure patient efficacy and improve their quality of life.

Raisbildeng® WJCC | https://www.wjgnet.com

Table 1 Efficacy of Botox combined with exercise therapy in neck dystonia						
Ref.	Groups	Specific content of intervention	Intervention time	Evaluation indicators	Research conclusion	
Hu <i>et al</i> [ <mark>10</mark> ]	C: Botox group; E: Botox group + exercise therapy group	Stretching exercises, active range of motion, and isometric exercises of the neck muscles	15 min/time, 5 d/wk for 6 wk	a, b, c	The spasticity and pain symptoms improved significantly	
Tassor <i>et al</i> [11]	C: Botox group; E: Botox group + exercise therapy group	Massage, stretch, educate posture, strengthen axial muscles	For 60-90 min/time, for 2 wk	c, d, e	Remission of spasticity symptoms was similar in both groups, and pain and daily living ability were significantly improved in group E	
van den Dool <i>et al</i> [ <mark>12</mark> ]	C: Botox + conven- tional PT group E: Botox + standardized PT group	Muscle stretching, passive neck activity, and training that has been found to be related to neurorehabilitation and motor learning	Family exercise: 10-15 min/time, 5-10 times/d. PT sessions twice/wk and once/wk in later stages. One year	b, c, d, f, g, h, i	Both groups were effective for spasticity and pain. Group E reduced patients' dependence on healthcare providers and reduced medical costs for this patient group	
Queiroz et al[13]	C: Botox group; E: Botox + exercise therapy group	Motor stretching, passive, and active cervical loosening of the cervical spine	25 min/time, 5 d/wk for 4 wk	b, c	Group E showed significant improvement in daily living capacity and subjective pain	
Stankovic et al[14]	C: Exercise therapy group, E: Botox + exercise therapy group	Increase range of motion exercise, muscle stretching, occupational and functional therapy	5 d/wk for 2 wk	c, d	Group E showed significant improvement in pain, torticollis and disability scales	
Guo et al [15]	C: Botox group, E: Botox + rehabil- itation treatment group	Muscle pulling and relaxation training, exercise relearning training, lateral flexion movement in each horizontal axis direction and vertical axis expansion movement in passive and active modes	757 min/time, 5 d/wk, for 4 wk	c, d	In group E, the scores of physiology, physiological function, physical pain, vitality and mental health were significantly improved compared with those before treatment	

"E" represents the test group; "C" represents the control group. a: The Visual Analogue Scale; b: SF-36 Quality of Life Scale; c: Toronto Western Torticollis Rating Scale; d: Torticollis Assessment scale; e: Daily Performance Scale; f: Functional Disability Questionnaire; g: Overall Clinical Impression-Disease severity Scale; h: Overall Clinical Impression-Improvement Scale; i: Cranio-cervical Dystonia Questionnaire.

# Table 2 Efficacy of rTMS therapy for dystonia

Ref.	Transcranial magnetic prescription	Evaluation method	Curative effect
Bradnam <i>et al</i> [ <mark>24</mark> ]	Five d/wk from Monday to Friday, 10 times	a, b	Cervical dystonia symptoms, pain and quality of life
Hao <i>et al</i> [25]	Frequency was 10 Hz, 30 min/time for 7 d	c, d, e, f	Significantly relieved muscle tension spasm and improved quality of life
Şan et al[26]	Frequency 5 Hz, 15 min/time, 10 for 2 wk	c, g, h, i	Spasticity and the frequency of the seizures improved significantly
Zhao et al[27]	Frequency 1 Hz, 1/d, 6/wk and continuous treatment for 4 wk	c, f, j	Improved limb spasm, restored limb movement function, and improved daily living ability
Cheng et al[28]	The stimulation frequency was 1 Hz, each sequence with 10 pulses at 2 s intervals for a total of 1200 pulses, 1/d, 5 times/wk, and 8 wk as 1 session	c, j, k	Improved the degree of muscle tone control, alleviated muscle spasm, and improved motor dysfunction
Yuan et al[ <mark>29</mark> ]	The stimulation frequency was 1 Hz with 120 stimulus trains of 2 s between 10 pulses for a total of 1200 pulses. Once/d for 5 d per wk for 8 wk	c, j, k	Effectively relieved the degree of spasm, improved the motor function and daily life activities

a: The Toronto Western Torticollis Rating Scale; b: Cranio-cervical Dystonia Questionnaire; c: Modified Ashworth Scale; d: Uniform Wilson's Disease Rating Scale; e: Dystonia Scale; f: Activities of daily living; g: Penn Spasticity Frequency Scale; h: Visual analogue scale; i: Quality of Life 54; j: Fugl-Meyer motor function rating; k: Modified Barthel quantity scale.

Zaisbideng® WJCC | https://www.wjgnet.com

# EXTRACORPOREAL SHOCK WAVE THERAPY

Extracorporeal Shock Wave Therapy (ESWT) is a series of high-energy single-pulse mechanical waves, notable for high pressure, fast rise, short action period, and three-dimensional propagation, the speed of which increases with pressure. It can be divided into focused ESWT (fESWT) and radial ESWT (rESWT)[32]. Although the mechanism of ESWT in spasmodic treatment is still unclear, it may be related to the following aspects: (1) ESWT induces the production of nitric oxide, which increases neovascularization of muscles and tendons, thereby improving muscle stiffness; (2) ESWT may reduce the excitability of motor neurons by vibration stimulation of the tendons, thereby reducing muscle tone; (3) ESWT can reduce the amount of acetylcholine at the neuromuscular junction, causing nerve conduction disorders; and (4) ESWT can cause alternating biochemical reactions between metabolism and proliferation, affecting muscle fibrosis and rheological properties[33]. Recent research has revealed that radial shock wave therapy can effectively reduce the degree of neck muscle spasm in patients with ST, maintain neck position activity, and enhance their daily living ability[34]. Currently, the mechanism of ESWT in the treatment of spasms has been posited from perspectives such as neuromuscular junction and muscle alterations; however, there is still a lack of concrete molecular biological evidence such as signal pathways and genes[35]. Although there is still no definite plan for precise positioning and energy level/density for ESWT, its effectiveness in treating spasms has been confirmed, as shown in Table 3[36-42]. This necessitates further clinical exploration to provide more precise treatment for patients.

# NEUROMUSCULAR ELECTRICAL STIMULATION

Neuromuscular Electrical Stimulation (NMES) is a non-invasive treatment technique, which when applied to spastic muscles, causes them to contract violently. To prevent muscle damage, the Golgi sensory organs are stimulated; impulses enter the spinal cord, then via interneurons to the corresponding anterior horn cells, reflexively inducing inhibition of the spastic muscle itself, thereby relieving spasms[43]. Studies have shown that the treatment of ST patients with NMES is significantly effective, and its short-term effect is superior to local injection of type A botulinum toxin[44]. Although research on NMES treatment for ST is currently limited, it has achieved significant efficacy in treating post-stroke spasm and spastic cerebral palsy [45-48], providing more options for clinical spasm treatment.

# VIBRATION THERAPY

Vibration is a type of sinusoidal mechanical oscillation characterized by amplitude, frequency, and phase angle and is categorized into whole-body vibration training (WBV) and focal muscle vibration (FMV) based on the area of effect[49]. During WBV, the patient is positioned on the vibration platform, and repetitive perturbations are allowed to be transferred, enabling modification of the vibration's frequency, amplitude, and direction (vertical displacement or a backand-forth vertical sinusoidal oscillation)[50]. Although the operating mode of this therapy remains unclear, it appears to display its function by stimulating muscle spindles and alpha motor neurons along with short-term metabolic activities [51]. FMV can activate muscle spindles, thereby stimulating the input of Ia fibers, altering the cortical spinal pathways [52]. It can lead to the inhibition of spinal reflexes, thereby suppressing antagonistic muscle neuron circuits, diminishing the excitability of antagonistic muscles, activating the functional muscle, and mitigating symptoms of spasms[53]. Vibration therapy studies for ST patients are limited, but it has maturely evolved in improving spasticity, particularly in patients with post-stroke spasms and cerebral palsy (Table 4)[54-58]. Studies have shown that the application of vibration to head and neck muscles in ST patients resulted in an immediate and significant reduction in involuntary muscle activity [59]. Recent research indicates that following vibration therapy in patients with right torticollis, the frequency and severity of the disturbances in tension posture significantly reduced, suggesting its effectiveness in spasm treatment[60]. In clinical practice, during the early stages of treatment, patients may experience muscle tension triggered by vibratory stimulation. At this juncture, an assistant is required to help the patient maintain the original passive stretching position and angle, whilst informing the patient to relax as much as possible. The vibratory therapy is highly operational and yields minimal adverse reactions, thus it is considered to be greatly promotable in the clinical context. It aims at alleviating as much pain and spasms in patients as possible and enhancing the patient's quality of life.

# ELECTROMYOGRAPHIC BIOFEEDBACK THERAPY

Electromyographic biofeedback is a method of retraining muscles by transforming the electromyographic signals in muscles into visual and auditory cues to form a new feedback system[61]. Electromyographic biofeedback therapy uses specific low-frequency pulses exciting the nerves or muscles under a preset stimulation program. Interacting with the central nervous system, endocrine system, and immune system, it consciously controls psychological processes to influence physiological processes, thereby causing muscle relaxation and reducing tension in spastic muscles[62]. With its non-invasive nature, high sensitivity, and objectivity, electromyographic biofeedback is often used in rehabilitation of muscles with weakened force or impaired control, such as damage to upper motor neurons (Table 5)[63-68], and has shown promising results. Training can improve muscle tension and activity, and enhance coordination between muscles. Studies have shown[69] that motor learning techniques focused on biofeedback can restore sensorimotor, altered body



# Table 3 Efficacy of shock wave treatment on post-stroke spasticity

Ref.	Position	Prescription	Evaluation method	Curative effect
Li et al <mark>[36</mark> ]	Upper limb	5 consecutive intervals with 4 d. 6000 pulses at pressure 1.2-1.4 bar at a frequency of 18 Hz	a, b, c, d	Clearly improved upper limb flexion spasm and significantly reduced pain
Yoldas <i>et al</i> [37]	Lower limbs	Twice/wk for 2 wk (4 times in total), 1500 pulses, pressure of 2 bar, frequency of 10 Hz	a, b, e, f, g	A marked reduction in plantarflexor spasticity, no significant improvement in the 6-minute walk test and activity range
Mihai et al [ <mark>38</mark> ]	Lower limbs	Once/wk for 2 wk, 2000 pulses at 10 Hz and energy density of 60 mJ	a, c, h, i, j	The degree of spasticity and pain intensity were significantly reduced, improved ankle range of motion, balance and gait
Wang et al [ <mark>39</mark> ]	Upper and lower limbs	Twice/wk for 4 wk, 1500-2500 pulses/site, pressure of 2.0-2.5 bar, frequency of 5-8 Hz	a, d, i	Relieved muscle tension of hemiplegic limb and improved interjoint coordination
Li et al[40]	Lower limbs	Three times/wk (every other day) for 4 wk, 2000 pulses per site with impact intensity of 1.5 bar at 10 Hz	a, d, k	Improved the degree of muscle spasm in the lower limbs, and improved lower limb function and balance function
He <i>et a</i> l[ <mark>41</mark> ]	Lower limbs	3 times/wk for a total of 3 wk. Calf muscle: 2000 shocks, shock wave strength 3 bar, frequency 6 Hz; Achilles tendon: 2000 shocks, shock wave strength 2 bar, frequency 9 Hz	a, d, l, k, m, n	Effectively inhibited calf muscle spasm, improved the active range and motor function, and improved the balance function, walking function and postural control ability
Zhao et al [ <mark>42</mark> ]	Upper limb	Once every 3 d, twice/wk for 3 wk at 8 Hz, pressure 1.5-2.0 bar, 1000 pulses per site	a, d, f	Improved the degree of upper limb spasm, improved the motor function and daily living ability

a: Modified Ashworth Scale; b: The modified Tardieu scale; c: Visual analog scale score; d: Fugl-Meyer Motor Function Rating Scale; e: The 6-min walk test; f: Modified Barthel Index scale; g: Measurement of ankle mobility of the affected side; h: Tinetti Assessment tool; i: Functional activity classification assessment; j: A handheld anglometer evaluated the ankle passive range of motion; k: Berg Balance scale; l: Passive and active motion of ankle dorsiflexion; m: 10 m Walking inspection; n: Time in the stand-up-walk test.

# Table 4 Efficacy of vibration therapy on spasticity

Ref.	Prescription	Evaluation method	Curative effect
Bao <i>et al</i> [54]	The frequency was 36.7 Hz, and the total duration of treatment was 10 min, once/d, 5 times/wk for 3 consecutive wk	b, c, d	Effectively reduced the spasm muscle tension, improved its elasticity and hardness, improved limb movement function
Yin <i>et al</i> [55]	Frequency 12 Hz, amplitude 4 mm, each lasting 15 min, 5 times/wk for 12 wk	e, f, g	Effectively improved ankle motion, balance function and lower limb movement ability in children with spastic diplegia
Li et al[ <mark>56</mark> ]	Frequency 30 Hz, 0.5 mm amplitude once/d 2 min each time and 2 min rest interval. Four-week treatment course with a total of two courses	a, d, h, i	Reduced ankle plantar flexor spasm, improved walking ability, and improved activities of daily living
Katusic et al[57]	Frequency 40 Hz, 20 min each time, twice/wk for 12 wk	a, j	Reduced the level of spasticity and improved the gross motor capacity in children with spastic cerebral palsy
Cai <i>et al</i> [58]	Treatment frequency: First 12 Hz, second 18 Hz, and third 21 Hz. 3 min/time for 9 min, training for 5 d/wk and 2 d rest for 8 wk	f, j	Effectively reduced muscle tension, improved muscle strength, abnormal posture and postural control ability in spastic cerebral paralysis

a: Modified Ashworth scale; b: Modified Tardieu scale; c: Myoton-3 muscle state detector assessment; d: Fugl-Meyer motor function rating scale; e: Scale of ankle dorsiflexion; f: Berg, balance scale; g: Gross motor function for cerebral palsy; h: Gait analysis; i: Modified Barthel index scale; j: Gross motor function (GMFM-88).

perception and motor control, and improve quality of life in patients with ST. It provides an effective supplementary regime for the treatment of ST.

# OTHER TREATMENTS

Other treatments regarding affected muscle groups include acupuncture therapy, as a single target adverse stimulus, might induce immediate facilitation in spasmodic muscles. However, when considered holistically, it acts on multiple



Raishideng® WJCC | https://www.wjgnet.com

### Table 5 Efficacy of electromyography biofeedback therapy on muscle spasticity

Ref.	Disease name	Intervention study	Frequency and duration of the intervention	Outcome indicators	Curative effect
Gulseren <i>et al</i> [63]	Post-stroke spasm	E: EMG-BF + conventional treatment; C: Conventional therapy	1/d, 5/wk, 20 min/time for 3 wk	a, d, e	Improvements in joint mobility, muscle strength, muscle tone, and function
Xie <i>et al</i> [64]	Post-stroke spasm	E: EMG-BF + conventional treatment; C: Conventional therapy	2 times/d, 5 times/wk, 15-20 min/time for 12 wk	b, c	Improved neurological function, motor function and quality of life in patients
Wang et al[ <mark>65</mark> ]	Post-stroke spasm	E: EMG-BF + rehabilitation training; C: Rehabilitation training	1 time/d, 5 times/wk, 30 min/time for 8 wk	a, c, d	Effectively reduced wrist spasm in patients with early cerebral infarction
Shi <i>et al</i> [ <mark>66</mark> ]	Post-stroke spasm	E: EMG-BF + rehabilitation training; C: Rehabilitation training	2 times/d, 30 min/time for 8 wk	a, c	Reduced patient spasticity level and improved motor function
Xu et al[67]	Spastic cerebral palsy	E: EMG-BF + rehabilitation training; C: Rehabilitation training	20-30 min/time, once/d, 6 times/wk for 12 wk	f, j, h, i	Improved gross motor function and serological indicators in children, and improved the development quotient and self-care ability score
Zhang et al[68]	Cerebral palsy	E: EMG-BF + rehabilitation training; C: rehabilitation training	20 min/time, once/d, 5 times/wk for 12 wk	a	Significantly improved exercise capacity and quality of life

"E" represents the test group; "C" represents the control group; EMG-BF: Electromyographic biofeedback; CPM: Continuous passive movement; a: Modified Ashworth score; b: Clinical spasm index; c: Fugl-Meyer motor function score; d: Degree of active joint activity; e: Surface EMG value; f: Gross motor function assessment table (GMFM-88); g: Daily life performance scale (ADL); h: Gesell pediatric neuropsychological development scale; i: Serological index.

targets in the body, which might elicit corresponding feedback and adjustment relative to the local spasm, and the longterm effect might inhibit the spasm[70]. Studies have shown that acupuncture could serve as an adjuvant therapy, helping ST patients reduce the frequency of botulinum toxin usage and alleviate muscle tension[71,72]. Multiple studies suggest that acupuncture therapy can relieve muscle spasm in ST patients, and as it has no side effects, it provides a new treatment option for ST treatment[73-76].

High-intensity laser treatment has photodynamic, photothermal, and photochemical properties, with several therapeutic effects including bio-stimulation, pain relief, anti-inflammatory effects, surface thermal effects, and muscle relaxation[77]. In one case of acute torticollis in a child caused by rotatory atlantoaxial subluxation, high-intensity laser therapy significantly alleviated muscle spasms and neck pain. This is mainly because high-power pulse emissions, believed to have photodynamic effects, can relax muscle spasms by massaging soft tissue structures, thereby stimulating deeper tissues[78]. Although there are few ST cases treated with high-intensity laser therapy, its functions of muscle relaxation and pain relief provide a new adjunct treatment for clinical practice.

# OUTLOOK

The etiology of ST remains unclear, and is currently believed to be associated with various factors including genetics and the environment, and their interactions. In the epidemiology of muscle tension disorders, idiopathic isolated dystonia is the most common, with neck dystonia being the most prevalent form (with 3 to 13 cases per 100000 persons)[79]. Genetically, dystonia presents in multiple forms with inherited isolated dystonia predominately showing an autosomal dominant inheritance (such as DYT6, DYT1)[80]. DYT-THAP1 (DYT6) is a common early-onset isolated dystonia, typically manifesting in children and adolescents[81], primarily displaying as cranio-cervical dystonia, with slow or no progress[82]. DYT-TOR1A (DYT1) is the most common early-onset generalized dystonia, involving the GAG — internal deletion in TOR1A, particularly common in individuals of German-Jewish descent[83]. Early symptoms often involve the leg or later the arm, then rapidly generalize while 20% may still remain localized, usually involving writer's cramp[83].

Evidence suggests that functional changes in the dopamine signaling process may be associated with the onset of dystonia[84]. Positron emission tomography showed that the binding rate of dopamine D2 receptors and labeled ligands in the shell nucleus of dystonic patients is lower compared to normal individuals[83,85]. Moreover, two genetic defects have been discovered leading to abnormal dopamine synthesis: Deficiencies in Guanosine-triphosphate Cyclase can result in autosomal dominant inherited dopamine-responsive dystonia (such as DYT5a/GCH1 mutations, DYT5b/TH defects), while Tyrosine Hydroxylase deficiency can result in autosomal recessive inherited dopamine-responsive dystonia[86]. Advances in genetic technology have led to a rapid increase in the number of dystonia genes. The increase in gene numbers is significant as they assist in understanding the genetic causes of inherited dystonia and help link them

Raishidena® WJCC | https://www.wjgnet.com

to shared biological pathways. It appears unlikely there could be a final universal pathway to explain all types of dystonia; instead, multiple pathways can explain specific dystonia. For inherited dystonia, drugs, surgery, and the aforementioned rehabilitation treatments have limited therapeutic efficacy. Future treatments, based on pathological mechanisms, tend to lean more towards gene therapy.

# CONCLUSION

The etiology and mechanism of ST are not yet clear. Currently, there are few articles on the independent use of the above rehabilitation treatments. Most are used in conjunction with botulinum toxin therapy or other rehabilitation measures. In clinical practice, when we encounter patients with complex conditions or poor symptom control, attention should also be paid to whether the patient has genetic factors or abnormal genes. In addition to the use of drugs and surgery, we should also try to combine various rehabilitation methods to solve the patient's problems as much as possible. It is hoped that in the future, gene therapy can be placed on the agenda as soon as possible to solve fundamental problems for patients.

# ACKNOWLEDGEMENTS

We would like to express our gratitude to Xin-Rui Li for their valuable feedback and constructive suggestions on the manuscript.

# FOOTNOTES

Author contributions: Zhang S wrote the manuscript; Kong MW provided crucial suggestions and guidance for the writing; Wu S, Zeng N and Wu HH reviewed and revised the manuscript; all authors read and approved the final manuscript.

**Conflict-of-interest statement:** All the authors declare that they have no competing interests.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

# Country/Territory of origin: China

ORCID number: Shuang Zhang 0009-0003-8865-784X; Mo-Wei Kong 0000-0002-1214-164X.

S-Editor: Liu JH L-Editor: Webster JR P-Editor: Zheng XM

# REFERENCES

- Simpson DM, Hallett M, Ashman EJ, Comella CL, Green MW, Gronseth GS, Armstrong MJ, Gloss D, Potrebic S, Jankovic J, Karp BP, 1 Naumann M, So YT, Yablon SA. Practice guideline update summary: Botulinum neurotoxin for the treatment of blepharospasm, cervical dystonia, adult spasticity, and headache: Report of the Guideline Development Subcommittee of the American Academy of Neurology. Neurology 2016; 86: 1818-1826 [PMID: 27164716 DOI: 10.1212/WNL.00000000002560]
- 2 Erro R, Picillo M, Pellecchia MT, Barone P. Improving the Efficacy of Botulinum Toxin for Cervical Dystonia: A Scoping Review. Toxins (Basel) 2023; 15 [PMID: 37368692 DOI: 10.3390/toxins15060391]
- Awan KH. The therapeutic usage of botulinum toxin (Botox) in non-cosmetic head and neck conditions An evidence based review. Saudi 3 Pharm J 2017; 25: 18-24 [PMID: 28223858 DOI: 10.1016/j.jsps.2016.04.024]
- Boyce MJ, McCambridge AB, Bradnam LV, Canning CG, Verhagen AP. The barriers and facilitators to satisfaction with botulinum 4 neurotoxin treatment in people with cervical dystonia: a systematic review. Neurol Sci 2022; 43: 4663-4670 [PMID: 35593979 DOI: 10.1007/s10072-022-06114-8]
- 5 Solish N, Carruthers J, Kaufman J, Rubio RG, Gross TM, Gallagher CJ. Overview of DaxibotulinumtoxinA for Injection: A Novel Formulation of Botulinum Toxin Type A. Drugs 2021; 81: 2091-2101 [PMID: 34787840 DOI: 10.1007/s40265-021-01631-w]
- Li J, Wang L. Clinical treatment of spasmodic torticollis. Linchuang Waike Zazhi 2019; 27: 840-843 6
- 7 Cheng R, Han YS, Han YZ, Hu JY, Xue BC, Yang RM. Clinical characteristics and therapeutic effect of botulinum toxin type A in 104 patients with spastic torticollis. Zhongyiyao Linchuang Zazhi 2021; 304-309
- 8 Hull M, Anupindi VR, DeKoven M, He J, Bouchard J. Botulinum Toxin Utilization, Treatment Patterns, and Healthcare Costs Among 8 Patients with Spasticity or Cervical Dystonia in the US. Adv Ther 2023; 40: 3986-4003 [PMID: 37414904 DOI: 10.1007/s12325-023-02563-5]
- 9 Feng SJ, Gao H, Chen J, LI YX, Hao YF, Chen SY. Exercise therapy in the application of the motor system related diseases treatment. Zhongguo Yundong Yixue zhazhi 2023; 74-79 [DOI: 10.3760/cma.j.issn.0254-1424.2020.10.025]



- Hu W, Rundle-Gonzalez V, Kulkarni SJ, Martinez-Ramirez D, Almeida L, Okun MS, Wagle Shukla A. A randomized study of botulinum 10 toxin versus botulinum toxin plus physical therapy for treatment of cervical dystonia. Parkinsonism Relat Disord 2019; 63: 195-198 [PMID: 30837195 DOI: 10.1016/j.parkreldis.2019.02.035]
- 11 Tassorelli C, Mancini F, Balloni L, Pacchetti C, Sandrini G, Nappi G, Martignoni E. Botulinum toxin and neuromotor rehabilitation: An 11 integrated approach to idiopathic cervical dystonia. Mov Disord 2006; 21: 2240-2243 [PMID: 17029278 DOI: 10.1002/mds.21145]
- van den Dool J, Visser B, Koelman JH, Engelbert RH, Tijssen MA. Cervical dystonia: effectiveness of a standardized physical therapy 12 program; study design and protocol of a single blind randomized controlled trial. BMC Neurol 2013; 13: 85 [PMID: 23855591 DOI: 10.1186/1471-2377-13-85
- 13 Queiroz MA, Chien HF, Sekeff-Sallem FA, Barbosa ER. Physical therapy program for cervical dystonia: a study of 20 cases. Funct Neurol 2012; 27: 187-192 [PMID: 23402680]
- 14 Stankovic I, Colovic H, Vesna ZD, Stamenovic J, Stanković A, Zlatanović D, Zivkovic D, Stanković T. The effect of physical therapy in the treatment of patients with cervical dystonia with or without concomitant use of botulinum toxin. Vojnosanitetski Pregled. Military-medical and Pharmaceutical Review 2017; 75: 16-16 [DOI: 10.2298/VSP161115016S]
- Guo GH, Zhang QZ, Li Z. Effect of botulinum toxin type A injection combined with rehabilitation intervention on spastic torticollis. 15 Zhonghua Wuli Yixue Yu Kangfu Zazhi 2014; 2: 111-114 [DOI: 10.3760/cma.J.issn.0254-1424.2014.02.008]
- 16 Ramdharry G. Case report: physiotherapy cuts the dose of botulinum toxin. Physiother Res Int 2006; 11: 117-122 [PMID: 16808092 DOI: 10.1002/pri.326]
- Odorfer TM, Homola GA, Reich MM, Volkmann J, Zeller D. Increased Finger-Tapping Related Cerebellar Activation in Cervical Dystonia, 17 Enhanced by Transcranial Stimulation: An Indicator of Compensation? Front Neurol 2019; 10: 231 [PMID: 30930842 DOI: 10.3389/fneur.2019.00231]
- Frey J, Ramirez-Zamora A, Wagle Shukla A. Applications of Transcranial Magnetic Stimulation for Understanding and Treating Dystonia. 18 *Adv Neurobiol* 2023; **31**: 119-139 [PMID: 37338699 DOI: 10.1007/978-3-031-26220-3\_7]
- Grimm K, Prilop L, Schön G, Gelderblom M, Misselhorn J, Gerloff C, Zittel S. Cerebellar Modulation of Sensorimotor Associative Plasticity 19 Is Impaired in Cervical Dystonia. Mov Disord 2023; 38: 2084-2093 [PMID: 37641392 DOI: 10.1002/mds.29586]
- 20 Brighina F, Romano M, Giglia G, Saia V, Puma A, Giglia F, Fierro B. Effects of cerebellar TMS on motor cortex of patients with focal dystonia: a preliminary report. Exp Brain Res 2009; 192: 651-656 [PMID: 18815775 DOI: 10.1007/s00221-008-1572-9]
- Teo JT, van de Warrenburg BP, Schneider SA, Rothwell JC, Bhatia KP. Neurophysiological evidence for cerebellar dysfunction in primary 21 focal dystonia. J Neurol Neurosurg Psychiatry 2009; 80: 80-83 [PMID: 19091711 DOI: 10.1136/jnnp.2008.144626]
- Koch G, Porcacchia P, Ponzo V, Carrillo F, Cáceres-Redondo MT, Brusa L, Desiato MT, Arciprete F, Di Lorenzo F, Pisani A, Caltagirone C, 22 Palomar FJ, Mir P. Effects of two weeks of cerebellar theta burst stimulation in cervical dystonia patients. Brain Stimul 2014; 7: 564-572 [PMID: 24881805 DOI: 10.1016/j.brs.2014.05.002]
- Bologna M, Paparella G, Fabbrini A, Leodori G, Rocchi L, Hallett M, Berardelli A. Effects of cerebellar theta-burst stimulation on arm and 23 neck movement kinematics in patients with focal dystonia. Clin Neurophysiol 2016; 127: 3472-3479 [PMID: 27721106 DOI: 10.1016/j.clinph.2016.09.008]
- Bradnam LV, McDonnell MN, Ridding MC. Cerebellar Intermittent Theta-Burst Stimulation and Motor Control Training in Individuals with 24 Cervical Dystonia. Brain Sci 2016; 6 [PMID: 27886079 DOI: 10.3390/brainsci6040056]
- Hao W, Wei T, Yang W, Yang Y, Cheng T, Li X, Dong W, Jiang H, Qian N, Wang H, Wang M. Effects of High-Frequency Repetitive 25 Transcranial Magnetic Stimulation on Upper Limb Dystonia in Patients With Wilson's Disease: A Randomized Controlled Trial. Front Neurol 2021; 12: 783365 [PMID: 34970214 DOI: 10.3389/fneur.2021.783365]
- San AU, Yılmaz B, Kesikburun S. The Effect of Repetitive Transcranial Magnetic Stimulation on Spasticity in Patients with Multiple 26 Sclerosis. J Clin Neurol 2019; 15: 461-467 [PMID: 31591833 DOI: 10.3988/jcn.2019.15.4.461]
- Zhao J. Effect of low-frequency repetitive transcranial magnetic stimulation on hemiplegic upper limb spasm and limb motor function in 27 stroke patients. Yiyao Yu Baojian 2021; 029: 83-85
- Cheng R, Tang HH, Zhang Y, Chen W. Effect of low-frequency repetitive transcranial magnetic stimulation on spastic dyskinesia after 28 ischemic stroke. Beihua Daxue Xuebao (Natural Science Edition) 2022; 23: 79-83
- 29 Yuan MZ. Multimodal magnetic resonance study of low-frequency repetitive transcranial magnetic stimulation in the treatment of post-29 stroke spasm. Fujian Zhongyiyao Daxue 2020; 8
- Wang L. The study of repetitive transcranial magnetic stimulation in the treatment of focal dystonia. Beijing Xiehe Yixueyuan 2014; 11 30
- Kaji R, Murase N, Urushihara R, Asanuma K. Sensory deficits in dystonia and their significance. Adv Neurol 2004; 94: 11-17 [PMID: 31 145096491
- Kong M, Pan Q, Cheng X, Li J, Gao Y, Tian X. Anthracycline-induced delayed-onset cardiac toxicity: A case report and literature review. Exp 32 Ther Med 2023; 26: 505 [PMID: 37822590 DOI: 10.3892/etm.2023.12204]
- Yang E, Lew HL, Özçakar L, Wu CH. Recent Advances in the Treatment of Spasticity: Extracorporeal Shock Wave Therapy. J Clin Med 33 2021; 10 [PMID: 34682846 DOI: 10.3390/jcm10204723]
- Liang CP, Huang GL, Ding WJ, Wang F, Su B. Radiation type extracorporeal shock wave on the clinical effect of the treatment of type 34 rotating spasmodic torticollis. Zhongguo Kangfu Yixue Zazhi 2023; 38: 105-107 [DOI: 10.3969/j.issn.1001-1242.2023.01.020]
- Xiao J, Liang HJ, Ji BS, Xing GY. Research progress on effects of extracorporeal shock wave on central muscle spasm and muscle 35 metabolism. Zhongguo Yixue Qianyan Zazhi (Electronic Version) 2022; 14: 7-14
- Li G, Yuan W, Liu G, Qiao L, Zhang Y, Wang Y, Wang W, Zhao M, Wang J. Effects of radial extracorporeal shockwave therapy on spasticity 36 of upper-limb agonist/antagonist muscles in patients affected by stroke: a randomized, single-blind clinical trial. Age Ageing 2020; 49: 246-252 [PMID: 31846499 DOI: 10.1093/ageing/afz159]
- Yoldaş Aslan Ş, Kutlay S, Düsünceli Atman E, Elhan AH, Gök H, Küçükdeveci AA. Does extracorporeal shock wave therapy decrease 37 spasticity of ankle plantar flexor muscles in patients with stroke: A randomized controlled trial. Clin Rehabil 2021; 35: 1442-1453 [PMID: 33906450 DOI: 10.1177/02692155211011320]
- Mihai EE, Berteanu M. Early Individualized Approach for a Patient with Spasticity of Stroke Origin. Curr Health Sci J 2021; 47: 608-611 38 [PMID: 35444829 DOI: 10.12865/CHSJ.47.04.20]
- 39 Wang ZY, Ma JJ, Tian XY, Tang S, Liu WW. Effect of extracorporeal shock wave on muscle spasm after stroke. Dianxian Yu Shenjingdian Shenglixue Zazhi 2021; 30: 284-289



- Li YF, Wu TL, Shen J. Extracorporeal shock wave in the application of lower limb spasm after cerebral apoplexy therapy. Anmo Yu Kangfu 40 Yixue 2020; 11: 28 and 30 [DOI: 10.19787/j.issn.1008-1879.2020.03.011]
- Duan H, Lian Y, Jing Y, Xing J, Li Z. Research progress in extracorporeal shock wave therapy for upper limb spasticity after stroke. Front 41 Neurol 2023; 14: 1121026 [PMID: 36846123 DOI: 10.3389/fneur.2023.1121026]
- 42 Zhao HY, Wang LS. Clinical observation of acupuncture combined with extracorporeal shock wave in the treatment of upper limb spasm after stroke. Xiandai Zhongxiyi Jiehe Zazhi 2021; 30: 3162-3165 [DOI: 10.3969/j.issn.1008-8849.2021.28.019]
- Zhou YP, Xu XL, Li MY. Electrical stimulation in the treatment of cerebral palsy application. Beijing Shengwu Yixue Gongcheng 2009; 28: 43 103 [DOI: 10.3969/j.issn.1002-3208.2009.01.025]
- Song Y. A comparative study of transcranial magnetic stimulation, electrical stimulation and botulinum toxin in the treatment of spastic 44 torticollis. Hebei Yike Daxue 2008
- Almutairi SM, Khalil ME, Almutairi N, Alenazi AM. Effects of Neuromuscular Electrical Stimulation on Plantarflexors Spasticity, Gait 45 Performance, and Self-Reported Health Outcomes in People With Chronic Stroke: A Study Protocol for a Double-Blinded Randomized Clinical Trial. Front Neurol 2021; 12: 770784 [PMID: 34925217 DOI: 10.3389/fneur.2021.770784]
- 46 Liu H, Xu DM, Deng XY, Yuan J, Rao A. Effect of low-frequency neuromuscular electrical stimulation on muscle tone recovery in elderly patients with hemiplegic muscle spasm after stroke. Zhongguo Laonianxue Zazhi 2023; 4: 1114-1117 [DOI: 10.3969/j.issn.1005-9202.2023.05.027
- Yang FN. Effect of neuromuscular electrical stimulation combined with head on lower limb muscle tone in patients with spastic hemiplegia 47 after ischemic stroke. Shiyong Zhongxiyi Jiehe Linchuang 2021; 21: 102-103 [DOI: 10.13638/j.issn.1671-4040.2021.11.051]
- 48 Xu DD, Cao HY, Fan YJ, Yan DM, Su M. Comparative Analysis of the Effect of Low-Frequency Repeated Transcranial Magnetic Stimulation and Extracorporeal Shock Wave on Improving the Spasm of Flexor after Stroke. Xunzheng Buchong Yu Tidai Yixue 2021; 2021: 7769581 [DOI: 10.1155/2021/7769581]
- 49 Xiao Y, Xu GX. Application of vibration therapy in limb spasm after stroke. Zhongguo Kangfu Yixue Zazhi 2019; 742-746 [DOI: 10.3969/j.issn.1001-1242.2019.06.026]
- Moggio L, de Sire A, Marotta N, Demeco A, Ammendolia A. Vibration therapy role in neurological diseases rehabilitation: an umbrella review 50 of systematic reviews. Disabil Rehabil 2022; 44: 5741-5749 [PMID: 34225557 DOI: 10.1080/09638288.2021.1946175]
- Naro A, Leo A, Russo M, Casella C, Buda A, Crespantini A, Porcari B, Carioti L, Billeri L, Bramanti A, Bramanti P, Calabrò RS. 51 Breakthroughs in the spasticity management: Are non-pharmacological treatments the future? J Clin Neurosci 2017; 39: 16-27 [PMID: 28262404 DOI: 10.1016/j.jocn.2017.02.044]
- Alashram AR, Padua E, Romagnoli C, Annino G. Effectiveness of focal muscle vibration on hemiplegic upper extremity spasticity in 52 individuals with stroke: A systematic review. NeuroRehabilitation 2019; 45: 471-481 [PMID: 31868686 DOI: 10.3233/NRE-192863]
- Lei J, Zhang C, Gai J, Fan X, Tang J. Deep muscle stimulator in the treatment of post-stroke spasticity: A protocol for systematic review and 53 meta-analysis. Medicine (Baltimore) 2023; 102: e33602 [PMID: 37115051 DOI: 10.1097/MD.00000000033602]
- Bao SR, Lin LH, Shan SR, Yang XP, Liu CL. Effect of electrical deep muscle stimulate on muscle tone, elasticity, and stiffness of biceps 54 brachii in stroke patients. Zhongguo Zuzhi Gongcheng Yanjiu 2021; 20: 3138-3143
- Yin HW, Li HF, Zhang X, Wang H, Ruan WC, Du Y, Che YP. Whole body vibration therapy on children with spastic type double collapsed 55 the impact of lower limb movement function. Zhonghua Wuliyixue Yu Kangfu Zazhi 2019; 9: 752-756
- 56 Li Z, Wang GS, Guo GH. Whole body vibration therapy to treat hemiplegia patients curative effect observation of ankle plantar flexion spasm. Zhongguo Kangfu Yixue Zazhi 2014; 29: 451-454 [DOI: 10.3969/j.issn.1001-1242.2014.05.011]
- Katusic A, Alimovic S, Mejaski-Bosnjak V. The effect of vibration therapy on spasticity and motor function in children with cerebral palsy: a 57 randomized controlled trial. Neuro Rehabilitation 2013; 32: 1-8 [PMID: 23422453 DOI: 10.3233/NRE-130817]
- Cai WL, He QY Chen XF, Zeng XL, Zhang XF. Effect of whole body vibration therapy on gross motor function and balance function of 58 spastic diplegia. Anmo Yu Kangfu Yixue 2018; 9: 8-9
- Leis AA, Dimitrijevic MR, Delapasse JS, Sharkey PC. Modification of cervical dystonia by selective sensory stimulation. J Neurol Sci 1992; 59 110: 79-89 [PMID: 1506873 DOI: 10.1016/0022-510X(92)90013-B]
- 60 Zhu Y, Mahnan A, Konczak J. Vibro-Tactile Stimulation as a Non-Invasive Neuromodulation Therapy for Cervical Dystonia: A Case Study. Annu Int Conf IEEE Eng Med Biol Soc 2021; 2021: 6314-6317 [PMID: 34892557 DOI: 10.1109/EMBC46164.2021.9629717]
- Giggins OM, Persson UM, Caulfield B. Biofeedback in rehabilitation. J Neuroeng Rehabil 2013; 10: 60 [PMID: 23777436 DOI: 61 10.1186/1743-0003-10-60]
- Su HP, Wu JL, Hu ZY. Electromyographic biofeedback therapy in the application of rehabilitation therapy. Shiyong Fuke Neifenmi Dianzi 62 Zazhi 2017; 4: 74-75 [DOI: 10.16484/j.cnki.issn2095-8803.2017.23.050]
- 63 Dost Sürücü G, Tezen Ö. The effect of EMG biofeedback on lower extremity functions in hemiplegic patients. Acta Neurol Belg 2021; 121: 113-118 [PMID: 31898758 DOI: 10.1007/s13760-019-01261-w]
- Xie BL, Zou Q, Tian Y. Clinical study of myoelectric biofeedback combined with rehabilitation training on motor function and quality of life 64 in stroke patients with hemiplegia. Chuanbei Yixueyuan Xuebao 2020; 35: 350-353 [DOI: 10.3969/j.issn.1005-3697.2020.02.045]
- Wang JM, Wang Y, Li KP, Jiang J, Pei S, Zhou DL, Qin HW, Zhu Y, Xu Y, Li F. Effect of myoelectric biofeedback on wrist spasm after 65 early cerebral infarction. Zhonghua Wuli Yixue Yu Kangfu Zazhi 2020: 709-711 [DOI: 10.3760/cma.j.issn.0254-1424.2020.08.009]
- Shi YB, Song DT, Zhang YR, Zhu J. Effect of myoelectric biofeedback therapy combined with PNF training on limb function recovery in 66 stroke patients. Zhonghua Wuli Yixue Yu Kangfu Zazhi 2020: 926-928 [DOI: 10.3760/cma.j.issn.0254-1424.2020.10.016]
- Xu WJ, Yang LL. Application of myoelectric biofeedback therapy in rehabilitation of children with spastic cerebral palsy. Zhongguo Minkang 67 *Yixue* 2022; 82-85 [DOI: 10.3969/j.issn.1002-1949.2018.z1.163]
- Zhang LJ, Wang YG. Electromyographic biofeedback therapy in children with cerebral palsy rehabilitation applications. Shenzhen Zhongxiyi 68 Jiehe Zazhi 2019; 29: 153-154
- Castagna A, Saibene E, Ramella M. How Do I Rehabilitate Patients with Cervical Dystonia Remotely? Mov Disord Clin Pract 2021; 8: 820-69 821 [PMID: 34295949 DOI: 10.1002/mdc3.13212]
- 70 Wang J Cui X. Progress in studies of acupuncture treatment of spastic. Zhongguo Kangfu Yixue Zazhi 2012; 27: 191-193 [DOI: 10.3969/j.issn.1001-1242.2012.02.027]
- Bega D, Park K, Grimone A, Lin F, Ring M. Acupuncture as Adjuvant Therapy for the Management of Cervical Dystonia. Med Acupunct 71 2018; **30**: 198-203 [PMID: 30147821 DOI: 10.1089/acu.2018.1291]



- Yu ZH, Xu SY, Li JH. Therapeutic effect of botulinum toxin type A combined with acupuncture on rotary spastic torticollis. Zhejiang 72 Zhongxiyi Jiehe Zazhi 2017; 27: 675-678 [DOI: 10.3969/j.issn.1005-4561.2017.08.013]
- 73 Huang CY, Zhang QP, Zha BX, Liu XJ, Xing J. Yang Jun using the combination of acupuncture with medicine in the treatment of spasmodic torticollis experience. Anhui Zhongyiyao Daxue Xuebao 2021; 1: 42-44 [DOI: 10.3969/j.issn.2095-7246.2021.01.012]
- Liu H. Spasmodic torticollis acupuncture therapy experience. Guangming Zhongyi 2023; 38: 1971-1973 [DOI: 74 10.3969/j.issn.1003-8914.2023.10.043]
- Xing HZ, Xiao WX, Guo Y, Wen YL, Li B. Treatment of spasmodic torticolium by differential acupuncture. Zhongguo Zhenjiu 2023; 43: 807-75 808 [DOI: 10.13703/J.0255-2930.20221011-0006]
- Liu DZ, Zhang X, Sun N, Zhang R, Feng CW, Wang YL, Yang TS. Cure method of anchor and "five heart hole" in the treatment of spasmodic 76 torticollis clinical research. Zhenjiu Linchuang Zazhi 2021; 5: 4-9
- 77 Szabo DA, Neagu N, Teodorescu S, Predescu C, Sopa IS, Panait L. TECAR Therapy Associated with High-Intensity Laser Therapy (Hilt) and Manual Therapy in the Treatment of Muscle Disorders: A Literature Review on the Theorised Effects Supporting Their Use. J Clin Med 2022; **11** [PMID: 36294470 DOI: 10.3390/jcm11206149]
- Tuan SH, Sun SF, Huang WY, Chen GB, Li MH, Liou IH. Effect of high intensity laser therapy in the treatment of acute atlantoaxial rotatory 78 subluxation: A case report. J Back Musculoskelet Rehabil 2022; 35: 963-969 [PMID: 35068439 DOI: 10.3233/BMR-210133]
- 79 Steeves TD, Day L, Dykeman J, Jette N, Pringsheim T. The prevalence of primary dystonia: a systematic review and meta-analysis. Mov *Disord* 2012; **27**: 1789-1796 [PMID: 23114997 DOI: 10.1002/mds.25244]
- 80 Stephen CD. The Dystonias. Continuum (Minneap Minn) 2022; 28: 1435-1475 [PMID: 36222773 DOI: 10.1212/CON.000000000001159]
- Almasy L, Bressman SB, Raymond D, Kramer PL, Greene PE, Heiman GA, Ford B, Yount J, de Leon D, Chouinard S, Saunders-Pullman R, 81 Brin MF, Kapoor RP, Jones AC, Shen H, Fahn S, Risch NJ, Nygaard TG. Idiopathic torsion dystonia linked to chromosome 8 in two Mennonite families. Ann Neurol 1997; 42: 670-673 [PMID: 9382482 DOI: 10.1002/ana.410420421]
- Cheng FB, Wan XH, Feng JC, Wang L, Yang YM, Cui LY. Clinical and genetic evaluation of DYT1 and DYT6 primary dystonia in China. 82 *Eur J Neurol* 2011; **18**: 497-503 [PMID: 20825472 DOI: 10.1111/j.1468-1331.2010.03192.x]
- 83 Ozelius LJ, Bressman SB. Genetic and clinical features of primary torsion dystonia. Neurobiol Dis 2011; 42: 127-135 [PMID: 21168499 DOI: 10.1016/j.nbd.2010.12.012]
- Ma L, Wan X. Spasmodic torticollis and its diagnosis and treatment. Xiehe Medical Journal 2012; 3: 332-336 [DOI: 84 10.3760/cma.j.issn.1673-4904.2008.29.029]
- Naumann M, Pirker W, Reiners K, Lange KW, Becker G, Brücke T. Imaging the pre- and postsynaptic side of striatal dopaminergic synapses 85 in idiopathic cervical dystonia: a SPECT study using [123I] epidepride and [123I] beta-CIT. Mov Disord 1998; 13: 319-323 [PMID: 9539347 DOI: 10.1002/mds.870130219]
- Perlmutter JS, Stambuk MK, Markham J, Black KJ, McGee-Minnich L, Jankovic J, Moerlein SM. Decreased [18F]spiperone binding in 86 putamen in idiopathic focal dystonia. J Neurosci 1997; 17: 843-850 [PMID: 8987805 DOI: 10.1523/JNEUROSCI.17-02-00843.1997]





# Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

