

World Journal of *Clinical Cases*

World J Clin Cases 2024 March 6; 12(7): 1196-1381



EDITORIAL

- 1196 Relevance of sleep for wellness: New trends in using artificial intelligence and machine learning
Nag DS, Swain A, Sahu S, Chatterjee A, Swain BP

MINIREVIEWS

- 1200 Expect the unexpected: Brown tumor of the mandible as the first manifestation of primary hyperparathyroidism
Majic Teng A, Cigrovski Berkovic M, Zajc I, Salaric I, Müller D, Markota I
- 1205 Research progress in spasmodic torticollis rehabilitation treatment
Zhang S, Zeng N, Wu S, Wu HH, Kong MW

ORIGINAL ARTICLE**Clinical and Translational Research**

- 1215 Investigating the causal associations between five anthropometric indicators and nonalcoholic fatty liver disease: Mendelian randomization study
Xiao XP, Dai YJ, Zhang Y, Yang M, Xie J, Chen G, Yang ZJ
- 1227 Causal role of immune cells in obstructive sleep apnea hypopnea syndrome: Mendelian randomization study
Zhao HH, Ma Z, Guan DS

Case Control Study

- 1235 Significant risk factors for intensive care unit-acquired weakness: A processing strategy based on repeated machine learning
Wang L, Long DY

Retrospective Cohort Study

- 1243 Perioperative and long-term results of ultrasonography-guided single- and multiple-tract percutaneous nephrolithotomy for staghorn calculi
Cheng RX, Dai N, Wang YM, Qi P, Chen F

Retrospective Study

- 1251 Clinical characteristics of testicular torsion and factors influencing testicular salvage in children: A 12-year study in tertiary center
Gang XH, Duan YY, Zhang B, Jiang ZG, Zhang R, Chen J, Teng XY, Zhang DB

META-ANALYSIS

- 1260** Effectiveness of sensory integration therapy in children, focusing on Korean children: A systematic review and meta-analysis
Oh S, Jang JS, Jeon AR, Kim G, Kwon M, Cho B, Lee N
- 1272** Safety and efficacy comparison of remimazolam and propofol for intravenous anesthesia during gastroenteroscopic surgery of older patients: A meta-analysis
Li FZ, Zhao C, Tang YX, Liu JT

CASE REPORT

- 1284** Sporadic gastrinoma with refractory benign esophageal stricture: A case report
Chen QN, Bai BQ, Xu Y, Mei Q, Liu XC
- 1290** Efficacy of borneol-gypsum in skin regeneration and pain control in toxic epidermal necrolysis: A case report
Yang LW, Zhang LJ, Zhou BB, Lin XY, Chen YT, Qin XY, Tian HY, Ma LL, Sun Y, Jiang LD
- 1296** Extended survival with metastatic pancreatic cancer under fruquintinib treatment after failed chemotherapy: Two case reports
Wu D, Wang Q, Yan S, Sun X, Qin Y, Yuan M, Wang NY, Huang XT
- 1305** Reconstruction of cervical necrotizing fasciitis defect with the modified keystone flap technique: Two case reports
Cho W, Jang EA, Kim KN
- 1313** Reversal of complete atrioventricular block in dialysis patients following parathyroidectomy: A case report
Xu SS, Hao LH, Guan YM
- 1320** Treatment of bilateral developmental dysplasia of the hip joint with an improved technique: A case report
Yu XX, Chen JY, Zhan HS, Liu MD, Li YF, Jia YY
- 1326** Misdiagnosis of synovial sarcoma - cellular myofibroblastoma with *SRF-RELA* gene fusion: A case report
Zhou Y, Sun YW, Liu XY, Shen DH
- 1333** Heterochronous multiple primary prostate cancer and lymphoma: A case report
Liang JL, Bu YQ, Peng LL, Zhang HZ
- 1339** Cardiac remodeling in patients with atrial fibrillation reversing bradycardia-induced cardiomyopathy: A case report
Gao DK, Ye XL, Duan Z, Zhang HY, Xiong T, Li ZH, Pei HF
- 1346** Microsurgical management of radicular cyst using guided tissue regeneration technique: A case report
Gómez Mireles JC, Martínez Carrillo EK, Alcalá Barbosa K, Gutiérrez Cortés E, González Ramos J, González Gómez LA, Bayardo González RA, Lomeli Martínez SM
- 1356** Delayed neurological dysfunction following posterior laminectomy with lateral mass screw fixation: A case report and review of literature
Yan RZ, Chen C, Lin CR, Wei YH, Guo ZJ, Li YK, Zhang Q, Shen HY, Sun HL

1365 Translocation of a fish spike from the pharynx to the thyroid gland: A case report

Li D, Zeng WT, Jiang JG, Chen JC

1371 Double plasma molecular adsorption system for Stevens–Johnson syndrome/toxic epidermal necrolysis: A case report

Tan YW, Liu LP, Zhang K

LETTER TO THE EDITOR

1378 Enhancing competency of clinical research nurses: A comprehensive training and evaluation framework

Liu YX, Xu Y

ABOUT COVER

Peer Reviewer of *World Journal of Clinical Cases*, Narendra Pamidi, PhD, Assistant Professor, Department of Anatomy, Melaka Manipal Medical College, Karnataka 576104, India. narendra.pamidi@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJCC* as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Si Zhao*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Li Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

March 6, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Reconstruction of cervical necrotizing fasciitis defect with the modified keystone flap technique: Two case reports

Wonseok Cho, Eun A Jang, Kyu Nam Kim

Specialty type: Medicine, research and experimental

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B, B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Boffano P, Italy;
Hrgovic Z, Germany

Received: October 9, 2023

Peer-review started: October 9, 2023

First decision: January 12, 2024

Revised: January 25, 2024

Accepted: February 6, 2024

Article in press: February 6, 2024

Published online: March 6, 2024



Wonseok Cho, Eun A Jang, Kyu Nam Kim, Department of Plastic and Reconstructive Surgery, Kangbuk Samsung Hospital, Sungkyunkwan University School of Medicine, Seoul 03181, South Korea

Corresponding author: Kyu Nam Kim, MD, PhD, Full Professor, Department of Plastic and Reconstructive Surgery, Kangbuk Samsung Hospital, Sungkyunkwan University School of Medicine, No. 29 Saemunan-ro, Jongno-gu, Seoul 03181, South Korea.

manabear77@naver.com

Abstract

BACKGROUND

Cervical necrotizing fasciitis (CNF) is a rare, aggressive form of deep neck space infection with significant morbidity and mortality rates. Serial surgical debridement acts as the cornerstone of CNF treatment; however, it often results in defects requiring complex reconstructions.

CASE SUMMARY

We report two cases in which the keystone flap (KF) was used for CNF defect coverage: Case 1, an 85-year-old patient with CNF in the anterior neck, and Case 2, a 54-year-old patient with CNF in the posterior neck. Both patients received empirical intravenous antibiotic therapy and underwent serial debridement, enabling adequate wound preparation and stabilization. The final defect size measured 5.5 cm × 12 cm in Case 1 and 6 cm × 11 cm in Case 2. For defect coverage, we employed an 8 cm × 19 cm type II KF based on perforators from the superior thyroid artery in Case 1 and a 9 cm × 18 cm type II KF based on perforators from the transverse cervical artery in Case 2. Both flaps showed complete survival. No postoperative complications occurred in both cases, and favorable outcomes were observed at 7- and 6-month follow-ups in case 1 and 2, respectively.

CONCLUSION

We effectively treated CNF-associated defects using the KF technique; KF is viable for covering CNF defects in carefully selected cases.

Key Words: Fasciitis; Necrotizing; Debridement; Plastic surgery; Dermatologic surgical procedures; Case report

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Cervical necrotizing fasciitis (CNF) is associated with rapid tissue destruction and mortality; hence, early diagnosis and prompt treatment are necessary. However, serial debridement of necrotic tissues often leads to defects that complicate reconstructions, necessitating the selection of an appropriate reconstruction method. This study, therefore, reports our experiences using the modified keystone flap (KF) technique for CNF defect coverage in two patients. The modified KF technique may serve as an effective reconstructive option for neck defects, owing to its reliable and favorable outcomes in selected cases with small- to moderate-sized defects.

Citation: Cho W, Jang EA, Kim KN. Reconstruction of cervical necrotizing fasciitis defect with the modified keystone flap technique: Two case reports. *World J Clin Cases* 2024; 12(7): 1305-1312

URL: <https://www.wjgnet.com/2307-8960/full/v12/i7/1305.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v12.i7.1305>

INTRODUCTION

Necrotizing fasciitis (NF) is a relatively uncommon but life-threatening soft tissue infection with significant mortality, even if optimal treatment is administered[1-6]. Cervical NF (CNF) is a severe soft tissue infection characterized by rapidly progressive necrosis of the fascia and subcutaneous tissues along cervical fascial planes, predominantly affecting the face and neck areas[1-4,7]. This condition can lead to extensive gangrene and abscess formation in the deep neck space[1-7]. Because of the rapid tissue destruction associated with CNF, early diagnosis and prompt management are crucial. Interventions such as securing the airway, administering empirical broad-spectrum intravenous antibiotics, and performing aggressive surgical debridement are essential to achieve improved outcomes and overall survival[1-7].

The indispensable serial debridement of necrotic tissues often results in defects that present challenging reconstruction scenarios. To address these issues, surgeons must carefully select appropriate reconstructive options based on the size and depth of the defects[8]. Skin grafts may be considered for superficial defects, whereas local flaps are suitable for small- to moderate-sized defects with exposed underlying structures[8,9]. In cases with extensive complex defects, free flaps are often preferred; however, all decisions are made on a case-by-case basis[8,9]. In this context, we report our experiences with the modified keystone flap (KF) technique for CNF defect coverage in two cases. We theorize that this technique can be a valuable option in the management of such challenging scenarios.

This study was approved by the Institutional Review Board of Konyang University Hospital (approval number: 2020-07-015-01). All research procedures were performed in accordance with the ethical guidelines of the 1975 Declaration of Helsinki. Written informed consent was obtained from all patients for the publication of this case report and accompanying images.

CASE PRESENTATION

Chief complaints

Case 1: An 85-year-old man presented with CNF in the anterior neck area.

Case 2: A 54-year-old man presented with CNF in the posterior neck area.

History of present illness

Case 1: The patient exhibited rapidly progressive swelling and inflammation in the right cheek and neck following a tooth extraction.

Case 2: The condition rapidly progressed, following the occurrence of a tiny folliculitis on the posterior neck area.

History of past illness

Case 1: The patient reported multiple comorbidities, including diabetes mellitus, hypertension, and chronic renal failure.

Case 2: The patient had several underlying conditions, including hypertension, diabetes mellitus, and chronic renal failure.

Personal and family history

There was no family history of similar diseases.

Imaging examinations

Case 1: Contrast-enhanced computed tomography (CT) scan showed diffuse thickening of the platysma muscle and subcutaneous fat stranding, as well as significant swelling in the right mastication space and parotid space, indicative of CNF (Figure 1).




Figure 1 The contrast-enhanced computed tomography scan of the neck for **Case 1**. The scan demonstrates diffuse thickening of the platysma muscle and subcutaneous fat stranding, along with significant swelling in the right mastication space and parotid space, indicative of cervical necrotizing fasciitis.



Figure 2 Contrast-enhanced computed tomography scan of the neck for **Case 2**. The scan illustrated heterogeneous enhancement in the right occipital area and the adjacent posterior neck region. It also reveals diffuse thickening of the surrounding subcutaneous tissue and fat stranding all of which are consistent with the presence of cervical necrotizing fasciitis.

Case 2: Contrast-enhanced CT scan showed heterogeneous enhancement in the right occipital area and adjacent posterior neck region. In addition, it revealed diffuse thickening of the surrounding subcutaneous tissue and fat stranding; these findings are consistent with the presence of CNF (Figure 2).

FINAL DIAGNOSIS

Case 1

Based on these examinations, we considered a diagnosis of CNF in the anterior neck area.

Case 2

Based on these examinations, we considered a diagnosis of CNF in the posterior neck area.

TREATMENT

Case 1

Empirical broad-spectrum intravenous antibiotic treatment was administered, and serial debridement was performed



Figure 3 Clinical photographs of Case 1. A: Skin necrosis caused by cervical necrotizing fasciitis in the right anterior neck area; B: The final post-debridement defect measuring 5.5 cm × 12 cm, along with the design of a keystone flap (KF) sized 8 cm × 19 cm, based on perforators from the superior thyroid artery; C: Successful coverage of the defect with the modified type II KF; D: Postoperative photographs taken after a 3-month follow-up.

over a 2-wk period. Following adequate wound preparation, a flap surgery was planned for defect coverage. The final size of the defect post-debridement was 5.5 cm × 12 cm. We designed an 8 cm × 19 cm KF, based on perforators from the superior thyroid artery (Figure 3). The defect was effectively managed using the modified Type II KF, which involved division of the deep fascia along the entire circumference line of the flap. The flap survived completely, without any complications.

Case 2

Empirical broad-spectrum intravenous antibiotic treatment was administered, and serial debridement performed over a 2-wk period. Following adequate wound preparation, flap surgery was planned to address the defect. The final size of the defect post-debridement was 6 cm × 11 cm, and a deep dead space was identified inside the defect. We designed a 9 cm × 18 cm-sized KF, based on perforators from the transverse cervical artery (Figure 4). The defect was effectively covered with the modified Type II KF, and the flap showed complete survival without any complications.

OUTCOME AND FOLLOW-UP

Case 1

At the 7-month follow-up, the patient demonstrated a favorable functional outcome, showing no limitation in neck movement. Moreover, the patient expressed satisfaction with the final aesthetic result (Figure 5).

Case 2

At the 6-month follow-up, the patient demonstrated a favorable functional outcome, with no limitations in neck movement. In addition, the patient expressed satisfaction with the overall outcome.



Figure 4 Clinical photographs of Case 2[8]. A: Skin necrosis resulting from cervical necrotizing fasciitis in the posterior neck area; B: The final post-debridement defect measuring 6 cm × 11 cm, alongside the design of a keystone flap (KF) sized 9 cm × 18 cm, based on perforators from the transverse cervical artery; C: Successful coverage of the defect with the modified Type II KF; D: Postoperative photographs taken after a 6-month follow-up. **Citation:** Kong YT, Kim J, Shin HW, Kim KN. Keystone flaps for coverage of defects in the posterior neck and lower occipital scalp: a retrospective clinical study. *J Craniofac Surg* 2021; **32**: 1813-1816. Copyright ©The Authors 2021. Published by Wolters Kluwer Health, Inc.

DISCUSSION

NF predominantly occurs in the extremities, perineum, or trunk, with occurrence in the head and neck region accounting for approximately 10% of all cases[4,6,7]. CNF in the head and neck is often caused by odontogenic abscesses resulting from dental caries or chronic periodontitis[1-4,7]. Additionally, CNF may develop as a secondary infection following trauma or upper respiratory infections[1-4,7]. In the cases presented in this study, one CNF case had an odontogenic cause, whereas the other was secondary to a skin infection.

Several risk factors have been associated with NF, including diabetes mellitus, hypertension, myxedema, alcoholism, arteriosclerosis, advanced age, chronic renal failure, malnutrition, and metastatic cancer[1-4,6,7]. Notably, both cases in this report showed some of these risk factors, including diabetes mellitus, hypertension, and chronic renal failure.

CNF management depends on the early initiation of broad-spectrum antibiotics and aggressive serial surgical debridement[4,6,7]. These strategies can decelerate the rate of tissue destruction and enhance the overall survival rate. However, serial debridement often results in defects that require reconstructive procedures, including skin grafts and flap coverage[4,5,7].

For superficial defects of any size that do not expose underlying structures, skin grafting is a straightforward and effective reconstructive option[8,9]. Nevertheless, the procedure has some limitations, such as potential skin color and texture mismatches, vulnerability to shearing and friction forces, contour deformity, and potential donor site morbidity[8,9]. In cases of deep defects involving exposed underlying structures, flap coverage is imperative for providing protective, durable tissue layers[8,9].

As previously mentioned, local flaps are suitable for small- to moderate-sized defects, as in our cases. Some studies have demonstrated the application of various local flaps, including the trapezius, pectoralis major muscle, deltopectoral,



Figure 5 Follow-up clinical photographs of Case 1. A-D: At the 7-month follow-up, no limitation of neck movement was observed.

bilobed fasciocutaneous, and perforator-based fasciocutaneous flaps[8,10].

The KF has emerged as a widely utilized reconstructive modality for various defects, serving as both a primary and alternative option, regardless of the defect's location[8,9,11,12]. This preference can be attributed to its definite advantages, including a simple and easily adjustable design to match specific defects, ease of reproducibility, and minimal learning curve for surgeons[8,9,11,12].

Numerous studies have demonstrated the effectiveness of KF reconstruction in the neck region[8,13-15]. Notably, our senior author has previously demonstrated successful application of KF for defect coverage in the posterior neck and lower occipital scalp[8].

The neck area is rich in perforators, providing an excellent resource for KF advancement flaps[8]. The anterior neck has perforators arising from the superior and inferior thyroid arteries, while the posterior neck has perforators originating from the occipital and transverse cervical arteries. Owing to its multi-perforator-based nature, the KF can take full advantage of these abundant perforators in the neck region, making it a safe and reliable choice for reconstructive purposes.

Figure 6 illustrates the regions of perforators in the neck area and representative design of the KF for neck defects, including the cases presented in this study. For anterior neck defects, KF based on perforators from the superior thyroid, inferior thyroid, and transverse cervical arteries can be employed, whereas for posterior neck defects, KF based on perforators from the occipital and transverse cervical arteries is utilized.

In our cases, we utilized the modified Type II KF, which involves a complete division of the whole circumferential deep fascia along the flap boundary[9,16]. This modified approach offers enhanced flap movement compared to the original Type II KF, where deep fascia division is limited along the outer curvilinear line of the flap[9,16]. By dividing the investing layer of deep cervical fascia along the entire circumference of the flap, we achieved sufficient flap movement to effectively cover the defects[9,16].

Previous studies have demonstrated the tension-reducing effect of KF reconstruction, primarily attributed to deep fascia division[11,12]. We believe that this tension-reducing effect facilitated wound closure with minimal tension, which is crucial for neck defect coverage. Therefore, no postoperative wound complications or limitations in neck movement were observed in the cases reported here.

CONCLUSION

This study presents successful cases of CNF defect coverage using the modified KF technique and provides illustrative figures (Figure 6) to enhance reader comprehension. To the best of our knowledge, this study is the first case report of CNF defect reconstruction using the modified KF in Korea. We suggest that the modified KF technique may serve as an effective reconstructive option for neck defects owing to its reliable and favorable outcomes in selected cases with small- to moderate-sized defects.




Figure 6 Illustrations of the regions of perforators in the neck area and the representative design of keystone flaps for neck defects. A: Perforator territories of the anterior neck area. The chartreuse-colored territory represents the perforator-area from the superior thyroid artery, the mustard-colored territory represents the perforator-area from the inferior thyroid artery, and the sky-blue-colored territory represent the perforator-area from the transverse cervical artery; B: Representative designs of keystone flaps (KFs) for anterior neck defects. The magenta-colored ellipses represent defects, and the apricot-colored figures represent the design of KFs; C: Perforator territories of the posterior neck area. The lemon-colored territory represents the perforator-area from the occipital artery and the sky-blue-colored territory represent the perforator-area from the transverse cervical artery; D: Representative designs of KFs for posterior neck defects. The magenta-colored ellipses represent defects and the apricot-colored figures represent the design of KFs.

FOOTNOTES

Author contributions: Cho W and Jang EA contributed equally to this work; Kim KN contributed to conceptualization and methodology, reviewed and edited the manuscript, and performed supervision; Jang EA contributed to software, provided resources and performed data curation; Kim KN and Cho W performed validation; Cho W performed formal analysis, and wrote the original draft; Cho W and Jang EA performed the research; Jang EA performed visualization. All authors have read and approve the final version of the manuscript.

Informed consent statement: Written informed consent was obtained from all patients for the publication of this case report and accompanying images.

Conflict-of-interest statement: The authors declare that they have no conflict of interest to disclose.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised

according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: South Korea

ORCID number: Wonseok Cho 0009-0007-0203-3059; Eun A Jang 0009-0007-2878-917X; Kyu Nam Kim 0000-0001-9998-8722.

S-Editor: Fan JR

L-Editor: A

P-Editor: Zheng XM

REFERENCES

- 1 **Yang YS**, Lee HU, Kim JS, Lee JK, Hong KH. A clinical study of the cervical necrotizing fasciitis. *Korean J Otorhinolaryngol-Head Neck Surg* 2005; **48**: 1020-1026
- 2 **Kim CH**, Choung YH, Oh JH, Lee JW. Management of cervical necrotizing fasciitis. *Korean J Otorhinolaryngol-Head Neck Surg* 2005; **48**: 771-777
- 3 **Hua J**, Friedlander P. Cervical Necrotizing Fasciitis, Diagnosis and Treatment of a Rare Life-Threatening Infection. *Ear Nose Throat J* 2023; **102**: NP109-NP113 [PMID: 33570428 DOI: 10.1177/0145561321991341]
- 4 **Gupta V**, Sidam S, Behera G, Kumar A, Mishra UP. Cervical Necrotizing Fasciitis: An Institutional Experience. *Cureus* 2022; **14**: e32382 [PMID: 36632261 DOI: 10.7759/cureus.32382]
- 5 **Choi JA**, Kwak JH, Yoon CM. Life-threatening necrotizing fasciitis of the posterior neck. *J Trauma Inj* 2020; **33**: 260-263 [DOI: 10.20408/jti.2020.0025]
- 6 **Misiakos EP**, Bagias G, Patapis P, Sotiropoulos D, Kanavidis P, Machairas A. Current concepts in the management of necrotizing fasciitis. *Front Surg* 2014; **1**: 36 [PMID: 25593960 DOI: 10.3389/fsurg.2014.00036]
- 7 **Gunaratne DA**, Tseros EA, Hasan Z, Kudpaje AS, Suruliraj A, Smith MC, Riffat F, Palme CE. Cervical necrotizing fasciitis: Systematic review and analysis of 1235 reported cases from the literature. *Head Neck* 2018; **40**: 2094-2102 [PMID: 29934952 DOI: 10.1002/hed.25184]
- 8 **Kong YT**, Kim J, Shin HW, Kim KN. Keystone Flaps for Coverage of Defects in the Posterior Neck and Lower Occipital Scalp: A Retrospective Clinical Study. *J Craniofac Surg* 2021; **32**: 1813-1816 [PMID: 33654029 DOI: 10.1097/SCS.00000000000007581]
- 9 **Yoo BW**, Oh KS, Kim J, Shin HW, Kim KN. Modified Keystone Perforator Island Flap Techniques for Small- to Moderate-Sized Scalp and Forehead Defect Coverage: A Retrospective Observational Study. *J Pers Med* 2023; **13** [PMID: 36836563 DOI: 10.3390/jpm13020329]
- 10 **Lee JH**, Foo FJ, Wong AW. Posterior cervical necrotizing fasciitis: a multidisciplinary endeavour in surgery. *J Surg Case Rep* 2023; **2023**: rjad127 [PMID: 36969714 DOI: 10.1093/jscr/rjad127]
- 11 **Yoon CS**, Kim HB, Kim YK, Kim H, Kim KN. Keystone-design perforator island flaps for the management of complicated epidermoid cysts on the back. *Sci Rep* 2019; **9**: 14699 [PMID: 31605009 DOI: 10.1038/s41598-019-51289-4]
- 12 **Yoon CS**, Kong YT, Lim SY, Kim J, Shin HW, Kim KN. A comparative study for tension-reducing effect of Type I and Type II keystone perforator island flap in the human back. *Sci Rep* 2021; **11**: 16699 [PMID: 34404867 DOI: 10.1038/s41598-021-96272-0]
- 13 **Behan FC**, Rozen WM, Wilson J, Kapila S, Sizeland A, Findlay MW. The cervico-submental keystone island flap for locoregional head and neck reconstruction. *J Plast Reconstr Aesthet Surg* 2013; **66**: 23-28 [PMID: 22974756 DOI: 10.1016/j.bjps.2012.08.027]
- 14 **Sun Y**. Keystone Flap for Large Posterior Neck Defect. *Indian J Surg* 2016; **78**: 321-322 [PMID: 27574353 DOI: 10.1007/s12262-015-1406-5]
- 15 **Pelissier P**, Gardet H, Pinsolle V, Santoul M, Behan FC. The keystone design perforator island flap. Part II: clinical applications. *J Plast Reconstr Aesthet Surg* 2007; **60**: 888-891 [PMID: 17493885 DOI: 10.1016/j.bjps.2007.03.023]
- 16 **Kim KH**, Yoo BW, Lim SY, Oh KS, Kim J, Shin HW, Kim KN. Modified Keystone Perforator Island Flap for Tension-Reducing Coverage of Axillary Defects Secondary to Radical Excision of Chronic Inflammatory Skin Lesions: A Retrospective Case Series. *Biomed Res Int* 2022; **2022**: 5600450 [PMID: 36212718 DOI: 10.1155/2022/5600450]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

