

World Journal of *Clinical Cases*

World J Clin Cases 2024 March 6; 12(7): 1196-1381



Contents

Thrice Monthly Volume 12 Number 7 March 6, 2024

EDITORIAL

- 1196 Relevance of sleep for wellness: New trends in using artificial intelligence and machine learning
Nag DS, Swain A, Sahu S, Chatterjee A, Swain BP

MINIREVIEWS

- 1200 Expect the unexpected: Brown tumor of the mandible as the first manifestation of primary hyperparathyroidism
Majic Teng A, Cigrovski Berkovic M, Zajc I, Salaric I, Müller D, Markota I
- 1205 Research progress in spasmodic torticollis rehabilitation treatment
Zhang S, Zeng N, Wu S, Wu HH, Kong MW

ORIGINAL ARTICLE

Clinical and Translational Research

- 1215 Investigating the causal associations between five anthropometric indicators and nonalcoholic fatty liver disease: Mendelian randomization study
Xiao XP, Dai YJ, Zhang Y, Yang M, Xie J, Chen G, Yang ZJ
- 1227 Causal role of immune cells in obstructive sleep apnea hypopnea syndrome: Mendelian randomization study
Zhao HH, Ma Z, Guan DS

Case Control Study

- 1235 Significant risk factors for intensive care unit-acquired weakness: A processing strategy based on repeated machine learning
Wang L, Long DY

Retrospective Cohort Study

- 1243 Perioperative and long-term results of ultrasonography-guided single- and multiple-tract percutaneous nephrolithotomy for staghorn calculi
Cheng RX, Dai N, Wang YM, Qi P, Chen F

Retrospective Study

- 1251 Clinical characteristics of testicular torsion and factors influencing testicular salvage in children: A 12-year study in tertiary center
Gang XH, Duan YY, Zhang B, Jiang ZG, Zhang R, Chen J, Teng XY, Zhang DB

META-ANALYSIS

- 1260 Effectiveness of sensory integration therapy in children, focusing on Korean children: A systematic review and meta-analysis
Oh S, Jang JS, Jeon AR, Kim G, Kwon M, Cho B, Lee N
- 1272 Safety and efficacy comparison of remimazolam and propofol for intravenous anesthesia during gastroenteroscopic surgery of older patients: A meta-analysis
Li FZ, Zhao C, Tang YX, Liu JT

CASE REPORT

- 1284 Sporadic gastrinoma with refractory benign esophageal stricture: A case report
Chen QN, Bai BQ, Xu Y, Mei Q, Liu XC
- 1290 Efficacy of borneol-gypsum in skin regeneration and pain control in toxic epidermal necrolysis: A case report
Yang LW, Zhang LJ, Zhou BB, Lin XY, Chen YT, Qin XY, Tian HY, Ma LL, Sun Y, Jiang LD
- 1296 Extended survival with metastatic pancreatic cancer under fruquintinib treatment after failed chemotherapy: Two case reports
Wu D, Wang Q, Yan S, Sun X, Qin Y, Yuan M, Wang NY, Huang XT
- 1305 Reconstruction of cervical necrotizing fasciitis defect with the modified keystone flap technique: Two case reports
Cho W, Jang EA, Kim KN
- 1313 Reversal of complete atrioventricular block in dialysis patients following parathyroidectomy: A case report
Xu SS, Hao LH, Guan YM
- 1320 Treatment of bilateral developmental dysplasia of the hip joint with an improved technique: A case report
Yu XX, Chen JY, Zhan HS, Liu MD, Li YF, Jia YY
- 1326 Misdiagnosis of synovial sarcoma - cellular myofibroma with *SRF-RELA* gene fusion: A case report
Zhou Y, Sun YW, Liu XY, Shen DH
- 1333 Heterochronous multiple primary prostate cancer and lymphoma: A case report
Liang JL, Bu YQ, Peng LL, Zhang HZ
- 1339 Cardiac remodeling in patients with atrial fibrillation reversing bradycardia-induced cardiomyopathy: A case report
Gao DK, Ye XL, Duan Z, Zhang HY, Xiong T, Li ZH, Pei HF
- 1346 Microsurgical management of radicular cyst using guided tissue regeneration technique: A case report
Gómez Mireles JC, Martínez Carrillo EK, Alcalá Barbosa K, Gutiérrez Cortés E, González Ramos J, González Gómez LA, Bayardo González RA, Lomeli Martínez SM
- 1356 Delayed neurological dysfunction following posterior laminectomy with lateral mass screw fixation: A case report and review of literature
Yan RZ, Chen C, Lin CR, Wei YH, Guo ZJ, Li YK, Zhang Q, Shen HY, Sun HL

- 1365** Translocation of a fish spike from the pharynx to the thyroid gland: A case report

Li D, Zeng WT, Jiang JG, Chen JC

- 1371** Double plasma molecular adsorption system for Stevens–Johnson syndrome/toxic epidermal necrolysis: A case report

Tan YW, Liu LP, Zhang K

LETTER TO THE EDITOR

- 1378** Enhancing competency of clinical research nurses: A comprehensive training and evaluation framework

Liu YX, Xu Y

ABOUT COVER

Peer Reviewer of *World Journal of Clinical Cases*, Narendra Pamidi, PhD, Assistant Professor, Department of Anatomy, Melaka Manipal Medical College, Karnataka 576104, India. narendra.pamidi@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Si Zhao; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

March 6, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Microsurgical management of radicular cyst using guided tissue regeneration technique: A case report

Juan Carlos Gómez Mireles, Eugenia Karina Martínez Carrillo, Katia Alcalá Barbosa, Evangelina Gutiérrez Cortés, Javier González Ramos, Luis Antonio González Gómez, Rubén Alberto Bayardo González, Sarah Monserrat Lomelí Martínez

Specialty type: Medicine, research and experimental

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): B, B

Grade C (Good): 0

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Ghannam WM, Egypt; He CY, China

Received: December 12, 2023

Peer-review started: December 12, 2023

First decision: December 31, 2023

Revised: January 9, 2024

Accepted: February 6, 2024

Article in press: February 6, 2024

Published online: March 6, 2024



Juan Carlos Gómez Mireles, Luis Antonio González Gómez, Periodontics Specialty, Department of Integral Dentistry Clinics, Centro Universitario de Ciencias de la Salud, Universidad de Guadalajara, Guadalajara 44340, Mexico

Eugenia Karina Martínez Carrillo, Endodontics Specialty, Department of Integral Dentistry Clinics, Centro Universitario de Ciencias de la Salud, Universidad de Guadalajara, Guadalajara 44340, Mexico

Katia Alcalá Barbosa, Endodontics Specialty, Department of Clinics, Centro Universitario de Los Altos, Universidad de Guadalajara, Tapatitlán de Morelos 47620, Mexico

Evangelina Gutiérrez Cortés, Specialty in Oral Pathology, Surgical-Stomatological Department, Autonomous University of Guadalajara, Zapopan 45129, Mexico

Javier González Ramos, Independent Researcher, Guadalajara 44340, Mexico

Rubén Alberto Bayardo González, Sarah Monserrat Lomelí Martínez, Department of Integral Dentistry Clinics, Centro Universitario de Ciencias de la Salud, Universidad de Guadalajara, Guadalajara 44340, Mexico

Sarah Monserrat Lomelí Martínez, Department of Medical and Life Sciences, Centro Universitario de la Ciénega, Universidad de Guadalajara, Ocotlán 47810, Mexico

Sarah Monserrat Lomelí Martínez, Master of Public Health, Department of Well-being and Sustainable Development, Centro Universitario del Norte, Universidad de Guadalajara, Colotlán 46200, Mexico

Corresponding author: Sarah Monserrat Lomelí Martínez, PhD, Academic Research, Adjunct Associate Professor, Department of Medical and Life Sciences, Centro Universitario de la Ciénega, Universidad de Guadalajara, Av. Universidad 1115, Col. Lindavista, Ocotlán 47810, Mexico. sarah.lomeli@academicos.udg.mx

Abstract

BACKGROUND

Radicular cyst is a lesion of odontogenic origin that arises from epithelial remains due to periapical periodontitis caused by inflammatory reactions generated at the apex of affected teeth with infected or necrotic pulps. The therapeutic mana-

gement of radicular cysts is controversial. There is only one case report of enucleation of a radicular cyst managed with microsurgery and apicoectomy, but without the use of the guided tissue regeneration (GTR) technique in the same surgical procedure. The present clinical case describes the management of a radicular cyst with microsurgical approach, performance of an apicoectomy of the tooth associated with the entity, application of GTR technique, use of a resorbable membrane of type I bovine collagen, and bovine xenograft.

CASE SUMMARY

A 68-year-old patient presented with a radicular cyst from an upper lateral incisor. The microsurgical management used was aimed at enucleating the chemical membrane, performing apicoectomy of the tooth along with careful and precise retrograde filling, and implementing GTR technique using a resorbable collagen membrane and bovine xenograft. The diagnosis of radicular cyst was confirmed using histopathological analysis. The patient underwent follow-up evaluations at 10 and 30 d postoperatively. At 4 months postoperative evaluation, she remained asymptomatic, and radiographs showed significant periapical healing with adequate bone formation.

CONCLUSION

These results suggest that microsurgical management using the GTR technique with collagen membrane and xenograft, contributes to bone regeneration.

Key Words: Odontogenic cysts; Apical periodontal cyst; Dental root end cyst; Case report

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The radicular cyst is a lesion of odontogenic origin whose therapeutic management is controversial. However, the present clinical case describes the management of a radicular cyst using a microsurgical approach, and the performance of an apicoectomy of the tooth associated with the entity, followed by the guided tissue regeneration technique, implementation of a resorbable type I bovine collagen membrane, and bovine xenograft. The diagnosis was based on clinical findings that were confirmed with histopathology analysis. This case underlines the importance of clinical, periapical radiographic, and cone-beam computed tomography evaluation as guides for an accurate diagnosis in order to enhance the development of an appropriate treatment plan for a radicular cyst.

Citation: Gómez Mireles JC, Martínez Carrillo EK, Alcalá Barbosa K, Gutiérrez Cortés E, González Ramos J, González Gómez LA, Bayardo González RA, Lomeli Martínez SM. Microsurgical management of radicular cyst using guided tissue regeneration technique: A case report. *World J Clin Cases* 2024; 12(7): 1346-1355

URL: <https://www.wjgnet.com/2307-8960/full/v12/i7/1346.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v12.i7.1346>

INTRODUCTION

Radicular cysts, also known as apical periodontal cysts, are the most common inflammatory odontogenic cysts of the jaw [1]. They are chronic lesions arising from pre-existing periapical granulomas triggered by inflammatory reactions generated in the apex of the affected teeth due to infected or necrotic pulps, although they may also be located on the lateral surfaces of roots associated with accessory canals [2]. Generally, radicular cysts have a predilection for the male gender, and they occur between the third and sixth decades of life [1]. A radicular cyst is clinically characterized by a slight increase in volume that is soft and fluctuating during palpation [1]. The lesion may present as a hard and irregular bony swelling [3]. It is usually asymptomatic, and sometimes presents mild pain and pain on percussion [4]. Extra-orally, an oval, diffuse, painless swelling is observed in the affected area. Radicular cysts are often slow-growing, and may cause sensitivity, mobility, root resorption and displacement of the affected teeth. When the cystic entity is infected, it can also cause pain and inflammation [1].

The treatment of radicular cyst is aimed at eradicating the lesion and enhancing the healing of the periapical tissue so as to improve the outcome. The modality chosen is usually the one that grants the lowest possible risk of recurrence and minimum morbidity. Some of the factors that determine the treatment modality for radicular cysts are the origin and extension of the lesion, its relationship with important anatomical structures, clinical characteristics, and patient collaboration, amongst others [5,6]. When the lesions are small, the first line of treatment is endodontic therapy which is a conservative method. In contrast, marsupialization and enucleation have been used as surgical treatment methods of choice when the cysts are large and/or there are invariable changes in the peri-radicular tissue that cannot be treated with non-surgical endodontic procedure [4,5]. Moreover, different conventional or microsurgical endodontic and periodontal surgical techniques such as apicectomy and guided tissue regeneration (GTR) have been described for complementing the enucleation treatment. However, despite the different modalities and techniques, the management of radicular cysts is controversial. The present clinical case describes the management of a radicular cyst using a microsurgical approach, and the performance of an apicectomy of the tooth associated with the entity, followed by GTR technique, implementation of

a resorbable membrane of type I bovine collagen, and bovine xenograft.

CASE PRESENTATION

Chief complaints

A 68-year-old female patient without systemic engagement, was referred by a general dentist to a private practice with finding of a radiolucent zone in the periapical area of the dental organ (OD) 22 and displacement of the adjacent roots.

History of present illness

The patient mentioned having received endodontic treatment in the previous 6 years, without clinical or radiographic follow-up.

History of past illness

The patient did not mention presence of any systemic disease, and she was not undergoing any medical treatment.

Personal and family history

Mother was diagnosed with high blood pressure in 2017.

Physical examination

During the clinical examination, a bulge was identified in the vestibular area corresponding to OD 22, with a soft, painless consistency (Figure 1). This OD did not present grade of mobility, supra- and sub-gingival calculus or probing depth.

Laboratory examinations

Hemoglobin, neutrophil and platelet levels were within normal limits.

Imaging examinations

Periapical radiograph revealed a well-defined, completely asymptomatic and circumferential radiolucent area (Figure 2). Evaluation using computed tomography revealed the absence of vestibular cortex, with the dimensions of the loss being 6 mm wide × 16 mm high and 5.5 mm deep (Figure 3). In the three-dimensional reconstruction, the extent and extension of the lesion can be observed. This evaluation was performed with the OnDemand computed tomography Viewer (Figure 4).

MULTIDISCIPLINARY EXPERT CONSULTATION

After the evaluation, analysis and interdisciplinary discussion were carried out with specialists in endodontics, periodontics and oral pathology, with respect to the possible treatment options. However, based on the patient's clinical history and her detachment from treatment, as evidenced by not attending any dental consultation for 6 years, including the recommended radiographic follow-up after root canal treatment, a micro-surgical approach was decided upon. This approach would enucleate the radicular cyst and carry out GTR to improve the treatment prognosis with the least possible trauma to the soft tissues. The treatment of choice and the associated risks were thoroughly explained to the patient, and she gave signed informed consent. Surgical enucleation was the chosen treatment, as well as apicoectomy of the piece involved, followed finally with GTR.

FINAL DIAGNOSIS

The results from histological analysis revealed a cavity lined by non-keratinized odontogenic epithelium with intracapsular projections and acanthosis. Moreover, intense mixed inflammatory infiltrates and foamy macrophages were present in the fibrous capsule, in addition to erythrocyte extravasation. These features were consistent with diagnosis of radicular cyst (Figure 5).

TREATMENT

The patient was scheduled for the surgery, during which specialists in the three previously mentioned specialties were involved. The procedure was performed under the magnification of a Zeiss Extaro 300 clinical microscope. The area was anesthetized with articaine: Epinephrine bi-tartrate (1:100000). Following anesthesia, a time lag of 20 min was allowed for adequate vasoconstriction. Then, initial intra-surgical incisions and a vertical incision were made in order to have a clear view of the lesion. A full-thickness flap was raised to expose the lesion which was carefully dissected completely using



Figure 1 Preoperative clinical view. An increase in volume corresponding to the circumscribed, well delimited lesion, between 2.2 and 2.3.



Figure 2 Preoperative periapical radiograph. A: X-ray prior to root canal treatment, 6 years ago; B: Current initial radiograph showing a well-defined and circumscribed radiolucent area at the apex of tooth 22.




Figure 3 Cone-beam computed tomography view. A: Sagittal; B: Axial; C: Coronal.

curettes. The lesion was placed in a container with 40% formalin (universal percentage for fixing histopathological samples). Following enucleation, the cavity was examined, washed and enlarged by means of a high-speed electric piece in order to have better control of the instruments during the apicoectomy. Using methylene blue, the root was inspected for possible fractures. The root apex was cut with an endo zekrya bur in a 45° turbine at 3 mm coronal to the apex, and the apex was again inspected with the help of magnification and micromirrors to locate any microfractures in the root canal. Retro-instrumentation was performed with diamond ultrasound tips. The cavity was dried and inspected for the presence of internal debris. Then, biodentine (Septodont) retro-preparation was placed in the canal (Figure 6). The excess was removed with the help of a carrier and burnisher, and using indirect vision, effective sealing of the canal was confirmed. Once the apicoectomy was completed, it was washed with cold saline, and a xenograft (Biooss by Geistlich) previously hydrated with physiological saline was used to fill the cavity. Then, a Bio-Gide® collagen membrane (Geistlich) was




Figure 4 Three-dimensional model reconstruction. Note the absence of vestibular wall.




Figure 5 Histological examination. A: Hematoxylin and eosin-stained histological sections exhibiting cavity lined by non-keratinized odontogenic epithelium with intracapsular projections beneath [Histological examination (H&E); $\times 20$]; B: Photomicrograph showing intense, mixed inflammatory infiltrates and foamy macrophages in the fibrous capsule (H&E; $\times 40$).

placed and adapted over the bone window (Figure 7). The flap was repositioned, making sure that the closure was first intention and tension-free (Figure 8A). The suture chosen was 6-0 polypropylene due to ease of handling and excellent biocompatibility. Conventional post-operative care was used. The antibiotic used was Amoxicillin (750 mg; 1 tablet every 12 h), while the analgesics were sodium naproxen 275 mg; Paracetamol 300 mg (2 tablets every 8 h). Additionally, 0.20% chlorhexidine gel was applied in the area for 2 wk. Ice application in the area was done for the first 72 h.

OUTCOME AND FOLLOW-UP

At the 4-month follow-up, gingival clinical examination revealed that the affected area which was subjected to GTR recovered its normal volume and color (Figure 8B and C). The periapical radiograph demonstrated total bone filling in the area adjacent to the dental piece up to the apicoectomy level of the tooth root. Two-dimensional (2D) radiographic healing was judged incomplete by two independent reviewers and calibrated according to the Molven criteria. Tomographic evaluation showed complete healing in the area, with beginnings of vestibular cortical formation (Figures 9 and 10).




Figure 6 Intraoperative photography. A: Complete enucleation of the lesion; B: Adequate sealing of the canal with biodentine, after apicectomy.



Figure 7 Immediate postoperative periapical radiograph.



Figure 8 Postoperative clinical view. A: Immediate; B: After 10 d; C: After 4 months.

DISCUSSION

The radicular cyst represents 55% of odontogenic cysts, and 52%-68% of all cysts of the jaws[7]. This chronic entity is the result of multiple inflammatory reactions initially guided by various proinflammatory interleukins and growth factors which cause proliferation of epithelial cells derived from epithelial rests of Malassez in granulomatous tissue, followed by cyst formation and growth[2,8]. However, since events associated with the molecular biology of the radicular cyst are still unclear, there is need for more research in this area[2]. The therapeutic management of radicular cysts is controversial. To our knowledge, in the literature, there is only one case report of enucleation of a radicular cyst managed with



Figure 9 6 months postoperative cone-beam computed tomography view. A: Sagittal; B: Axial; C: Coronal.




Figure 10 Six months postoperative three-dimensional model reconstruction.

microsurgery and apicoectomy, but without the use of the GTR technique in the same surgical procedure. The present clinical case describes the management of a radicular cyst with microsurgical approach, performance of an apicoectomy of the tooth associated with the entity, application of GTR technique, use of a resorbable membrane of type I bovine collagen, and bovine xenograft.

In recent years, the incorporation of magnification into dental practice has resulted in improved visual acuity, ergonomic benefits, decreased patient morbidity, rapid healing, and increased patient acceptance[4,9]. Endodontic microsurgery becomes the recommended procedure when initial root canal therapy or subsequent retreatments fail to prevent or heal apical periodontitis, and the lesion near the tooth's apex remains unrepaired[4]. In the meta-analysis of Setzer *et al*[10], the probability of success of conventional endodontic surgery *vs* microsurgery, and the probability of success for the 2 techniques were compared. The authors concluded that the success rates of conventional surgeries and microsurgeries were 59% and 94%, respectively[11]. In a review article, it was reported that endodontic microsurgery had an overall success rate of 91.5% at 5 years, and 93.3% at 10 years. These data indicate that the microsurgical approach has resulted in a significant increase in the success rate of endodontic surgeries, since this technique is minimally invasive, and it allows for greater precision. In the present case, the apicoectomy with retrodental filling through a microsurgical approach using biodentine was carried out with the aim of removing the periapical lesion and generating an environment conducive to healing in both hard and soft tissues. The use of biodentine for retrograde obturation after apicoectomy has been described by other authors[12,13]. Biodentine is a biologically active cement based on calcium silicate. It has a variety of dental applications due to its physicochemical characteristics, biocompatibility and ease of handling which account for its current wide use, particularly in the endodontic area[14,15]. For several years, Mineral Trioxide Aggregate (MTA) was considered the first treatment option in many surgical and non-surgical dental procedures such as root perforations, direct pulp capping, apexification processes, periapical lesions, retrograde obturation, and filling/temporary padding. This ceramic material possesses biocompatibility and sealing capacity, and it is not adversely affected by contact with blood and peri-radicular fluids. However, it has the disadvantage of long setting time, high cost and difficulty in handling[11]. To compensate for the deficiencies of MTA, biodentine gained prominence due to its excellent sealing capacity. It does not shrink or reabsorb, and it hardens quickly and is easy to handle. Biodentine has several advantages over MTA, such as greater antibacterial and antifungal activity, greater adhesion strength, and greater biocompatibility[12,14,15]. Due to these benefits, biodentine is an excellent option for use as a retro-filling material.

Reports have shown that GTR has been implemented as an adjunct to radicular cyst enucleation and apicoectomy for enhancement of periapical tissue healing, prevention of soft tissue collapse, and improvement of treatment outcomes[11, 16,17]. Full recovery of the area around the apex of a tooth includes the regeneration of alveolar bone, periodontal ligament cells and cementum. However, within the bone defect, the surrounding connective tissues may grow, thereby

preventing bone healing. GTR is a technique that uses a bone substitute material along with a membrane that serves both as a biological and physical shield to prevent cells that do not contribute to bone growth, such as epithelial cells, from invading the area. Moreover, GTR allows the migration of bone-forming cells from slow migration to bad sites, thereby ensuring selective cell re-population and guiding the proliferation of various tissues during the healing process[17,18]. Numerous studies have described this technique, with respect to the application of a wide range of materials and treatment protocols, all of which achieved clinical effectiveness. Zubizarreta-Macho *et al*[16] have described a meta-analysis of clinical evidence on the efficacy of the GTR technique in bone defects after endodontic surgical treatment[16]. These authors highlighted the importance of membranes and demonstrated that employing a membrane by itself, without the addition of a bone graft, was 1.02 times more effective compared to endodontic surgery without the aid of GTR techniques. It was also found to surpass the efficacy of methods involving platelet-rich plasma. Nonetheless, the synergy of both membrane and bone grafting proved to be substantially more advantageous, achieving a success rate that was 3.6 times higher than using the membrane in isolation. On the other hand, the success rate of the combination of membrane and bone graft was 3.7 times higher than that of endodontic surgery alone (control). It was concluded in the meta-analysis that GTR techniques in which bone grafts and membranes were used as complements to surgical endodontic treatment, promoted complete healing of the periapical bone with enhanced success rate, and improved the prognosis of endodontic surgery. The determinants of the success of GTR are the type of bone regenerative biomaterial and the barriers applied. In a systematic review and meta-analysis, Sumangali *et al*[19] analyzed the success of GTR in periapical surgeries based on clinical, histological, and radiographic evaluations with different materials, and reported that the use of xenograft in combination with collagen membrane produced a superior result. In the present case, in order to improve the healing of the periapical tissue and avoid tissue collapse due to bone defect, a resorbable type I bovine collagen membrane was used to cover the particulate bovine xenograft after the enucleation of the cyst and the apicoectomy. The membrane is 100% natural and multi-porous. The GTR was performed by Morillo *et al*[20] using cortico-cancellous bone substitute and fibrin-rich plasma (PRF) membrane, after the removal of a radicular cyst[20]. The latter is a natural blood-derived structure associated with the immune system and platelet concentrate that exhibits several vital properties such as cell attachment, cell migration, cell proliferation, and cell differentiation. Additionally, it contains multiple growth factor-like platelet-derived growth factor, transforming growth factor β 1, insulin-like growth factor, and vascular endothelial growth factor, all of which enhance stimulation of the healing of surgical wounds in big bones[17]. The PRF was also implemented as a post-enucleation membrane of a radicular cyst by Govindaraju *et al*[6], but without the addition of bone filler biomaterials[6]. Despite the multiple and beneficial properties of PRF, its main disadvantage is that it resorbs within approximately 7-28 d, which is substantially shorter than the 4-6 wk required for most periodontal regeneration applications[17], and also shorter than the 16 to 24 wk required for bone enhancement[18]. Furthermore, its capacity to maintain space is compromised by its high rate of reabsorption[17]. In the present study, the use of PRF membrane was ruled out in favor of a collagen membrane. Tseng *et al*[21] have reported that after enucleation of radicular cyst with apicoectomy and retrograde obturation of the apical root canal, micro-macroporous biphasic calcium phosphate bio-ceramic was placed in the bone defect. However, they did not use any membrane. On the other hand, there are reports of clinical cases in which no biomaterials were used for bone regeneration after the enucleation of the radicular cyst[4,7,22,23]. For instance, Elhakim *et al*[4] made the complete enucleation of a periapical cyst without sacrificing the vitality of the adjacent teeth, and used no bone filler material or membrane: They argued that the bone graft material might have compromised the integrity of the neurovascular bundles of vital teeth and limited their repair potential.

The microsurgical approach with GTR is associated with better illumination and enlargement of the surgical field, thereby providing adequate access to the intraosseous defect and debridement thereof with greater precision and minimal trauma[4,9,24]. This approach allows for obtaining and maintaining primary closure at the edges of the wound, and minimizing bacterial contamination, thereby ensuring favorable conditions for regeneration[4]. Likewise, this microsurgical modality is characterized by minimal marginal tissue recession and very limited intraoperative and postoperative morbidity[4,9,25]. Due to these fundamental factors, the microsurgical approach for the treatment of intra-bony defects has been proposed by several authors[26]. In the present case, the magnification allowed for a greater precision in the management of the enucleation of the lesion, as well as primary closure without tension, thereby favoring an adequate evolution of the wound with minimal discomfort. Additionally, at 6 months of follow-up in the clinical and radiographic evaluation, successful results were observed which were attributed to adequate surgical management and the use of correct regeneration biomaterials.

Finally, it is important to highlight that the management of radicular cysts requires a multidisciplinary approach that involves several dental specialists, in order to obtain accurate diagnosis and safe and effective treatment plans. Regular follow-up of endodontically treated teeth is essential for verification of apical healing, and if a periapical lesion develops, it is identified at an early stage. Early detection and accurate diagnosis are essential for adequate treatment and prevention of possible complications.

CONCLUSION

The microsurgical management of a radicular cyst using the GTR technique, resorbable membrane of type I bovine collagen and bovine xenograft after apicoectomy of the associated tooth with retrograde biodentine seal, allowed for successful removal of the periapical lesion and generated an environment conducive to healing in both hard and soft tissues. The successful outcome of this case confirms the efficacy of microsurgical management in GTR using correctly-chosen biomaterials for the enucleation of the radicular cyst. Therefore, dentists should be aware of the potential benefits

of these biomaterials that may be used to provide quality treatments for their patients.

FOOTNOTES

Author contributions: Gómez Mireles JC, Martínez Carrillo EK, Alcalá Barbosa K, Gutiérrez Cortés E and Lomeli Martínez SM contributed to diagnosis, surgical treatment and manage of patients; González Ramos J, González Gómez LA and Bayardo González RA contributed to acquisition and analysis of data; Gómez Mireles JC, Lomeli Martínez SM, Martínez Carrillo EK, Alcalá Barbosa K drafted manuscript and critically revised manuscript; Gómez Mireles JC, Martínez Carrillo EK and Lomeli Martínez SM contributed to conceptualization and supervision; all authors have read and approved the final manuscript.

Informed consent statement: Informed written consent was obtained from the patients for publication of this report and any accompanying images.

Conflict-of-interest statement: The authors declare that they have no conflict of interest to disclose.

CARE Checklist (2016) statement: The authors have read the CARE checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Mexico

ORCID number: Juan Carlos Gómez Mireles 0009-0001-7790-5591; Eugenia Karina Martínez Carrillo 0000-0002-4475-3552; Katia Alcalá Barbosa 0009-0007-0604-0272; Evangelina Gutiérrez Cortés 0000-0003-0115-1110; Javier González Ramos 0009-0004-2574-8354; Luis Antonio González Gómez 0009-0003-7179-7133; Rubén Alberto Bayardo González 0000-0002-0851-348X; Sarah Monserrat Lomeli Martínez 0000-0002-0569-1387.

S-Editor: Liu H

L-Editor: A

P-Editor: Li X

REFERENCES

- 1 Deshmukh J, Shrivastava R, Bharath KP, Mallikarjuna R. Giant radicular cyst of the maxilla. *BMJ Case Rep* 2014; **2014** [PMID: 24792022 DOI: 10.1136/bcr-2014-203678]
- 2 Rios Osorio N, Caviedes-Bucheli J, Mosquera-Guevara L, Adames-Martinez JS, Gomez-Pinto D, Jimenez-Jimenez K, Avendano Maz H, Bornacelly-Mendoza S. The Paradigm of the Inflammatory Radicular Cyst: Biological Aspects to be Considered. *Eur Endod J* 2023; **8**: 20-36 [PMID: 36748442 DOI: 10.14744/eej.2022.26918]
- 3 Shetty S, Angadi PV, Rekha K. Radicular cyst in deciduous maxillary molars: a rarity. *Head Neck Pathol* 2010; **4**: 27-30 [PMID: 20237985 DOI: 10.1007/s12105-009-0152-z]
- 4 Elhakim A, Kim S, Kim E, Elshazli AH. Preserving the vitality of teeth adjacent to a large radicular cyst in periapical microsurgery: a case report with 4-year follow-up. *BMC Oral Health* 2021; **21**: 382 [PMID: 34344347 DOI: 10.1186/s12903-021-01738-2]
- 5 Vega Llauradó A, Ayuso Montero R, Teixidor Olmo I, Salas Enric J, Mari Roig A, López López J. Opcionesterapéuticas en quistes odontogénicos. Revisión. *Av Odontoestomatol* 2013; **29**: 81-93
- 6 Govindaraju L, Antony DP, S P. Surgical Management of Radicular Cyst With the Application of a Natural Platelet Concentrate: A Case Report. *Cureus* 2023; **15**: e33992 [PMID: 36811049 DOI: 10.7759/cureus.33992]
- 7 Noda A, Abe M, Shinozaki-Ushiku A, Ohata Y, Zong L, Abe T, Hoshi K. A Bilocular Radicular Cyst in the Mandible with Tooth Structure Components Inside. *Case Rep Dent* 2019; **2019**: 6245808 [PMID: 31565444 DOI: 10.1155/2019/6245808]
- 8 Lin LM, Huang GT, Rosenberg PA. Proliferation of epithelial cell rests, formation of apical cysts, and regression of apical cysts after periapical wound healing. *J Endod* 2007; **33**: 908-916 [PMID: 17878074 DOI: 10.1016/j.joen.2007.02.006]
- 9 Yadav VS, Salaria SK, Bhatia A, Yadav R. Periodontal microsurgery: Reaching new heights of precision. *J Indian Soc Periodontol* 2018; **22**: 5-11 [PMID: 29568165 DOI: 10.4103/jisp.jisp_364_17]
- 10 Setzer FC, Shah SB, Kohli MR, Karabucak B, Kim S. Outcome of endodontic surgery: a meta-analysis of the literature--part 1: Comparison of traditional root-end surgery and endodontic microsurgery. *J Endod* 2010; **36**: 1757-1765 [PMID: 20951283 DOI: 10.1016/j.joen.2010.08.007]
- 11 Montero-Mirallés P, Ibáñez-Barranco R, Cabanillas-Balsera D, Areal-Quecuty V, Sánchez-Domínguez B, Martín-González J, Segura-Egea JJ, Jiménez-Sánchez MC. Biomaterials in periapical regeneration after microsurgical endodontics: A narrative review. *J Clin Exp Dent* 2021; **13**: e935-e940 [PMID: 34603623 DOI: 10.4317/jced.58651]
- 12 Barzuna M, Cárdenas T. Use of a bioceramic (Biodentine™) in retrodental filling in apicoectomy report of a clinical case. *Vital Dent* 2014; **20**: 69-74
- 13 Alvarez HA, Santillana CH, Portocarrero JP. Quiste odontogénico inflamatorio: Reporte de caso. *Salud Vida Sipanense* 2020; 132-143 [DOI: 10.26495/svs.v7i2.1473]
- 14 Malkondur Ö, Karapinar Kazandağ M, Kazazoğlu E. A review on biodentine, a contemporary dentine replacement and repair material. *Biomed Res Int* 2014; **2014**: 160951 [PMID: 25025034 DOI: 10.1155/2014/160951]

- 15 **Rajasekharan S**, Martens LC, Cauwels RGE, Anthonappa RP. Biodentine™ material characteristics and clinical applications: a 3 years literature review and update. *Eur Arch Paediatr Dent* 2018; **19**: 1-22 [PMID: [29372451](#) DOI: [10.1007/s40368-018-0328-x](#)]
- 16 **Zubizarreta-Macho Á**, Tosin R, Tosin F, Velasco Bohórquez P, San Hipólito Marín L, Montiel-Company JM, Mena-Álvarez J, Hernández Montero S. Influence of Guided Tissue Regeneration Techniques on the Success Rate of Healing of Surgical Endodontic Treatment: A Systematic Review and Network Meta-Analysis. *J Clin Med* 2022; **11** [PMID: [35207335](#) DOI: [10.3390/jcm11041062](#)]
- 17 **Agrawal AA**. Platelet rich fibrin is not a barrier membrane! Or is it? *World J Clin Cases* 2023; **11**: 2396-2404 [PMID: [37123322](#) DOI: [10.12998/wjcc.v11.i11.2396](#)]
- 18 **Ren Y**, Fan L, Alkildani S, Liu L, Emmert S, Najman S, Rimashevskiy D, Schnettler R, Jung O, Xiong X, Barbeck M. Barrier Membranes for Guided Bone Regeneration (GBR): A Focus on Recent Advances in Collagen Membranes. *Int J Mol Sci* 2022; **23** [PMID: [36499315](#) DOI: [10.3390/ijms232314987](#)]
- 19 **Sumangali A**, Naik AC, Mohan N, Gautam N, Abrol S, Mustafa M, Tiwari H. Bone Regenerative Biomaterials in Periapical Surgery: A Systemic Review and Meta-Analysis. *J Pharm Bioallied Sci* 2021; **13**: S933-S937 [PMID: [35017901](#) DOI: [10.4103/jpbs.jpbs_386_21](#)]
- 20 **Morillo ML**, Peralta HV, Vásquez VM, Guzmán DJ, Harris RJ. Guided bone regeneration in the treatment of radicular cyst: Case report. *Den J* 2023; **26**: 99-103 [DOI: [10.29166/odontologia.vol26.n1.2023-e4025](#)]
- 21 **Tseng SK**, Tsai YL, Li UM, Jeng JH. Radicular cyst with actinomycotic infection in an upper anterior tooth. *J Formos Med Assoc* 2009; **108**: 808-813 [PMID: [19864202](#) DOI: [10.1016/S0929-6646\(09\)60409-5](#)]
- 22 **Molgatini S**, Rey E, Basilaki J, Mosca C, Galante R, Gliosca L. [Isolation of *Pseudomonas stutzeri* from an odontogenic inflammatory cyst: Diagnostic relevance]. *Rev Argent Microbiol* 2017; **49**: 32-38 [PMID: [28188092](#) DOI: [10.1016/j.ram.2016.10.006](#)]
- 23 **Nilesh K**, Dadhich A. Unusually large radicular cyst presenting in the maxillary sinus. *BMJ Case Rep* 2020; **13** [PMID: [32907869](#) DOI: [10.1136/bcr-2020-236582](#)]
- 24 **Cortellini P**, Tonetti MS. Minimally invasive surgical technique and enamel matrix derivative in intra-bony defects. I: Clinical outcomes and morbidity. *J Clin Periodontol* 2007; **34**: 1082-1088 [PMID: [17953696](#) DOI: [10.1111/j.1600-051X.2007.01144.x](#)]
- 25 **Ribeiro FV**, Nociti Júnior FH, Sallum EA, Sallum AW, Casati MZ. Use of enamel matrix protein derivative with minimally invasive surgical approach in intra-bony periodontal defects: clinical and patient-centered outcomes. *Braz Dent J* 2010; **21**: 60-67 [PMID: [20464323](#) DOI: [10.1590/s0103-64402010000100010](#)]
- 26 **Aimetti M**, Ferrarotti F, Mariani GM, Romano F. A novel flapless approach versus minimally invasive surgery in periodontal regeneration with enamel matrix derivative proteins: a 24-month randomized controlled clinical trial. *Clin Oral Investig* 2017; **21**: 327-337 [PMID: [27044318](#) DOI: [10.1007/s00784-016-1795-2](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

