World Journal of *Clinical Cases*

World J Clin Cases 2024 March 16; 12(8): 1382-1548





Published by Baishideng Publishing Group Inc

W J C C World Journal of Clinical Cases

Contents

Thrice Monthly Volume 12 Number 8 March 16, 2024

EDITORIAL

1382 Latest updates on structure and recommendations of cardiac rehabilitation programs in chronic heart failure

Kourek C, Briasoulis A, Magouliotis DE, Skoularigis J, Xanthopoulos A

1388 Immunoglobulin A glomerulonephropathy: A review

El Labban M, Surani S

ORIGINAL ARTICLE

Retrospective Study

1395 Serum proteins differentially expressed in gestational diabetes mellitus assessed using isobaric tag for relative and absolute quantitation proteomics

Cao WL, Yu CP, Zhang LL

Observational Study

1406 Effect of comprehensive nursing on the quality of life and swallowing function in individuals diagnosed with ischemic stroke

Hu HF, Sang YF, Xiao YQ

META-ANALYSIS

Safety and effectiveness of butorphanol in epidural labor analgesia: A protocol for a systematic review and 1416 meta-analysis

Tang GC, He M, Huang ZZ, Cheng Y

CASE REPORT

1422 Left lower lobe sleeve resection for the clear cell variant of pulmonary mucoepidermoid carcinoma: A case report

Yu XH, Wang WX, Yang DS, Gong LH

1430 Lower extremity peripherally inserted central catheter placement ectopic to the ascending lumbar vein: A case report

Zhu XJ, Zhao L, Peng N, Luo JM, Liu SX

- Effect of foot reflexology on an infant with sensorineural hearing loss: A case report 1437 Lee YJ, Chen MQ, Dong J
- 1442 Development of immature ovarian teratoma after mature teratoma in a girl with familial ovarian teratoma: A case report

Ju UC, Kang WD, Kim SM



.	World Journal of Clinical Cases	
Conten	Thrice Monthly Volume 12 Number 8 March 16, 2024	
1448	Metastatic clear cell sarcoma of the pancreas: A rare case report	
	Liu YJ, Zou C, Wu YY	
1454	3M syndrome patient with a novel mutation: A case report	
	Luo MR, Dai SM, Li Y, Wang Q, Liu H, Gao P, Liu JY, Chen J, Zhao SJ, Yin GY	
1461	Appendiceal intussusception complicated by adenocarcinoma of the cecum: A case report	
	Long Y, Xiang YN, Huang F, Xu L, Li XY, Zhen YH	
1467	Malignant triton tumor in the abdominal wall: A case report	
	Yang HJ, Kim D, Lee WS, Oh SH	
1474	Multilocular thymic cysts can be easily misdiagnosed as malignant tumor on computer tomography: A case report	
	Sun J, Yang QN, Guo Y, Zeng P, Ma LY, Kong LW, Zhao BY, Li CM	
1481	Diagnosis of poorly differentiated adenocarcinoma of the stomach by confocal laser endomicroscopy: A case report	
	Lou JX, Wu Y, Huhe M, Zhang JJ, Jia DW, Jiang ZY	
1487	High-grade pancreatic intraepithelial neoplasia diagnosed based on changes in magnetic resonance cholangiopancreatography findings: A case report	
	Furuya N, Yamaguchi A, Kato N, Sugata S, Hamada T, Mizumoto T, Tamaru Y, Kusunoki R, Kuwai T, Kouno H, Kuraoka K, Shibata Y, Tazuma S, Sudo T, Kohno H, Oka S	
1497	Chronic infectious unilateral giant thyroid cyst related to diabetes mellitus: A case report	
	Liu JB, Zhang SL, Jiang WL, Sun HK, Yang HC	
1504	Multiple thoracic and abdominal foregut duplication cysts: A case report	
	Alsinan TA, Altokhais TI	
1510	Advanced cervix cancer patient with chemotherapy-induced grade IV myelosuppression achieved complete remission with cadonilimab: A case report	
	Zhu R, Chen TZ, Sun MT, Zhu CR	
1517	Detection of 4p16.3 deletion and 11p15.5p15.4 gain in a boy by comparative genomic hybridization array: A case report	
	Kaya I	
1523	Eccrine porocarcinoma in the tempus of an elderly woman: A case report	
	Wu ZW, Zhu WJ, Huang S, Tan Q, You C, Hu DG, Li LN	
1530	Analysis and monitoring of drug therapy in a patient with peptic ulcer complicated by infection: A case report	
	Gou YP, Huang Y, Chen L, Zheng W, Zheng YH	
1536	Primary gastroduodenal tuberculosis presenting as gastric outlet obstruction: A case report and review of literature	
	Ali AM, Mohamed YG, Mohamud AA, Mohamed AN, Ahmed MR, Abdullahi IM, Saydam T	



Conton	World Journal of Clinical Cases
Conten	Thrice Monthly Volume 12 Number 8 March 16, 2024
1544	Clinical manifestations and the prenatal diagnosis of trisomy 7 mosaicism: Two case reports <i>Hou F, Li Y, Jin H</i>

Contents

Thrice Monthly Volume 12 Number 8 March 16, 2024

ABOUT COVER

Peer Reviewer of World Journal of Clinical Cases, Makram Koubaa, MD, Assistant Professor, Department of Infectious Diseases, Hedi Chaker University Hospital, Sfax 3029, Tunisia. makram.koubaa@gmail.com

AIMS AND SCOPE

The primary aim of World Journal of Clinical Cases (WJCC, World J Clin Cases) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Hua-Ge Yu; Production Department Director: Xu Guo; Editorial Office Director: Jin-Lei Wang,

NAME OF JOURNAL World Journal of Clinical Cases	INSTRUCTIONS TO AUTHORS https://www.wjgnet.com/bpg/gerinfo/204
ISSN ISSN 2307-8960 (online)	GUIDELINES FOR ETHICS DOCUMENTS
LAUNCH DATE April 16, 2013	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
FREQUENCY Thrice Monthly	PUBLICATION ETHICS
EDITORS-IN-CHIEF Bao Gan Bang Salim Sugari, Jaray Tadayar Chudak, Gaorga Kantagaaraa	PUBLICATION MISCONDUCT
Maurizio Serati	ADTICLE DOCESSING CHADGE
https://www.wjgnet.com/2307-8960/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS
March 16, 2024	https://www.wjgnet.com/bpg/GerInfo/239
COPYRIGHT © 2024 Baishideng Publishing Group Inc	ONLINE SUBMISSION https://www.f6publishing.com

© 2024 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: office@baishideng.com https://www.wjgnet.com



W J C C World Journal of Clinical Cases

Submit a Manuscript: https://www.f6publishing.com

World J Clin Cases 2024 March 16; 12(8): 1523-1529

DOI: 10.12998/wjcc.v12.i8.1523

ISSN 2307-8960 (online)

CASE REPORT

Eccrine porocarcinoma in the tempus of an elderly woman: A case report

Zhi-Wei Wu, Wen-Jie Zhu, Shan Huang, Qi Tan, Cong You, Dian-Gui Hu, Long-Nian Li

Specialty type: Medicine, research and experimental

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): 0 Grade C (Good): C, C Grade D (Fair): 0 Grade E (Poor): 0

P-Reviewer: Cassell III AK, Liberia; Sira AM, Egypt

Received: December 16, 2023 Peer-review started: December 16, 2023

First decision: January 10, 2024 Revised: January 19, 2024 Accepted: February 22, 2024 Article in press: February 22, 2024 Published online: March 16, 2024



Zhi-Wei Wu, Wen-Jie Zhu, Cong You, Long-Nian Li, Department of Dermatology, First Affiliated Hospital of Gannan Medical University, Ganzhou 341000, Jiangxi Province, China

Shan Huang, Department of Laser and Cosmetic Dermatology, Ganzhou Dermatology Hospital, Ganzhou 341000, Jiangxi Province, China

Qi Tan, Department of Dermatology, Tongxiang Dermatology Hospital, Jiaxing 314000, Zhejiang Province, China

Dian-Gui Hu, Department of Infectious Diseases, The Fifth people's Hospital of Ganzhou, Ganzhou 341000, Jiangxi Province, China

Corresponding author: Long-Nian Li, PhD, Associate Chief Physician, Department of Dermatology, First Affiliated Hospital of Gannan Medical University, No. 23 Qingnian Road, Zhanggong District, Ganzhou 341000, Jiangxi Province, China. li longnian@foxmail.com

Abstract

BACKGROUND

Eccrine porocarcinoma (EPC) is a rare skin tumor that mainly affects the elderly population. Tumors often present with slow growth and a good prognosis. EPCs are usually distinguished from other skin tumors using histopathology and immunohistochemistry. However, surgical management alone may be inadequate if the tumor has metastasized. However, currently, surgical resection is the most commonly used treatment modality.

CASE SUMMARY

A seventy-four-year-old woman presented with a slow-growing nodule in her left temporal area, with no obvious itching or pain, for more than four months. Histopathological examination showed small columnar and short spindle-shaped cells; thus, basal cell carcinoma was suspected. However, immunohistochemical analysis revealed the expression of cytokeratin 5/6, p63 protein, p16 protein, and Ki-67 antigen (40%), and EPC was taken into consideration. The skin biopsy was repeated, and hematoxylin and eosin staining revealed ductal differentiation in some cells. Finally, the patient was diagnosed with EPC, and Mohs micrographic surgery was performed. We adapted follow-up visits in a year and not found any recurrence of nodules.

CONCLUSION

This case report emphasizes the diagnosis and differentiation of EPC.



Key Words: Eccrine porocarcinoma; Sweat gland carcinoma; Histopathology; Immunohistochemistry; Prognosis; Case report

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Eccrine porocarcinomas (EPCs) are rare skin tumors that grow slowly with no obvious discomfort. They are difficult to distinguish from basal cell carcinomas, particularly those that occur on the face. Histopathological examination and immunohistochemistry are necessary to diagnose atypical lesions, their importance is demonstrated in our case. Multiple biopsies at different times are necessary to visualize ductal differentiation in cells. Our case reported a brief period of EPC in the temporal region of an elderly woman. We hope that our case findings will help in better understanding of EPCs.

Citation: Wu ZW, Zhu WJ, Huang S, Tan Q, You C, Hu DG, Li LN. Eccrine porocarcinoma in the tempus of an elderly woman: A case report. World J Clin Cases 2024; 12(8): 1523-1529 URL: https://www.wjgnet.com/2307-8960/full/v12/i8/1523.htm DOI: https://dx.doi.org/10.12998/wjcc.v12.i8.1523

INTRODUCTION

The skin is the largest organ in the human body, and its tissue structure is complex. Skin tumors can be identified through the actions of various pathogenic factors. The diversity of skin tumors makes their diagnoses difficult. Here, we report a case of eccrine porocarcinoma (EPC) of the tempus in an elderly woman. EPC is a rare skin tumor which originates from sweat glands. It has the high misdiagnosis rate due to the rarity of the disease and the atypicality of the site of occurrence[1,2]. Our case described the diagnostic methods of EPC from the histopathological examination and immunohistochemistry, and detailed explanation of the current treatment methods for EPC.

CASE PRESENTATION

Chief complaints

A nodule on the left tempus, which was growing for more than four months.

History of present illness

A soybean-sized papule had developed on the left tempus of a seventy-four-year-old woman, four months prior to presentation. The nodule developed slowly with the accumulation of adherent grevish brown oily scales on the surface; the patient experienced slight itching without pain. Because the number of nodules affected her facial appearance, she sought help from a local hospital. Based on clinical manifestations, basal cell carcinoma was the primary diagnosis. A biopsy and histopathological examination were performed as additional diagnostic evaluations.

History of past illness

The patient had no history of tumor and immune disease.

Personal and family history

Similar symptoms were not reported in her family.

Physical examination

An approximately 1.5 cm × 2 cm nodule was observed on the left tempus of the patient's face. The nodule had a rough surface and an irregular boundary (Figure 1). A black suture line was present at the center of the lesion. No intumescent lymph nodes were palpable in the neck.

Laboratory examinations

Skin biopsy was performed. The first histopathological examination revealed small columnar and short spindle-shaped cells (Figure 2A). The later one revealed part of the cells appearing to be ductally differentiated. It can be seen many pore structures (Figure 2B). Some tumor cells have large and hyperchromatic nuclei, and it can be seen lots of apoptotic bodies (Figure 2C). Immunohistochemistry revealed the expression of cytokeratin (CK) 5/6, p63 protein (p63), p16 protein (p16), and Ki-67 antigen (40%) (Figure 3). Blood biochemistry showed no significant abnormalities.

Imaging examinations

The computed tomography of the skull, and ultrasound of superficial lymph glands showed no remarkable findings.



WJCC | https://www.wjgnet.com



Figure 1 The clinical features of the patient: The broad bean-sized greyish nodule with a greasy crust.



Figure 2 Pathological examination. A: Hematoxylin and eosin (HE) (× 100); B: HE (× 100); C: HE (× 200).

 Jaishideng®
 WJCC
 https://www.wignet.com
 1525



Figure 3 Immunohistochemical examination. A: Cytokeratin5/6: (+); B: Ki-67 antigen: (40%+); C: P16 protein: (+); D: P63 Protein: (+).

FINAL DIAGNOSIS

The presence of ductal differentiation in cells ruled out basal cell carcinoma. The small cell size, ductal differentiation, and lumen formation excluded squamous cell carcinoma (SCC). Combined with clinical manifestations and pathological findings, EPC was diagnosed.

TREATMENT

After obtaining informed consent from the patient, Mohs micrographic surgery was performed (Figure 4).

OUTCOME AND FOLLOW-UP

No recurrence of nodules was noted during the follow-up visits in the next year.

DISCUSSION

EPC is a rare pernicious tumor originating from the sweat glands of the skin. It was first reported by Pinkus and Mehregan^[3]. Studies have shown that from 2000 to 2018, the incidence of EPC in the United States was 0.045 per 100000



Baishideng® WJCC | https://www.wjgnet.com



Figure 4 The clinical features of the patient: The incision after the resection of the lesion.

individuals[4]; with the improvement of diagnostic technology, the diagnostic rate of EPC has also increased annually. Furthermore, the elderly population is at a higher risk of EPCs. However, there have been reports of young women with EPC whose conditions progressed significantly during pregnancy[5]. No significant sex differences have been noted for this disease[6]. The most common location of presentation is the head and neck (39.9%), followed by the lower extremities (33.9%)[6]. Similar to our case, EPC has been reported in the left temporal area, with the appearance of a verrucous lesion [7]. However, the tumors had a longer growth time (one year to twenty years)[7] than that in our case (four months). What makes it special is that the EPC in our case seemed growing more quickly than the reported ones[7], which needs more study to clarify it. EPCs usually appear as hard, purple or red nodular plaques with smooth warty or ulcerated surfaces[8,9], and are often associated with a malignant transformation that occurs over a previous poroma originating from the intraepidermal components of the sweat glands[10]. The pathogenesis of EPC is complex and varied, and includes benign tumor transformation, sun exposure, and immunosuppression[11]. In the present case, the patient was an elderly woman living in a rural area, who was exposed to ultraviolet radiation.

Currently, surgical resection is the most commonly accepted treatment option for EPC especially the Mohs micrographic surgery. Mohs micrographic surgery is widely used in melanoma and some infrequent skin tumors, it has the high cure rate and low recurrence rate in EPC patients with Mohs micrographic surgery[12,13]. Upon assessing the metastasis of EPC, radiotherapy, chemotherapy and targeted therapy are considered to have an adjuvant effect on EPC. The treatment of paclitaxel pus cetuximab was reported to have an almost complete response for EPC[14]. EPCs are divided into "pushing" and "infiltrating" with predictable propulsion margins; the latter have a greater risk of local metastasis. The rate of mitosis, lymphovascular invasion, and tumor invasion depth (> 7 mm) can increase the risk of metastasis[10]. The most common metastatic organs for EPC are the nearby lymph nodes (57.7%), followed by the lungs (12.8%) and the liver and brain (both 9%)[6].

Currently, there is no clear diagnostic protocol for EPCs. Hematoxylin and eosin (HE) staining and immunohistochemistry have a high diagnostic value for EPCs. HE staining of EPC tissue sections reveals irregular tumors consisting of characteristic porous basal-like cells, with part of the cells appearing to be ductally differentiated at the least, in addition to the characteristics of typical tumor cells. The presence of a tumor duct is necessary for the diagnosis of EPC[9,10]. Additionally, immunochemical markers play a significant role in the diagnosis. One study investigated whether the immunohistochemical expression of p53 protein, retinoblastoma protein, and p16 could differentiate EPCs from poromas and found that six EPC cases (43%) showed an abnormal expression of p16[15]. The strong expression of p63 and CK5/6 were detected in EPCs and p63 could distinguish EPCs from metastatic adenocarcinomas[16,17]. EPCs often need to be distinguished from basal cell carcinomas, which consist of morphologically basaloid cells with scant cytoplasm, elongated hyperchromatic nuclei, intercellular bridges, and peripheral cells that are not fence-like [18]. Compared to SCCs, EPCs have a small cell size, ductal differentiation, and lumen formation. The expression of engrailed homeobox 1 and CK19 were found to be able to improve the accuracy of histologic diagnosis of EPC, and these two markers are independent that distinguish EPCs from SCCs[2]. Furthermore, the express of mast/stem cell growth factor receptor also become a more and more crucial role to make a distinction between EPCs and SCCs[19]. EPC is also often misdiagnosed as amelanotic melanoma, granuloma, telangiectasia, or fibroma^[20]. Additionally, we need to summarize more EPC cases in this area of skin to summarize the clinical characteristics of the disease better. We also need to follow up for a longer time period to evaluate the recurrence.

Zaishidena® WJCC | https://www.wjgnet.com

CONCLUSION

EPC is a rare skin tumor that grows slowly and is heterogeneous and nonspecific. EPCs in the temporal area are even rarer. Owing to its similar clinical presentation with those of SCC and basal cell carcinoma, its differential diagnosis is warranted. Diagnosis is often made through histopathological examination and immunohistochemistry, and specific immunochemical markers still need to be studied to help distinguish EPCs from other skin tumors. Surgical resection is currently the most commonly used mode of treatment, especially Mohs micrographic surgery. We hope that this report will provide a better understanding of EPCs.

FOOTNOTES

Author contributions: Li LN and You C conceptualized the design; Wu ZW, Zhu WJ, Huang S, and Tan Q conducted an investigation; You C, Wu ZW and Li LN perform software operations; Li LN, You C, and Hu DG to supervise; Li LN and Zhu WJ do the writing; Li LN conducts resource collection, writing and editing.

Informed consent statement: Informed written consent was obtained from the patient and her families for publication of this report and any accompanying images.

Conflict-of-interest statement: The authors declare that they have no competing interests.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: China

4631-0905; Cong You 0000-0002-0314-5884; Dian-Gui Hu 0009-0000-4627-1701; Long-Nian Li 0000-0002-3441-037X.

S-Editor: Liu H L-Editor: A P-Editor: Zheng XM

REFERENCES

- Li YX, Gudi M, Yan Z. Primary Eccrine Porocarcinoma of the Breast: A Case Report and Review of Literature. Case Rep Oncol Med 2022; 1 2022: 4042298 [PMID: 35685061 DOI: 10.1155/2022/4042298]
- 2 Miura K, Akashi T, Namiki T, Hishima T, Bae Y, Sakurai U, Murano K, Shiraishi J, Warabi M, Tanizawa T, Tanaka M, Bhunchet E, Kumagai J, Ayabe S, Sekiya T, Ando N, Shintaku H, Kinowaki Y, Tomii S, Kirimura S, Kayamori K, Yamamoto K, Ito T, Eishi Y. Engrailed Homeobox 1 and Cytokeratin 19 Are Independent Diagnostic Markers of Eccrine Porocarcinoma and Distinguish It From Squamous Cell Carcinoma. Am J Clin Pathol 2020; 154: 499-509 [PMID: 32556098 DOI: 10.1093/ajcp/agaa066]
- Pinkus H, Mehregan AH. Epidermotropic errccrine carcinoma. A case combining features of errccrine poroma and Paget's dermatosis. Arch 3 Dermatol 1963; 88: 597-606 [PMID: 14060075 DOI: 10.1001/archderm.1963.01590230105015]
- Gibbs DC, Yeung H, Blalock TW. Incidence and trends of cutaneous adnexal tumors in the United States in 2000-2018: A population-based 4 study. J Am Acad Dermatol 2023; 88: 226-228 [PMID: 35525505 DOI: 10.1016/j.jaad.2022.04.052]
- Jeon SP, Kang SJ, Jung SJ. Rapidly growing eccrine porocarcinoma of the face in a pregnant woman. J Craniofac Surg 2014; 25: 715-717 5 [PMID: 24621740 DOI: 10.1097/01.scs.0000436739.52418.97]
- Salih AM, Kakamad FH, Baba HO, Salih RQ, Hawbash MR, Mohammed SH, Othman S, Saeed YA, Habibullah IJ, Muhialdeen AS, Nawroly 6 RO, Hammood ZD, Abdulkarim NH. Porocarcinoma; presentation and management, a meta-analysis of 453 cases. Ann Med Surg (Lond) 2017; 20: 74-79 [PMID: 28721214 DOI: 10.1016/j.amsu.2017.06.027]
- Mulinari-Brenner FA, Mukai MM, Bastos CA, Filho EA, Santamaria JR, Neto JF. [Eccrine porocarcinoma: report of four cases and literature 7 review]. An Bras Dermatol 2009; 84: 519-523 [PMID: 20098856 DOI: 10.1590/s0365-05962009000500012]
- Sawaya JL, Khachemoune A. Poroma: a review of eccrine, apocrine, and malignant forms. Int J Dermatol 2014; 53: 1053-1061 [PMID: 8 24697501 DOI: 10.1111/ijd.12448]
- Meriläinen AS, Pukkala E, Böhling T, Koljonen V. Malignant Eccrine Porocarcinoma in Finland During 2007 to 2017. Acta Derm Venereol 9 2021; 101: adv00363 [PMID: 33313937 DOI: 10.2340/00015555-3718]
- Robson A, Greene J, Ansari N, Kim B, Seed PT, McKee PH, Calonje E. Eccrine porocarcinoma (malignant eccrine poroma): a 10 clinicopathologic study of 69 cases. Am J Surg Pathol 2001; 25: 710-720 [PMID: 11395548 DOI: 10.1097/00000478-200106000-00002]
- Puttonen M, Isola J, Ylinen O, Böhling T, Koljonen V, Sihto H. UV-induced local immunosuppression in the tumour microenvironment of 11 eccrine porocarcinoma and poroma. Sci Rep 2022; 12: 5529 [PMID: 35365704 DOI: 10.1038/s41598-022-09490-5]



- Wittenberg GP, Robertson DB, Solomon AR, Washington CV. Eccrine porocarcinoma treated with Mohs micrographic surgery: A report of 12 five cases. Dermatol Surg 1999; 25: 911-913 [PMID: 10594609 DOI: 10.1046/j.1524-4725.1999.99121.x]
- 13 Tidwell WJ, Mayer JE, Malone J, Schadt C, Brown T. Treatment of eccrine porocarcinoma with Mohs micrographic surgery: a cases series and literature review. Int J Dermatol 2015; 54: 1078-1083 [PMID: 26205087 DOI: 10.1111/ijd.12997]
- Godillot C, Boulinguez S, Riffaud L, Sibaud V, Chira C, Tournier E, Paul C, Meyer N. Complete response of a metastatic porocarcinoma 14 treated with paclitaxel, cetuximab and radiotherapy. Eur J Cancer 2018; 90: 142-145 [PMID: 29233613 DOI: 10.1016/j.ejca.2017.11.009]
- Zahn J, Chan MP, Wang G, Patel RM, Andea AA, Bresler SC, Harms PW. Altered Rb, p16, and p53 expression is specific for porocarcinoma 15 relative to poroma. J Cutan Pathol 2019; 46: 659-664 [PMID: 31012122 DOI: 10.1111/cup.13480]
- Qureshi HS, Ormsby AH, Lee MW, Zarbo RJ, Ma CK. The diagnostic utility of p63, CK5/6, CK 7, and CK 20 in distinguishing primary 16 cutaneous adnexal neoplasms from metastatic carcinomas. J Cutan Pathol 2004; 31: 145-152 [PMID: 14690459 DOI: 10.1111/j.0303-6987.2004.00147.x]
- Ivan D, Nash JW, Prieto VG, Calonje E, Lyle S, Diwan AH, Lazar AJ. Use of p63 expression in distinguishing primary and metastatic 17 cutaneous adnexal neoplasms from metastatic adenocarcinoma to skin. J Cutan Pathol 2007; 34: 474-480 [PMID: 17518775 DOI: 10.1111/j.1600-0560.2006.00644.x]
- Grimme H, Petres A, Bergen E, Wiemers S, Schöpf E, Vanscheidt W. Metastasizing porocarcinoma of the head with lethal outcome. 18 Dermatology 1999; 198: 298-300 [PMID: 10393458 DOI: 10.1159/000018135]
- Joshy J, Mistry K, Levell NJ, van Bodegraven B, Vernon S, Rajan N, Craig P, Venables ZC. Porocarcinoma: a review. Clin Exp Dermatol 19 2022; 47: 1030-1035 [PMID: 35149987 DOI: 10.1111/ced.15126]
- Snow SN, Reizner GT. Eccrine porocarcinoma of the face. J Am Acad Dermatol 1992; 27: 306-311 [PMID: 1325487 DOI: 20 10.1016/0190-9622(92)70187-k]





Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

