

World Journal of *Clinical Cases*

World J Clin Cases 2024 March 26; 12(9): 1549-1713



Contents

Thrice Monthly Volume 12 Number 9 March 26, 2024

EDITORIAL

- 1549 Multidisciplinary approach toward enhanced recovery after surgery for total knee arthroplasty improves outcomes

Nag DS, Swain A, Sahu S, Sahoo A, Wadhwa G

- 1555 Using clinical cases to guide healthcare

Colwill M, Baillie S, Pollok R, Poullis A

ORIGINAL ARTICLE

Retrospective Study

- 1560 Analysis of the causes of primary revision after unicompartmental knee arthroplasty: A case series

Zhao JL, Jin X, Huang HT, Yang WY, Li JH, Luo MH, Liu J, Pan JK

- 1569 Efficacy and safety of minimally invasive laparoscopic surgery under general anesthesia for ovarian cancer

Qin X, Chen C, Liu Y, Hua XH, Li JY, Liang MJ, Wu F

- 1578 Factors influencing Frey syndrome after parotidectomy with acellular dermal matrix

Chai XD, Jiang H, Tang LL, Zhang J, Yue LF

Clinical Trials Study

- 1585 Allogeneic mesenchymal stem cells may be a viable treatment modality in cerebral palsy

Boyalı O, Kabatas S, Civelek E, Özdemir O, Bahar-Ozdemir Y, Kaplan N, Savrunlu EC, Karaöz E

Observational Study

- 1597 Clinical characteristics of acute non-varicose upper gastrointestinal bleeding and the effect of endoscopic hemostasis

Wang XJ, Shi YP, Wang L, Li YN, Xu LJ, Zhang Y, Han S

Clinical and Translational Research

- 1606 Construction of the underlying circRNA-miRNA-mRNA regulatory network and a new diagnostic model in ulcerative colitis by bioinformatics analysis

Yuan YY, Wu H, Chen QY, Fan H, Shuai B

- 1622 Exploring the autophagy-related pathogenesis of active ulcerative colitis

Gong ZZ, Li T, Yan H, Xu MH, Lian Y, Yang YX, Wei W, Liu T

CASE REPORT

- 1634 Low-molecular-weight heparin and preeclampsia — does the sword cut both ways? Three case reports and review of literature

Shan D, Li T, Tan X, Hu YY

- 1644** Pulmonary alveolar proteinosis induced by X-linked agammaglobulinemia: A case report
Zhang T, Li M, Tan L, Li X
- 1649** Gradient inflammation in the pancreatic stump after pancreaticoduodenectomy: Two case reports and review of literature
Wang TG, Tian L, Zhang XL, Zhang L, Zhao XL, Kong DS
- 1660** Low interleukin-10 level indicates a good prognosis in *Salmonella enterica* serovar typhimurium-induced pediatric hemophagocytic lymphohistiocytosis: A case report
Chen YY, Xu XZ, Xu XJ
- 1669** Multi-systemic melioidosis in a patient with type 2 diabetes in non-endemic areas: A case report and review of literature
Ni HY, Zhang Y, Huang DH, Zhou F
- 1677** Endoscopic ultrasound-guided tissue sampling induced pancreatic duct leak resolved by the placement of a pancreatic stent: A case report
Kim KH, Park CH, Cho E, Lee Y
- 1685** Upadacitinib for refractory ulcerative colitis with primary nonresponse to infliximab and vedolizumab: A case report
Xu X, Jiang JW, Lu BY, Li XX
- 1691** Exogenous insulin autoimmune syndrome: A case report and review of literature
Xu LL, Chen JX, Cheng JP, Luo N
- 1698** Unexplained fetal tachycardia: A case report
Wang H, Duan RZ, Bai XJ, Zhang BT, Wang J, Song WX
- 1704** Challenging anticoagulation therapy for multiple primary malignant tumors combined with thrombosis: A case report and review of literature
Chen JX, Xu LL, Cheng JP, Xu XH

LETTER TO THE EDITOR

- 1712** Epinephrine also acts on beta cells and insulin secretion
Zabulienė L, Ilias I

ABOUT COVER

Peer Reviewer of *World Journal of Clinical Cases*, Luca Mezzetto, MD, Surgeon, Department of Vascular Surgery, University Hospital of Verona, Verona 37126, Italy. luca.mezzetto@aovr.veneto.it

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Zi-Hang Xu, Production Department Director: Xiang Li, Cover Editor: Jin-Lai Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

March 26, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Endoscopic ultrasound-guided tissue sampling induced pancreatic duct leak resolved by the placement of a pancreatic stent: A case report

Ki-Hyun Kim, Chang Hwan Park, Eunae Cho, Yohan Lee

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Nagaya M, Japan;
Tang W, China

Received: December 4, 2023

Peer-review started: December 4, 2023

First decision: January 9, 2024

Revised: February 3, 2024

Accepted: March 5, 2024

Article in press: March 5, 2024

Published online: March 26, 2024



Ki-Hyun Kim, Eunae Cho, Yohan Lee, Department of Internal Medicine, Chonnam National University Hospital, Gwangju 61469, South Korea

Chang Hwan Park, Department of Gastroenterology, Chonnam National University Hospital, Gwangju 61469, South Korea

Corresponding author: Chang Hwan Park, MD, PhD, Professor, Department of Gastroenterology, Chonnam National University Hospital, No. 42 Jebong-ro Donggu, Gwangju 61469, South Korea. p1052ccy@hanmail.net

Abstract

BACKGROUND

Pancreatic ductal leaks complicated by endoscopic ultrasonography-guided tissue sampling (EUS-TS) can manifest as acute pancreatitis.

CASE SUMMARY

A 63-year-old man presented with persistent abdominal pain and weight loss. Diagnosis: Laboratory findings revealed elevated carbohydrate antigen 19-9 (5920 U/mL) and carcinoembryonic antigen (23.7 ng/mL) levels. Magnetic resonance imaging of the pancreas revealed an approximately 3 cm ill-defined space-occupying lesion in the inferior aspect of the head, with severe encasement of the superior mesenteric artery. Pancreatic ductal adenocarcinoma was confirmed after pathological examination of specimens obtained by EUS-TS using the fanning method. Interventions and outcomes: The following day, the patient experienced severe abdominal pain with high amylase (265 U/L) and lipase (1173 U/L) levels. Computed tomography of the abdomen revealed edematous wall thickening of the second portion of the duodenum with adjacent fluid collections and a suspicious leak from either the distal common bile duct or the main pancreatic duct in the head. Endoscopic retrograde cholangiopancreatography revealed dye leakage in the head of the main pancreatic duct. Therefore, a 5F 7 cm linear plastic stent was deployed into the pancreatic duct to divert the pancreatic juice. The patient's abdominal pain improved immediately after pancreatic stent insertion, and amylase and lipase levels normalized within a week. Neoadjuvant chemotherapy was then initiated.

CONCLUSION

Using the fanning method in EUS-TS can inadvertently cause damage to the pan-

creatic duct and may lead to clinically significant pancreatitis. Placing a pancreatic stent may immediately resolve acute pancreatitis and shorten the waiting time for curative therapy. When using the fanning method during EUS-TS, ductal structures should be excluded to prevent pancreatic ductal leakage.

Key Words: Endoscopic ultrasound-guided tissue sampling; Pancreatitis; Pancreatic duct leak; Pancreatic stent; Case report

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Endoscopic ultrasound guided tissue sampling is a crucial procedure for histological diagnosis of pancreatic lesions, but it may occasionally be accompanied by unforeseen complications. The Fanning method is a technique that can enhance diagnostic accuracy, but it also carries the risk of unintended ductal injury. Here, we report a case where the use of the Fanning method during endoscopic ultrasound guided tissue sampling resulted in leakage of the pancreatic duct. We successfully managed this complication by performing pancreatic duct stent *via* endoscopic retrograde cholangiopancreatography.

Citation: Kim KH, Park CH, Cho E, Lee Y. Endoscopic ultrasound-guided tissue sampling induced pancreatic duct leak resolved by the placement of a pancreatic stent: A case report. *World J Clin Cases* 2024; 12(9): 1677-1684

URL: <https://www.wjgnet.com/2307-8960/full/v12/i9/1677.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v12.i9.1677>

INTRODUCTION

Endoscopic ultrasound-guided tissue sampling (EUS-TS) has become the gold standard for histological diagnosis of solid pancreatic lesions[1,2]. Although EUS-TS is considered relatively safe, unexpected complications inevitably arise. Acute pancreatitis is one of the more severe complications of EUS-TS, as it can delay the surgical schedule of resectable lesions and can even make curative treatment impossible[3]. Pancreatic ductal leaks often complicate the situation, while recent treatment guidelines for acute pancreatitis, such as early feeding, can be counterproductive. EUS-TS puts patients at risk of developing pancreatic ductal leaks if normal pancreatic ducts are damaged during the procedure[4]. Fortunately, pancreatic ductal leaks can be adequately treated by placing a pancreatic stent to divert the leaking pancreatic juice into the duodenal lumen[5]. According to our review of English-language scientific literature, this is the first recorded case of a pancreatic ductal leak developing after timely EUS-TS treatment with pancreatic stent placement. Therefore, clinical suspicion is key to promptly diagnosing and treating pancreatic ductal leaks in patients who present with acute pancreatitis after EUS-TS. Herein, we present a case of acute pancreatitis caused by a pancreatic ductal leak after EUS-TS that was treated with a pancreatic stent.

CASE PRESENTATION

Chief complaints

A 63-year-old man visited the outpatient department with a weight loss.

History of present illness

The weight loss has been significant, with a decrease of 14 kg over the past two months, and upper abdominal pain was occurred 10 d ago.

History of past illness

He had been diagnosed with hypertension and diabetes mellitus four years earlier.

Personal and family history

He was a chronic alcoholic and a current smoker.

Physical examination

The physical examination showed tenderness in the epigastric area without rebound tenderness.

Laboratory examinations

His white blood cell count was 3500/mm³ (reference value: 6000–10000/mm³), lipase level was 120 U/L (reference value: 7–60 U/L), carbohydrate antigen 19-9 level was 5920 U/mL (reference value: < 37 U/mL), and carcinoembryonic antigen level was 23.7 ng/mL (reference value: 0–4.7 ng/mL). All other laboratory test results were within the normal ranges.




Figure 1 Magnetic resonance imaging. A and B: An approximately 3 cm, ill-defined, space-occupying lesion in the inferior aspect of the head with high signal intensity on T1- and T2-weighted imaging (orange arrows).




Figure 2 Magnetic resonance imaging. A: An ill-defined space-occupying lesion invading the distal common bile duct and main pancreatic duct, both of which were narrowed, and the upstream main pancreatic duct dilated (orange arrow); B: The lesion severely encased the superior mesenteric artery (orange arrow).

Imaging examinations

Magnetic resonance imaging of the pancreas showed an approximately 3 cm-sized, ill-defined space-occupying lesion in the inferior aspect of the head, with high signal intensity on T1- and T2-weighted imaging (Figure 1). The distal common bile duct and main pancreatic duct adjacent to the lesion were narrowed, the upstream main pancreatic duct was dilated, and the superior mesenteric artery was severely encased (Figure 2).

FINAL DIAGNOSIS

EUS-TS was performed for pathologic diagnosis, which demonstrated a 36.0 mm × 32.9 mm sized ill-defined, relatively heterogeneous, hypoechoic mass in the pancreatic head. EUS-TS was performed with a 22-gauge EZ shot (EZ shot 3 plus; Olympus, Tokyo, Japan) using the fanning method (Figure 3). Pathological examination confirmed a pancreatic ductal adenocarcinoma. The day after the procedure, the patient experienced severe abdominal pain in the right upper quadrant and epigastric areas. His follow-up amylase, lipase, and C-reactive protein levels were 265 U/L (reference range: 43–116 U/L), 1173 U/L (reference range: 7–60 U/L), and 27.7 mg/dL (reference range: 0–3 mg/dL), respectively. Computed tomography (CT) of the abdomen two days after EUS-TS revealed edematous wall thickening of the second portion of the duodenum with adjacent fluid collections and a suspicious leak from either the distal common bile duct or the main pancreatic duct in the head (Figure 4).

TREATMENT

During the initial endoscopic retrograde cholangiopancreatography (ERCP), selective biliary cannulation could only be achieved through the ampulla of Vater. Although no definitive dye leak from the distal CBD was observed, a 7F 7 cm double pigtail plastic stent (ZSO-7-7; Cook Medical, Indiana, United States) was deployed over the stricture to allow for effective biliary drainage (Figure 5). Selective pancreatic cannulation was attempted several times without success. Despite biliary diversion, the patient continued to experience severe abdominal pain for four days. Follow-up abdominal CT showed no resolution of the suspected leak, suggesting a pancreatic ductal leak. During the second ERCP, a separate orifice was observed below the biliary orifice, which was not observed in the first ERCP. Selective pancreatic cannulation was easily achieved through the orifice (Figure 6). As the pancreatogram showed a dye leak in the head portion of the




Figure 3 Endoscopic ultrasound-guided tissue sampling. Endoscopic ultrasound (EUS) showed a 36.0 mm × 32.9 mm sized ill-defined relative heterogeneous hypoechoic mass in the pancreatic head. EUS-guided tissue sampling (EUS-TS) was performed using a 22-gauge EZ Shot (EZ Shot 3 Plus; Olympus, Tokyo, Japan) using the fanning method.




Figure 4 Computed tomography of the abdomen two days after endoscopic ultrasound-guided tissue sampling. A and B: Edematous wall thickening of the second portion of the duodenum with adjacent fluid collection was detected and a leak was suspected (orange arrows).

main pancreatic duct, a 5F 7 cm linear plastic stent was deployed to divert the pancreatic juice (Figure 7).

OUTCOME AND FOLLOW-UP

Immediately after insertion of the pancreatic stent, his abdominal pain markedly improved, and amylase and lipase levels normalized within a week. Two weeks after the pancreatic stent insertion procedure, follow-up abdominal CT imaging revealed the disappearance of adjacent fluid collection and edematous change of head of pancreas findings and healing of the duct leakage site (Figure 8). Based on these observations, neoadjuvant chemotherapy was initiated.




Figure 5 First endoscopic retrograde cholangiopancreatography. A: Selective biliary cannulation was achieved only through the ampulla of Vater; B: There is no definitive dye leakage from the distal common bile duct; C: A 7 F 7 cm-sized double pigtail plastic stent was deployed for effective biliary drainage.




Figure 6 Second endoscopic retrograde cholangiopancreatography. A: A separate orifice below the biliary orifice that had not been observed during the first endoscopic retrograde cholangiopancreatography is noted (white arrow); B: Selective pancreatic cannulation was easily achieved through a separate orifice.

DISCUSSION

EUS-TS is a revolutionary technique that shifts the narrative surrounding the diagnosis of pancreatic lesions, such as cysts and solid lesions, and can target small pancreatic tumors. Before the advent of EUS-TS, ultrasonography-guided percutaneous transabdominal TS played a major role in significant seeding risk[5]. However, despite the many benefits of EUS-TS, it can also cause adverse side effects such as bleeding, pancreatitis, infection, duodenal perforation, abscesses, and sepsis[6]. Pancreatic ductal leaks are also a rare complication of EUS-TS, with only two cases reported thus far in the English-language scientific literature[1,7]. Unlike the current case, these cases involved pancreatic juice collected as a pseudocyst or as pancreatic ascites.

The patient in this case study was diagnosed with acute pancreatitis as he had suffered persistent severe pain over 24 h and had amylase and lipase levels more than three times the upper normal limit immediately after EUS-TS. Sequential abdominal CT imaging studies suggested a pancreatic ductal leak from the dilated main pancreatic duct, which a second ERCP confirmed. The patient recovered immediately after pancreatic stent placement to divert the leaking pancreatic juice from the intraperitoneal cavity to the duodenal lumen.

Acute pancreatitis (AP) is a rare EUS-TS complication. Recent research has demonstrated that a recent history of acute pancreatitis and undertaking EUS-TS through more than 5 mm of the normal parenchyma can lead to a much greater risk of acute pancreatitis[3]. However, a history of acute pancreatitis was not found in the current case, and the EUS-TS fine needle passed through less than 5 mm of the normal parenchyma. Another study reported that patients with a branch




Figure 7 Second endoscopic retrograde cholangiopancreatography. A: A pancreatogram showing dye leakage in the head of the main pancreatic duct (orange arrow); B and C: A 5F 7 cm linear plastic stent was deployed to divert the pancreatic juice.



Figure 8 Computed tomography of the abdomen two weeks after pancreatic stent placement. Disappearance of adjacent fluid collection and edematous change of head of pancreas, and healing of the duct leakage site finding were seen.

duct-type intraductal papillary mucin-secreting neoplasm are at risk of developing acute pancreatitis after EUS-TS[2]. However, unlike that study, the current case involved pancreatic ductal adenocarcinoma (PDAC).

In the present case, the fanning method was used to achieve a high diagnostic accuracy. The fanning method in EUS-TS involves sampling by cycles of in-and-out needle passes addressed in multiple directions within a solid lesion to obtain more target tissue[8]. While blood vessels were visualized and excluded by color Doppler during EUS-TS in the current case, pancreatic ductal structures without color Doppler might have a greater chance of being damaged by the fanning technique than the standard technique. A retrospective analysis of the EUS images in the current case revealed a suspicious ductal structure within the target fanning area (Figure 9). When using the fanning method during EUS-TS, ductal structures without color Doppler imaging should be excluded to prevent pancreatic ductal leaks. To avoid pancreatic ductal injury, a thorough evaluation of the ductal structures should be performed before EUS-TS with Doppler imaging in the pancreatic mass, and the fanning method range should be limited in the field without suspicious ductal structures.

Although PDAC generally has a poor prognosis because it is frequently detected at an advanced stage, recent studies demonstrate that innovations in neoadjuvant chemotherapy have provided opportunities to proceed to curative surgery, thereby improving long-term outcomes in patients with borderline resectable and locally advanced PDAC[9]. Therefore, prompt treatment with neoadjuvant chemotherapy is critical for improving prognosis, while a delay in treatment could cause borderline resectable and locally advanced PDAC to become distant and metastatic. Acute pancreatitis initiates a hypercatabolic state caused by elevated protein catabolism and an inflammatory cytokine storm[10]. Recent guidelines dictate that patients with acute pancreatitis may benefit from early oral or enteral nutrition, which is in contrast to managing pancreatic ductal leaks as they require prolonged fasting and parenteral nutrition[4]. The hypercatabolic state and prolonged fasting in acute pancreatitis due to pancreatic ductal leaks could make PDAC patients unsuitable for toxic chemotherapy. In the current case, placement of the pancreatic stent helped to reduce the waiting interval for neoadjuvant chemotherapy, as the patient's acute pancreatitis was immediately resolved.




Figure 9 Retrospective analysis of endoscopic ultrasound imaging in the current case. Suspected ductal structure within the fanning area of the target (orange arrow).

CONCLUSION

In conclusion, pancreatic ductal leakage caused by EUS-TS can lead to acute pancreatitis. Therefore, imaging studies should be immediately performed after EUS-TS to rule out pancreatic ductal leaks in patients with clinical signs and symptoms of AP. If a pancreatic ductal leak is suspected, ERCP must confirm the leak. Pancreatic stent placement may allow patients to immediately recover from acute pancreatitis and shorten the waiting time for curative therapy. When using the fanning method during EUS-TS, ductal structures without color Doppler imaging should be excluded to prevent pancreatic ductal leaks.

FOOTNOTES

Author contributions: Kim KH, Cho E, Lee Y designed the research study; Kim KH and Park CH analyzed the data and wrote the manuscript; all authors have read and approve the final manuscript.

Informed consent statement: Written informed consent was obtained from the patient for publication of this report and all accompanying images.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: South Korea

ORCID number: Ki-Hyun Kim 0009-0007-8558-0540; Chang Hwan Park 0000-0002-2995-8779; Eunae Cho 0000-0001-5931-4643; Yohan Lee 0000-0002-0643-3304.

S-Editor: Gong ZM

L-Editor: A

P-Editor: Xu ZH

REFERENCES

- 1 Reddymasu S, Oropeza-Vail MM, Williamson S, Jafri F, Olyae M. Pancreatic leak after endoscopic ultrasound guided fine needle aspiration managed by transpapillary pancreatic duct stenting. *JOP* 2011; **12**: 489-490 [PMID: 21904078]
- 2 Siddiqui AA, Shahid H, Shah A, Khurana T, Huntington W, Ghumman SS, Loren DE, Kowalski TE, Laique S, Hayat U, Eloubeidi MA. High risk of acute pancreatitis after endoscopic ultrasound-guided fine needle aspiration of side branch intraductal papillary mucinous neoplasms. *Endosc Ultrasound* 2015; **4**: 109-114 [PMID: 26020044 DOI: 10.4103/2303-9027.156728]
- 3 Ribeiro A, Goel A. The Risk Factors for Acute Pancreatitis after Endoscopic Ultrasound Guided Biopsy. *Korean J Gastroenterol* 2018; **72**:

- 135-140 [PMID: [30270595](#) DOI: [10.4166/kjg.2018.72.3.135](#)]
- 4 **Larsen M**, Kozarek R. Management of pancreatic ductal leaks and fistulae. *J Gastroenterol Hepatol* 2014; **29**: 1360-1370 [PMID: [24650171](#) DOI: [10.1111/jgh.12574](#)]
- 5 **Micames C**, Jowell PS, White R, Paulson E, Nelson R, Morse M, Hurwitz H, Pappas T, Tyler D, McGrath K. Lower frequency of peritoneal carcinomatosis in patients with pancreatic cancer diagnosed by EUS-guided FNA vs. percutaneous FNA. *Gastrointest Endosc* 2003; **58**: 690-695 [PMID: [14595302](#) DOI: [10.1016/s0016-5107\(03\)02009-1](#)]
- 6 **Tian G**, Ye Z, Zhao Q, Jiang T. Complication incidence of EUS-guided pancreas biopsy: A systematic review and meta-analysis of 11 thousand population from 78 cohort studies. *Asian J Surg* 2020; **43**: 1049-1055 [PMID: [31974051](#) DOI: [10.1016/j.asjsur.2019.12.011](#)]
- 7 **Ikarashi S**, Tsuchiya A, Hayashi K, Terai S. Delayed pancreatic ductal leakage after EUS-FNA for autoimmune pancreatitis. *Endosc Ultrasound* 2019; **8**: 277-278 [PMID: [30719995](#) DOI: [10.4103/eus.eus_55_18](#)]
- 8 **Bang JY**, Magee SH, Ramesh J, Trevino JM, Varadarajulu S. Randomized trial comparing fanning with standard technique for endoscopic ultrasound-guided fine-needle aspiration of solid pancreatic mass lesions. *Endoscopy* 2013; **45**: 445-450 [PMID: [23504490](#) DOI: [10.1055/s-0032-1326268](#)]
- 9 **Versteijne E**, van Dam JL, Suker M, Janssen QP, Groothuis K, Akkermans-Vogelaar JM, Besselink MG, Bonsing BA, Buijsen J, Busch OR, Creemers GM, van Dam RM, Eskens FALM, Festen S, de Groot JWB, Groot Koerkamp B, de Hingh IH, Homs MYV, van Hooft JE, Kerver ED, Luelmo SAC, Neelis KJ, Nuyttens J, Paardekooper GMRM, Patijn GA, van der Sangen MJC, de Vos-Geelen J, Wilmink JW, Zwinderman AH, Punt CJ, van Tienhoven G, van Eijck CHJ; Dutch Pancreatic Cancer Group. Neoadjuvant Chemoradiotherapy Versus Upfront Surgery for Resectable and Borderline Resectable Pancreatic Cancer: Long-Term Results of the Dutch Randomized PREOPANC Trial. *J Clin Oncol* 2022; **40**: 1220-1230 [PMID: [35084987](#) DOI: [10.1200/jco.21.02233](#)]
- 10 **Rinninella E**, Annetta MG, Serricchio ML, Dal Lago AA, Miggiano GA, Mele MC. Nutritional support in acute pancreatitis: from physiopathology to practice. An evidence-based approach. *Eur Rev Med Pharmacol Sci* 2017; **21**: 421-432 [PMID: [28165542](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

