

World Journal of *Clinical Cases*

World J Clin Cases 2018 November 6; 6(13): 577-715





REVIEW

- 577 Role of bile acids in colon carcinogenesis
Nguyen TT, Ung TT, Kim NH, Jung YD

MINIREVIEWS

- 589 Update on global epidemiology of viral hepatitis and preventive strategies
Jefferies M, Rauff B, Rashid H, Lam T, Rafiq S

ORIGINAL ARTICLE

Case Control Study

- 600 Iron metabolism disorders in patients with hepatitis B-related liver diseases
Gao YH, Wang JY, Liu PY, Sun J, Wang XM, Wu RH, He XT, Tu ZK, Wang CG, Xu HQ, Niu JQ

Retrospective Cohort Study

- 611 Impact of an acute hemodynamic response-guided protocol for primary prophylaxis of variceal bleeding
Forteza JI, Puente Á, Ruiz P, Ezcurra I, Vaquero J, Cuadrado A, Arias-Loste MT, Cabezas J, Llerena S, Iruzubieta P, Rodríguez-Lope C, Huelin P, Casafont F, Fábrega E, Crespo J

Retrospective Study

- 624 Effect of a region-wide incorporation of an algorithm based on the 2012 international consensus guideline on the practice pattern for the management of pancreatic cystic neoplasms in an integrated health system
Nguyen AK, Girg A, Tekeste T, Chang K, Adeyemo M, Eskandari A, Alonso E, Yaramada P, Chaya C, Ko A, Burke E, Roggow I, Butler R, Kawatkar A, Lim BS

- 632 Usefulness of colonic tattooing using indocyanine green in patients with colorectal tumors
Park JH, Moon HS, Kwon IS, Yun GY, Lee SH, Park DH, Kim JS, Kang SH, Lee ES, Kim SH, Sung JK, Lee BS, Jeong HY

Randomized Clinical Trial

- 641 *Helicobacter pylori* may be an initiating factor in newly diagnosed ulcerative colitis patients: A pilot study
Mansour L, El-Kalla F, Kobtan A, Abd-Elsalam S, Yousef M, Soliman S, Ali LA, Elkhawany W, Amer I, Harras H, Hagar MM, Elhendawy M

META-ANALYSIS

- 650 Photodynamic therapy for middle-advanced stage upper gastrointestinal carcinomas: A systematic review and meta-analysis
Chen B, Xiong L, Chen WD, Zhao XH, He J, Zheng YW, Kong FH, Liu X, Zhang ZJ, Miao XY

CASE REPORT

- 659** Successful rescue of acute liver failure and hemophagocytic lymphohistiocytosis following varicella infection: A case report and review of literature
Zhang LN, Guo W, Zhu JH, Guo Y
- 666** Bilateral thoracic kidneys combined with inferior vena cava located behind the anterior abdominal wall: A case report and review of literature
Peng XX, Cheng SA, Liang QL, Luo XB, Hong XC, Yuan GL, Zhang HJ
- 671** Incident hepatocellular carcinoma developing during tenofovir alafenamide treatment as a rescue therapy for multi-drug resistant hepatitis B virus infection: A case report and review of the literature
Lu JC, Liu LG, Lin L, Zheng SQ, Xue Y
- 675** Possible connection between elevated serum α -fetoprotein and placental necrosis during pregnancy: A case report and review of literature
Yu MY, Xi L, Zhang JX, Zhang SC
- 679** Laparoscopic pancreatic duct incision and stone removal and T-type tube drainage for pancreatic duct stone: A case report and review of literature
Bai Y, Yu SA, Wang LY, Gong DJ
- 683** Detection of a unicentric type of Castleman-like mass at the site of adrenal gland: A case report and review of literature
Chen J, Yang C, Liang CZ
- 688** Systemic lupus erythematosus complicated by noncirrhotic portal hypertension: A case report and review of literature
Yang QB, He YL, Peng CM, Qing YF, He Q, Zhou JG
- 694** Natural killer/T-cell lymphoma with concomitant syndrome of inappropriate antidiuretic hormone secretion: A case report and review of literature
Liu QB, Zheng R
- 703** Successful treatment of pyoderma gangrenosum with concomitant immunoglobulin A nephropathy: A case report and review of literature
Li XL, Ma ZG, Huang WH, Chai EQ, Hao YF



- 707 Highlighting the importance of early diagnosis in progressive multi-organ involvement of IgG4-related disease: A case report and review of literature

Xue J, Wang XM, Li Y, Zhu L, Liu XM, Chen J, Chi SH

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Byung-Wook Kim, MD, PhD, Professor, Division of Gastroenterology, Department of Internal Medicine, Incheon St. Mary's Hospital, the Catholic University of Korea, Incheon 21431, South Korea

AIM AND SCOPE

World Journal of Clinical Cases (*World J Clin Cases*, *WJCC*, online ISSN 2307-8960, DOI: 10.12998) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

The primary task of *WJCC* is to rapidly publish high-quality Autobiography, Case Report, Clinical Case Conference (Clinicopathological Conference), Clinical Management, Diagnostic Advances, Editorial, Field of Vision, Frontier, Medical Ethics, Original Articles, Clinical Practice, Meta-Analysis, Minireviews, Review, Therapeutics Advances, and Topic Highlight, in the fields of allergy, anesthesiology, cardiac medicine, clinical genetics, clinical neurology, critical care, dentistry, dermatology, emergency medicine, endocrinology, family medicine, gastroenterology and hepatology, geriatrics and gerontology, hematology, immunology, infectious diseases, internal medicine, obstetrics and gynecology, oncology, ophthalmology, orthopedics, otolaryngology, pathology, pediatrics, peripheral vascular disease, psychiatry, radiology, rehabilitation, respiratory medicine, rheumatology, surgery, toxicology, transplantation, and urology and nephrology.

INDEXING/ABSTRACTING

World Journal of Clinical Cases (*WJCC*) is now indexed in PubMed, PubMed Central, Science Citation Index Expanded (also known as SciSearch®), and Journal Citation Reports/Science Edition. The 2018 Edition of Journal Citation Reports cites the 2017 impact factor for *WJCC* as 1.931 (5-year impact factor: N/A), ranking *WJCC* as 60 among 154 journals in Medicine, General and Internal (quartile in category Q2).

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Yun-XiaoJian Wu*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Ying Dou*
Proofing Editorial Office Director: *Jin-Lei Wang*

NAME OF JOURNAL
World Journal of Clinical Cases

ISSN
ISSN 2307-8960 (online)

LAUNCH DATE
April 16, 2013

FREQUENCY
Semimonthly

EDITORS-IN-CHIEF
Sandro Vento, MD, Department of Internal Medicine, University of Botswana, Private Bag 00713, Gaborone, Botswana

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com/2307-8960/editorialboard.htm>

EDITORIAL OFFICE
Jin-Lei Wang, Director

World Journal of Clinical Cases
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
November 6, 2018

COPYRIGHT

© 2018 Baishideng Publishing Group Inc. Articles published by this Open Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT

All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS

<http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION

<http://www.f6publishing.com>

Case Control Study

Iron metabolism disorders in patients with hepatitis B-related liver diseases

Yan-Hang Gao, Jing-Yun Wang, Pei-Yan Liu, Jing Sun, Xiao-Mei Wang, Rui-Hong Wu, Xiu-Ting He, Zheng-Kun Tu, Chun-Guang Wang, Hong-Qin Xu, Jun-Qi Niu

Yan-Hang Gao, Jing-Yun Wang, Pei-Yan Liu, Jing Sun, Xiao-Mei Wang, Rui-Hong Wu, Hong-Qin Xu, Jun-Qi Niu, Department of Hepatology, The First Hospital of Jilin University, Jilin University, Changchun 130021, Jilin Province, China

Jing-Yun Wang, Department of Gastroenterology, Weihaiwei People's Hospital, Weihai 264200, Shandong Province, China

Jing Sun, Department of Gastroenterology, Heping Hospital, Changzhi Medical College, Changzhi 046011, Shanxi Province, China

Xiu-Ting He, Department of Geriatrics, The First Hospital of Jilin University, Changchun 130021, Jilin Province, China

Zheng-Kun Tu, Department of Translational Medicine, The First Hospital of Jilin University, Changchun 130021, Jilin Province, China

Chun-Guang Wang, Department of Surgery, The Second Hospital of Jilin University, Changchun 130021, Jilin Province, China

Jun-Qi Niu, Jilin Provincial Key Laboratory of Infectious Diseases, Laboratory of Molecular Virology, Changchun 130021, Jilin Province, China

ORCID number: Yan-Hang Gao (0000-0002-7248-8290); Jing-Yun Wang (0000-0003-0987-8322); Pei-Yan Liu (0000-0002-4616-7012); Jing Sun (0000-0001-8910-5965); Xiao-Mei Wang (0000-0003-2606-4013); Rui-Hong Wu (0000-0001-7585-0059); Xiu-Ting He (0000-0002-6110-9531); Zheng-Kun Tu (0000-0001-6803-2015); Hong-Qin Xu (0000-0002-7022-7732); Jun-Qi Niu (0000-0002-5187-7297).

Author contributions: Niu JQ, Wang CG and Xu HQ designed the research; Niu JQ, Gao YH, Wang CG, Liu PY, Sun J, and He XT treated the patients and collected the materials and clinical data from patients; Wang JY, Wang XM, and Tu ZK performed the assays; Xu HQ analyzed the data; Gao YH and Xu HQ wrote the paper. Wang CG, Xu HQ, and Niu JQ are co-corresponding authors. Each author contributed important intellectual content during manuscript drafting or revision and accepts accountability

for the overall work by ensuring that questions pertaining to the accuracy or integrity of any portion of the work are appropriately investigated and resolved.

Supported by the National Science and Technology Major Project, No. 2014ZX10002002 and No. 2017ZX10202202; the National Key Research Plan "Precision Medicine Research" Key Project, No. 2017YFC0908103; the National Natural Science Foundation of China, No. 81700534; and Program for JLU Science and Technology Innovative Research Team, No. 2017TD-08.

Institutional review board statement: The study was reviewed and approved by the First Hospital Ethical Committee of Jilin University.

Informed consent statement: All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

Conflict-of-interest statement: The authors declare that they have no competing interests.

Data sharing statement: Dataset available from the corresponding author at junqi_niu@163.com or hongqinxu86@jlu.edu.cn.

STROBE Statement: The authors have read the STROBE Statement-checklist of items, and the manuscript was prepared and revised according to the STROBE Statement-checklist of items

Open-Access: This is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Correspondence to: Jun-Qi Niu, PhD, Professor, Department of Hepatology, First Hospital of Jilin University, No. 71, Xinmin Street, Changchun 130021, Jilin Province, China. junqi_niu@163.com
Telephone: +86-431-88783486
Fax: +86-431-85612468

Received: July 17, 2018
Peer-review started: July 17, 2018
First decision: August 25, 2018
Revised: September 16, 2018
Accepted: October 16, 2018
Article in press: October 16, 2018
Published online: November 6, 2018

Abstract

AIM

To investigate the relationship between levels of iron metabolism markers and hepatitis B virus (HBV)-related chronic liver diseases.

METHODS

This case-control study with 318 participants included 78 cases of chronic hepatitis B, 85 cases of HBV-related liver cirrhosis, 77 cases of HBV-related hepatocellular carcinoma, and 78 healthy controls. Markers of iron metabolism were detected in participants. Hematological and biochemical parameters and HBV-DNA were assessed. Child-Pugh grade and Barcelona Clinic Liver Cancer stage were determined for each hepatocellular carcinoma patient. Perls' staining was performed on liver sections. The SPSS program was used for all statistical analyses, and statistical significance was considered if a *P*-value < 0.05.

RESULTS

Significantly higher serum ferritin and lower serum hepcidin levels were detected in all groups of HBV-infected patients compared with healthy controls. Serum iron, total iron binding capacity, and serum transferrin levels were significantly lower in patients with cirrhosis and hepatocellular carcinoma, whereas the hepcidin level was higher than that in chronic hepatitis B patients. Correlation analysis indicated that serum hepcidin was negatively correlated with HBV-DNA load (*P* < 0.01). Serum ferritin and transferrin saturation levels increased proportionally to the extent of liver cirrhosis and poorer Child-Pugh scores (*P* < 0.05). The decreased serum iron and transferrin saturation levels were significantly correlated with a smaller hepatocellular carcinoma tumor burden according to Barcelona Clinic Liver Cancer staging. Liver histology showed a clearly increasing trend in iron deposition in the liver tissues with increased fibrosis, which became prominent at stages 3 (severe liver fibrosis) and 4 (cirrhosis).

CONCLUSION

Iron metabolism disorders occur in patients with HBV-related liver diseases. The serum markers of iron me-

tabolism disorders vary in different stages of HBV-related liver diseases.

Key words: Hepatocellular carcinoma; Hepcidin; Iron staining; Hepatitis B; Liver fibrogenesis

© **The Author(s) 2018.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: The relationship between hepatitis B viruses (HBV) related chronic liver diseases and levels of components in iron metabolism and the corresponding impact on liver disease severity have not been clearly described. In our study, we found that significantly higher serum ferritin and lower serum hepcidin levels were detected in all groups of HBV-infected patients compared with healthy controls. Serum iron, total iron binding capacity, and serum transferrin levels were significantly lower in patients with cirrhosis and hepatocellular carcinoma, whereas the hepcidin level was higher than that in chronic hepatitis B patients. In conclusion, iron metabolism disorders are present in patients with HBV-related liver diseases. The characteristics of iron metabolism disorders in different development stages of HBV-related liver diseases varied.

Gao YH, Wang JY, Liu PY, Sun J, Wang XM, Wu RH, He XT, Tu ZK, Wang CG, Xu HQ, Niu JQ. Iron metabolism disorders in patients with hepatitis B-related liver diseases. *World J Clin Cases* 2018; 6(13): 600-610 Available from: URL: <http://www.wjgnet.com/2307-8960/full/v6/i13/600.htm> DOI: <http://dx.doi.org/10.12998/wjcc.v6.i13.600>

INTRODUCTION

Chronic hepatitis B virus (HBV) infection affects 240 million people worldwide and can lead to chronic liver diseases including chronic hepatitis B (CHB), cirrhosis, and hepatocellular carcinoma (HCC). An estimated 788000 patients die of HBV-related liver diseases each year^[1]. In managing chronic HBV infection, the objective is to detect liver injury early and then stop the progression of liver disease through effective treatment. To achieve this goal, a clear understanding of various factors involved in HBV pathogenesis is required.

Maintenance of iron metabolism within the physiological range is essential to human health. The liver synthesizes several proteins that are required for iron metabolism in addition to iron storage. Chronic liver injury and histological alteration impact iron metabolism, and altered iron metabolism may aggravate liver injury^[2-4]. For instance, both experimental and clinical studies have suggested that sustained iron retention in hepatocytes aggravates liver injury and is associated with higher risks of developing fibrosis, cirrhosis, and HCC in CHB patients^[5-7]. In 1981, Blumberg *et al*^[8] reported that serum iron indices tend to be higher in CHB patients than in those who had HBV infection cl-

eared. Moderate iron accumulation in liver cells has been observed in patients with nonalcoholic fatty liver disease, and the down-regulation of the iron exporter ferroportin-1 in affected cells was suggested to be responsible for the reduced iron clearance from the liver^[9]. Hepcidin, a homeostatic regulator, modulates iron absorption in the intestinal mucosal epithelial cells, iron recycle in macrophages, and iron storage in liver cells. The serum level of hepcidin is decreased in patients with chronic hepatitis C^[10,11]. Advanced alcoholic liver disease is associated with abnormally accumulated iron in the liver^[12]. Wang *et al.*^[13] found that the mean serum hepcidin level was significantly higher in non-cirrhotic CHB patients than in healthy controls. Another study found that successful treatment with lamivudine may lead to a reduction in the serum ferritin level and hepatic siderosis in CHB, implying that hepatic siderosis is secondary to HBV-related liver injury or may contribute to liver injury^[14].

The relationship between HBV-related chronic liver diseases and levels of components in iron metabolism and the corresponding impact on liver disease severity have not been clearly described. The aim of this study was to characterize changes in serum iron components in patients with HBV-related diseases and identify correlations between changes in iron metabolism and HBV-related liver injury.

MATERIALS AND METHODS

Participants

This study included 78 healthy controls, 78 CHB patients including 25 diagnosed by biopsy, 85 patients with HBV-related liver cirrhosis (LC) including 5 diagnosed by biopsy, and 77 with HBV-related HCC. All patients were ethnic Han Chinese recruited between January 2012 and December 2014 in The First Hospital of Jilin University in Changchun, China. The groups were matched for age and sex. CHB, cirrhosis, and HCC were diagnosed based on the criteria outlined in the Guidelines for the Prevention and Treatment of CHB (2010 version adapted at the 10th National Conference on Viral Hepatitis and Hepatology 2011, China). Briefly, CHB was diagnosed when an individual had been HBsAg positive for more than 6 mo and had clinical manifestations of hepatitis and abnormal liver biochemistry or histological evidence of liver injury. The diagnosis of LC was made for cases meeting one of the following terms: (1) histological evidence based on liver biopsy; and (2) biochemical/hematological and imaging evidence of liver dysfunction or portal hypertension accompanied by hypersplenism/gastroesophageal varices/hepatic encephalopathy/ascites. LC cases were divided into subgroups according to the Child-Pugh classification^[15]. HBV-related cirrhosis was defined as cirrhosis with a history of CHB or chronic HBV infection. Fibrosis in patients with biopsy was staged according to the METAVIR scoring system^[16]. HBV-related HCC was diagnosed based on: (1) a solid mass detected

by one of the imaging modalities (computed tomography, magnetic resonance imaging, or ultrasonography) or (2) cytological or histological evidence of liver cancer cells appearing in the liver of a chronic HBV-infected patient. All patients were positive for one or multiple HBV serological markers. Patients were excluded if they: (1) were co-infected with hepatitis A, C, or D or had alcoholic liver disease or drug-induced hepatitis; (2) had diseases of the blood system including hereditary hemochromatosis (types I, II, III, and IV) and chronic anemia; or (3) had an iron overload disease.

The healthy controls were negative for hepatitis B surface antigen and hepatitis C antibody and had normal liver transaminase levels and abdominal ultrasound examinations. Individuals who manifested iron metabolism disorders were excluded. Written informed consent was obtained from all participants, and the study was approved by The First Hospital Ethical Committee of Jilin University.

Laboratory analysis

Blood samples were collected from all patients after overnight fasting. Serum iron, ferritin, and total iron binding capacity were analyzed with an automatic clinical chemistry analyzer (Hitachi 7600 D-210, Japan). Transferrin (TF) was determined using an automatic protein analyzer (BN II System, SIEMENS, Germany). HBV serological markers were measured with a ThermoScientific MultiskanGo (Kehua, Shanghai, China). The Elecsys 2010 and Roche COBAS e 411 immunoassay systems (Roche Diagnostics, Grenzach, Germany) were used to quantitate serum HBsAg, and the COBAS AmpliPrep/COBAS TaqMan assay (Roche Molecular Diagnostics, Grenzach, Germany) was used for quantitating HBV DNA in serum, which had a lower limit of detection at 15 U/mL. Biochemical markers were determined with the Synchron LX120 Autoanalyzer (Beckman Coulter, Brea, CA, United States). Hepcidin was measured using an enzyme-linked immuno sorbent assay kit (IBL, Germany), and transferrin saturation was calculated as a percentage [serum iron/total iron binding capacity (TIBC) × 100%].

Histological assessment

Liver biopsy was performed with a 16 G Tru-Cut needle guided by color Doppler ultrasound. A biopsy that was 1.5 cm or longer or a liver section that contained at least five portal tracts was considered eligible for diagnosis. The liver sections from each biopsy were stained with hematoxylin and eosin and Perls' stain, respectively. Liver fibrosis was staged using the METAVIR scoring system, which consists of five stages: S0 (no fibrosis), S1 (portal fibrosis without septa), S2 (portal fibrosis with rare septa), S3 (portal fibrosis with many septa), and S4 (cirrhosis). Two pathologists at the Diagnostic Center of the First Hospital of Jilin University, who were blinded to patients' clinical data, independently scored sections, and the resultant scores for each section were

Table 1 Demographic and clinical characteristics of the participants in the study groups

Characteristic	HC (<i>n</i> = 78)	CHB (<i>n</i> = 78)	LC (<i>n</i> = 85)	HCC (<i>n</i> = 77)	<i>P</i> -value
Gender (M/F)	48/30	53/25	54/31	62/15	0.05
Age (yr)	44 (25-67)	46 (30-61)	49 (27-65)	49 (22-57)	0.137
ALT (U/L)	14.95 (4.4-564)	122.9 (7-1906)	64 (8-855)	41 (15-450.2)	< 0.01
AST (U/L)	21.55 (11.0-38.5)	75.5 (14-1499)	56 (16-845)	56 (19-634)	< 0.01
ALB (g/L)	48.22 ± 2.51	38.01 ± 5.82	32.10 ± 6.87	34.56 ± 6.56	< 0.01
HGB (g/L)	147.79 ± 13.80	145.41 ± 17.74	126.33 ± 22.70	133.58 ± 22.5	< 0.01
PLT (× 10 ⁹ /L)	217 (117-328)	152.5 (55-298)	79 (28-303)	124 (17-515)	< 0.01
APRI	0.24 (0.1-0.5)	1.25 (0.16-68.14)	2.05 (0.24-42.25)	1.49 (0.24-9.43)	< 0.01
HBV-DNA (Log ₁₀)	0 (0-0)	6.64 (1.3-8.91)	4.96 (1.7-8.35)	4.61 (1.7-7.88)	< 0.01

M: Male; F: Female; HC: Healthy controls; CHB: Chronic hepatitis B; LC: Liver cirrhosis; HCC: Hepatocellular carcinoma; ALT: Alanine aminotransferase; AST: Aspartate transaminase; ALB: Albumin; HGB: Hemoglobin; PLT: Platelets; APRI: AST to platelet ratio index; HBV: Hepatitis B virus.

averaged in the analysis.

Statistical analysis

Mean ± SEM or median (range) was calculated for all numerical variables. Percentages are used to express categorical data. Differences between groups were determined by analysis of variance and Student's *t*-tests (normal distribution), the Kruskal-Wallis and Mann-Whitney *U*-tests (non-normal distribution), or the χ^2 -test or Fisher's exact test for categorical data. Correlations between variables were computed based on Spearman's correlation coefficients. The identified factors were further subjected to multiple linear regression analysis. The SPSS program (version 18.0, SPSS Inc., Chicago, IL, United States) was used for all statistical analyses, and statistical significance was considered if a *P*-value < 0.05.

RESULTS

Baseline characteristics

The baseline demographic, clinical and laboratory features of the 318 participants are summarized in Table 1. Among the participants, there were 78 cases of CHB (53 males), 85 cases of LC (54 males), 77 cases of HCC (62 males), and 78 healthy controls (48 males). The median (P25-P75) ages of the CHB, LC, and HCC patients as well as the control participants were 46 years (30-61 years), 49 years (27-65 years), 49 years (22-57 years), and 44 years (22-57 years), respectively, and there was no significant difference in age among the groups (*P* = 0.137). The proportions of male participants were significantly higher than those of female counterparts among all four groups (*P* = 0.049). Multiple comparisons showed that the male percentage in the HCC group was significantly higher than those in the CHB and LC groups (both *P* < 0.05), while there was no significant difference among the CHB, LC, and control groups. Higher alanine transaminase (ALT), aspartate transaminase, and HBV-DNA levels and lower serum albumin level, hemoglobin, and platelets (PLT) were detected in the CHB, LC, and HCC patients compared with controls.

Comparison of iron markers among different groups

As shown in Figure 1A-F, the serum ferritin levels were significantly higher in the CHB, LC, and HCC groups compared with the control group. The median TIBC level was significantly lower in the CHB group (59.63 ± 15.07), LC (49.29 ± 15.37), and HBV-HCC patients (54.16 ± 12.86) than in healthy controls (65.81 ± 9.18), as was the mean transferrin level in CHB (2.30 ± 0.59), LC (2.37 ± 3.42), and HCC (2.11 ± 0.52) patients relative to the values in controls (2.72 ± 0.47). Among the three liver disease groups, serum iron, total iron binding capacity, and serum transferrin levels were significantly lower, and the hepcidin level was significantly higher in the patients with LC or HCC (*P* < 0.05) than in the patients with CHB. However, the mean serum iron level in LC patients (20.99 ± 9.51 μmol/L) was comparable to that in the controls (20.05 ± 6.12 μmol/L). Increased transferrin saturation was observed in the CHB and LC groups, but not in HCC patients compared with the control group (*P* < 0.05 and *P* = 0.731).

Serum iron metabolite levels at different stages of LC and HCC

The increases in serum ferritin and transferrin saturation were proportional to the extent of LC and differed significantly among patients with different Child-Pugh scores (*P* < 0.05; Figure 2C and E). The TIBC and TF levels significantly decreased with increasing Child-Pugh scores (Figure 2F and D). There was no significant difference in serum iron or hepcidin among patients with different Child-Pugh scores (*P* > 0.05; Figure 2A and B).

The decreased serum iron and transferrin saturation levels significantly correlated with the smaller tumor burden in the first three stages of HCC by BCLC staging. Unexpectedly, they were significantly higher at stage D compared with stage B or C (Figure 3B and E). However, the mean transferrin level in patients with stage D was significantly lower than that in patients with stage A or B disease (Figure 3D). The serum ferritin level was significantly higher at stage D than




Figure 1 Mean levels of serum iron markers and their standard errors among the four groups. A: Hepcidin; B: Serum iron; C: Serum ferritin; D: Transferrin; E: Transferrin saturation; F: Total iron binding capacity. ^a $P < 0.05$ vs HC; ^b $P < 0.05$ vs CHB; ^c $P < 0.05$ vs LC; ^d $P < 0.05$ vs HCC. HC: Healthy controls; CHB: Chronic hepatitis B; LC: Liver cirrhosis; HCC: Hepatocellular carcinoma.

during the other three early stages (Figure 3C). There was no significant difference in hepcidin or total iron binding capacity among HCC patients at different BCLC stages ($P > 0.05$; Figure 3A and F).

Iron deposition in liver tissues with LC of varying severity

The extent of iron deposition in liver cells was investigated in patients with fibrosis at different stages as indicated by histology. A total of 29 patients were selected for iron staining, including 5 CHB patients (4 males) at S0, 8 (5 males) at S1, 4 (3 males) at S2, 7 (6 males) at S3, and 5 (2 males) at S4. There was no

significant difference in the median (P25-P75) ages among the different stage groups ($P = 0.122$). There was a clear trend of an increase in iron deposition in the liver tissues with increased fibrosis, which became significant at stages 3 and 4 (Figure 4A-F; $P < 0.05$). The iron level was barely detectable at stage 0 (Figure 4A), while prominent iron retention in hepatocytes was detected at S3 and S4 (Figure 4D and E). Iron deposition in liver cells was observed in 10%–50% of CHB patients. The mean iron level was significantly higher in severe fibrosis (S3: 23.7%) and cirrhosis (S4: 43.6%) compared with that in tissues with no or mild fibrosis (S0: 5.2%; S1: 7.9%; S2: 8.5%).




Figure 2 Serum iron markers among liver cirrhosis patients with different Child-Pugh classes. A: Hepcidin; B: Serum iron; C: Serum ferritin; D: Transferrin; E: Transferrin saturation; F: Total iron binding capacity. ^a $P < 0.05$ vs Child A; ^b $P < 0.05$ vs Child B; ^c $P < 0.05$ vs Child C. Child A: Child-Pugh class A; Child B: Child-Pugh class B; Child C: Child-Pugh class C.

Correlations between hepcidin level and other variables

The relationships of serum hepcidin levels and demographic and clinical findings are summarized in Table 2. Univariate analysis showed that serum hepcidin was significantly correlated with age ($\rho = 0.3$, $P < 0.001$), total bile acid (TBA; $\rho = 0.196$, $P = 0.002$), and international normalized ratio (INR; $\rho = 0.188$, $P = 0.003$). The serum hepcidin level was inversely correlated with HBV-DNA ($\rho = -0.347$, $P < 0.001$), red blood cell count (RBC; $\rho = -0.153$, $P = 0.018$), platelet count (PLT; $\rho = -0.195$, $P = 0.002$), hemoglobin (HGB; $\rho = -0.158$, $P = 0.014$), and hematocrit (HCT,

$\rho = -0.162$, $P = 0.012$). Subsequently, these factors were subjected to multiple linear regression analysis, which identified age (adjusted effect size = 0.213, $P = 0.001$), INR (adjusted effect size = 0.198, $P \leq 0.001$), and HBV-DNA (adjusted effect size = -0.282, $P < 0.001$) as independent factors associated with higher serum hepcidin. Notably, HBV-DNA was ranked as the most significant among all independent factors.

DISCUSSION

Our results showed that serum iron level was elevated




Figure 3 Serum iron markers among hepatocellular carcinoma patients with different Barcelona Clinic Liver Cancer stages. A: Hepcidin; B: Serum iron; C: Serum ferritin; D: Transferrin; E: Transferrin saturation; F: Total iron binding capacity. ^a $P < 0.05$ vs Stage A; ^b $P < 0.05$ vs Stage B; ^c $P < 0.05$ vs Stage C; ^d $P < 0.05$ vs Stage D. Stage A: Stage A of BCLC; Stage B: Stage B of BCLC; Stage C: Stage C of BCLC; Stage D: Stage D of BCLC.

ed in CHB but not in LC and HCC. The elevated serum iron level has been described in other chronic liver diseases, including chronic hepatitis C and alcoholic liver disease^[17-19]. However, cancer cells consume a large amount of iron to sustain their rapid proliferation^[20], resulting in a reduced serum iron level as shown in this study (Figure 1B). In addition, the serum ferritin level was significantly higher in hepatitis B-related liver diseases including LC and HCC (Figure 1C), likely reflecting the increased release of ferritin from hepatocytes that had increased iron deposition and were destroyed as a result of HBV replication and/or iron

retention-related injury. This suggestion was supported by our finding that the serum ferritin level was higher in CHB patients than in LC and HCC patients, because a more prominent liver injury, as indicated by ALT level, occurred in CHB^[21]. Sebastiani *et al.*^[22] reported that serum ferritin is an independent protein associated with hepatic iron deposition, while serum iron and transferrin saturation were not^[23].

In the present study, decreased serum transferrin and total iron binding capacity accompanied by increased transferrin saturation was detected in patients with HBV-related liver diseases compared with controls (Figure




Figure 4 Iron deposition in liver tissues with fibrosis of different stages. Perl's staining of iron appears as red granular particles in liver cells ($\times 400$ magnification.) A-E: Different stages of liver fibrosis, respectively. Liver fibrosis was staged using the METAVIR scoring system, which consists of five stages: S0 (no fibrosis, $n = 5$), S1 (portal fibrosis without septa, $n = 8$), S2 (portal fibrosis with rare septa, $n = 4$), S3 (portal fibrosis with many septa, $n = 7$), and S4 (cirrhosis, $n = 5$). Markedly increased iron deposition was observed in the severe liver fibrosis (S3) and cirrhosis (S4) groups, but not in groups S0-S2. F: Significantly higher average iron retention (mean \pm SEM) with severe fibrosis (S3: 23.7%) and cirrhosis (S4: 43.6%) compared to that with no or mild fibrosis (S0: 5.2%, S1: 7.9%; S2: 8.5%). A statistically significant difference in iron staining was observed among patients with severe fibrosis and cirrhosis ($P < 0.05$). ^a $P < 0.05$ vs S0; ^b $P < 0.05$ vs S1; ^c $P < 0.05$ vs S2; ^d $P < 0.05$ vs S3.

1D-F). Similar findings were reported by two studies. One study found that both serum transferrin saturation and ferritin concentration were elevated in patients with HBV-related liver disease^[24]. The other showed that 27.1% of the study participants experienced elevated transferrin saturation and 48.7% experienced increased liver iron deposition^[25]. However, we found no significant increase in the serum iron level, and in fact, the serum iron was reduced in HCC patients. Similar findings to ours were also reported. For instance, previous studies

also found there was no significant change in the serum iron level^[26] and no reduction of the serum iron level in patients with LC and HCC^[27]. The diverse stages of liver diseases among different studies may have contributed to the discrepancies in the serum iron level.

Intracellular iron deposition may trigger the cellular oxidative stress that may lead to liver injury. Liver iron accumulation significantly correlated with the severity of fibrosis based on METAVIR classification and cirrhosis^[28]. Our results also suggested the following relationship:

Table 2 Factors associated with serum hepcidin concentration in multiple linear regression models

Characteristic	Correlation analysis		Linear regression analysis	
	rho	P-value	Effect size	P-value
Age (yr)	0.3	< 0.01	0.213	0.001
Gender	0.01	0.986	-	-
ALT (U/L)	-0.095	0.141	-	-
AST (U/L)	-0.013	0.842	-	-
GGT (U/L)	0.011	0.871	-	-
AKP (U/L)	0.123	0.058	-	-
ALB (g/L)	-0.055	0.394	-	-
TBIL ($\mu\text{mol/L}$)	0.108	0.096	-	-
TBA ($\mu\text{mol/L}$)	0.196	0.002	-	-
HBV-DNA (Log_{10})	-0.374	< 0.01	-0.282	< 0.001
APRI	0.114	0.077	-	-
RBC ($\times 10^{12}/\text{L}$)	-0.153	0.018	-	-
WBC ($\times 10^9/\text{L}$)	-0.119	0.065	-	-
PLT ($\times 10^9/\text{L}$)	-0.195	0.002	-	-
HGB (g/L)	-0.158	0.014	-	-
MCV (fL)	0.01	0.879	-	-
HCT (L/L)	-0.162	0.012	-	-
INR	0.188	0.003	0.198	< 0.001
Serum iron ($\mu\text{mol/L}$)	0.027	0.672	-	-
Total iron binding capacity ($\mu\text{mol/L}$)	-0.073	0.261	-	-
Serum ferritin ($\mu\text{g/L}$)	-0.13	0.836	-	-
Transferrin saturation (%)	0.056	0.387	-	-
transferrin (g/L)	0.002	0.98	-	-

ALT: Alanine aminotransferase; AST: Aspartate transaminase; AKP: Alkaline phosphatase; ALB: Albumin; APRI: AST to platelet ratio index; HBV: Hepatitis B virus; TBIL: Total bilirubin; RBC: Red blood cell count; WBC: White blood cell count; TBA: Total bile acid; HGB: Hemoglobin; HCT: Hematocrit; PLT: Platelets; MCV: Mean corpuscular volume; INR: International normalized ratio.

Higher levels of serum ferritin and transferrin saturation were proportional to poorer Child-Pugh scores (Figure 2C and E). The extent of the intracellular iron staining was associated with the degree of liver fibrosis (Figure 4). We found cellular iron accumulation in more than 20% of patients with severe fibrosis and cirrhosis ($P < 0.05$), which was in line with the report that intracellular iron deposition was detected in chronic hepatitis C patients. Hepatic iron density was an independent predictor of advanced fibrosis in CHC^[29]. We did not observe coincident changes among different BCLC stages in the HCC group. The relatively small population in the HCC group may hinder more definitive results. Still, whether subgroups of HCC patients have coincidentally changed iron remains to be determined by further studies.

In addition to ferritin and transferrin, hepatocytes also produce and secrete hepcidin, an acute phase reactant protein that may negatively regulate the endogenous iron level and reduce the release of iron from cells by interacting with the cellular iron exporter ferroportin that leads to subsequent internalization and degradation^[30,31]. Our study further indicates that the major factors independently associated with the altered hepcidin levels were age, HBV-DNA, and INR in patients with HBV-related diseases. The cross-sectional study design limited the ability of the study to clarify causality between HBV-DNA load and altered hepcidin levels. Wang *et al.*^[32] indicated that hepcidin expression was regulated by iron and inflammatory factors in HBV-

infected patients, and that the virus accumulation in infected hepatocytes can affect hepcidin production. However, the molecular basis by which hepcidin may alter HBV replication is unknown. Interestingly, iron-induced hepcidin expression altered HCV replication in cultured cells^[33]. Of note, other studies found no link between HBV infection and hepcidin production^[34].

This study does have some limitations. We were unable to determine the cause-effect relationship between the increased iron accumulation and liver injury in HBV-infected patients because of the nature of this cross-sectional study. The relatively small size of the study population may render insufficient statistical power in the analysis. Our findings should be verified in future studies.

In conclusion, iron metabolism disorders can occur in patients with HBV-related liver disease. The serum markers of iron metabolism disorders vary in different stages of HBV-related liver diseases.

ARTICLE HIGHLIGHTS

Research background

Chronic hepatitis B virus (HBV) infection affects 240 million people worldwide and can lead to chronic liver diseases including chronic hepatitis B (CHB), cirrhosis, and hepatocellular carcinoma (HCC). In managing chronic HBV infection, the objective is to detect liver injury early and then stop the progression of liver disease through effective treatment. To achieve this goal, a clear understanding of various factors involved in HBV pathogenesis is required. The relationship between HBV-related chronic liver diseases and levels of serum markers of iron metabolism and their impact on liver disease severity

have not been well investigated. The aim of this study was to characterize changes in serum iron markers in patients with HBV-related diseases and to assess correlations between changes in iron metabolism and HBV-related liver injury.

Research motivation

In managing chronic HBV infection, the objective is to detect liver injury early and then stop the progression of liver disease through effective treatment. To achieve this goal, a clear understanding of various factors involved in HBV pathogenesis is required.

Research objectives

The aim of this study was to characterize changes in serum iron markers in patients with HBV-related diseases and assess correlations between changes in iron metabolism and HBV-related liver injury.

Research methods

This was a case-control study, which included 78 healthy controls, 78 CHB patients including 25 diagnosed by biopsy, 85 patients with HBV-related liver cirrhosis including 5 diagnosed by biopsy, and 77 with HBV-related HCC. Blood samples were collected from all patients after overnight fasting. Serum iron metabolism markers, HBV serological markers and HBV-DNA, and biochemical markers were determined with autoanalyzers. Liver biopsy was performed with a 16 G Tru-Cut needle guided by color Doppler ultrasound. A biopsy that was 1.5 cm or longer or a liver section that contained at least five portal tracts was considered eligible for diagnosis. The liver sections from each biopsy were stained with hematoxylin and eosin and Perls, respectively. Liver fibrosis was staged using the METAVIR scoring system. The SPSS program was used for all statistical analyses, and statistical significance was considered if a *P*-value < 0.05.

Research results

The serum ferritin levels were significantly higher in CHB, liver cirrhosis, and HCC groups compared with the control group. The median TIBC level was significantly lower in the CHB group, liver cirrhosis, and HBV-HCC patients than in healthy controls, as was the mean transferrin level in CHB, liver cirrhosis, and HCC patients relative to the values in controls. Among the three liver disease groups, serum iron, total iron binding capacity, and serum transferrin levels were significantly lower, and the hepcidin level was significantly higher in the patients with cirrhosis or HCC (*P* < 0.05) than in the patients with CHB. Serum ferritin and transferrin saturation levels increased proportionally to the extent of liver cirrhosis and poorer Child-Pugh scores (*P* < 0.05). The decreased serum iron and transferrin saturation levels significantly correlated with a smaller hepatocellular carcinoma tumor burden according to Barcelona Clinic Liver Cancer staging. There was a clear trend of an increase in iron deposition in the liver tissues with increased fibrosis, which became prominent at stages 3 and 4 (Figure 4A-F; *P* < 0.05). The iron level was barely detectable at stage 0 (Figure 4A), while prominent iron retention in hepatocytes was detected at S3 and S4 (Figure 4D and E). Iron deposition in liver cells was observed in 10%–50% of CHB patients. The mean iron level was significantly higher in severe fibrosis and cirrhosis compared with that in tissues with no or mild fibrosis. We also analyzed the relationships between serum hepcidin levels and demographic factors. Age, international normalized ratio, and HBV-DNA were independent factors associated with higher serum hepcidin. The cross-sectional study design limited the ability of this study to clarify causality between HBV-DNA load and altered hepcidin levels.

Research conclusions

Iron metabolism disorders occur in patients with HBV-related liver disease. The iron metabolism disorders vary in different stages of HBV-related liver diseases.

Research perspectives

The mechanism of iron metabolism disorders present in patients of HBV-related liver disease should be further studied in the future.

ACKNOWLEDGMENTS

We appreciate Li-Shan Su for assistance in revising this

manuscript.

REFERENCES

- 1 Stanaway JD, Flaxman AD, Naghavi M, Fitzmaurice C, Vos T, Abubakar I, Abu-Raddad LJ, Assadi R, Bhala N, Cowie B, Forouzanfar MH, Groeger J, Hanafiah KM, Jacobsen KH, James SL, MacLachlan J, Malekzadeh R, Martin NK, Mokdad AA, Mokdad AH, Murray CJL, Plass D, Rana S, Rein DB, Richardus JH, Sanabria J, Saylan M, Shahraz S, So S, Vlassov VV, Weiderpass E, Wiersma ST, Younis M, Yu C, El Sayed Zaki M, Cooke GS. The global burden of viral hepatitis from 1990 to 2013: findings from the Global Burden of Disease Study 2013. *Lancet* 2016; **388**: 1081-1088 [PMID: 27394647 DOI: 10.1016/S0140-6736(16)30579-7]
- 2 Hino K, Nishina S, Hara Y. Iron metabolic disorder in chronic hepatitis C: mechanisms and relevance to hepatocarcinogenesis. *J Gastroenterol Hepatol* 2013; **28** Suppl 4: 93-98 [PMID: 24251712 DOI: 10.1111/jgh.12243]
- 3 Wei Y, Ye W, Zhao W. Serum Iron Levels Decreased in Patients with HBV-Related Hepatocellular Carcinoma, as a Risk Factor for the Prognosis of HBV-Related HCC. *Front Physiol* 2018; **9**: 66 [PMID: 29467672 DOI: 10.3389/fphys.2018.00066]
- 4 Zou DM, Sun WL. Relationship between Hepatitis C Virus Infection and Iron Overload. *Chin Med J (Engl)* 2017; **130**: 866-871 [PMID: 28345552 DOI: 10.4103/0366-6999.202737]
- 5 Jaeschke H, Gores GJ, Cederbaum AI, Hinson JA, Pessayre D, Lemasters JJ. Mechanisms of hepatotoxicity. *Toxicol Sci* 2002; **65**: 166-176 [PMID: 11812920 DOI: 10.1093/toxsci/65.2.166]
- 6 Powell LW, Seckington RC, Deugnier Y. Haemochromatosis. *Lancet* 2016; **388**: 706-716 [PMID: 26975792 DOI: 10.1016/S0140-6736(15)01315-X]
- 7 Mao W, Hu Y, Lou Y, Chen Y, Zhang J. Abnormal serum iron markers in chronic hepatitis B virus infection may be because of liver injury. *Eur J Gastroenterol Hepatol* 2015; **27**: 130-136 [PMID: 25419642 DOI: 10.1097/MEG.0000000000000247]
- 8 Blumberg BS, Lustbader ED, Whitford PL. Changes in serum iron levels due to infection with hepatitis B virus. *Proc Natl Acad Sci USA* 1981; **78**: 3222-3224 [PMID: 6942427 DOI: 10.1073/pnas.78.5.3222]
- 9 Lecube A, Hernández C, Simó R. Glucose abnormalities in non-alcoholic fatty liver disease and chronic hepatitis C virus infection: the role of iron overload. *Diabetes Metab Res Rev* 2009; **25**: 403-410 [PMID: 19444865 DOI: 10.1002/dmrr.972]
- 10 Metwally MA, Zein CO, Zein NN. Clinical significance of hepatic iron deposition and serum iron values in patients with chronic hepatitis C infection. *Am J Gastroenterol* 2004; **99**: 286-291 [PMID: 15046219 DOI: 10.1111/j.1572-0241.2004.04049.x]
- 11 Nishina S, Hino K, Korenaga M, Vecchi C, Pietrangelo A, Mizukami Y, Furutani T, Sakai A, Okuda M, Hidaka I, Okita K, Sakaida I. Hepatitis C virus-induced reactive oxygen species raise hepatic iron level in mice by reducing hepcidin transcription. *Gastroenterology* 2008; **134**: 226-238 [PMID: 18166355 DOI: 10.1053/j.gastro.2007.10.011]
- 12 Varghese J, James JV, Sagi S, Chakraborty S, Sukumaran A, Ramakrishna B, Jacob M. Decreased hepatic iron in response to alcohol may contribute to alcohol-induced suppression of hepcidin. *Br J Nutr* 2016; **115**: 1978-1986 [PMID: 27080262 DOI: 10.1017/S0007114516001197]
- 13 Wang J, Dong A, Liu G, Anderson GJ, Hu TY, Shi J, Hu Y, Nie G. Correlation of serum hepcidin levels with disease progression in hepatitis B virus-related disease assessed by nanopore film based assay. *Sci Rep* 2016; **6**: 34252 [PMID: 27694815 DOI: 10.1038/srep34252]
- 14 Ohkoshi S, Yoshimura A, Yamamoto S, Yano M, Kurita S, Yamazaki K, Aoki YH, Yamagiwa S, Wakabayashi H, Sugiyama M, Takahashi T, Ishikawa T, Matsuda Y, Ichida T, Kamimura T, Aoyagi Y. Successful treatment with lamivudine may correlate with reduction of serum ferritin levels in the patients with chronic hepatitis and liver cirrhosis type B. *Hepatol Int* 2008; **2**: 382-387

- [PMID: 19669269 DOI: 10.1007/s12072-008-9084-z]
- 15 **Suk KT**, Baik SK, Yoon JH, Cheong JY, Paik YH, Lee CH, Kim YS, Lee JW, Kim DJ, Cho SW, Hwang SG, Sohn JH, Kim MY, Kim YB, Kim JG, Cho YK, Choi MS, Kim HJ, Lee HW, Kim SU, Kim JK, Choi JY, Jun DW, Tak WY, Lee BS, Jang BK, Chung WJ, Kim HS, Jang JY, Jeong SW, Kim SG, Kwon OS, Jung YK, Choe WH, Lee JS, Kim IH, Shim JJ, Cheon GJ, Bae SH, Seo YS, Choi DH, Jang SJ; Korean Association for the Study of the Liver. Revision and update on clinical practice guideline for liver cirrhosis. *Korean J Hepatol* 2012; **18**: 1-21 [PMID: 22511898 DOI: 10.3350/kjhep.2012.18.1.1]
 - 16 **Rozario R**, Ramakrishna B. Histopathological study of chronic hepatitis B and C: a comparison of two scoring systems. *J Hepatol* 2003; **38**: 223-229 [PMID: 12547412 DOI: 10.1016/S0168-8278(02)00357-4]
 - 17 **Di Bisceglie AM**, Axiotis CA, Hoofnagle JH, Bacon BR. Measurements of iron status in patients with chronic hepatitis. *Gastroenterology* 1992; **102**: 2108-2113 [PMID: 1587431 DOI: 10.1016/0016-5085(92)90339-Z]
 - 18 **Datz C**, Müller E, Aigner E. Iron overload and non-alcoholic fatty liver disease. *Minerva Endocrinol* 2017; **42**: 173-183 [PMID: 27834478 DOI: 10.23736/S0391-1977.16.02565-7]
 - 19 **Britton LJ**, Subramaniam VN, Crawford DH. Iron and non-alcoholic fatty liver disease. *World J Gastroenterol* 2016; **22**: 8112-8122 [PMID: 27688653 DOI: 10.3748/wjg.v22.i36.8112]
 - 20 **Ba Q**, Hao M, Huang H, Hou J, Ge S, Zhang Z, Yin J, Chu R, Jiang H, Wang F, Chen K, Liu H, Wang H. Iron deprivation suppresses hepatocellular carcinoma growth in experimental studies. *Clin Cancer Res* 2011; **17**: 7625-7633 [PMID: 22052937 DOI: 10.1158/1078-0432.CCR-10-3099]
 - 21 **Lustbader ED**, Hann HW, Blumberg BS. Serum ferritin as a predictor of host response to hepatitis B virus infection. *Science* 1983; **220**: 423-425 [PMID: 6301008 DOI: 10.1126/science.6301008]
 - 22 **Sebastiani G**, Tempesta D, Alberti A. Hepatic iron overload is common in chronic hepatitis B and is more severe in patients coinfecting with hepatitis D virus. *J Viral Hepat* 2012; **19**: e170-e176 [PMID: 22239515 DOI: 10.1111/j.1365-2893.2011.01508.x]
 - 23 **Torti FM**, Torti SV. Regulation of ferritin genes and protein. *Blood* 2002; **99**: 3505-3516 [PMID: 11986201 DOI: 10.1182/blood.V99.10.3505]
 - 24 **Yonal O**, Akyuz F, Demir K, Ciftci S, Keskin F, Pinarbasi B, Uyanikoglu A, Issever H, Ozdil S, Boztas G, Besisik F, Kaymakoglu S, Cakaloglu Y, Mungan Z, Okten A. Decreased prohepcidin levels in patients with HBV-related liver disease: relation with ferritin levels. *Dig Dis Sci* 2010; **55**: 3548-3551 [PMID: 20238243 DOI: 10.1007/s10620-010-1183-8]
 - 25 **Martinelli AL**, Filho AB, Franco RF, Tavella MH, Ramalho LN, Zucoloto S, Rodrigues SS, Zago MA. Liver iron deposits in hepatitis B patients: association with severity of liver disease but not with hemochromatosis gene mutations. *J Gastroenterol Hepatol* 2004; **19**: 1036-1041 [PMID: 15304122 DOI: 10.1111/j.1440-1746.2004.03410.x]
 - 26 **Boige V**, Castéra L, de Roux N, Ganne-Carrié N, Ducot B, Pelletier G, Beaugrand M, Buffet C. Lack of association between HFE gene mutations and hepatocellular carcinoma in patients with cirrhosis. *Gut* 2003; **52**: 1178-1181 [PMID: 12865278 DOI: 10.1136/gut.52.8.1178]
 - 27 **Büyükaşık NS**, Nadir I, Akin FE, Cakal B, Kav T, Ersoy O, Büyükaşık Y. Serum iron parameters in cirrhosis and chronic hepatitis: detailed description. *Turk J Gastroenterol* 2011; **22**: 606-611 [PMID: 22287406 DOI: 10.4318/tjg.2011.0323]
 - 28 **Hézode C**, Cazeneuve C, Coué O, Roudot-Thoraval F, Lonjon I, Bastie A, Duvoux C, Pawlotsky JM, Zafrani ES, Amselem S, Dhumeaux D. Liver iron accumulation in patients with chronic active hepatitis C: prevalence and role of hemochromatosis gene mutations and relationship with hepatic histological lesions. *J Hepatol* 1999; **31**: 979-984 [PMID: 10604569 DOI: 10.1016/S0168-8278(99)80308-0]
 - 29 **Missiha SB**, Ostrowski M, Heathcote EJ. Disease progression in chronic hepatitis C: modifiable and nonmodifiable factors. *Gastroenterology* 2008; **134**: 1699-1714 [PMID: 18471548 DOI: 10.1053/j.gastro.2008.02.069]
 - 30 **Nemeth E**, Tuttle MS, Powelson J, Vaughn MB, Donovan A, Ward DM, Ganz T, Kaplan J. Heparin regulates cellular iron efflux by binding to ferroportin and inducing its internalization. *Science* 2004; **306**: 2090-2093 [PMID: 15514116 DOI: 10.1126/science.1104742]
 - 31 **Singh B**, Arora S, Agrawal P, Gupta SK. Heparin: a novel peptide hormone regulating iron metabolism. *Clin Chim Acta* 2011; **412**: 823-830 [PMID: 21333642 DOI: 10.1016/j.cca.2011.02.014]
 - 32 **Wang XH**, Cheng PP, Jiang F, Jiao XY. The effect of hepatitis B virus infection on hepcidin expression in hepatitis B patients. *Ann Clin Lab Sci* 2013; **43**: 126-134 [PMID: 23694786]
 - 33 **Bartolomei G**, Cevik RE, Marcello A. Modulation of hepatitis C virus replication by iron and hepcidin in Huh7 hepatocytes. *J Gen Virol* 2011; **92**: 2072-2081 [PMID: 21593278 DOI: 10.1099/vir.0.032706-0]
 - 34 **Olmez OF**, Gurel S, Yilmaz Y. Plasma prohepcidin levels in patients with chronic viral hepatitis: relationship with liver fibrosis. *Eur J Gastroenterol Hepatol* 2010; **22**: 461-465 [PMID: 19940783 DOI: 10.1097/MEG.0b013e3283344708]

P-Reviewer: Shimizu Y, Trinder D **S-Editor:** Ma RY
L-Editor: Wang TQ **E-Editor:** Tan WW





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

