

World Journal of *Clinical Cases*

World J Clin Cases 2019 January 6; 7(1): 1-121



MINIREVIEWS

- 1 Role of endoscopy in the surveillance and management of colorectal neoplasia in inflammatory bowel disease
Manchanda S, Rizvi QUA, Singh R

ORIGINAL ARTICLE**Retrospective Cohort Study**

- 10 Risk factors for perforation during endoscopic retrograde cholangiopancreatography in post-reconstruction intestinal tract
Takano S, Fukasawa M, Shindo H, Takahashi E, Hirose S, Fukasawa Y, Kawakami S, Hayakawa H, Yokomichi H, Kadokura M, Sato T, Enomoto N

Retrospective Study

- 19 Instant evaluation of contrast enhanced endoscopic ultrasound helps to differentiate various solid pancreatic lesions in daily routine
Kannengiesser K, Mahlke R, Petersen F, Peters A, Kucharzik T, Maaser C
- 28 Correlation of serum albumin and prognostic nutritional index with outcomes following pancreaticoduodenectomy
Rungsakulkij N, Tangtawee P, Suragul W, Muangkaew P, Mingphruedhi S, Aeesoa S

Clinical Trials Study

- 39 Efficacy of 0.5-L vs 1-L polyethylene glycol containing ascorbic acid as additional colon cleansing methods for inadequate bowel preparation as expected by last stool examination before colonoscopy
Cho JH, Goo EJ, Kim KO, Lee SH, Jang BI, Kim TN

Observational Study

- 49 Value of contrast-enhanced ultrasound combined with elastography in evaluating cervical lymph node metastasis in papillary thyroid carcinoma
Jiang W, Wei HY, Zhang HY, Zhuo QL

CASE REPORT

- 58 Full-term pregnancy in breast cancer survivor with fertility preservation: A case report and review of literature
Garrido-Marín M, Argacha PM, Fernández L, Molfino F, Martínez-Soler F, Tortosa A, Gimenez-Bonafé P
- 69 Psoriatic fasciitis in a pediatric patient: A case report
Otar Yener G, Ekici Tekin Z, Yuksel S

- 73 Vertebrobasilar artery dissection manifesting as Millard-Gubler syndrome in a young ischemic stroke patient: A case report
Li XT, Yuan JL, Hu WL
- 79 Endodontic management of the maxillary first molars with two root canals: A case report and review of the literature
Liu J, Que KH, Xiao ZH, Wen W
- 89 Tegafur deteriorates established cardiovascular atherosclerosis in colon cancer: A case report and review of the literature
Zhang SC, Yu MY, Xi L, Zhang JX
- 95 Authenticity of pulmonary *Lophomonas blattarum* infection: A case report
Meng SS, Dai ZF, Wang HC, Li YX, Wei DD, Yang RL, Lin XH
- 102 Co-occurrence of IPMN and malignant IPNB complicated by a pancreatobiliary fistula: A case report and review of the literature
Ren X, Zhu CL, Qin XF, Jiang H, Xia T, Qu YP
- 109 Bilateral and symmetric C1-C2 dumbbell ganglioneuromas associated with neurofibromatosis type 1: A case report
Tan CY, Liu JW, Lin Y, Tie XX, Cheng P, Qi X, Gao Y, Guo ZZ
- 116 Follicular dendritic cell sarcoma detected in hepatogastric ligament: A case report and review of the literature
Yan WX, Yu YX, Zhang P, Liu XK, Li Y

ABOUT COVER

Editor-in-Chief of *World Journal of Clinical Cases*, Sandro Vento, MD, Full Professor, (E-mail: ventosandro@yahoo.it) Department of Medicine, Nazarbayev University School of Medicine and University Medical Center, Astana 010000, Kazakhstan

AIMS AND SCOPE

World Journal of Clinical Cases (World J Clin Cases, WJCC, online ISSN 2307-8960, DOI: 10.12998) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

The primary task of *WJCC* is to rapidly publish high-quality Autobiography, Case Report, Clinical Case Conference (Clinicopathological Conference), Clinical Management, Diagnostic Advances, Editorial, Field of Vision, Frontier, Medical Ethics, Original Articles, Clinical Practice, Meta-Analysis, Minireviews, Review, Therapeutics Advances, and Topic Highlight, in the fields of allergy, anesthesiology, cardiac medicine, clinical genetics, clinical neurology, critical care, dentistry, dermatology, emergency medicine, endocrinology, family medicine, gastroenterology and hepatology, geriatrics and gerontology, hematology, immunology, etc.

INDEXING/ABSTRACTING

World Journal of Clinical Cases (WJCC) is now indexed in PubMed, PubMed Central, Science Citation Index Expanded (also known as SciSearch®), and Journal Citation Reports/Science Edition. The 2018 Edition of Journal Citation Reports cites the 2017 impact factor for *WJCC* as 1.931 (5-year impact factor: N/A), ranking *WJCC* as 60 among 154 journals in Medicine, General and Internal (quartile in category Q2).

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: *Ying-Na Bian* Proofing Editorial Office Director: *Jin-Lei Wang*

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Semimonthly

EDITORS-IN-CHIEF

Dennis A Bloomfield, Sandro Vento

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/2307-8960/editorialboard.htm>

EDITORIAL OFFICE

Jin-Lei Wang, Director

PUBLICATION DATE

January 6, 2019

COPYRIGHT

© 2019 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Role of endoscopy in the surveillance and management of colorectal neoplasia in inflammatory bowel disease

Shaad Manchanda, Qurat-ul-ain Rizvi, Rajvinder Singh

ORCID number: Shaad Manchanda (0000-0002-3787-0558); Qurat-ul-ain Rizvi (0000-0002-5083-0227); Rajvinder Singh (0000-0001-9116-6054).

Author contributions: Manchanda S, Rizvi Q and Singh R contributed equally to this work.

Conflict-of-interest statement: The authors have no conflicts of interest.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Received: October 27, 2018

Peer-review started: October 29, 2018

First decision: November 27, 2018

Revised: December 19, 2018

Accepted: December 21, 2018

Article in press: December 21, 2018

Published online: January 6, 2019

Shaad Manchanda, Rajvinder Singh, Department of Gastroenterology, Lyell McEwin Hospital, South Australia 5112, Australia

Shaad Manchanda, Rajvinder Singh, School of Medicine, University of Adelaide, South Australia 5000, Australia

Qurat-ul-ain Rizvi, Eastern Health, Department of Gastroenterology, Box Hill Hospital, Victoria 3128, Australia

Corresponding author: Rajvinder Singh, FRACP, FRCP(C), MBBS, MPhil, MRCP, Professor, Department of Gastroenterology, Lyell McEwin Hospital, Haydown Road, Elizabeth Vale, Adelaide, South Australia 5112, Australia. rajvinder.singh@sa.gov.au

Telephone: +61-8-81829000

Fax: +61-8-82821764

Abstract

Endoscopy has become increasingly fundamental in the management of patients with inflammatory bowel disease (IBD). It is required for diagnosis, assessment of therapeutic response, postoperative follow up and in the surveillance of dysplasia. With rapid advances in technology, including high definition colonoscopy and chromoendoscopy, questions have arisen regarding the most appropriate surveillance and management strategies of colorectal neoplasia in IBD. We aim to review current surveillance strategies, explore the utility of new technologies, and examine the role of endoscopic resection, with the aim of clarifying these questions.

Key words: Inflammatory bowel disease; Colorectal cancer; High definition endoscopy; Chromoendoscopy

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: With the use of new generation, high definition endoscopy, most dysplasia is visually identifiable and hence targeted biopsies are advised. Random biopsies may be utilised in patients with a personal history of neoplasia, primary sclerosis cholangitis, and a tubular colon. Any lesion deemed to be endoscopically resectable should be referred to centres with expertise to do so whilst invisible dysplasia should prompt consideration towards a colectomy.

Citation: Manchanda S, Rizvi QUA, Singh R. Role of endoscopy in the surveillance

and management of colorectal neoplasia in inflammatory bowel disease. *World J Clin Cases* 2019; 7(1): 1-9

URL: <https://www.wjgnet.com/2307-8960/full/v7/i1/1.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v7.i1.1>

INTRODUCTION

Endoscopy has increasingly become a fundamental investigation in the management of patients with inflammatory bowel disease (IBD). It is required for initial diagnosis, assessment of therapeutic response, postoperative follow up and in the surveillance of dysplasia. It is estimated that up to 10% of deaths amongst those with IBD may be attributed to the development of colorectal cancer (CRC). The cumulative risk of CRC in IBD is approximately 2% at 10 years of disease duration, 5% to 8% after 20 years, and as high as 20% to 30% by the 30-year mark^[1-3]. The risk of developing CRC in ulcerative colitis (UC) is similar to that in Crohn's disease (CD) if adjusted for the area of colonic involvement^[4]. Improvements in therapy and refinements in surveillance methodology has seen an overall reduction in CRC in patients with IBD^[5]. With rapid advances in technology, including high definition colonoscopy and chromoendoscopy, questions have arisen regarding the most appropriate surveillance and management strategies of colorectal neoplasia in IBD. This article aims to review and clarify these questions.

SURVEILLANCE

According to a 2006 Cochrane review, there was no evidence to support surveillance programs conferring survival benefit^[6]. The authors attributed this to a lead-time bias, noting earlier malignancy detection and improved prognosis. Nevertheless, due to the inherent risk of developing CRC, many gastroenterological societies still endorse surveillance programs. The recognized risk factors for the development of malignancy include disease duration, extent of bowel involvement, early age of onset, severity of inflammation, family history of CRC in a first degree relative and presence of primary sclerosing cholangitis (PSC)^[1-3] (Table 1). Of these, the strongest predictor is duration of disease, although cancer rarely occurs within the first 7 years. Additional potential risk factors are post inflammatory polyps, backwash ileitis and the presence of strictures. Proctitis, where inflammation is limited to the rectum and anus, is thought to not confer increased risk of cancer, hence such patients should undergo normal CRC screening^[6]. All patients diagnosed with either UC or CD should be enrolled into a screening program. Neoplasia surveillance has three goals; first is to define the extent of mucosal healing and response to treatment; second is to identify premalignant lesions and facilitate endoscopic resection; and third is to allow for prompt diagnosis of colitis associated malignancy^[4].

There are several recognized guidelines regarding surveillance including the European Crohn's and Colitis Organization (2017) guideline, the American Gastroenterological Association (2010) guideline and the British Society of Gastroenterology (2010) guideline (Table 2). Timing and surveillance intervals vary between these. Typically, surveillance is recommended to begin approximately 8 to 10 years after either the beginning of symptoms or diagnosis. Surveillance intervals range from 1 to 5 years, and is often determined by an individual patient's risk profile for the development of colorectal neoplasia. High risk features, warranting annual surveillance, include active extensive disease, a history of dysplasia or stricture, a personal history of PSC and a strong family history of CRC. Intermediate risk patients may be screened every 3 years. Low risk features, allowing for up to 5 years of surveillance intervals, include endoscopic and histological remission, no history of neoplasia and no family history of CRC^[5,7]. Table 3 summarizes low, intermediate and high risk features. Pseudopolyps are not considered a risk factor for the development of neoplasia, however do reflect chronic inflammation which may impact upon the quality of mucosal visualization and examination during endoscopic surveillance^[4].

Ideal conditions for surveillance include thorough bowel preparation and a non-inflamed mucosa during a quiescent disease period given the difficulty in distinguishing inflammatory from early dysplastic changes both visually endoscopically and on histology^[4]. Inadequate bowel preparation may obscure non polypoid lesions and is associated with reduced dysplasia detection and increased

Table 1 Risk factors for the development of dysplasia in inflammatory bowel disease

Risk factors	Endoscopic factors
Disease duration	Active disease
Disease extent	Presence of strictures in ulcerative colitis
Disease severity	Post inflammatory polyps
Past dysplasia	Tubular appearance of colon with loss of colonic haustration
Primary sclerosing cholangitis	
Family history of colorectal cancer	

need for repeat procedures^[5]. It is important to note that surveillance should not be delayed indeterminately in waiting for disease quiescence. Some guidelines may recommend the use of random biopsies; however, the current gold standard is the use of targeted biopsies as discussed below.

IBD associated dysplasia is often visually flat and may be quite subtle^[3]. A fundamental new concept which has recently developed is that of visible and invisible dysplasia. Visible dysplasia, by definition, is histopathological proven dysplasia on a targeted biopsy of a specific, concerning area recognized on colonoscopic visualization. Invisible dysplasia is histopathological proven dysplasia on a random biopsy, from a visually unremarkable colonic region^[8].

With traditional standard definition white light colonoscopy, conventional screening recommendations incorporated random biopsies to increase the pick-up rate of endoscopically invisible dysplasia^[4,7]. Random 4 quadrant biopsies were performed every 10 cm from rectum to cecum, with a total of at least 33 samples. This benchmark was estimated to have approximately 90% confidence of detecting dysplasia in the presence of pancolitis. Two major drawbacks of random biopsies are the significant prolongation of procedure time as well as the relatively poor diagnostic yield. One trial comparing targeted and random biopsies in the surveillance of CRC in patients with UC noted the random group had a longer examination time of 41.7 min compared with 26.6 min in the targeted biopsy group^[9]. They also found the mean number of biopsies in the random group to be 34.8 compared with 3.1 amongst the targeted biopsy study arm. Targeted biopsies were also reported to have improved neoplastic detection, with improvements in neoplasia detection per colonoscopy (0.211 vs 0.168) as well as a 2.1% improvement in identification of patients with neoplasia. Van den Broek *et al.* demonstrated neoplasia detection in up to 85% of targeted biopsies, with neoplasia identified in 5.7% of random biopsies in only 7.5% of cases^[10]. They also demonstrated neoplasia to be macroscopically visible in 94% of cases. A review by Moussata and colleagues noted random biopsies added an additional 15% detection rate of neoplasia compared to chromoendoscopy with targeted biopsy, and that this was exclusively in patients with either a personal history of neoplasia (OR = 5.3; CI: 3.3-8.8, $P < 0.001$), concomitant PSC (OR = 2.3; CI: 1.2-4.2, $P = 0.006$) or had a tubular appearing colon (OR = 1.5, CI: 0.7-3.5, $P = 0.303$)^[11]. The authors also noted that when these 3 risk factors were absent, neoplasia was not detected on random biopsies. Taken together, this study suggests that judicious use of random biopsies has a role in detecting invisible dysplasia, and can be considered in patients with either a personal history of neoplasia, concomitant PSC, or noted to have a tubular appearing colon.

With the new era of high definition and image enhanced colonoscopy, the utility of random biopsies has come into question. It is now generally regarded that most colonic neoplasia may be identified through high definition or image enhanced endoscopy^[7]. In a retrospective analysis, comparing standard definition, the new generation endoscopes have been estimated to offer a higher ratio both in terms of prevalence (2.21, CI: 1.09-4.45) and in detecting dysplasia on targeted biopsy (2.99, CI: 1.16-7.79)^[12]. Large, prospective trials would be required to better appreciate the difference between standard definition and the new generation endoscopes, however such trials are unlikely given many centers have already shifted to high definition imaging modalities.

Chromoendoscopy is an endoscopic technique utilizing dyes to emphasize mucosal characteristics and highlight pathologically abnormal areas. The utility of dyes to enhance mucosal abnormalities was first developed in the 1990s^[3]. Initially contrast dyes were applied via a spraying catheter and foot pump^[4]. Chromoendoscopy with targeted biopsies is now regarded as the preferred surveillance technique^[7,13]. It has been reported to have up to a 3 fold increase in per patient dysplasia detection, and up to 5 fold increase in per lesion dysplasia detection. Dilution of contrast agents must

Table 2 Commonly used guidelines for the screening of neoplasia in inflammatory bowel disease

Society	Commencement	Risk stratification	Interval
ECCO, 2017	8 yr post symptom onset	Stricture or dysplasia, PSC, extensive colitis, severe active inflammation	Annual
		Mild to moderate active inflammation, post inflammatory polyps, or first degree relative with CRC	2-3 yr
		None of the above features	5 yr
AGA, 2010	8 yr post diagnosis	Active inflammation, stricture, post inflammatory polyps, history of dysplasia, first degree relative with CRC, PSC	Annual
ACG, 2010	8-10 yr post diagnosis	After 2 negative colonoscopies No risk stratification	1-3 yr 1-2 yr
BSG, 2010	10 yr post symptom onset	Moderate/severe active inflammation on the prior colonoscopy, stricture, dysplasia, PSC, first degree relative with CRC aged < 50 yr	Annual
		Mild active inflammation on prior colonoscopy, post inflammatory polyps, first degree relative with CRC aged > 50 yr	3 yr
		Nil prior inflammation, left sided colitis or CD colitis affecting > 50% surface area of the colon	5 yr

ECCO: European Crohn's and Colitis Organisation; AGA: American Gastroenterological Association; ACG: American College of Gastroenterology; BSG: British Society of Gastroenterology.

be appropriate to allow for adequate mucosal staining without being too intense and visually obscuring subtle mucosal abnormalities. Topical contrast agents commonly used include 0.1% methylene blue or 0.02% to 0.5% indigo carmine^[4,7]. The Surveillance for Colorectal Endoscopic Neoplasia Detection and Management in Inflammatory Bowel Disease Patients: International Consensus Recommendations (SCENIC) guidelines recommend the use of either 0.03% Indigocarmine or 0.04% Methylene blue applied through biopsy or water jet channels. Identified lesions should be first evaluated as resectable or non resectable. Resectable lesions should be referred for Endoscopic Mucosal Resection and/or Endoscopic Submucosal Dissection. Targeted biopsies should be performed on non resectable lesions and those of uncertain significance. The major limitations of chromoendoscopy are the prolonged procedure times, the additional training required on the part of the proceduralist, and the additional costs, all of which act as barriers to the widespread uptake of this technique^[3,5,8].

Random biopsies are not recommended from areas which appear macroscopically normal on chromoendoscopy^[4]. In addition to targeted biopsies, some guidelines recommend two biopsies from each colon segment to further define disease extent and the severity of inflammation^[7]. Chromoendoscopy has an 83.3% (CI: 35.9-99.6) sensitivity and a 91.3% (CI: 43.8-100) specificity for the detection of dysplasia^[14]. Its yield may be reduced by severe inflammation, significant pseudopolyposis or poor bowel preparation. In these conditions, or if chromoendoscopy is not available, previous recommendations of standard definition colonoscopy with combined targeted and random biopsies is regarded as an appropriate approach (Figure 1).

If chromoendoscopy is not available, high definition endoscopy is recommended over standard definition endoscopy^[7,8]. Iannone *et al*^[15] reviewed 10 randomized controlled trials and found chromoendoscopy to be superior to standard definition white light endoscopy, with a relative rate of 2.12 of detecting dysplasia. The major drawback of chromoendoscopy was the longer procedure times. The authors also commented there was no direct evidence of benefit in terms of all cause or cancer specific mortality, nor in the time interval to the development of neoplasia. With further technological improvements there have been significant advancement in endoscopy resolution however most trials have compared chromoendoscopy to standard definition endoscopy. Iacucci *et al*^[16] performed a randomised, non-inferiority study to compare high definition endoscopy, chromoendoscopy and iSCAN image enhanced virtual chromoendoscopy. This paper concluded that virtual

Table 3 Low, intermediate, and high risk features to risk stratify patients and guide surveillance intervals

Low risk	Intermediate risk	High risk
Quiescent disease, even with extensive colonic involvement; left sided IBD	Extensive colonic involvement with mild inflammation; post inflammatory polyps; CRC in 1 st degree relative aged > 50	Extensive colitis with moderate/severe inflammation; primary sclerosing cholangitis; colonic strictures ¹ ; dysplasia of any grade ¹ ; CRC in 1 st degree relative aged < 50

¹In particular if in the prior 5 years.

chromoendoscopy and high definition endoscopy were non inferior to chromoendoscopy in the detection of neoplasia during surveillance. In actual fact, they found high definition endoscopy alone to be adequate in detecting dysplasia without the drawback of the longer procedure times of chromoendoscopy.

An area of interest is the utility of virtual chromoendoscopy or the utility of digital image enhancement, such as narrow band imaging (NBI, Olympus, Japan), Fujifilm Intelligent Colour Enhancement (FICE, Fujifilm, Japan), i-scan OE (Pentax, Japan), CBI (Aohua Photoelectricity, China), VIST (Sonoscape, China), Storz Professional Image Enhancement System (SPIES, Storz, Germany). These optical techniques allow for visual enhancement of the mucosal images through the utility of optical filters, rather than the use of physical contrast dyes, however unlike dye chromoendoscopy, optical filters attempt to narrow the red light spectrum, which preferentially emphasizes vascular patterns^[3].

NBI has been demonstrated to have less withdrawal times compared to chromoendoscopy, with mean times of 15.74 min and 26.87 min respectively. However, although NBI has a similar true positive rate compared to chromoendoscopy and an inferior false positive biopsy rate, NBI has a higher percentage rate of missed dysplasia. As a consequence, at present, NBI is not recommended in place of standard or high definition endoscopy or chromoendoscopy^[7,8,17].

I-Scan is a new generation hybrid virtual chromoendoscopy system integrated with a EPKi high definition unit (Pentax, Japan) allowing for a combination of optical and virtual enhancements in the single entity. It can enhance surface characteristics through modification of light-dark contrast, can digitally enhance blue areas to emphasize vascular abnormalities, and can provide real time modification of the red-green-blue spectrum of the high definition images to highlight fine mucosal abnormalities^[18,19]. This is a digital technique of chromoendoscopy, without the need for contrast dyes. Other digital options include the FICE and SPIES systems. At present, there is no conclusive evidence for the utility of I-Scan, FICE or SPIES in the surveillance of colorectal neoplasia in IBD, however there is ongoing research which will have to be followed closely^[3]. Due to the inherent limitations of xenon light sources, novel approaches have been to utilize laser based systems, such as the Blue Laser Imaging (Fujifilm, Japan), or light-emitting diodes such as Blue Light Imaging (Fujifilm, Japan) or Linked Colour Imaging (Fujifilm, Japan). One report suggested Linked Colour Imaging technology to be superior in the diagnosis of residual inflammation, especially where the Mayo endoscopic subscore was 0^[5,20]. These are however novel approaches and further research and refinement will be required before they become standard of care. At present, virtual chromoendoscopy has not been demonstrated to be superior to conventional chromoendoscopy. This lack of superiority may perhaps be related to virtual techniques generating less light intensity as well as enhancing vasculature patterns whilst chromoendoscopy emphasizes changes in crypt patterns^[4].

Current evidence indicates optimum surveillance modality is through the use of conventional chromoendoscopy or high definition endoscopy with targeted biopsies of concerning mucosal areas. It must be acknowledged that with further technological advances and clinical trials, the preferred modality may change. It should be highlighted that irrespective of modality, the time for withdrawal when performing surveillance for neoplasia in IBD is often higher given dysplastic changes may be very subtle and elusive. Furthermore, it must be emphasized that irrespective of modality, quality is largely impacted upon by mucosal interrogation, with longer and more detailed examinations conferring improvements in neoplasia detection^[4].

ENDOSCOPIC RESECTION

Table 4 summarizes the SCENIC consensus nomenclature of dysplasia in




Figure 1 An approach to dysplasia. EMR: Endoscopic mucosal resection; ESD: Endoscopic submucosal dissection.

inflammatory bowel disease^[5]. On visualization of a lesion, it is imperative to first recognize whether it is within an area of colitis or not. Key features to appreciate include the morphology, margin definition, surface characteristics, size, background mucosa characteristics, and endoscopic access. The lesion should be categorized as either polypoid, *i.e.*, pedunculated or sessile, or non-polypoid, *i.e.*, slightly elevated, flat or depressed. Borders should be classified as distinct or indistinct. It is important to note any suggestions of submucosal invasion, such as the presence of ulceration, depression or failure to lift following submucosal injection^[7].

On identification of an area of dysplasia, it is important to determine if the lesion is endoscopically resectable. Performing a biopsy can result in fibrosis, a major risk factor for complications of subsequent future resection strategies such as endoscopic submucosal dissection^[21]. Therefore, if it is felt to be technically safe and appropriate, areas of dysplasia should be endoscopically resected⁴ rather than undergo unnecessary biopsy. A lesion should be biopsied if the lesion is considered inappropriate for immediate resection or from surrounding tissue post endoscopic resection to assess for invisible dysplasia^[9].

A lesion with distinct borders may be considered for endoscopic resection whilst

Table 4 SCENIC consensus nomenclature of dysplasia in inflammatory bowel disease

Term	Definition
Visible dysplasia	Dysplasia confirmed histologically on a targeted biopsy
Invisible dysplasia	Dysplasia on a random biopsy
Polypoid	Lesion protruding ≥ 2.5 mm into the lumen
Non polypoid	Lesion protruding < 2.5 mm into the lumen or not protruding
Superficial elevated	Protrusion < 2.5 mm
Pedunculated	Attached to mucosa via stalk
Sessile	Not attached via stalk; base contiguous with mucosa
Flat	No protrusion above mucosa
Depressed	At least a portion of lesion depressed below mucosa
Ulcerated	Fibrinous appearing base within lesion
Distinct border	Easily identified from surrounding mucosa
Indistinct border	Not discrete; difficult to distinguish from surrounding mucosa

indistinct borders are preferably surgically managed. An important caveat is that inflammation can impact upon margin visualization. Nevertheless, any well-defined lesion should be endoscopically resected, regardless of the grade of dysplasia^[4]. Surface features should be categorized according to standard grading systems such as the Kudo Pit pattern, JNET or MS classification. Lesions identified in areas distinct from areas of colitis should be treated as per standard practice for spontaneous adenomas.

The risk and benefit profile of an individual case will guide the appropriateness of endoscopic management against definitive colectomy. Features favorable for colectomy include ill-defined margins, submucosal invasion, asymmetrical lift not attributable to fibrosis, ulceration or large depression and flat neoplastic changes adjacent to the lesion. It is also important to highlight that if performing endoscopic resection in inflammatory bowel disease, submucosal injection may be difficult, hence careful marking (APC/tip of snare) is essential and one may require a stiffer snare such as a braided or "Histolock" snare. Following endoscopic resection, biopsies of surrounding tissues help evaluate the margins.

Standard snaring is acceptable for small, polypoid or protuberant lesions less than 10 mm in size. Extra care must be taken to ensure complete resection and retrieval. For larger lesions up to 2 cm in size, one may consider *en bloc* endoscopic mucosal resection. If it is larger than 2 cm, then one may consider piecemeal endoscopic mucosal resection unless there is suspicion of submucosal invasion. Endoscopic submucosal dissection (ESD) is an alternative however this may be technically very challenging. Current guidelines recommend random biopsies of peri-resection areas to ensure complete resection, however the diagnostic yield of this is questionable given the development of high definition and chromoendoscopy, as previously discussed^[5].

Following endoscopic resection, current guidelines recommend surveillance high definition chromoendoscopy by a specialized IBD proceduralist, rather than colectomy, annually for 5 years. If surveillance is not an option, colectomy may be considered^[4].

With regards to invisible dysplasia, once confirmed by a specialist gastrointestinal pathologist, cases should be referred to an IBD specialist and surveillance performed with chromoendoscopy with high definition colonoscopy^[3]. Low grade endoscopically invisible dysplasia should be referred for multidisciplinary team review. Given factors such as disease activity and overall patient specific CRC risk factors, a patient centered management plan must be determined and may involve 6 monthly to yearly surveillance or may warrant referral for colectomy. Before the development of high definition and chromoendoscopy, high grade invisible dysplasia would warrant prompt referral for colectomy^[22-24]. With the new generation of endoscopic techniques, the management of invisible high grade dysplasia is similar to that of invisible low grade dysplasia, with emphasis being on a multidisciplinary team, patient centered approach to determine whether close surveillance or colectomy is more appropriate^[5].

CONCLUSION

With the improvement in endoscopic imaging technologies, surveillance of patients

with IBD have reached new frontiers. Huge progress has also been made in the management of dysplastic lesions. Many of the flat or 'targeted biopsy-only' detected dysplasia that had historically warranted a colectomy can now be visualised as circumscribed lesions with the help of high definition scopes and chromoendoscopy. These lesions are now amenable to endoscopic excision with close subsequent endoscopic follow-up. However, management decisions should be balanced with the risks and benefits of endoscopic, medical, and surgical treatments, during clinical decision making.

Key points

Most colonic dysplasia is visually identifiable with current high definition endoscopic imaging modalities and hence targeted biopsies is advised. Random biopsies may be utilized in patients with a personal history of neoplasia, PSC, or a tubular appearing colon. Lesions deemed to be endoscopically resectable should be referred to centers with expertise to do so and be endoscopically resected rather than undergo unnecessary biopsy. Invisible dysplasia should be referred to an IBD specialist and surveillance performed with chromoendoscopy with high definition colonoscopy

REFERENCES

- 1 **Danese S, Fiocchi C.** Ulcerative colitis. *N Engl J Med* 2011; **365**: 1713-1725 [PMID: [22047562](#) DOI: [10.1056/NEJMra1102942](#)]
- 2 **Ten Hove JR, Bernstein CN, Oldenburg B.** Putting Evidence into Practice: IBD Surveillance, Chromoendoscopy and Future Directions. *Am J Gastroenterol* 2017 [PMID: [29087392](#) DOI: [10.1038/ajg.2017.390](#)]
- 3 **Sivanathan V, Tontini GE, Möhler M, Galle PR, Neumann H.** Advanced endoscopic imaging for diagnosis of inflammatory bowel diseases: Present and future perspectives. *Dig Endosc* 2018; **30**: 441-448 [PMID: [29360261](#) DOI: [10.1111/den.13023](#)]
- 4 **Goetz M.** Endoscopic Surveillance in Inflammatory Bowel Disease. *Visc Med* 2018; **34**: 66-71 [PMID: [29594172](#) DOI: [10.1159/000485019](#)]
- 5 **Ibraheim H, Dhillon AS, Koumoutsos I, Gulati S, Hayee B.** Curriculum review: colorectal cancer surveillance and management of dysplasia in IBD. *Frontline Gastroenterol* 2018; **9**: 271-277 [PMID: [30245789](#) DOI: [10.1136/flgastro-2017-100919](#)]
- 6 **Collins PD, Mpofu C, Watson AJ, Rhodes JM.** Strategies for detecting colon cancer and/or dysplasia in patients with inflammatory bowel disease. *Cochrane Database Syst Rev* 2006; CD000279 [PMID: [16625534](#) DOI: [10.1002/14651858.CD000279.pub3](#)]
- 7 **American Society for Gastrointestinal Endoscopy Standards of Practice Committee, Shergill AK, Lightdale JR, Bruining DH, Acosta RD, Chandrasekhara V, Chathadi KV, Decker GA, Early DS, Evans JA, Fanelli RD, Fisher DA, Fonkalsrud L, Foley K, Hwang JH, Jue TL, Khashab MA, Muthusamy VR, Pasha SF, Saltzman JR, Sharaf R, Cash BD, DeWitt JM.** The role of endoscopy in inflammatory bowel disease. *Gastrointest Endosc* 2015; **81**: 1101-21.e1-13 [PMID: [25800660](#) DOI: [10.1016/j.gie.2014.10.030](#)]
- 8 **Laine L, Kaltenbach T, Barkun A, McQuaid KR, Subramanian V, Soetikno R; SCENIC Guideline Development Panel.** SCENIC international consensus statement on surveillance and management of dysplasia in inflammatory bowel disease. *Gastroenterology* 2015; **148**: 639-651.e28 [PMID: [25702852](#) DOI: [10.1053/j.gastro.2015.01.031](#)]
- 9 **Watanabe T, Ajioka Y, Mitsuyama K, Watanabe K, Hanai H, Nakase H, Kunisaki R, Matsuda K, Iwakiri R, Hida N, Tanaka S, Takeuchi Y, Ohtsuka K, Murakami K, Kobayashi K, Iwao Y, Nagahori M, Iizuka B, Hata K, Igarashi M, Hirata I, Kudo SE, Matsumoto T, Ueno F, Watanabe G, Ikegami M, Ito Y, Oba K, Inoue E, Tomotsugu N, Takebayashi T, Sugihara K, Suzuki Y, Watanabe M, Hibi T.** Comparison of Targeted vs Random Biopsies for Surveillance of Ulcerative Colitis-Associated Colorectal Cancer. *Gastroenterology* 2016; **151**: 1122-1130 [PMID: [27523980](#) DOI: [10.1053/j.gastro.2016.08.002](#)]
- 10 **van den Broek FJ, Stokkers PC, Reitsma JB, Boltjes RP, Ponsioen CY, Fockens P, Dekker E.** Random biopsies taken during colonoscopic surveillance of patients with longstanding ulcerative colitis: low yield and absence of clinical consequences. *Am J Gastroenterol* 2014; **109**: 715-722 [PMID: [21427710](#) DOI: [10.1038/ajg.2011.93](#)]
- 11 **Moussata D, Allez M, Cazals-Hatem D, Treton X, Laharie D, Reimund JM, Bertheau P, Bourrille A, Lavergne-Slove A, Brixi H, Branche J, Gornet JM, Stefanescu C, Moreau J, Marteau P, Pelletier AL, Carbonnel F, Seksik P, Simon M, Fléjou JF, Colombel JF, Charlois AL, Roblin X, Nancey S, Bouhnik Y, Berger F, Flourié B; the GETAID.** Are random biopsies still useful for the detection of neoplasia in patients with IBD undergoing surveillance colonoscopy with chromoendoscopy? *Gut* 2018; **67**: 616-624 [PMID: [28115492](#) DOI: [10.1136/gutjnl-2016-311892](#)]
- 12 **Subramanian V, Ramappa V, Telakis E, Mannath J, Jawhari AU, Hawkey CJ, Ragunath K.** Comparison of high definition with standard white light endoscopy for detection of dysplastic lesions during surveillance colonoscopy in patients with colonic inflammatory bowel disease. *Inflamm Bowel Dis* 2013; **19**: 350-355 [PMID: [22552948](#) DOI: [10.1002/ibd.23002](#)]
- 13 **Kaltenbach T, Shergill AK, Wallace MB.** How to obtain and use chromoendoscopy dyes for surveillance colonoscopy in inflammatory bowel disease: a technical guide. *Gastrointest Endosc* 2017; **86**: 949-951 [PMID: [28974381](#) DOI: [10.1016/j.gie.2017.09.022](#)]
- 14 **Wu L, Li P, Wu J, Cao Y, Gao F.** The diagnostic accuracy of chromoendoscopy for dysplasia in ulcerative colitis: meta-analysis of six randomized controlled trials. *Colorectal Dis* 2012; **14**: 416-420 [PMID: [21073646](#) DOI: [10.1111/j.1463-1318.2010.02505.x](#)]
- 15 **Iannone A, Ruospo M, Wong G, Principi M, Barone M, Strippoli GFM, Di Leo A.** Chromoendoscopy for Surveillance in Ulcerative Colitis and Crohn's Disease: A Systematic Review of Randomized Trials. *Clin Gastroenterol Hepatol* 2017; **15**: 1684-1697.e11 [PMID: [27890853](#)]

- DOI: [10.1016/j.cgh.2016.11.021](https://doi.org/10.1016/j.cgh.2016.11.021)]
- 16 **Iacucci M**, Kaplan GG, Panaccione R, Akinola O, Lethebe BC, Lowerison M, Leung Y, Novak KL, Seow CH, Urbanski S, Minoo P, Gui X, Ghosh S. A Randomized Trial Comparing High Definition Colonoscopy Alone With High Definition Dye Spraying and Electronic Virtual Chromoendoscopy for Detection of Colonic Neoplastic Lesions During IBD Surveillance Colonoscopy. *Am J Gastroenterol* 2018; **113**: 225-234 [PMID: [29134964](https://pubmed.ncbi.nlm.nih.gov/29134964/) DOI: [10.1038/ajg.2017.417](https://doi.org/10.1038/ajg.2017.417)]
 - 17 **Pellisé M**, López-Cerón M, Rodríguez de Miguel C, Jimeno M, Zabalza M, Ricart E, Aceituno M, Fernández-Esparrach G, Ginès A, Sendino O, Cuatrecasas M, Llach J, Panés J. Narrow-band imaging as an alternative to chromoendoscopy for the detection of dysplasia in long-standing inflammatory bowel disease: a prospective, randomized, crossover study. *Gastrointest Endosc* 2011; **74**: 840-848 [PMID: [21802681](https://pubmed.ncbi.nlm.nih.gov/21802681/) DOI: [10.1016/j.gie.2011.05.013](https://doi.org/10.1016/j.gie.2011.05.013)]
 - 18 **Neumann H**, Fujishiro M, Wilcox CM, Mönkemüller K. Present and future perspectives of virtual chromoendoscopy with i-scan and optical enhancement technology. *Dig Endosc* 2014; **26** Suppl 1: 43-51 [PMID: [24373000](https://pubmed.ncbi.nlm.nih.gov/24373000/) DOI: [10.1111/den.12190](https://doi.org/10.1111/den.12190)]
 - 19 **Iacucci M**, Kiesslich R, Gui X, Panaccione R, Heatherington J, Akinola O, Ghosh S. Beyond white light: optical enhancement in conjunction with magnification colonoscopy for the assessment of mucosal healing in ulcerative colitis. *Endoscopy* 2017; **49**: 553-559 [PMID: [28315280](https://pubmed.ncbi.nlm.nih.gov/28315280/) DOI: [10.1055/s-0042-124363](https://doi.org/10.1055/s-0042-124363)]
 - 20 **Uchiyama K**, Takagi T, Kashiwagi S, Toyokawa Y, Tanaka M, Hotta Y, Dohi O, Okayama T, Yoshida N, Katada K, Kamada K, Ishikawa T, Handa O, Konishi H, Kishimoto M, Yagi N, Naito Y, Itoh Y. Assessment of Endoscopic Mucosal Healing of Ulcerative Colitis Using Linked Colour Imaging, a Novel Endoscopic Enhancement System. *J Crohns Colitis* 2017; **11**: 963-969 [PMID: [28333209](https://pubmed.ncbi.nlm.nih.gov/28333209/) DOI: [10.1093/ecco-jcc/jjx026](https://doi.org/10.1093/ecco-jcc/jjx026)]
 - 21 **Lee SP**, Kim JH, Sung IK, Lee SY, Park HS, Shim CS, Han HS. Effect of submucosal fibrosis on endoscopic submucosal dissection of colorectal tumors: pathologic review of 173 cases. *J Gastroenterol Hepatol* 2015; **30**: 872-878 [PMID: [25641510](https://pubmed.ncbi.nlm.nih.gov/25641510/) DOI: [10.1111/jgh.12886](https://doi.org/10.1111/jgh.12886)]
 - 22 **Ignjatovic A**, East JE, Subramanian V, Suzuki N, Guenther T, Palmer N, Bassett P, Ragnath K, Saunders BP. Narrow band imaging for detection of dysplasia in colitis: a randomized controlled trial. *Am J Gastroenterol* 2012; **107**: 885-890 [PMID: [22613903](https://pubmed.ncbi.nlm.nih.gov/22613903/) DOI: [10.1038/ajg.2012.67](https://doi.org/10.1038/ajg.2012.67)]
 - 23 **Ayoola R**, Mohanty M, Lee JE, Sifuentes H. The Role of Chromoendoscopy and Enhanced Imaging Techniques in Inflammatory Bowel Disease Colorectal Cancer Colonoscopy Surveillance. Sridhar S, Wu G. *Diagnostic and Therapeutic Procedures in Gastroenterology*. Cham: Humana Press; 2018; 331-337 [DOI: [10.1007/978-3-319-62993-3_25](https://doi.org/10.1007/978-3-319-62993-3_25)]
 - 24 **Castañó-Milla C**, Chaparro M, Gisbert JP. Systematic review with meta-analysis: the declining risk of colorectal cancer in ulcerative colitis. *Aliment Pharmacol Ther* 2014; **39**: 645-659 [PMID: [24612141](https://pubmed.ncbi.nlm.nih.gov/24612141/) DOI: [10.1111/apt.12651](https://doi.org/10.1111/apt.12651)]

P- Reviewer: Gazouli M, Papamichail K, Yuksel I

S- Editor: Ma YJ **L- Editor:** A **E- Editor:** Bian YN





Published By Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

