

World Journal of *Clinical Cases*

World J Clin Cases 2019 December 6; 7(23): 3915-4171



**REVIEW**

- 3915** Overview of organic anion transporters and organic anion transporter polypeptides and their roles in the liver
Li TT, An JX, Xu JY, Tuo BG

ORIGINAL ARTICLE**Observational Study**

- 3934** Value of early diagnosis of sepsis complicated with acute kidney injury by renal contrast-enhanced ultrasound
Wang XY, Pang YP, Jiang T, Wang S, Li JT, Shi BM, Yu C
- 3945** Value of elastography point quantification in improving the diagnostic accuracy of early diabetic kidney disease
Liu QY, Duan Q, Fu XH, Fu LQ, Xia HW, Wan YL
- 3957** Resection of recurrent third branchial cleft fistulas assisted by flexible pharyngotomy
Ding XQ, Zhu X, Li L, Feng X, Huang ZC
- 3964** Therapeutic efficacy of acupuncture combined with neuromuscular joint facilitation in treatment of hemiplegic shoulder pain
Wei YH, Du DC, Jiang K
- 3971** Comparison of intra-articular injection of parecoxib *vs* oral administration of celecoxib for the clinical efficacy in the treatment of early knee osteoarthritis
Lu L, Xie Y, Gan K, Huang XW

Retrospective Study

- 3980** Celiomesenteric trunk: New classification based on multidetector computed tomography angiographic findings and probable embryological mechanisms
Tang W, Shi J, Kuang LQ, Tang SY, Wang Y

Prospective Study

- 3990** Interaction of arylsulfatases A and B with maspin: A possible explanation for dysregulation of tumor cell metabolism and invasive potential of colorectal cancer
Kovacs Z, Jung I, Szalman K, Baniás L, Bara TJ, Gurzu S

CASE REPORT

- 4004** Recuperation of severe tumoral calcinosis in a dialysis patient: A case report
Westermann L, Isbell LK, Breitenfeldt MK, Arnold F, Röthele E, Schneider J, Widmeier E

- 4011** Robotic wedge resection of a rare gastric perivascular epithelioid cell tumor: A case report
Marano A, Maione F, Woo Y, Pellegrino L, Geretto P, Sasia D, Fortunato M, Orcioni GF, Priotto R, Fasoli R, Borghi F
- 4020** Primary parahiatal hernias: A case report and review of the literature
Preda SD, Pătrașcu Ș, Ungureanu BS, Cristian D, Bințișan V, Nica CM, Calu V, Strâmbu V, Sapalidis K, Șurlin VM
- 4029** Diagnosis of Laron syndrome using monoplex-polymerase chain reaction technology with a whole-genome amplification template: A case report
Neumann A, Alcántara-Ortigoza MÁ, González-del Ángel A, Camargo-Diaz F, López-Bayghen E
- 4036** *In-vitro* proliferation assay with recycled ascitic cancer cells in malignant pleural mesothelioma: A case report
Anayama T, Taguchi M, Tatenuma T, Okada H, Miyazaki R, Hirohashi K, Kume M, Matsusaki K, Orihashi K
- 4044** Distant metastasis in choroidal melanoma with spontaneous corneal perforation and intratumoral calcification: A case report
Wang TW, Liu HW, Bee YS
- 4052** Secondary Parkinson disease caused by breast cancer during pregnancy: A case report
Li L
- 4057** Pulmonary embolism and deep vein thrombosis caused by nitrous oxide abuse: A case report
Sun W, Liao JP, Hu Y, Zhang W, Ma J, Wang GF
- 4063** Micronodular thymic tumor with lymphoid stroma: A case report and review of the literature
Wang B, Li K, Song QK, Wang XH, Yang L, Zhang HL, Zhong DR
- 4075** Diffuse large B cell lymphoma with bilateral adrenal and hypothalamic involvement: A case report and literature review
An P, Chen K, Yang GQ, Dou JT, Chen YL, Jin XY, Wang XL, Mu YM, Wang QS
- 4084** Urethral pressure profilometry in artificial urinary sphincter implantation: A case report
Meng LF, Liu XD, Wang M, Zhang W, Zhang YG
- 4091** Hydroxyurea-induced cutaneous squamous cell carcinoma: A case report
Xu Y, Liu J
- 4098** Recurrent hypotension induced by sacubitril/valsartan in cardiomyopathy secondary to Duchenne muscular dystrophy: A case report
Li JM, Chen H
- 4106** Complete duodenal obstruction induced by groove pancreatitis: A case report
Wang YL, Tong CH, Yu JH, Chen ZL, Fu H, Yang JH, Zhu X, Lu BC

- 4111** Radiological aspects of giant hepatocellular adenoma of the left liver: A case report
Zheng LP, Hu CD, Wang J, Chen XJ, Shen YY
- 4119** Mixed serous-neuroendocrine neoplasm of the pancreas: A case report and review of the literature
Xu YM, Li ZW, Wu HY, Fan XS, Sun Q
- 4130** Rigid esophagoscopy combined with angle endoscopy for treatment of superior mediastinal foreign bodies penetrating into the esophagus caused by neck trauma: A case report
Wang D, Gao CB
- 4137** Left armpit subcutaneous metastasis of gastric cancer: A case report
He FJ, Zhang P, Wang MJ, Chen Y, Zhuang W
- 4144** Bouveret syndrome: A case report
Wang F, Du ZQ, Chen YL, Chen TM, Wang Y, Zhou XR
- 4150** Fatal complications in a patient with severe multi-space infections in the oral and maxillofacial head and neck regions: A case report
Dai TG, Ran HB, Qiu YX, Xu B, Cheng JQ, Liu YK
- 4157** Management of massive fistula bleeding after endoscopic ultrasound-guided pancreatic pseudocyst drainage using hemostatic forceps: A case report
Ge N, Sun SY
- 4163** Pure squamous cell carcinoma of the gallbladder locally invading the liver and abdominal cavity: A case report and review of the literature
Jin S, Zhang L, Wei YF, Zhang HJ, Wang CY, Zou H, Hu JM, Jiang JF, Pang LJ

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Consolato M Sergi, FRCP (C), MD, PhD, Professor, Department of Lab Medicine and Pathology, University of Alberta, Edmonton T6G 2B7, Canada

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now indexed in PubMed, PubMed Central, Science Citation Index Expanded (also known as SciSearch®), and Journal Citation Reports/Science Edition. The 2019 Edition of Journal Citation Reports cites the 2018 impact factor for WJCC as 1.153 (5-year impact factor: N/A), ranking WJCC as 99 among 160 journals in Medicine, General and Internal (quartile in category Q3).

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: *Yan-Xia Xing*

Proofing Production Department Director: *Xiang Li*

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Semimonthly

EDITORS-IN-CHIEF

Dennis A Bloomfield, Bao-Gan Peng, Sandro Vento

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/2307-8960/editorialboard.htm>

EDITORIAL OFFICE

Jin-Lei Wang, Director

PUBLICATION DATE

December 6, 2019

COPYRIGHT

© 2019 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Observational Study

Therapeutic efficacy of acupuncture combined with neuromuscular joint facilitation in treatment of hemiplegic shoulder pain

Yan-Hui Wei, De-Chao Du, Ke Jiang

ORCID number: Yan-Hui Wei (0000-0002-3455-9836); De-Chao Du (0000-0003-2068-3008); Ke Jiang (0000-0002-5946-215X).

Author contributions: Wei YH and Du DC contributed equally to this manuscript; Wei YH, Du DC, and Jiang K designed the research and wrote and revised the manuscript.

Institutional review board

statement: The study was approved by the Ethics Committee of the Affiliated Hospital of Nanjing University of Chinese Medicine.

Informed consent statement: All patients in this study gave informed consent.

Conflict-of-interest statement: The authors declare that they have no competing interests.

Data sharing statement: No additional data are available.

STROBE statement: The authors have read the STROBE statement, and the manuscript was prepared and revised according to the STROBE statement.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the

Yan-Hui Wei, Department of Rehabilitation, Nanjing University of Chinese Medicine, Nanjing 210028, Jiangsu Province, China

De-Chao Du, Department of Rehabilitation, Nanjing Red Cross Hospital, Nanjing 210028, Jiangsu Province, China

Ke Jiang, Department of Rehabilitation Physiotherapy, Jiangsu Provincial Hospital of Integration of Chinese and Western Medicine, Nanjing 210028, Jiangsu Province, China

Corresponding author: Ke Jiang, MD, Chief Physician, Department of Rehabilitation Physiotherapy, Jiangsu Provincial Hospital of Integration of Chinese and Western Medicine, No. 100, Cross Street, Hongshan Road, Nanjing 210028, Jiangsu Province, China. shanchuan-baobei@163.com

Abstract

BACKGROUND

Stroke is a type of cerebrovascular disease with high prevalence, mortality, and onset of disability. As a neurodevelopmental therapy, neuromuscular joint facilitation (NJF) is widely used in the treatment of orthopedic and neurological disorders in the clinical practice. It is mainly used for central nervous system diseases or orthopedic diseases, movement disorders, and pain rehabilitation. According to related studies, NJF can also be used as a rehabilitation treatment in patients with hemiplegic shoulder pain (HSP).

AIM

To investigate the clinical efficacy of acupuncture combined with NJF in patients with HSP.

METHODS

Forty patients with HSP were randomly divided into a treatment group and a control group. The treatment group was treated with acupuncture combined with NJF and the control group was treated with acupuncture alone. All patients were assessed by using the visual analogue scale (VAS), Fugl-Meyer assessment (FMA), Barthel index (BI), and passive range of motion (PROM) before and after the training. All the clinical data were analyzed using SPSS 20.0 statistical software.

RESULTS

There was no statistical difference in the general characteristics between the two groups. In the terms of duration of treatment, age, and pre-treatment indicators, the two groups were comparable ($P > 0.05$). After the treatment, VAS, PROM, BI,

original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Received: September 8, 2019

Peer-review started: September 8, 2019

First decision: November 11, 2019

Revised: October 19, 2019

Accepted: November 23, 2019

Article in press: November 23, 2019

Published online: December 6, 2019

P-Reviewer: Munjal K, Ugezu C

S-Editor: Wang JL

L-Editor: Wang TQ

E-Editor: Wu YXJ



and FMA scores were significantly improved in the two groups of patients ($P < 0.05$). The VAS, PROM and FMA scores were significantly higher in the treatment group than in the control group ($P < 0.05$). However, there was no significant difference in BI scores between the two groups ($P > 0.05$).

CONCLUSION

Both acupuncture alone and acupuncture combined with NJF in the treatment of HSP are effective, and can improve the clinical symptoms of patients.

Acupuncture combined with NJF can improve the upper limb motor function, relieve pain, and increase joint mobility in patients with HSP. The combination therapy is better than acupuncture alone. However, there is no significant difference in improving the score of patients' self-care ability.

Key words: Acupuncture; Neuromuscular joint facilitation; Hemiplegic shoulder pain; Visual analogue scale; Fugl-Meyer assessment; Therapy

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Shoulder pain is one of the common complications in stroke patients.

Neuromuscular joint facilitation (NJF) is a therapeutic exercise which is used to improve the movement of the joint. Acupuncture is a key component of traditional treatment to relieve pain. The present research studied whether the efficacy of acupuncture combined with NJF was superior to acupuncture alone for hemiplegic shoulder pain (HSP). The results showed that the combined therapy of NJF and acupuncture is better than acupuncture alone in the treatment of HSP.

Citation: Wei YH, Du DC, Jiang K. Therapeutic efficacy of acupuncture combined with neuromuscular joint facilitation in treatment of hemiplegic shoulder pain. *World J Clin Cases* 2019; 7(23): 3964-3970

URL: <https://www.wjgnet.com/2307-8960/full/v7/i23/3964.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v7.i23.3964>

INTRODUCTION

Stroke is a type of cerebrovascular disease with high prevalence, mortality, and onset of disability. It has become the second leading cause of death in human beings today^[1]. According to the statistics, the incidence of new-onset stroke in China exceeds 2 million per year^[2], and its incidence is increasing year by year. With the increase in the level of social medical care, the mortality of stroke has decreased, but the disability rate of up to 80% has always been a serious burden for individuals, families, and the society. Shoulder pain is one of the common stroke sequelae, and has an incidence rate as high as 84%^[3], two thirds of which occur within 4 wk to 2 mo after stroke, which is the best period for rehabilitation of stroke patients. It is often presented as a common mode of motion. However, the occurrence of hemiplegic shoulder pain (HSP) can easily affect the upper limb training, hinder the movements, and have a certain effect on the recovery of joint function, the daily living ability, and the quality of life in the patients. It also causes harm to the mental health of patients, resulting in the occurrence and aggravation of adverse psychological reactions such as depression and anxiety after stroke^[4,5].

As a neurodevelopmental therapy, NJF is widely used in the treatment of orthopedic and neurological disorders in the clinical practice^[6-9]. In particular, it is mainly used for central nervous system diseases or orthopedic diseases, movement disorders, and pain rehabilitation. According to related studies, NJF can also be used as a rehabilitation treatment in patients with HSP. However, the research is still insufficient. The present study combined acupuncture with NJF technology to treat HSP, with an aim to provide a new treatment choice for this disease.

MATERIALS AND METHODS

Participants

From January 2016 to February 2017, 40 patients with HSP were selected at the Department of Acupuncture and Rehabilitation and the Department of Neurology, Integration of Chinese and Western Medicine.

Diagnostic criteria

The criteria for traditional Chinese medicine diagnosis used in the present study were the Standards for Diagnosis and Efficacy of Stroke Diseases compiled by the State Administration of Traditional Chinese Medicine in 1996^[10].

The criteria for Western medicine diagnosis used in the present study were the Guidelines for the Diagnosis and Treatment of Acute Ischemic Stroke in China published by the Cerebrovascular Diseases Group of the Chinese Medical Association Neurology Branch in 2014^[11].

Inclusion and exclusion criteria

To be included, patients should meet the following criteria: (1) Diagnosed as having cerebral hemorrhage or cerebral infarction according to the diagnostic criteria of Chinese and Western medicine, and by head computed tomography or magnetic resonance imaging (MRI); (2) Aged between 35 and 85 years; (3) First seizure with a disease course of 2 wk to 6 mo; (4) Hemiplegia and shoulder pain after stroke; (5) Sober-minded, with stable vital signs, and without communication barriers or other serious disorders; and (6) Willing to receive the treatment.

Patients were excluded if they clearly met one or more of the following exclusion criteria: (1) With a history of progressive stroke or multiple episodes, or having undergone a previous thrombolysis; (2) With aphasia, sensory processing disorder, or cognitive dysfunction; (3) With pre-existing orthopedic or neurological diseases which affect upper limb functional activities such as cervical spondylosis, frozen shoulder, upper limb fracture, thalamic lesion, and peripheral nerve injury; and (4) With a history of brain trauma, tumor, or surgery, or other heart disease with atrial fibrillation or pacemaker installed.

In addition, patients who cannot complete the study and naturally dropped out during the observation period, patients who took other drugs without permission during the study, which affects the validity of efficacy, patients whose condition was unstable and whose condition was aggravated during the treatment, and patients in whom adverse reactions or other serious complications occur were also excluded.

Treatment

The 40 patients selected were numbered according to the date of the visit, and were randomly assigned to a treatment group and a control group according to the random number table method, with 20 patients in each group.

Hua Tuo brand disposable acupuncture needles (0.30 mm × 40 mm) were used as acupuncture needles in the study. Our acupuncturists mainly selected the following acupuncture points to perform the acupuncture procedure: Jianyu (LI15), punctured obliquely 0.8–1.5 inches on the deltoid muscle; Jianzhen (SI9), punctured obliquely 1–1.5 inch; Jianliao (SJ14), punctured straightly 1–1.5 inch; Tianzong (SI11), punctured obliquely 0.5–1 inch; Jianqian (EX-UE), punctured straightly 1–1.5 inches; Binao (LI14), punctured straightly 0.8–1 inch; Quchi (LI11), punctured straightly 0.5–1 inch; Waiguan (SJ5), punctured straightly 0.5–1 inch; Hegu (LI4), straightly pricked 0.5–1 inch; and Shousanli (LI10), punctured straightly 0.8–1.2 inches. We suggest using the method of inserting and twisting, mild reinforcing and attenuating acupuncture to achieve arrival of the qi. Each time the needle was left for 30 min, once a day and six times a week, for a total of three weeks.

For the control group, the following acupuncture points were selected on the affected side of the shoulder: LI15, SI9, SJ14, SI11, EX-UE, LI14, LI11, LI10, SJ5, and LI4. For the treatment group, the acupuncture treatment was the same as the control group. In addition, NJF was applied for rehabilitation training.

Patients in the treatment group were also trained to do exercises mainly involving articular surface movement including extension-adduction-internal rotation, flexion-abduction-external rotation, flexion-adduction-external rotation, and extension-abduction-internal rotation. Opposite joint motion was also performed, which involved the upper limb-shoulder, including extension-adduction-internal rotation + shoulder abduction-upper rotation, flexion-abduction-external rotation + shoulder adduction-lower rotation, extension-abduction-internal rotation + shoulder abduction-above rotation, and flexion-adduction-external rotation + shoulder extension-lower rotation.

Effectiveness evaluation

Patients were assessed for rehabilitation before and after treatment. The Clinical Diagnosis and Treatment Guide, Physical Medicine and Rehabilitation compiled by the Chinese Medical Association in 2005^[12] was used to evaluate the functional activity of the shoulder joint. The relevant rehabilitation indexes were used to evaluate the rehabilitation effect of HSP in the following four aspects: (1) Visual analogue scale (VAS); (2) Passive range of shoulder joints (PROM); (3) Barthel index score (BI); and (4) Fugl-Meyer assessment (FMA) score for upper limb motor function.

Statistical analysis

Statistical analyses were performed on relevant clinical data with IBM SPSS Statistics 20.0. Student's *t*-test was used for the data measured which conform to a normal distribution. Student's *t*-test was also used to compare the difference before and after treatment. Independent *t*-test was used to compare the difference between groups. χ^2 test was used to compare the difference in quantitative data, and nonparametric test was used for data which do not conform to a normal distribution. $P < 0.05$ indicated that there was a statistical difference.

RESULTS

Patient characteristics

There was no significant difference in age or duration of disease between the treatment group and the control group ($P > 0.05$, Table 1).

VAS scores

Table 2 shows that there was no significant difference in VAS scores between the two groups before treatment ($P > 0.05$). However, after the treatment, the VAS scores were significantly reduced in the two groups ($P < 0.05$). The VAS score was significantly better in the treatment group than in the control group ($P < 0.05$).

PROM scores

Table 3 shows that there was no significant difference in the PROM scores between the two groups before treatment ($P > 0.05$). However, after the treatment, the PROM scores were significantly increased in the two groups ($P < 0.05$). The PROM score was significantly better in the treatment group than in the control group ($P < 0.05$).

FMA scores

There was no significant difference in FMA scores between the two groups before treatment ($P > 0.05$). However, after the treatment, the FMA scores of the two groups were significantly increased ($P < 0.05$). The FMA score was significantly better in the treatment group than in the control group ($P < 0.05$, Table 2).

BI scores

There was no significant difference in BI scores between the two groups before treatment ($P > 0.05$, Table 2). After the treatment, the BI scores of the two groups were significantly increased ($P < 0.05$). However, there was no significant difference in BI scores between the two groups after the treatment ($P > 0.05$).

DISCUSSION

Acupuncture has a positive effect on shoulder joint activity in patients with hemiplegia^[13]. The acupuncture points we selected based on the disease are mainly on the extensor muscle. For example, LI15, SI9, SJ14, EX-UE and LI14 are located on the deltoid muscle; SI11 is located on the infraorbital muscle; LI11 is located on the radial extensor muscle. LI4 is located between the 1st and 2nd metacarpal bones, and at its deep layer there is the elbow extensor muscle. The acupuncture points are mainly on antagonistic muscle groups, the upper limbs contain lots of extensor muscle groups, and the lower limbs contain lots of flexor muscle groups. The acupuncture points near the extensor muscle group can improve muscle strength^[14]. This helps to improve motor dysfunction and limb function activities in patients after stroke. The present study found that acupuncture has a certain effect on shoulder mobility and upper limb motor function in patients with hemiplegia.

Acupuncture can change the microenvironment of damaged muscles^[15-20]. Moreover, it promotes the recovery of muscle fiber structures by improving arrangement of filament in muscle, eliminating mitochondrial swelling, dilating

Table 1 Comparison of basic conditions between the two groups

Group	Age (yr)	Disease course
Treatment	63.85 ± 11.07	2.50 ± 1.54
Control	65.55 ± 13.30	2.25 ± 1.68

sarcoplasmic reticulum and so on; it also improves intracellular calcium metabolism by promoting cell membrane sodium and calcium exchange, promotes the recovery of resting potential of skeletal muscle, accelerates muscle regeneration by promoting the proliferation of muscle satellite cells, and accelerates polymerization and recombination by inhibiting myosin degradation. By inhibiting the production of myogenic IL-6, it has an effect on skeletal muscle inflammation. Acupuncture can play a protective role in the treatment of muscle injury through a series of microstructural changes of skeletal muscle. For the effect of acupuncture on pain relief, studies^[21] have shown that acupuncture can not only activate the endogenous opioid peptide system in the body, result in the release of related transmitter products, increase pain threshold, and relieve pain, but also stimulate the transmission of pain-induced fibers and C fibers through the excitatory and central nervous system inhibition of the spinal cord. This may ultimately influence the interaction of various functional intervals of the cortex and achieve analgesic purposes, result in the release of amine and enkephalin-like neurotransmitters, and promote the decline of striatal adrenaline and cortical adrenaline^[22]. A study shows that electro-acupuncture for rabbits at LI10, LI4, ST36, and LI14 can evoke the caudate nucleus and increase the pain threshold^[23]. Using fMRI to observe the brain function image of the patients after needling at LI4 on the right side, these authors found that the signal intensity of many functional areas of the brain can be changed by needling. It was also observed that the signal intensity of the gray matter and lenticular nucleus on the left side was enhanced, and the signal intensity of the occipital lobe and anterior cingulate gyrus on both sides was weakened^[24]. Acupuncture alone and acupuncture combined with NJF for HSP can achieve appreciated clinical efficacy.

NJF is a movement pattern that is diagonal and spiral in nature and applies the concave and convex rule in joint mobilization. It is a new type of exercise therapy that can achieve therapeutic effects on joints, muscles, and nerves^[25,26]. Joint pain and limited activity, from the perspective of anatomy, generally involve internal or external causes of joint capsule. The internal capsule includes joint capsule, ligament, bone, and articular cartilage. The external joint capsule mainly includes skin, muscle, tendon, and nerve factors, and hemiplegia and shoulder pain are just a kind of disorders caused by changes in joint structure stability and tissue. Shoulder joint is a triaxial joint. Different from head and the joint iliac crest, it can perform multi-directional movement such as flexion, extension stretching, and rotation on three mutually perpendicular axes. In terms of the concave and convex rule, the axis for the rotation at the shoulder joint is on the humerus, and the movement of the articular surface is opposite to the direction of bone movement. In the normal mode of motion, patients with hemiplegia will exhibit flexor-based mode motion for their upper limbs, mainly manifested as retraction and pull-up of scapular band, extension, abduction, and rotation of the shoulder joint, flexion of the elbow joint, forearm rotation, wrist flexion, and finger flexion and adduction. Generally, shoulder pain makes the patient unable to lift the upper limb. Long-term pain can affect the patient's psychology. Loss of scapular rhythm and inadequate external rotation of the humerus are the major mechanism for shoulder pain^[27]. NJF can assist and resist the movement of the articular surface, and add spiral diagonal motion, which can reduce joint impact and pain, correct force line, and improve motion ability. The muscle shuttle can reflexively contract by extension. Traction can increase joint clearance and increase mobility^[28]. Extrusion can make the joint gap smaller and increase joint stability. Active resistance motion can promote the excitability of proprioceptors and promote the production of muscle, muscle strength, and tendon tension^[29,30]. This study found that acupuncture combined with NJF has clinical efficacy in the treatment of limb movement disorders and joint pain in patients with hemiplegia.

Table 2 Comparison of scores between the two groups

Group	VAS score		FMA score		BI score	
	Before treatment	After treatment	Before treatment	After treatment	Before treatment	After treatment
Treatment	5.52 ± 2.02	3.62 ± 1.52	18.05 ± 6.70	23.30 ± 7.03	49.00 ± 21.44	54.75 ± 22.39
Control	5.45 ± 1.99	4.71 ± 1.76	17.45 ± 4.21	19.40 ± 4.06	50.00 ± 23.90	54.00 ± 23.15

VAS: Visual analogue scale; FMA: Fugl-Meyer assessment; BI: Barthel index.

Table 3 Passive range of motion scores in the treatment group vs the control group

	PROM (flexion)	PROM (stretch)	PROM (outreach)	PROM (pronation)	PROM (external rotation)
Treatment group					
Before treatment	111.96 ± 17.44	21.99 ± 6.83	114.58 ± 16.75	29.80 ± 6.27	30.40 ± 6.83
After treatment	135.98 ± 16.61	32.42 ± 6.28	136.32 ± 17.14	44.88 ± 8.53	43.60 ± 9.87
Control group					
Before treatment	113.85 ± 15.30	22.25 ± 6.20	112.65 ± 13.43	27.26 ± 6.42	27.71 ± 5.79
After treatment	125.66 ± 12.55	27.87 ± 6.91	125.88 ± 14.22	39.18 ± 7.67	37.61 ± 8.09

PROM: Passive range of motion.

ARTICLE HIGHLIGHTS

Research background

Hemiplegic shoulder pain (HSP), a common sequela of stroke, causes damage to patients' movements and quality of life. Both traditional acupuncture and neuromuscular joint facilitation (NJF) help alleviate shoulder pain and inflammation associated with bursitis and/or tendonitis. However, the combination of the two therapies was seldom applied in the clinical practice.

Research motivation

The present study combined acupuncture with NJF technology to treat HSP, with an aim to provide a new treatment choice for this disease.

Research objectives

To study whether the efficacy of acupuncture combined with NJF is superior to acupuncture alone.

Research methods

Forty patients with HSP were randomly divided into a treatment group and a control group. The treatment group was treated with acupuncture combined with NJF and the control group was treated with acupuncture alone. In addition, patients in the treatment group were also trained to do exercises including articular surface movement and extension-abduction-internal rotation. The efficacy of the therapies was assessed by using the visual analogue scale (VAS), Fugl-Meyer assessment (FMA), Barthel index (BI), and passive range of motion (PROM) before and after the training.

Research results

After the treatment, the scores of VAS, FMA, and PROM were better in patients receiving neuromuscular joint facilitation combined with acupuncture than in patients only receiving acupuncture. However, there was no significant difference in BI scores between the two groups after the treatment.

Research conclusions

Both acupuncture alone and acupuncture combined with NJF in the treatment of HSP are effective, and can improve the clinical symptoms of patients. Acupuncture combined with NJF can improve the upper limb motor function, relieve pain, and increase joint mobility in patients with HSP. The combination therapy is better than acupuncture alone. However, there is no significant difference in improving the score of patients' self-care ability.

Research perspectives

Multidisciplinary treatment plan should be further studied in patients with HSP.

REFERENCES

- 1 **He J**, Gu D, Wu X, Reynolds K, Duan X, Yao C, Wang J, Chen CS, Chen J, Wildman RP, Klag MJ, Whelton PK. Major causes of death among men and women in China. *N Engl J Med* 2005; **353**: 1124-1134 [PMID: [16162883](#) DOI: [10.1056/NEJMsa050467](#)]
- 2 **Li Z**. The Effect of SES and Hypertension Control on the Incidence and Mortality of Stroke in the Elderly. M.Sc. Thesis, Anhui Medical University. 2013
- 3 **Zhao T**. Modern Hemiplegia Treatment. Beijing: People's Military Medical Press 1996; 190-265
- 4 **Harrison RA**, Field TS. Post stroke pain: identification, assessment, and therapy. *Cerebrovasc Dis* 2015; **39**: 190-201 [PMID: [25766121](#) DOI: [10.1159/000375397](#)]
- 5 **Snels IAK**, Beckerman H, Lankhorst GJ, Bouter LM. Risk factors for hemiplegic shoulder pain: A systematic review. *Crit Rev Phys Rehabil Med* 2002; **14**: 223-233
- 6 **Zheng T**, Huo M, Maruyama H, Kurosawa K, Hiiragi Y, Huang Q, Li D, Zhou B, Yin L, Wang H. Effects on ROM and joint position sense of the neck of two different interventions. *J Phys Ther Sci* 2015; **27**: 1041-1043 [PMID: [25995551](#) DOI: [10.1589/jpts.27.1041](#)]
- 7 **Huang Q**, Li D, Zhang J, Yang D, Huo M, Maruyama H. Comparison of the Efficacy of Different Long-term Interventions on Chronic Low Back Pain Using the Cross-sectional Area of the Multifidus Muscle and the Thickness of the Transversus Abdominis Muscle as Evaluation Indicators. *J Phys Ther Sci* 2014; **26**: 1851-1854 [PMID: [25540480](#) DOI: [10.1589/jpts.26.1851](#)]
- 8 **Huo M**, Wang H, Ge M, Huang Q, Li D, Maruyama H. The Immediate Effect of Neuromuscular Joint Facilitation (NJF) Treatment on Electromechanical Reaction Times of Hip Flexion. *J Phys Ther Sci* 2013; **25**: 1463-1465 [PMID: [24396211](#) DOI: [10.1589/jpts.25.1463](#)]
- 9 **Yin LQ**, Yin K, Hao XB. The treatment of 30 cases of frozen shoulder with neuromuscular joint promotion method. *Zhongguo Laonianxue Zazhi* 2010; **30**: 2366-2367
- 10 **State Administration of Traditional Chinese Medicine**. Encephalopathy Emergency Collaboration Group. Diagnostic and Efficacy Scoring Criteria for Stroke. *Beijing Zhongyiyao Daxue Xuebao* 1996; **16**: 55-56
- 11 **Chinese Medical Association Neurology Branch**. Chinese Medical Association Neurology Branch Cerebrovascular Disease Group. Guidelines for the Diagnosis and Treatment of Acute Ischemic Stroke in China 2014. *Zhonghua Shenjingke Zazhi* 2015; **48**: 246-257
- 12 **Chinese Medical Association**. Clinical Technical Practices - Physical Medicine and Rehabilitation. Beijing: People's Military Medical Press 2004; 12-58
- 13 **Zhao JX**, Zheng WY. Acupuncture Treatment of 30 Cases of Hemiplegia and Shoulder Pain after Stroke. *Zhongguo Minjian Liaofo* 2012; **18**: 13
- 14 **Tong YD**, Wang GX, Wei D, Xie YQ, Wu YL, Lai M. Preliminary Study on the Influence of Acupuncture and Moxibustion on Individual Muscle Mechanical Properties. *Shenyang Tiyu Xueyuan Xuebao* 1996; **4**: 9-10
- 15 **Duan CP**, Lu DH, Fu XQ. Effects of Acupuncture and Static Stretch on the Ultrastructure of Skeletal Muscle during Delayed Soreness. *Beijing Tiyu Daxue Xuebao* 1984; **4**: 12-23
- 16 **Qu ZQ**, Lu DH, Yi R. Effects of Acupuncture on Intracellular Calcium Distribution during Skeletal Muscle Injury and its Mechanism. *Zhongguo Yundong Yixue Zazhi* 1995; **4**: 7-12
- 17 **Xu YM**, Wang RY. Effects of Acupuncture and Magnetic Extraction on Resting Potential Changes of Skeletal Muscle after Long-term Contraction. *Tiyu Kexue* 1999; **1**: 75-76
- 18 **Li JH**, Wang ZZ, Tang YF. Effect of Acupuncture on Muscle Regeneration after Skeletal Muscle Contusion in Rats. *Beijing Tiyu Daxue Xuebao* 2012; **6**: 61-65
- 19 **Qin CJ**, Wang RY. Changes of Myosin after Skeletal Muscle Stimulation by Electric Stimulation and the Influence of Acupuncture on it. *Guangzhou Tiyu Xueyuan Xuebao* 2001; **1**: 50-53
- 20 **Bai SC**. Effect of Acupuncture on Inflammation of Bone and Muscle after an Eccentric Exercise. M.Sc. Thesis, Beijing Sport University. 2015
- 21 **Guo G**. Experimental Acupuncture and Moxibustion. Beijing: China Traditional Chinese Medicine Press 2008; 263-275
- 22 **Guo CQ**. Modern Research and Application of Acupuncture and Moxibustion. Beijing: Xueyuan Publishing House 1998; 205
- 23 **Wang MJ**, Zhu J, Yan YQ. Chinese Acupuncture and Moxibustion Book. Zhengzhou: Henan Science and Technology Press 2012; 158
- 24 **Hou JW**, Huang WW, Wang Q. MRI Study of Brain Function in Acupuncture Analgesia. *Zhonghua Fangshexue Zazhi* 2002; **36**: 206-210
- 25 **Wu P**, Huo M, Maruyama H. Effects of neuromuscular joint facilitation on baseball pitching velocity and electromechanical reaction times of the teres major of young amateur baseball players. *J Phys Ther Sci* 2013; **25**: 1459-1461 [PMID: [24396210](#) DOI: [10.1589/jpts.25.1459](#)]
- 26 **Huang Q**, Li D, Yokotsuka N, Zhang Y, Ubukata H, Huo M, Maruyama H. The Intervention Effects of Different Treatment for Chronic Low Back Pain as Assessed by the Cross-sectional Area of the Multifidus Muscle. *J Phys Ther Sci* 2013; **25**: 811-813 [PMID: [24259859](#) DOI: [10.1589/jpts.25.811](#)]
- 27 **Wang G**, Liu Q. Treatment of Neurological Rehabilitation. Beijing: People's Medical Publishing House 2015; 458-459
- 28 **Onoda K**, Huo M, Maruyama H. The immediate effect of neuromuscular joint facilitation (NJF) treatment on the standing balance in younger persons. *J Phys Ther Sci* 2015; **27**: 1481-1483 [PMID: [26157245](#) DOI: [10.1589/jpts.27.1481](#)]
- 29 **Wang H**, Huo M, Huang Q, Li D, Maruyama H. The Immediate Effect of Neuromuscular Joint Facilitation (NJF) Treatment on Hip Muscle Strength. *J Phys Ther Sci* 2013; **25**: 1455-1457 [PMID: [24396209](#) DOI: [10.1589/jpts.25.1455](#)]
- 30 **Huo M**, Maruyama H, Kaneko T, Naito D, Koiso Y. The immediate effect of lumbar spine patterns of neuromuscular joint facilitation in young amateur baseball players. *J Phys Ther Sci* 2013; **25**: 1523-1524 [PMID: [24409011](#) DOI: [10.1589/jpts.25.1523](#)]



Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-2238242
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

