

World Journal of *Clinical Cases*

World J Clin Cases 2020 October 26; 8(20): 4688-5069



MINIREVIEWS

- 4688 Relationship between non-alcoholic fatty liver disease and coronary heart disease
Arslan U, Yenercağ M

ORIGINAL ARTICLE

Retrospective Cohort Study

- 4700 Remission of hepatotoxicity in chronic pulmonary aspergillosis patients after lowering trough concentration of voriconazole
Teng GJ, Bai XR, Zhang L, Liu HJ, Nie XH

Retrospective Study

- 4708 Endoscopic submucosal dissection as alternative to surgery for complicated gastric heterotopic pancreas
Noh JH, Kim DH, Kim SW, Park YS, Na HK, Ahn JY, Jung KW, Lee JH, Choi KD, Song HJ, Lee GH, Jung HY
- 4719 Observation of the effects of three methods for reducing perineal swelling in children with developmental hip dislocation
Wang L, Wang N, He M, Liu H, Wang XQ
- 4726 Predictive value of serum cystatin C for risk of mortality in severe and critically ill patients with COVID-19
Li Y, Yang S, Peng D, Zhu HM, Li BY, Yang X, Sun XL, Zhang M
- 4735 Sleep quality of patients with postoperative glioma at home
Huang Y, Jiang ZJ, Deng J, Qi YJ
- 4743 Early complications of preoperative external traction fixation in the staged treatment of tibial fractures: A series of 402 cases
Yang JZ, Zhu WB, Li LB, Dong QR
- 4753 Retroperitoneal vs transperitoneal laparoscopic lithotripsy of 20-40 mm renal stones within horseshoe kidneys
Chen X, Wang Y, Gao L, Song J, Wang JY, Wang DD, Ma JX, Zhang ZQ, Bi LK, Xie DD, Yu DX
- 4763 Undifferentiated embryonal sarcoma of the liver: Clinical characteristics and outcomes
Zhang C, Jia CJ, Xu C, Sheng QJ, Dou XG, Ding Y
- 4773 Cerebral infarct secondary to traumatic internal carotid artery dissection
Wang GM, Xue H, Guo ZJ, Yu JL
- 4785 Home-based nursing for improvement of quality of life and depression in patients with postpartum depression
Zhuang CY, Lin SY, Cheng CJ, Chen XJ, Shi HL, Sun H, Zhang HY, Fu MA

Observational Study

- 4793** Cost-effectiveness of lutetium (^{177}Lu) oxodotreotide *vs* everolimus in gastroenteropancreatic neuroendocrine tumors in Norway and Sweden
Palmer J, Leeuwenkamp OR
- 4807** Factors related to improved American Spinal Injury Association grade of acute traumatic spinal cord injury
Tian C, Lv Y, Li S, Wang DD, Bai Y, Zhou F, Ma QB
- 4816** Intraoperative systemic vascular resistance is associated with postoperative nausea and vomiting after laparoscopic hysterectomy
Qu MD, Zhang MY, Wang GM, Wang Z, Wang X

META-ANALYSIS

- 4826** Underwater *vs* conventional endoscopic mucosal resection in treatment of colorectal polyps: A meta-analysis
Ni DQ, Lu YP, Liu XQ, Gao LY, Huang X

CASE REPORT

- 4838** Dehydrated patient without clinically evident cause: A case report
Palladino F, Fedele MC, Casertano M, Liguori L, Esposito T, Guarino S, Miraglia del Giudice E, Marzuillo P
- 4844** Intracranial malignant solitary fibrous tumor metastasized to the chest wall: A case report and review of literature
Usuda D, Yamada S, Izumida T, Sangen R, Higashikawa T, Nakagawa K, Iguchi M, Kasamaki Y
- 4853** End-of-life home care of an interstitial pneumonia patient supported by high-flow nasal cannula therapy: A case report
Goda K, Kenzaka T, Kuriyama K, Hoshijima M, Akita H
- 4858** Rupture of carotid artery pseudoaneurysm in the modern era of definitive chemoradiation for head and neck cancer: Two case reports
Kim M, Hong JH, Park SK, Kim SJ, Lee JH, Byun J, Ko YH
- 4866** Unremitting diarrhoea in a girl diagnosed anti-N-methyl-D-aspartate-receptor encephalitis: A case report
Onpoaree N, Veeravigrom M, Sanpavat A, Suratannon N, Sintusek P
- 4876** Paliperidone palmitate-induced facial angioedema: A case report
Srifueungfung M, Sukakul T, Liangcheep C, Viravan N
- 4883** Improvement of lenvatinib-induced nephrotic syndrome after adaptation to sorafenib in thyroid cancer: A case report
Yang CH, Chen KT, Lin YS, Hsu CY, Ou YC, Tung MC
- 4895** Adult metaplastic hutch diverticulum with robotic-assisted diverticulectomy and reconstruction: A case report
Yang CH, Lin YS, Ou YC, Weng WC, Huang LH, Lu CH, Hsu CY, Tung MC

- 4902** Thrombus straddling a patent foramen ovale and pulmonary embolism: A case report
Huang YX, Chen Y, Cao Y, Qiu YG, Zheng JY, Li TC
- 4908** Therapeutic experience of an 89-year-old high-risk patient with incarcerated cholecystolithiasis: A case report and literature review
Zhang ZM, Zhang C, Liu Z, Liu LM, Zhu MW, Zhao Y, Wan BJ, Deng H, Yang HY, Liao JH, Zhu HY, Wen X, Liu LL, Wang M, Ma XT, Zhang MM, Liu JJ, Liu TT, Huang NN, Yuan PY, Gao YJ, Zhao J, Guo XA, Liao F, Li FY, Wang XT, Yuan RJ, Wu F
- 4917** Woven coronary artery: A case report
Wei W, Zhang Q, Gao LM
- 4922** Idiopathic multicentric Castleman disease with pulmonary and cutaneous lesions treated with tocilizumab: A case report
Han PY, Chi HH, Su YT
- 4930** Perianorectal abscesses and fistula due to ingested jujube pit in infant: Two case reports
Liu YH, Lv ZB, Liu JB, Sheng QF
- 4938** Forniceal deep brain stimulation in severe Alzheimer's disease: A case report
Lin W, Bao WQ, Ge JJ, Yang LK, Ling ZP, Xu X, Jiang JH, Zuo CT, Wang YH
- 4946** Systemic autoimmune abnormalities complicated by cytomegalovirus-induced hemophagocytic lymphohistiocytosis: A case report
Miao SX, Wu ZQ, Xu HG
- 4953** Nasal mucosa pyoderma vegetans associated with ulcerative colitis: A case report
Yu SX, Cheng XK, Li B, Hao JH
- 4958** Amiodarone-induced hepatotoxicity — quantitative measurement of iodine density in the liver using dual-energy computed tomography: Three case reports
Lv HJ, Zhao HW
- 4966** Multisystem involvement Langerhans cell histiocytosis in an adult: A case report
Wang BB, Ye JR, Li YL, Jin Y, Chen ZW, Li JM, Li YP
- 4975** New mutation in EPCAM for congenital tufting enteropathy: A case report
Zhou YQ, Wu GS, Kong YM, Zhang XY, Wang CL
- 4981** Catastrophic vertebral artery and subclavian artery pseudoaneurysms caused by a fishbone: A case report
Huang W, Zhang GQ, Wu JJ, Li B, Han SG, Chao M, Jin K
- 4986** Anastomosing hemangioma arising from the left renal vein: A case report
Zheng LP, Shen WA, Wang CH, Hu CD, Chen XJ, Shen YY, Wang J
- 4993** Bladder perforation caused by long-term catheterization misdiagnosed as digestive tract perforation: A case report
Wu B, Wang J, Chen XJ, Zhou ZC, Zhu MY, Shen YY, Zhong ZX

- 4999** Primary pulmonary plasmacytoma accompanied by overlap syndrome: A case report and review of the literature
Zhou Y, Wang XH, Meng SS, Wang HC, Li YX, Xu R, Lin XH
- 5007** Gastrointestinal stromal tumor metastasis at the site of a totally implantable venous access port insertion: A rare case report
Yin XN, Yin Y, Wang J, Shen CY, Chen X, Zhao Z, Cai ZL, Zhang B
- 5013** Massive gastrointestinal bleeding caused by a Dieulafoy's lesion in a duodenal diverticulum: A case report
He ZW, Zhong L, Xu H, Shi H, Wang YM, Liu XC
- 5019** Plastic bronchitis associated with *Botrytis cinerea* infection in a child: A case report
Liu YR, Ai T
- 5025** Chest, pericardium, abdomen, and thigh penetrating injury by a steel rebar: A case report
Yang XW, Wang WT
- 5030** Monocular posterior scleritis presenting as acute conjunctivitis: A case report
Li YZ, Qin XH, Lu JM, Wang YP
- 5036** Choriocarcinoma with lumbar muscle metastases: A case report
Pang L, Ma XX
- 5042** Primary chondrosarcoma of the liver: A case report
Liu ZY, Jin XM, Yan GH, Jin GY
- 5049** Successful management of a tooth with endodontic-periodontal lesion: A case report
Alshawwa H, Wang JF, Liu M, Sun SF
- 5057** Rare imaging findings of hypersensitivity pneumonitis: A case report
Wang HJ, Chen XJ, Fan LX, Qi QL, Chen QZ
- 5062** Effective administration of cranial drilling therapy in the treatment of fourth degree temporal, facial and upper limb burns at high altitude: A case report
Shen CM, Li Y, Liu Z, Qi YZ

ABOUT COVER

Peer-reviewer of *World Journal of Clinical Cases*, Dr. Aleem Ahmed Khan is a Distinguished Scientist and Head of The Central Laboratory for Stem Cell Research and Translational Medicine, Centre for Liver Research and Diagnostics, Deccan College of Medical Sciences, Kanchanbagh, Hyderabad (India). Dr. Aleem completed his Doctorate from Osmania University, Hyderabad in 1998 and has since performed pioneering work in the treatment of acute liver failure and decompensated cirrhosis using hepatic stem cell transplantation. During his extensive research career he supervised 10 PhD students and published > 150 research articles, 7 book chapters, and 2 patents. His ongoing research involves developing innovative technologies for organ regeneration and management of advanced cancers. (L-Editor: Filipodia)

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, PubMed, and PubMed Central. The 2020 Edition of Journal Citation Reports® cites the 2019 impact factor (IF) for *WJCC* as 1.013; IF without journal self cites: 0.991; Ranking: 120 among 165 journals in medicine, general and internal; and Quartile category: Q3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Ji-Hong Liu; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lai Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Semimonthly

EDITORS-IN-CHIEF

Dennis A Bloomfield, Sandro Vento, Bao-Gan Peng

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

October 26, 2020

COPYRIGHT

© 2020 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Retrospective Study

Predictive value of serum cystatin C for risk of mortality in severe and critically ill patients with COVID-19

Yan Li, Shuang Yang, Ding Peng, Hong-Ming Zhu, Bang-Yi Li, Xiaojiao Yang, Xue-Lian Sun, Mei Zhang

ORCID number: Yan Li 0000-0002-2340-1937; Shuang Yang 0000-0003-0652-9266; Ding Peng 0000-0003-3614-5538; Hong-Ming Zhu 0000-0002-8596-177x; Bang-Yi Li 0000-0002-5944-9641; Xiaojiao Yang 0000-0003-3667-1239; Xue-Lian Sun 0000-0003-3372-0410; Mei Zhang 0000-0002-8596-1868.

Author contributions: Li Y wrote the paper; Zhang M designed and supervised the research; Zhu HM and Yang S performed the research; Peng D and Yang X performed data and statistical analyses; Peng D and Li BY analyzed the data; Sun XL collected the data; all authors approved the final version of the article.

Institutional review board

statement: This survey is a retrospective study, only collecting the clinical data of patients. Since it will not bring risks to patients' physiology and do not interfere with patients' treatment plan, and researchers will protect patients' information from disclosure, Xuanwu Hospital of Capital Medical University agreed to exempt this study from ethical review.

Informed consent statement:

Patients were not required to give informed consent to the study

Yan Li, Department of Pulmonology, Xuanwu Hospital of Capital Medical University, Beijing 100053, China

Shuang Yang, Ding Peng, Hong-Ming Zhu, Bang-Yi Li, Mei Zhang, Department of Gastroenterology, Xuanwu Hospital of Capital Medical University, Beijing 100053, China

Xiaojiao Yang, School of Human Nutrition, Faculty of Agricultural and Environmental Sciences, McGill University, Montreal H9X 3V9, Quebec, Canada

Xue-Lian Sun, Department of Emergency Medicine, Xuanwu Hospital of Capital Medical University, Beijing 100053, China

Corresponding author: Mei Zhang, MD, Chief Physician, Professor, Department of Gastroenterology, Xuanwu Hospital of Capital Medical University, No. 45 Changchun Street, Xicheng District, Beijing 100053, China. zhang2955@sina.com

Abstract

BACKGROUND

The outbreak of coronavirus disease 2019 (COVID-19) has rapidly evolved into a global pandemic. COVID-19 is clinically categorized into mild, moderate, severe, and critical illness. Acute kidney injury is an independent risk factor for poor prognosis in patients with. Serum cystatin C (sCys C) is considered a more sensitive biomarker for early renal insufficiency than conventional indicators of renal function. Early detection of risk factors that affect the prognosis of severe and critically ill patients while using active and effective treatment measures is very important and can effectively reduce the potential mortality rate.

AIM

To determine the predictive value of sCys C for the prognosis of patients with COVID-19.

METHODS

The clinical data of 101 severe and critically ill patients with COVID-19 at a designated hospital in Wuhan, Hubei Province, China were analyzed retrospectively. According to the clinical outcome, the patients were divided into a discharge group (64 cases) and a death group (37 cases). The general information, underlying diseases, and laboratory examination indexes of the two groups were compared. Multivariate Cox regression was used to explore the

because the analysis used anonymous clinical data.

Conflict-of-interest statement: The authors declare no conflict of interest regarding the manuscript.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Received: July 4, 2020

Peer-review started: July 4, 2020

First decision: July 24, 2020

Revised: August 5, 2020

Accepted: September 3, 2020

Article in press: September 3, 2020

Published online: October 26, 2020

P-Reviewer: Yeh YC

S-Editor: Wang JL

L-Editor: Wang TQ

P-Editor: Wang LL



relationship between sCys C and prognosis. The receiver operating characteristic (ROC) curve was used to demonstrate the sensitivity and specificity of sCys C and its optimal cut-off value for predicting death.

RESULTS

There were significant differences in age, sCys C, creatinine, C-reactive protein, serum albumin, creatine kinase-MB, alkaline phosphatase, lactate dehydrogenase, neutrophil count, and lymphocyte count between the two groups ($P < 0.001$). Multivariate logistic regression analysis showed that sCys C was an independent risk factor for death in patients with COVID-19 (Odds ratio = 1.812, 95% confidence interval [CI]: 1.300-2.527, $P < 0.001$). The area under the ROC curve was 0.755 (95% CI: 1.300-2.527), the cut-off value was 0.80, the specificity was 0.562, and the sensitivity was 0.865.

CONCLUSION

sCys C is an independent risk factor for death in patients with COVID-19. Patients with a sCys C level of 0.80 mg/L or greater are at a high risk of death.

Key Words: COVID-19; Cystatin C; Acute kidney injury; Renal function; Coronavirus infections; Multiple organ dysfunction syndrome

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The novel coronavirus epidemic has become a major public health event in the world. The mortality rate of severe and critical patients is high. It is very important to find an available, cheap, and sensitive biomarker to predict mortality. Cystatin C is a predictor of renal injury. We compared 101 severe and critical patients with coronavirus disease 2019 (COVID-19) in Wuhan, China, and found that patients with a cystatin C level of 0.80 mg/L or greater were at a high risk of death. Therefore, cystatin C may have an important role in the prognosis of severe and critical patients with COVID-19.

Citation: Li Y, Yang S, Peng D, Zhu HM, Li BY, Yang X, Sun XL, Zhang M. Predictive value of serum cystatin C for risk of mortality in severe and critically ill patients with COVID-19. *World J Clin Cases* 2020; 8(20): 4726-4734

URL: <https://www.wjgnet.com/2307-8960/full/v8/i20/4726.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v8.i20.4726>

INTRODUCTION

The outbreak of coronavirus disease 2019 (COVID-19) has rapidly evolved into a global pandemic. As of June 17, 2020, more than 8 million people have been infected worldwide. The World Health Organization has reported more than 440000 deaths and 130000 new confirmed cases daily. Most patients with COVID-19 have mild symptoms, but approximately 5%-20%^[1] have severe symptoms, including acute respiratory distress syndrome, septic shock, and multiple organ failure. The mortality rate of patients with severe COVID-19 was higher than that of patients with mild COVID-19. The later the diagnosis is made, the higher the risk of death.

In addition to the impacts of COVID-19 on the respiratory system and immune system, the kidney is also one of the main organs being affected. Whether COVID-19 patients will develop acute kidney injury (AKI), which may affect their prognosis, is an important issue and worthy of clinical attention. Some studies have shown that AKI is an independent risk factor for poor prognosis in patients with COVID-19^[2]. In addition to the direct toxicity of coronaviruses, factors attributed to AKI include systemic hypoxia, coagulation abnormalities, and rhabdomyolysis which may be related to drugs or hyperventilation^[3]. Early detection of AKI in patients with COVID-19 is conducive to establishing standardized AKI cluster prevention and treatment (5R principle): Risk screening, early recognition, timely response, renal replacement therapy, and renal recovery, which are key to improve the cure rate and reduce the

mortality rate of patients with COVID-19.

Serum cystatin C (sCys C) is a newly developed index with a high sensitivity and specificity for evaluating renal function, which is not affected by age, sex, weight, inflammation, and other factors. It is considered an endogenous marker superior to serum creatinine (sCr)^[4]. sCys C is a cysteine protease inhibitor, also known as γ -trace basic protein or post-gamma globulin, which is encoded by the *Cys C* gene that can be continuously transcribed and expressed in all nucleated cells at a constant speed. It has no tissue specificity with a small molecular weight (13 kDa), being positively charged under physiological conditions. It is freely filtered in the glomeruli, completely reabsorbed by the renal tubular epithelial cells, and degraded within cells without returning to the blood. Renal tubular epithelial cells do not secrete sCys C into the lumen, therefore its serum concentration is mainly determined by glomerular filtration rate which is an important indicator of glomerular filtration^[5]. In addition, it has been reported that sCys C can be used to evaluate the survival rate of patients with sepsis^[6].

The purpose of this study was to investigate the effect of COVID-19 on renal function and assess the predictive value of sCys C for risk of mortality in critically ill patients with COVID-19.

MATERIALS AND METHODS

Study design and patient population

This single-center retrospective study included adults diagnosed with COVID-19 admitted to the ward of a designated hospital in Wuhan, Hubei Province, from January to March 2020 without any history of chronic kidney disease. The diagnostic criteria were based on the COVID-19 Diagnosis and Treatment Protocol (trial version 7) formulated by the National Health Commission of China, with the epidemiological history and clinical expression of COVID-19, and the etiological or serological evidence (positive novel coronavirus nucleic acid test or serum specific antibody test) considered. Patients recruited in this study were all severe or critically ill patients. Severe condition was determined when subjects had any of the following conditions: (1) Shortness of breath with a respiration rate ≥ 30 times/min; (2) Oxygen saturation $\leq 93\%$ in the resting state; and (3) $\text{PaO}_2/\text{FiO}_2 \leq 300$ mmHg. Subjects were considered critical if any of the following conditions arose: (1) Respiratory failure and need for mechanical ventilation; (2) Shock; and (3) Patients with other organ failure that should be monitored in the intensive care unit (ICU).

This survey was a retrospective study collecting only the clinical data of patients. Since it did not bring risks to patients' physiology and did not interfere with patients' treatment plan, and researchers protected patients' information from disclosure, Xuanwu Hospital of Capital Medical University agreed to exempt this study from ethical review.

Data collection

All subjects enrolled in our study had a case report form, and data were collected within 12 h after hospital admission, including sex, age, medical history, complications, sCys C, creatine kinase-MB (CK-MB), serum creatinine (sCr), aspartate aminotransferase (AST), alanine aminotransferase (ALT), gamma-glutamyl transpeptidase (GGTP), alkaline phosphatase (ALP), lactate dehydrogenase (LDH), total bilirubin, C-reactive protein (CRP), neutrophil count, albumin, uric acid, lymphocyte count, hemoglobin, platelet count, and blood calcium. All data were obtained from the first laboratory examination after admission. Discharge or death was considered the study outcome.

Statistical analysis

All data were analyzed using SPSS version 22.0. Continuous data (sCr, AST, ALT, *etc.*) with a normal distribution are presented as the mean \pm SD. Non-normally distributed variables are presented as medians with interquartile ranges and analyzed using a non-parametric test. Categorical data (sex, medical history, *etc.*) were analyzed by the χ^2 test. The independent effect of sCys C was calculated using Cox forward stepwise multivariate proportional hazards regression analysis, incorporating classic risk factors. All tests were two-sided, and a *P* value of < 0.05 was considered statistically significant. The receiver operating characteristic (ROC) curve was used to demonstrate the sensitivity and specificity of sCys C and its optimal cut-off value for predicting death.

RESULTS

There were a total of 101 cases including 18 critically ill patients and 83 severe patients at baseline, and they were divided into two groups: Discharge group (64 cases) and death group (37 cases). All data were obtained from the first laboratory examination after admission within 12 h after the diagnosis of severe and critical COVID-19. The median length of hospital stay was 15 d. Comparison of the baseline data between the two groups showed that the group of patients who died had the following characteristics: Advanced age ($P < 0.001$), higher levels of sCys C ($P < 0.001$), CK-MB ($P < 0.001$), sCr ($P = 0.001$), AST ($P < 0.001$), ALT ($P < 0.05$), GGTP ($P < 0.05$), ALP ($P < 0.05$), LDH ($P < 0.001$), total bilirubin ($P < 0.05$), CRP ($P < 0.001$), and neutrophil count ($P < 0.05$), and lower levels of albumin ($P < 0.001$), calcium ion concentration ($P < 0.001$), lymphocyte count ($P < 0.001$), and platelet count ($P < 0.05$) (Table 1).

Predictive value of renal function

By using sCys C, sCr, and the differences in factors as explanatory variables in forward Cox regression analysis to explore the independent factors (Table 2), sCys C was found to be an independent risk factor for death, but the sCr level was not. The ROC curve was used to determine the optimal threshold for predicting the death of severe and critical COVID-19 patients. The results are shown in Figure 1. The area under the curve, optimal threshold, specificity, and sensitivity were 0.755, 0.80, 0.562, and 0.865, respectively. Patients with a sCys C level of 0.80 mg/L or greater were at a high risk of death. The patients were divided into two groups based on a cut-off of 0.80 mg/L, and the survival curve was obtained, as shown in Figure 2. The results demonstrated that this threshold was effective in differentiating high-risk groups (log-rank = 13.375, $P < 0.001$).

DISCUSSION

From January to March 2020, the clinical data of 101 severe and critically ill patients in a designated hospital in Wuhan, Hubei Province, China were analyzed retrospectively, including 37 dead patients (death group) and 64 surviving patients (discharge group). The risk factors for prognosis were assessed by comparing the laboratory examination results of the discharge and death groups. Multivariate analysis showed that sCys C, CK-MB, AST, albumin, and serum calcium were independent risk factors for death in severe and critically ill patients. The predictive value of CK-MB in patients with COVID-19 has been confirmed by many studies. Wang *et al*^[7] showed that serum cardiac troponin (cTnI) and CK-MB levels in ICU patients were significantly higher than those in non-ICU patients ($P = 0.004$), suggesting that increased CK-MB and cTnI levels represent a poor prognosis in patients with COVID-19. Zhou *et al*^[1] also illustrated that an increase in CK-MB and cTnI levels in COVID-19 patients was significantly correlated with hospital death ($P = 0.043$). AST and albumin are indicators for liver function, and previous literature supported the predictive value of abnormal liver function in the prognosis of COVID-19 patients. A meta-analysis showed that abnormal liver function indicators such as AST, ALT, bilirubin, and albumin can predict the severity and prognosis of COVID-19 patients^[8]. Lei *et al*^[9] included 5771 adult COVID-19 patients from Hubei Province, and retrospectively analyzed the time distribution of liver function indicators in patients and correlated them with death. The results indicated that abnormal liver function, especially elevated AST, was closely related to the risk of death, which was consistent with the results of this study^[9]. In addition, many studies have shown that a decrease in serum calcium is a risk factor for disease severity and death in patients with COVID-19^[10,11].

Presently, the predictive value of sCys C in the prognosis of patients with COVID-19 is rarely reported, but studies have confirmed that sCys C can predict the risk of AKI and death in critically ill patients such as those with acute cerebral infarction, acute myocardial infarction, heart failure, and sepsis^[12,13]. This indicates that sCys C has a predictive value in the prognosis of critically ill patients. However, the predictive value of sCys C for severe and critically ill COVID-19 patients has not been confirmed. In this study, the sCys C level in the death group was significantly higher than that in the discharge group, suggesting that the sCys C level was related to the risk of death in COVID-19. Moreover, multivariate regression analysis showed that the sCys C level was an independent risk factor for the death of patients with COVID-19. To further clarify the correlation between sCys C value and prognosis of COVID-19, we used the ROC curve to assess the cut-off value of sCys C in predicting the death of patients. The

Table 1 Baseline clinical characteristics

Characteristic	Discharged	Dead	P value
Age, yr	54.09 ± 14.95	71.76 ± 10.012	0.000
Male sex, <i>n</i> (%)	30 (46.9)	23 (62.2)	0.138
Critically ill patients, <i>n</i> (%)	6 (9.4)	12 (32.4)	0.004
Hypertension, <i>n</i> (%)	26 (40.6)	20 (54.1)	0.192
Diabetes, <i>n</i> (%)	20 (31.3)	7 (18.9)	0.177
Combined with renal insufficiency, <i>n</i> (%)	9 (14.1)	3 (8.1)	0.373
Coronary heart disease, <i>n</i> (%)	5 (7.8)	7 (18.9)	0.097
sCys C, mg/L	0.8 (0.7, 0.9)	1 (0.9, 1.3)	0.000
CK-MB activity, U/L	12 (9, 14)	19(13, 35)	0.000
Creatinine, μmol/L	64.5 (57.5, 79.1)	78.2 (70, 94.2)	0.001
Aspartate aminotransferase, U/L	29 (19, 40)	44 (31, 73)	0.000
Alanine aminotransferase, U/L	29 (17, 52)	44 (23, 58)	0.028
Glutamyl transpeptidase, U/L	25 (17, 49)	37 (26, 86)	0.005
Alkaline phosphatase, U/L	47 (41, 58)	68 (52, 95)	0.000
Lactic acid dehydrogenase, U/L	228 (188, 283)	548 (399, 623)	0.000
Total bilirubin, μmol/L	9.6 (7.4, 13.5)	13.4 (8.5, 18.6)	0.024
Albumin, g/L	32.7 (30.4, 35.9)	28 (24.8, 29.4)	0.000
Uric acid, μmol/L	229.2 (176, 290.8)	233.8 (177.2, 366.7)	0.521
Calcium, mmol/L	1.97 (1.91, 2.04)	1.73 (1.03, 1.88)	0.000
C reactive protein, mg/L	11.1 (3.55,29.13)	92.29 (42.57, 116.45)	0.000
Neutrophils, G/L	3 (2.21, 4.27)	7.28 (5.63, 9.35)	0.000
Lymphocytes, G/L	1.02 (0.79, 1.39)	0.58 (0.42, 0.88)	0.000
Hemoglobin, g/L	128 (119, 139)	127 (105, 138)	0.540
Platelets, G/L	211 (159, 277)	147(109, 229)	0.011

sCys C: Serum cystatin C; CK-MB: Creatine kinase-MB.

Table 2 Independent predictors identified by univariate Cox regression analysis

	B	SE	Wald	P value	Exp(B)	95%CI
sCys C	0.595	0.17	12.279	0.000	1.812	1.300, 2.527
CK-MB	0.037	0.014	7.149	0.008	1.037	1.010, 1.066
AST	0.026	0.008	10.875	0.001	1.027	1.011, 1.043
Albumin	-0.221	0.041	28.531	0.000	0.802	0.739, 0.869
Calcium	-3.157	0.555	32.347	0.000	0.043	0.014, 0.126

sCys C: Serum cystatin C; CK-MB: Creatine kinase-MB; AST: Aspartate aminotransferase.

results showed that sCys C had an AUC of 0.755, sensitivity of 0.865, and specificity of 0.562, which indicated that sCys C had appreciated value in predicting the death of patients with severe and critical COVID-19. The cut-off value was 0.80 mg/L, that is, patients with sCys C ≥ 0.80 mg/L are at a high risk of death. The low specificity may be related to the low sample size and retrospective study. Clinicians should be vigilant when patients have sCys C ≥ 0.80 mg/L so that early intervention could be given to improve the prognosis.




Figure 1 Receiver operating characteristic curve of serum cystatin C for predicting the risk of mortality in severe and critically ill patients with coronavirus disease 2019. Area under the curve, optimal threshold, specificity, and sensitivity were 0.755, 0.80, 0.562, and 0.865, respectively.




Figure 2 Effect of serum cystatin C level on patient survival. The Kaplan-Meier survival curves for severe and critically ill coronavirus disease 2019 patients with a cut-off value of serum cystatin C at 0.80 mg/L are shown.

In this study, the sCr level of the dead patients was significantly higher than that of the survivors ($P < 0.01$). However, sCr was not an independent risk factor for death when other different factors were included in the Cox regression equation. Accordingly, sCys C was deemed superior to sCr in predicting the risk of death in severe and critically ill patients with COVID-19. sCys C is a member of the cystatin family. It is freely filtered by the glomerulus, reabsorbed by the proximal tubules, and completely catabolized without returning to the blood stream. Therefore, its

concentration in the blood is not affected by age, sex, or liver function^[14,15]. sCys C is generally considered an earlier and more sensitive marker of renal insufficiency than the conventional indicators of renal function such as sCr and blood urea nitrogen (BUN). Numerous studies have shown that there is no significant increase in sCr or BUN levels in the early stage of renal injury, and their elevations can only be observed in the middle and late stages of renal injury. Abnormal levels of indicators usually indicate that renal damage is serious and irreversible^[16]. Xiang *et al*^[17] showed that BUN, sCr, and sCys C (biochemical indicators of renal function) were significantly increased in patients with severe COVID-19, indicating that severe acute respiratory syndrome coronavirus 2 infection may damage the kidney. To date, a number of studies have revealed that patients with COVID-19 may have renal function damage, and the clinical manifestations include rise in sCr, hematuria, albuminuria, and AKI^[18,19]. It has been reported that the incidence of AKI in patients with COVID-19 ranged from 0.9% to 29%^[20-22]. For patients who required intensive care, the incidence of AKI increased significantly to 50%^[4]. Autopsy results indicated that patients with COVID-19 had acute proximal renal tubular injury that was characterized by loss of brush margin, vacuolar degeneration, dilation of the tubular lumen with cell fragments, necrosis, and epithelial exfoliation^[9]. Pei *et al*^[23] showed that the total mortality rate of patients with renal involvement was 11.2%, while the mortality rate of patients without renal involvement was 1.2%, suggesting that renal complications were related to the mortality rate of COVID-19. In addition, another study^[24] found that the levels of sCys C were positively correlated with inflammatory reaction indexes such as interleukin-1 β (IL-1 β), IL-6, tumor necrosis factor- α , and hsCRP, and promoted the occurrence and development of inflammatory reaction. Therefore, in the pathological state of COVID-19 invading lung tissue, sCys C is synthesized and released in large quantities, and the level of sCys C is increased, which regulates the cathepsin activity released from necrotic or inflammatory cells. It is suggested that clinicians should pay close attention to the sCys C level and its changes.

Some limitations pertaining to this study

Since the specificity of the prediction was relatively low, interpretation with caution in clinical practice is suggested. This was a single-center retrospective study and the sample size was relatively small and we lacked clinical data of surviving patients after discharge, thus we were unable to evaluate the effect of sCys C on the long-term prognosis of patients with COVID-19. Our research population was Chinese, further research based on other populations is needed before we can infer the results and draw general conclusions.

CONCLUSION

In conclusion, sCys C can be used as a means for early diagnosis of severe and critical COVID-19 patients with acute renal function damage, providing predictive information for clinical prognosis. Clinicians should improve their understanding pertaining to kidney injury in severe and critically ill patients with COVID-19. Early detection and effective intervention of renal involvement may help reduce potential death in patients with COVID-19.

ARTICLE HIGHLIGHTS

Research background

Coronavirus disease 2019 (COVID-19) is an acute infectious disease caused by a new coronavirus, which is clinically categorized into mild, moderate, severe, and critical illness. Severe and critically ill patients progress rapidly with dyspnea, hypoxemia, and even life-threatening complications such as multiple organ dysfunction syndrome, sepsis, and shock. Early detection of risk factors that affect the prognosis of severe and critically ill patients while using active and effective treatment measures is very important and can effectively reduce the potential mortality rate.

Research motivation

A rapid, effective, and accurate biomarker is urgently needed to predict the prognosis of patients with COVID-19. Serum cystatin C (sCys C) has a predictive value in the prognosis of critically ill patients such as those with acute cerebral infarction, acute

myocardial infarction, heart failure, and sepsis. However, the predictive value of sCys C for severe and critically ill COVID-19 patients has not been confirmed. We designed this study to confirm whether sCys C can be used as a prognostic indicator for COVID-19 patients.

Research objectives

The main objective of this study was to determine the predictive value of sCys C for the prognosis of severe and critically ill COVID-19 patients.

Research methods

A total of 101 severe or critically ill patients with COVID-19 were divided into a discharge group (64 cases) and a death group (37 cases). We compared the general information, underlying diseases, and laboratory examination indexes of the two groups. Multivariate Cox regression was used to explore the relationship between sCys C and prognosis. In addition, we used the receiver operating characteristic curve to assess the cut-off value of sCys C in predicting the death of patients.

Research results

sCys C, creatine kinase-MB, aspartate aminotransferase, albumin, and serum calcium were independent risk factors for death in severe and critically ill patients. sCys C had an AUC of 0.755, sensitivity of 0.865, and specificity of 0.562. Patients with a sCys C level of 0.80 or greater were at a high risk of death.

Research conclusions

sCys C is superior to sCr in predicting the risk of death in severe and critically ill patients with COVID-19. Patients with sCys C ≥ 0.80 mg/L are at a high risk of death.

Research perspectives

sCys C could be a marker of mortality in severe and critically ill COVID-19 patients.

REFERENCES

- 1 Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, Xiang J, Wang Y, Song B, Gu X, Guan L, Wei Y, Li H, Wu X, Xu J, Tu S, Zhang Y, Chen H, Cao B. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020; **395**: 1054-1062 [PMID: 32171076 DOI: 10.1016/S0140-6736(20)30566-3]
- 2 Hirsch JS, Ng JH, Ross DW, Sharma P, Shah HH, Barnett RL, Hazzan AD, Fishbane S, Jhaveri KD; Northwell COVID-19 Research Consortium; Northwell Nephrology COVID-19 Research Consortium. Acute kidney injury in patients hospitalized with COVID-19. *Kidney Int* 2020; **98**: 209-218 [PMID: 32416116 DOI: 10.1016/j.kint.2020.05.006]
- 3 Adapa S, Aeddula NR, Konala VM, Chenna A, Naramala S, Madhira BR, Gayam V, Balla M, Muppidi V, Bose S. COVID-19 and Renal Failure: Challenges in the Delivery of Renal Replacement Therapy. *J Clin Med Res* 2020; **12**: 276-285 [PMID: 32489502 DOI: 10.14740/jocmr4160]
- 4 Delanaye P, Cavalier E, Morel J, Mehdi M, Maillard N, Claisse G, Lambermont B, Dubois BE, Damas P, Krzesinski JM, Lautrette A, Mariat C. Detection of decreased glomerular filtration rate in intensive care units: serum cystatin C versus serum creatinine. *BMC Nephrol* 2014; **15**: 9 [PMID: 24410757 DOI: 10.1186/1471-2369-15-9]
- 5 Liu J. Evaluation of serum cystatin C for diagnosis of acute rejection after renal transplantation. *Transplant Proc* 2012; **44**: 1250-1253 [PMID: 22663994 DOI: 10.1016/j.transproceed.2012.01.138]
- 6 Al-Beladi FI. Cystatin C is an early marker of contrast-induced nephropathy in patients with sepsis in the intensive care unit. *Saudi J Kidney Dis Transpl* 2015; **26**: 718-724 [PMID: 26178544 DOI: 10.4103/1319-2442.160170]
- 7 Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, Wang B, Xiang H, Cheng Z, Xiong Y, Zhao Y, Li Y, Wang X, Peng Z. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA* 2020; **323**: 1061-1069 [PMID: 32031570 DOI: 10.1001/jama.2020.1585]
- 8 Parohan M, Yaghoubi S, Seraji A. Liver injury is associated with severe coronavirus disease 2019 (COVID-19) infection: A systematic review and meta-analysis of retrospective studies. *Hepatol Res* 2020; **50**: 924-935 [PMID: 32386449 DOI: 10.1111/hepr.13510]
- 9 Lei F, Liu YM, Zhou F, Qin JJ, Zhang P, Zhu L, Zhang XJ, Cai J, Lin L, Ouyang S, Wang X, Yang C, Cheng X, Liu W, Li H, Xie J, Wu B, Luo H, Xiao F, Chen J, Tao L, Cheng G, She ZG, Zhou J, Wang H, Lin J, Luo P, Fu S, Zhou J, Ye P, Xiao B, Mao W, Liu L, Yan Y, Liu L, Chen G, Li H, Huang X, Zhang BH, Yuan Y. Longitudinal Association Between Markers of Liver Injury and Mortality in COVID-19 in China. *Hepatology* 2020; **72**: 389-398 [PMID: 32359177 DOI: 10.1002/hep.31301]
- 10 Cappellini F, Brivio R, Casati M, Cavallero A, Contro E, Brambilla P. Low levels of total and ionized calcium in blood of COVID-19 patients. *Clin Chem Lab Med* 2020 [PMID: 32459190 DOI: 10.1515/cclm-2020-0611]

- 11 **Lippi G**, South AM, Henry BM. Electrolyte imbalances in patients with severe coronavirus disease 2019 (COVID-19). *Ann Clin Biochem* 2020; **57**: 262-265 [PMID: [32266828](#) DOI: [10.1177/0004563220922255](#)]
- 12 **Zhi H**, Zhang M, Cui X, Li Y. [Renal echography and cystatin C for prediction of acute kidney injury: very different in patients with cardiac failure or sepsis]. *Zhonghua Wei Zhong Bing Ji Jiu Yi Xue* 2019; **31**: 1258-1263 [PMID: [31771725](#) DOI: [10.3760/cma.j.issn.2095-4352.2019.10.015](#)]
- 13 **Fu Z**, Yang X, Shen M, Xue H, Qian G, Cao F, Guo J, Dong W, Chen Y. Prognostic ability of cystatin C and homocysteine plasma levels for long-term outcomes in very old acute myocardial infarction patients. *Clin Interv Aging* 2018; **13**: 1201-1209 [PMID: [30013331](#) DOI: [10.2147/CIA.S151211](#)]
- 14 **Jung YJ**, Lee HR, Kwon OJ. Comparison of serum cystatin C and creatinine as a marker for early detection of decreasing glomerular filtration rate in renal transplants. *J Korean Surg Soc* 2012; **83**: 69-74 [PMID: [22880179](#) DOI: [10.4174/jkss.2012.83.2.69](#)]
- 15 **Simonsen O**, Grubb A, Thysell H. The blood serum concentration of cystatin C (gamma-trace) as a measure of the glomerular filtration rate. *Scand J Clin Lab Invest* 1985; **45**: 97-101 [PMID: [3923607](#) DOI: [10.3109/00365518509160980](#)]
- 16 **Jickling GC**, Sharp FR. Biomarker panels in ischemic stroke. *Stroke* 2015; **46**: 915-920 [PMID: [25657186](#) DOI: [10.1161/STROKEAHA.114.005604](#)]
- 17 **Xiang J**, Wen J, Yuan X, Xiong S, Zhou X, Liu C, Min X. Potential biochemical markers to identify severe cases among COVID-19 patients. 2020 Preprint. Available from: [medRxiv:2020.03.19.20034447](#) [DOI: [10.1101/2020.03.19.20034447](#)]
- 18 **Henry BM**, de Oliveira MHS, Benoit S, Plebani M, Lippi G. Hematologic, biochemical and immune biomarker abnormalities associated with severe illness and mortality in coronavirus disease 2019 (COVID-19): a meta-analysis. *Clin Chem Lab Med* 2020; **58**: 1021-1028 [PMID: [32286245](#) DOI: [10.1515/cclm-2020-0369](#)]
- 19 **Su H**, Yang M, Wan C, Yi LX, Tang F, Zhu HY, Yi F, Yang HC, Fogo AB, Nie X, Zhang C. Renal histopathological analysis of 26 postmortem findings of patients with COVID-19 in China. *Kidney Int* 2020; **98**: 219-227 [PMID: [32327202](#) DOI: [10.1016/j.kint.2020.04.003](#)]
- 20 **Huang C**, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z, Yu T, Xia J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo L, Xie J, Wang G, Jiang R, Gao Z, Jin Q, Wang J, Cao B. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; **395**: 497-506 [PMID: [31986264](#) DOI: [10.1016/S0140-6736\(20\)30183-5](#)]
- 21 **Chen N**, Zhou M, Dong X, Qu J, Gong F, Han Y, Qiu Y, Wang J, Liu Y, Wei Y, Xia J, Yu T, Zhang X, Zhang L. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020; **395**: 507-513 [PMID: [32007143](#) DOI: [10.1016/S0140-6736\(20\)30211-7](#)]
- 22 **Guan WJ**, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, Liu L, Shan H, Lei CL, Hui DSC, Du B, Li LJ, Zeng G, Yuen KY, Chen RC, Tang CL, Wang T, Chen PY, Xiang J, Li SY, Wang JL, Liang ZJ, Peng YX, Wei L, Liu Y, Hu YH, Peng P, Wang JM, Liu JY, Chen Z, Li G, Zheng ZJ, Qiu SQ, Luo J, Ye CJ, Zhu SY, Zhong NS; China Medical Treatment Expert Group for Covid-19. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* 2020; **382**: 1708-1720 [PMID: [32109013](#) DOI: [10.1056/NEJMoa2002032](#)]
- 23 **Pei G**, Zhang Z, Peng J, Liu L, Zhang C, Yu C, Ma Z, Huang Y, Liu W, Yao Y, Zeng R, Xu G. Renal Involvement and Early Prognosis in Patients with COVID-19 Pneumonia. *J Am Soc Nephrol* 2020; **31**: 1157-1165 [PMID: [32345702](#) DOI: [10.1681/ASN.2020030276](#)]
- 24 **Akerfeldt T**, Helmersson J, Larsson A. Postsurgical inflammatory response is not associated with increased serum cystatin C values. *Clin Biochem* 2010; **43**: 1138-1140 [PMID: [20627098](#) DOI: [10.1016/j.clinbiochem.2010.07.004](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

