

World Journal of *Clinical Cases*

World J Clin Cases 2020 October 26; 8(20): 4688-5069



MINIREVIEWS

- 4688 Relationship between non-alcoholic fatty liver disease and coronary heart disease
Arslan U, Yenercağ M

ORIGINAL ARTICLE

Retrospective Cohort Study

- 4700 Remission of hepatotoxicity in chronic pulmonary aspergillosis patients after lowering trough concentration of voriconazole
Teng GJ, Bai XR, Zhang L, Liu HJ, Nie XH

Retrospective Study

- 4708 Endoscopic submucosal dissection as alternative to surgery for complicated gastric heterotopic pancreas
Noh JH, Kim DH, Kim SW, Park YS, Na HK, Ahn JY, Jung KW, Lee JH, Choi KD, Song HJ, Lee GH, Jung HY
- 4719 Observation of the effects of three methods for reducing perineal swelling in children with developmental hip dislocation
Wang L, Wang N, He M, Liu H, Wang XQ
- 4726 Predictive value of serum cystatin C for risk of mortality in severe and critically ill patients with COVID-19
Li Y, Yang S, Peng D, Zhu HM, Li BY, Yang X, Sun XL, Zhang M
- 4735 Sleep quality of patients with postoperative glioma at home
Huang Y, Jiang ZJ, Deng J, Qi YJ
- 4743 Early complications of preoperative external traction fixation in the staged treatment of tibial fractures: A series of 402 cases
Yang JZ, Zhu WB, Li LB, Dong QR
- 4753 Retroperitoneal vs transperitoneal laparoscopic lithotripsy of 20-40 mm renal stones within horseshoe kidneys
Chen X, Wang Y, Gao L, Song J, Wang JY, Wang DD, Ma JX, Zhang ZQ, Bi LK, Xie DD, Yu DX
- 4763 Undifferentiated embryonal sarcoma of the liver: Clinical characteristics and outcomes
Zhang C, Jia CJ, Xu C, Sheng QJ, Dou XG, Ding Y
- 4773 Cerebral infarct secondary to traumatic internal carotid artery dissection
Wang GM, Xue H, Guo ZJ, Yu JL
- 4785 Home-based nursing for improvement of quality of life and depression in patients with postpartum depression
Zhuang CY, Lin SY, Cheng CJ, Chen XJ, Shi HL, Sun H, Zhang HY, Fu MA

Observational Study

- 4793** Cost-effectiveness of lutetium (^{177}Lu) oxodotreotide *vs* everolimus in gastroenteropancreatic neuroendocrine tumors in Norway and Sweden
Palmer J, Leeuwenkamp OR
- 4807** Factors related to improved American Spinal Injury Association grade of acute traumatic spinal cord injury
Tian C, Lv Y, Li S, Wang DD, Bai Y, Zhou F, Ma QB
- 4816** Intraoperative systemic vascular resistance is associated with postoperative nausea and vomiting after laparoscopic hysterectomy
Qu MD, Zhang MY, Wang GM, Wang Z, Wang X

META-ANALYSIS

- 4826** Underwater *vs* conventional endoscopic mucosal resection in treatment of colorectal polyps: A meta-analysis
Ni DQ, Lu YP, Liu XQ, Gao LY, Huang X

CASE REPORT

- 4838** Dehydrated patient without clinically evident cause: A case report
Palladino F, Fedele MC, Casertano M, Liguori L, Esposito T, Guarino S, Miraglia del Giudice E, Marzuillo P
- 4844** Intracranial malignant solitary fibrous tumor metastasized to the chest wall: A case report and review of literature
Usuda D, Yamada S, Izumida T, Sangen R, Higashikawa T, Nakagawa K, Iguchi M, Kasamaki Y
- 4853** End-of-life home care of an interstitial pneumonia patient supported by high-flow nasal cannula therapy: A case report
Goda K, Kenzaka T, Kuriyama K, Hoshijima M, Akita H
- 4858** Rupture of carotid artery pseudoaneurysm in the modern era of definitive chemoradiation for head and neck cancer: Two case reports
Kim M, Hong JH, Park SK, Kim SJ, Lee JH, Byun J, Ko YH
- 4866** Unremitting diarrhoea in a girl diagnosed anti-N-methyl-D-aspartate-receptor encephalitis: A case report
Onpoaree N, Veeravigrom M, Sanpavat A, Suratannon N, Sintusek P
- 4876** Paliperidone palmitate-induced facial angioedema: A case report
Srifuengfung M, Sukakul T, Liangcheep C, Viravan N
- 4883** Improvement of lenvatinib-induced nephrotic syndrome after adaptation to sorafenib in thyroid cancer: A case report
Yang CH, Chen KT, Lin YS, Hsu CY, Ou YC, Tung MC
- 4895** Adult metaplastic hutch diverticulum with robotic-assisted diverticulectomy and reconstruction: A case report
Yang CH, Lin YS, Ou YC, Weng WC, Huang LH, Lu CH, Hsu CY, Tung MC

- 4902** Thrombus straddling a patent foramen ovale and pulmonary embolism: A case report
Huang YX, Chen Y, Cao Y, Qiu YG, Zheng JY, Li TC
- 4908** Therapeutic experience of an 89-year-old high-risk patient with incarcerated cholecystolithiasis: A case report and literature review
Zhang ZM, Zhang C, Liu Z, Liu LM, Zhu MW, Zhao Y, Wan BJ, Deng H, Yang HY, Liao JH, Zhu HY, Wen X, Liu LL, Wang M, Ma XT, Zhang MM, Liu JJ, Liu TT, Huang NN, Yuan PY, Gao YJ, Zhao J, Guo XA, Liao F, Li FY, Wang XT, Yuan RJ, Wu F
- 4917** Woven coronary artery: A case report
Wei W, Zhang Q, Gao LM
- 4922** Idiopathic multicentric Castleman disease with pulmonary and cutaneous lesions treated with tocilizumab: A case report
Han PY, Chi HH, Su YT
- 4930** Perianorectal abscesses and fistula due to ingested jujube pit in infant: Two case reports
Liu YH, Lv ZB, Liu JB, Sheng QF
- 4938** Forniceal deep brain stimulation in severe Alzheimer's disease: A case report
Lin W, Bao WQ, Ge JJ, Yang LK, Ling ZP, Xu X, Jiang JH, Zuo CT, Wang YH
- 4946** Systemic autoimmune abnormalities complicated by cytomegalovirus-induced hemophagocytic lymphohistiocytosis: A case report
Miao SX, Wu ZQ, Xu HG
- 4953** Nasal mucosa pyoderma vegetans associated with ulcerative colitis: A case report
Yu SX, Cheng XK, Li B, Hao JH
- 4958** Amiodarone-induced hepatotoxicity — quantitative measurement of iodine density in the liver using dual-energy computed tomography: Three case reports
Lv HJ, Zhao HW
- 4966** Multisystem involvement Langerhans cell histiocytosis in an adult: A case report
Wang BB, Ye JR, Li YL, Jin Y, Chen ZW, Li JM, Li YP
- 4975** New mutation in EPCAM for congenital tufting enteropathy: A case report
Zhou YQ, Wu GS, Kong YM, Zhang XY, Wang CL
- 4981** Catastrophic vertebral artery and subclavian artery pseudoaneurysms caused by a fishbone: A case report
Huang W, Zhang GQ, Wu JJ, Li B, Han SG, Chao M, Jin K
- 4986** Anastomosing hemangioma arising from the left renal vein: A case report
Zheng LP, Shen WA, Wang CH, Hu CD, Chen XJ, Shen YY, Wang J
- 4993** Bladder perforation caused by long-term catheterization misdiagnosed as digestive tract perforation: A case report
Wu B, Wang J, Chen XJ, Zhou ZC, Zhu MY, Shen YY, Zhong ZX

- 4999** Primary pulmonary plasmacytoma accompanied by overlap syndrome: A case report and review of the literature
Zhou Y, Wang XH, Meng SS, Wang HC, Li YX, Xu R, Lin XH
- 5007** Gastrointestinal stromal tumor metastasis at the site of a totally implantable venous access port insertion: A rare case report
Yin XN, Yin Y, Wang J, Shen CY, Chen X, Zhao Z, Cai ZL, Zhang B
- 5013** Massive gastrointestinal bleeding caused by a Dieulafoy's lesion in a duodenal diverticulum: A case report
He ZW, Zhong L, Xu H, Shi H, Wang YM, Liu XC
- 5019** Plastic bronchitis associated with *Botrytis cinerea* infection in a child: A case report
Liu YR, Ai T
- 5025** Chest, pericardium, abdomen, and thigh penetrating injury by a steel rebar: A case report
Yang XW, Wang WT
- 5030** Monocular posterior scleritis presenting as acute conjunctivitis: A case report
Li YZ, Qin XH, Lu JM, Wang YP
- 5036** Choriocarcinoma with lumbar muscle metastases: A case report
Pang L, Ma XX
- 5042** Primary chondrosarcoma of the liver: A case report
Liu ZY, Jin XM, Yan GH, Jin GY
- 5049** Successful management of a tooth with endodontic-periodontal lesion: A case report
Alshawwa H, Wang JF, Liu M, Sun SF
- 5057** Rare imaging findings of hypersensitivity pneumonitis: A case report
Wang HJ, Chen XJ, Fan LX, Qi QL, Chen QZ
- 5062** Effective administration of cranial drilling therapy in the treatment of fourth degree temporal, facial and upper limb burns at high altitude: A case report
Shen CM, Li Y, Liu Z, Qi YZ

ABOUT COVER

Peer-reviewer of *World Journal of Clinical Cases*, Dr. Aleem Ahmed Khan is a Distinguished Scientist and Head of The Central Laboratory for Stem Cell Research and Translational Medicine, Centre for Liver Research and Diagnostics, Deccan College of Medical Sciences, Kanchanbagh, Hyderabad (India). Dr. Aleem completed his Doctorate from Osmania University, Hyderabad in 1998 and has since performed pioneering work in the treatment of acute liver failure and decompensated cirrhosis using hepatic stem cell transplantation. During his extensive research career he supervised 10 PhD students and published > 150 research articles, 7 book chapters, and 2 patents. His ongoing research involves developing innovative technologies for organ regeneration and management of advanced cancers. (L-Editor: Filipodia)

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, PubMed, and PubMed Central. The 2020 Edition of Journal Citation Reports® cites the 2019 impact factor (IF) for *WJCC* as 1.013; IF without journal self cites: 0.991; Ranking: 120 among 165 journals in medicine, general and internal; and Quartile category: Q3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Ji-Hong Liu; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lai Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Semimonthly

EDITORS-IN-CHIEF

Dennis A Bloomfield, Sandro Vento, Bao-Gan Peng

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

October 26, 2020

COPYRIGHT

© 2020 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

End-of-life home care of an interstitial pneumonia patient supported by high-flow nasal cannula therapy: A case report

Ken Goda, Tsuneaki Kenzaka, Kyosuke Kuriyama, Masahiko Hoshijima, Hozuka Akita

ORCID number: Ken Goda 0000-0002-2462-1157; Tsuneaki Kenzaka 0000-0002-3120-6605; Kyosuke Kuriyama 0000-0001-8854-3314; Masahiko Hoshijima 0000-0002-0616-7317; Hozuka Akita 0000-0002-7612-9703.

Author contributions: Goda K managed the case and prepared and revised the manuscript; Kenzaka T assisted with manuscript preparation, correction, and revision; Kuriyama K and Hoshijima M managed the case and assisted with manuscript preparation, correction, and revision; Akita H assisted with manuscript preparation, correction, and revision. All authors have read and approved the final version of the manuscript.

Informed consent statement:

Written informed consent was obtained from the patient's wife for the publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor of this journal.

Conflict-of-interest statement: The authors declare that they have no competing interests.

CARE Checklist (2016) statement:

The authors have read the CARE Checklist (2016), and the manuscript was prepared and

Ken Goda, Tsuneaki Kenzaka, Masahiko Hoshijima, Hozuka Akita, Department of Internal Medicine, Hyogo Prefectural Tamba Medical Center, Tamba 669-3495, Japan

Ken Goda, Tsuneaki Kenzaka, Division of Community Medicine and Career Development, Kobe University Graduate School of Medicine, Kobe 652-0032, Japan

Ken Goda, Tsuneaki Kenzaka, Department of Medicine, Tamba City mirune Clinic, Tamba 669-3464, Japan

Kyosuke Kuriyama, Clinical Engineer, Nursing Department, Hyogo Prefectural Tamba Medical Center, Tamba 669-3495, Japan

Corresponding author: Tsuneaki Kenzaka, MD, PhD, Professor, Division of Community Medicine and Career Development, Kobe University Graduate School of Medicine, 2-1-5, Arata-cho, Hyogo-ku, Kobe 652-0032, Japan. smile.kenzaka@jichi.ac.jp

Abstract

BACKGROUND

High-flow nasal cannula (HFNC) therapy and morphine continuous subcutaneous infusion (CSI) have been used to ameliorate dyspnea in non-cancer patients with end-stage respiratory diseases, including chronic obstructive pulmonary disease and interstitial pneumonia, primarily in hospital settings. However, it is rare to perform home-based medical treatment using these. We observe a case to assess the feasibility of this treatment strategy.

CASE SUMMARY

Here, we report a case of a 75-year-old man who was diagnosed with interstitial pneumonia 11 years ago and was successfully nursed at home during his terminal phase for over 10 mo without hospitalization, by introducing domiciliary uses of HFNC and morphine CSI with a patient-controlled analgesia device.

CONCLUSION

Active utilization of HFNC and morphine CSI with patient-controlled analgesia device would substantiate successful end-of-life palliative home care of idiopathic interstitial pneumonia patients.

Key Words: High flow nasal cannula; Continuous subcutaneous infusion; Morphine; Patient controlled analgesia; Home care; Interstitial pneumonia; Case report

revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Received: August 24, 2020

Peer-review started: August 24, 2020

First decision: September 13, 2020

Revised: September 18, 2020

Accepted: September 25, 2020

Article in press: September 25, 2020

Published online: October 26, 2020

P-Reviewer: Chan SM

S-Editor: Zhang L

L-Editor: A

P-Editor: Zhang YL



©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: We propose that active utilization of high-flow nasal cannula and morphine continuous subcutaneous infusion with a patient-controlled analgesia device would substantiate successful palliative care of patients with idiopathic interstitial pneumonia at home, nearing their end-of-life.

Citation: Goda K, Kenzaka T, Kuriyama K, Hoshijima M, Akita H. End-of-life home care of an interstitial pneumonia patient supported by high-flow nasal cannula therapy: A case report. *World J Clin Cases* 2020; 8(20): 4853-4857

URL: <https://www.wjgnet.com/2307-8960/full/v8/i20/4853.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v8.i20.4853>

INTRODUCTION

High-flow nasal cannula (HFNC) therapy has been widely used in hospital settings to treat various forms of respiratory disorders. These include acute hypoxemic respiratory failure of patients with congestive heart failure, exacerbation of chronic obstructive pulmonary disease (COPD), lung cancer, and interstitial pneumonia, postoperative respiratory support, post-extubation respiratory support^[1]. Recently, the use of HFNC has been extended to the palliative care of critically ill patients^[1]; however, there have been a limited number of reports of the domiciliary uses of HFNC for patients with COPD^[2] and chronic respiratory tract infection after gastrectomy^[3].

Morphine is commonly known to provide effective relief to dyspnea caused by malignant tumors^[4]. Similarly, it has been shown to be an effective home-based treatment for dyspnea in patients with COPD, at the end of life^[5]. In contrast, while both oral administration^[6] and continuous subcutaneous injection (CSI)^[7] have been used effectively in hospital settings to ease dyspnea in interstitial pneumonia, the domiciliary use of morphine in patients with terminal-stage interstitial pneumonia has not been reported.

Here, we report a case where a patient with severe interstitial pneumonia and frequent relapse of fatal type 1 respiratory failure spent his final 10 mo at home without hospitalization by utilizing HFNC and morphine CSI with a patient-controlled analgesia (PCA) device.

CASE PRESENTATION

Chief complaints

A 75-year-old man presented to our hospital with dyspnea.

History of present illness

He was diagnosed with interstitial pneumonia 11 years ago. He had been treated with prednisolone (5 mg/d) and home oxygen therapy (3 L/min) for 1 year. However, between November of the same year to January of the following year, his lung function deteriorated, leading to hospitalization for 10 times over the next 13 mo for acute type 1 respiratory failure. The maximum length of stay at home between hospitalization was 30 d. Before his final admission to our hospital (his eleventh admission), the patient was at home for 6 d. At that time, hospitalization until the very end of life was proposed to the patient.

History of past illness

The patient had type 2 diabetes.

Personal and family history

At home he lived with his wife, and while he could consume food independently, he required assistance in changing clothes. Furthermore, excretion was done on the floor, and bathing was not possible due to restricted movements caused by exertion

dyspnea. His modified Medical Research Council grade was 4. He smoked 80 cigarettes a day between the ages of 20 and 50.

Physical examination

The patient's was thin during the final admission. He was fully conscious, with a body temperature of 38.4 °C and a respiratory rate of 57 breaths/min. His SpO₂ was 60% at 15 L oxygen/min with a mask with reservoir and 94% with non-invasive positive pressure ventilation [inspiratory positive airway pressure, 8 cmH₂O; expiratory positive airway pressure, 4 cmH₂O, and fraction of inspired oxygen (FiO₂) 100%]. Fine crackles were heard bilaterally on the chest.

Laboratory examinations

Blood tests showed increases in his white blood cell count [32100 cells/μL, (neutrophils: 48.2%)] and C-reactive protein level (9.12 mg/L); no abnormalities were detected in the patient's urinalysis.

Imaging examinations

Chest X-rays showed reticular shadows on both sides of the lungs, and the computed tomography scan displayed a mixture of frosted glass shadows, reticular shadows, and tractive bronchodilation on both lungs (Figure 1).

FINAL DIAGNOSIS

We considered acute exacerbations of interstitial pneumonia and bacterial pneumonia.

TREATMENT

Although his general condition improved with piperazine/tazobactam treatment and the use of non-invasive positive pressure ventilation, the effect of orally administered codeine phosphate and morphine hydrochloride to suppress dyspnea and bouts of coughing was limited. Accordingly, morphine CSI (7.2 mg/d) was introduced. HFNC (FiO₂ 35%, 35 L/min) was also used to successfully assist his respiration.

OUTCOME AND FOLLOW-UP

Adhering to the patient's wish to stay home, we coordinated with home care support to establish a system at home to provide CSI service with PCA, as well as domiciliary HFNC with oxygen supplied through two oxygen concentrators (maximum productivity 7 L oxygen/min each); then, he was discharged. CSI used a CADD Legacy® (Smiths Medical, St. Paul, MN, United States) with a 50–100-mL disposable medication cassette reservoir, which was replaced weekly. Initially, HFNC was set to provide humidified air of FiO₂ 37% at the rate of 35 L/min. The patient developed pneumonia several times, which was treated with antimicrobials without hospitalization. He survived for 10 more mo within home care.

DISCUSSION

We present the case of a patient with advanced stage of interstitial pneumonia, who was treated at home for a long period of time through a combination of HFNC and CSI. To the best of our knowledge, there is no preceding report on the domiciliary use of HFNC for terminal care of patients with interstitial pneumonia, nor reports that successfully used CSI at home in such patients.

Compared to conventional oxygen therapy, HFNC provides more stable high-flow high-concentration oxygen^[8]. It has a beneficial effect in flushing anatomical dead spaces in the airway^[9] and improving airway health through heating and humidification^[10,11]. This has broad implications for inpatients for both acute and terminal stages of lung diseases. The exacerbating symptoms of COPD patients, who are hospitalized, are alleviated by the use of HFNC^[12]. The long-term high-flow oxygen administration facilitates home-based care for critically ill patients at their end of




Figure 1 Chest computed tomography at latest admission.

lives^[2,13]. HFNC also improves respiration in idiopathic interstitial pneumonia^[1]. Moreover, patients using HFNC can comfortably eat and converse while high-concentration oxygen is administered, maintaining better quality of life. Nonetheless, there have been a limited number of reports of using HFNC to treat critically ill lung disease patients at home at the end of life, mainly due to issues associated with oxygen supplies and securing purified water for humidification. For example, in the current case, the patient needed approximately 10 L of purified distilled water per week, which was supplied through home visiting service by a community pharmacist. The maximum supply of oxygen from the two oxygen concentrators connected together was 14 L/min. We also discussed rescue plans with the local fire department for issues such as unscheduled power outage. Solving these problems was crucial for the success of home care.

Morphine has been reported to be useful for respiratory distress in malignant tumors^[4]. In COPD, morphine is clearly effective in alleviating dyspnea symptoms in severely ill patients^[14,15]. It has central and peripheral antitussive effects (*i.e.* alleviation of anxiety and perceptual changes in dyspnea in the center), while decreasing systemic oxygen consumption in the periphery^[16]. The efficacy of CSI has been demonstrated for interstitial pneumonia^[7], which we also confirmed in the current case. CSI of morphine has the advantages over oral or intravascular administration as follows: it does not require a blood vessel, patients' movements are not restricted by infusion lines, and morphine can be administered at a stable rate^[17].

The majority of idiopathic interstitial pneumonia cases are progressive and do not respond to therapy. For patients who do not respond to disease specific therapy and who are not candidates for lung transplantation, the focus needs to shift towards end-of-life care^[18]. The symptoms of dyspnea, persistent cough, and general fatigue are difficult to manage. The prognosis of idiopathic interstitial pneumonia is variable and unpredictable^[19]. As effective remedies are limited, early introduction of palliative care is important to alleviate symptoms^[18]. However, this is not generally the case^[20]. While many patients with interstitial pneumonia wish to spend their last days at home^[21], only a small number are actually able to do so^[22,23]. Patient-related factors such as their own and their families' willingness, and the care environment they accept, are essential to facilitate spending their last days under home care^[24]. On the other hand, it is necessary to have a factor that can provide medical services including a home-visit medical system and home-visit nursing system^[24]. Here, we propose that active utilization of HFNC and morphine CSI with a PCA device would substantiate successful palliative care of patients with idiopathic interstitial pneumonia at home nearing their end of life.

CONCLUSION

We facilitated and examined a case where HFNC and morphine CSI with PCA device were introduced at home to a patient with interstitial pneumonia who had recurrent type 1 respiratory failure; this enabled long-term treatment and home care. Our observations supported the option of having end-of-life medical care at home even in patients with interstitial pneumonia.

REFERENCES

- 1 **Ito J**, Nagata K, Sato S, Shiraki A, Nishimura N, Izumi S, Tachikawa R, Morimoto T, Tomii K. The clinical practice of high-flow nasal cannula oxygen therapy in adults: A Japanese cross-sectional multicenter survey. *Respir Investig* 2018; **56**: 249-257 [PMID: [29773297](#) DOI: [10.1016/j.resinv.2018.02.002](#)]
- 2 **Nagata K**, Kikuchi T, Horie T, Shiraki A, Kitajima T, Kadowaki T, Tokioka F, Chohnabayashi N, Watanabe A, Sato S, Tomii K. Domiciliary High-Flow Nasal Cannula Oxygen Therapy for Patients with Stable Hypercapnic Chronic Obstructive Pulmonary Disease. *Ann Am Thorac Soc* 2018; **15**: 432-439 [PMID: [29283682](#) DOI: [10.1513/AnnalsATS.201706-425OC](#)]
- 3 **Hamada S**, Tsukino M. The clinical utility of domiciliary nocturnal high-flow nasal cannula in a post-gastrectomy patient with chronic lower respiratory tract infection. *Pulmonology* 2018; **24**: 312-313 [PMID: [30122335](#) DOI: [10.1016/j.pulmoe.2018.07.002](#)]
- 4 **Bruera E**, MacEachern T, Ripamonti C, Hanson J. Subcutaneous morphine for dyspnea in cancer patients. *Ann Intern Med* 1993; **119**: 906-907 [PMID: [8215003](#) DOI: [10.7326/0003-4819-119-9-199311010-00007](#)]
- 5 **Faes K**, Cohen J, Annemans L. Resource Use during the Last Six Months of Life among COPD Patients: A Population-Level Study. *J Pain Symptom Manage* 2018; **56**: 318-326.e7 [PMID: [29902554](#) DOI: [10.1016/j.jpainsymman.2018.06.002](#)]
- 6 **Allen S**, Raut S, Woollard J, Vassallo M. Low dose diamorphine reduces breathlessness without causing a fall in oxygen saturation in elderly patients with end-stage idiopathic pulmonary fibrosis. *Palliat Med* 2005; **19**: 128-130 [PMID: [15810751](#) DOI: [10.1191/0269216305pm998oa](#)]
- 7 **Matsuda Y**, Maeda I, Tachibana K, Nakao K, Sasaki Y, Sugimoto C, Arai T, Tokoro A, Akira M, Inoue Y. Low-Dose Morphine for Dyspnea in Terminally Ill Patients with Idiopathic Interstitial Pneumonias. *J Palliat Med* 2017; **20**: 879-883 [PMID: [28437202](#) DOI: [10.1089/jpm.2016.0432](#)]
- 8 **Dysart K**, Miller TL, Wolfson MR, Shaffer TH. Research in high flow therapy: mechanisms of action. *Respir Med* 2009; **103**: 1400-1405 [PMID: [19467849](#) DOI: [10.1016/j.rmed.2009.04.007](#)]
- 9 **Kernick J**, Magarey J. What is the evidence for the use of high flow nasal cannula oxygen in adult patients admitted to critical care units? *Aust Crit Care* 2010; **23**: 53-70 [PMID: [20206546](#) DOI: [10.1016/j.aucc.2010.01.001](#)]
- 10 **Hasani A**, Chapman TH, McCool D, Smith RE, Dilworth JP, Agnew JE. Domiciliary humidification improves lung mucociliary clearance in patients with bronchiectasis. *Chron Respir Dis* 2008; **5**: 81-86 [PMID: [18539721](#) DOI: [10.1177/1479972307087190](#)]
- 11 **Tomii K**. Indications and limitations of nasal high-flow therapy. *J Care Rehabil* 2015; **25**: 53-57 [DOI: [10.15032/jsr.25.1_53](#)]
- 12 **Storgaard LH**, Hockey HU, Laursen BS, Weinreich UM. Long-term effects of oxygen-enriched high-flow nasal cannula treatment in COPD patients with chronic hypoxemic respiratory failure. *Int J Chron Obstruct Pulmon Dis* 2018; **13**: 1195-1205 [PMID: [29713153](#) DOI: [10.2147/COPD.S159666](#)]
- 13 **Dolidon S**, Dupuis J, Molano Valencia LC, Salaün M, Thiberville L, Muir JF, Cuvelier A, Patout M. Characteristics and outcome of patients set up on high-flow oxygen therapy at home. *Ther Adv Respir Dis* 2019; **13**: 1753466619879794 [PMID: [31610722](#) DOI: [10.1177/1753466619879794](#)]
- 14 **Singh D**, Agusti A, Anzueto A, Barnes PJ, Bourbeau J, Celli BR, Criner GJ, Frith P, Halpin DMG, Han M, López Varela MV, Martínez F, Montes de Oca M, Papi A, Pavord ID, Roche N, Sin DD, Stockley R, Vestbo J, Wedzicha JA, Vogelmeier C. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Lung Disease: the GOLD science committee report 2019. *Eur Respir J* 2019; **53** [PMID: [30846476](#) DOI: [10.1183/13993003.00164-2019](#)]
- 15 **Rocker GM**, Sinuff T, Horton R, Hernandez P. Advanced chronic obstructive pulmonary disease: innovative approaches to palliation. *J Palliat Med* 2007; **10**: 783-797 [PMID: [17592991](#) DOI: [10.1089/jpm.2007.9951](#)]
- 16 **Kin SC**. Palliative medicine in malignant respiratory diseases. 4th ed. Hawks G, Cherhy NI, Christakis NA, editors. Oxford textbook of palliative medicine. Oxford: Oxford University Press, 2009: 1129-1130
- 17 **Koshi A**, Masaru H. Both switching from a winged needle to a small plastic intravenous catheter and adding dexamethasone to continuous subcutaneous infusion (CSCI) successfully treated inflammatory skin changes caused by CSCI. *Palliat Care Res* 2012; **7**: 112-120 [DOI: [10.2512/jspm.7.112](#)]
- 18 **Garibaldi BT**, Danoff SK. Symptom-based management of the idiopathic interstitial pneumonia. *Respirology* 2016; **21**: 1357-1365 [PMID: [26450007](#) DOI: [10.1111/resp.12649](#)]
- 19 **Wijsenbeek MS**, Holland AE, Swigris JJ, Renzoni EA. Comprehensive Supportive Care for Patients with Fibrosing Interstitial Lung Disease. *Am J Respir Crit Care Med* 2019; **200**: 152-159 [PMID: [31051080](#) DOI: [10.1164/rccm.201903-0614PP](#)]
- 20 **Ahmadi Z**, Wysham NG, Lundström S, Janson C, Currow DC, Ekström M. End-of-life care in oxygen-dependent ILD compared with lung cancer: a national population-based study. *Thorax* 2016; **71**: 510-516 [PMID: [26865603](#) DOI: [10.1136/thoraxjnl-2015-207439](#)]
- 21 **Skorstengaard MH**, Neergaard MA, Andreassen P, Brogaard T, Bendstrup E, Løkke A, Aagaard S, Wiggers H, Bech P, Jensen AB. Preferred Place of Care and Death in Terminally Ill Patients with Lung and Heart Disease Compared to Cancer Patients. *J Palliat Med* 2017; **20**: 1217-1224 [PMID: [28574737](#) DOI: [10.1089/jpm.2017.0082](#)]
- 22 **Rajala K**, Lehto JT, Saarinen M, Sutinen E, Saarto T, Myllärniemi M. End-of-life care of patients with idiopathic pulmonary fibrosis. *BMC Palliat Care* 2016; **15**: 85 [PMID: [27729035](#) DOI: [10.1186/s12904-016-0158-8](#)]
- 23 **Lindell KO**, Liang Z, Hoffman LA, Rosenzweig MQ, Saul MI, Pilewski JM, Gibson KF, Kaminski N. Palliative care and location of death in decedents with idiopathic pulmonary fibrosis. *Chest* 2015; **147**: 423-429 [PMID: [25187973](#) DOI: [10.1378/chest.14-1127](#)]
- 24 **Hattori A**, Uemura K, Masuda Y, Mogi N, Naito M, Iguchi A. [Factors contributing to dying at home in elderly patients who received home care service]. *Nihon Ronen Igakkai Zasshi* 2001; **38**: 399-404 [PMID: [11431898](#) DOI: [10.3143/geriatrics.38.399](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

