

World Journal of *Clinical Cases*

World J Clin Cases 2021 November 6; 9(31): 9320-9698



Contents

Thrice Monthly Volume 9 Number 31 November 6, 2021

FRONTIER

- 9320** Gut-liver axis in cirrhosis: Are hemodynamic changes a missing link?

Maslennikov R, Ivashkin V, Efremova I, Poluektova E, Shirokova E

REVIEW

- 9333** Pharmaconutrition strategy to resolve SARS-CoV-2-induced inflammatory cytokine storm in non-alcoholic fatty liver disease: Omega-3 long-chain polyunsaturated fatty acids

Jeyakumar SM, Vajreswari A

- 9350** Major depressive disorder: Validated treatments and future challenges

Karroui R, Hammani Z, Benjelloun R, Otheman Y

MINIREVIEWS

- 9368** Gene × environment interaction in major depressive disorder

Zhao MZ, Song XS, Ma JS

- 9376** Deep learning driven colorectal lesion detection in gastrointestinal endoscopic and pathological imaging

Cai YW, Dong FF, Shi YH, Lu LY, Chen C, Lin P, Xue YS, Chen JH, Chen SY, Luo XB

ORIGINAL ARTICLE

Case Control Study

- 9386** Cognitive behavioral therapy on personality characteristics of cancer patients

Yuan XH, Peng J, Hu SW, Yang Y, Bai YJ

Retrospective Cohort Study

- 9395** Extrapneumonic necrosis volume: A new tool in acute pancreatitis severity assessment?

Cucuteanu B, Negru D, Gavrilescu O, Popa IV, Floria M, Mihai C, Cijevschi Prelipcean C, Dranga M

- 9406** Establishment of a risk assessment score for deep vein thrombosis after artificial liver support system treatment

Ye Y, Li X, Zhu L, Yang C, Tan YW

Retrospective Study

- 9417** Clinical management and susceptibility of primary hepatic lymphoma: A cases-based retrospective study

Hai T, Zou LQ

- 9431** Association of serum pepsinogen with degree of gastric mucosal atrophy in an asymptomatic population

Cai HL, Tong YL

- 9440** Risk factors for relapse and nomogram for relapse probability prediction in patients with minor ischemic stroke

Yu XF, Yin WW, Huang CJ, Yuan X, Xia Y, Zhang W, Zhou X, Sun ZW

- 9452** Incidence, prognosis, and risk factors of sepsis-induced cardiomyopathy

Liang YW, Zhu YF, Zhang R, Zhang M, Ye XL, Wei JR

- 9469** Associations with pancreatic exocrine insufficiency: An United Kingdom single-centre study

Shandro BM, Chen J, Ritehnia J, Poullis A

- 9481** Retrospective analysis of influencing factors on the efficacy of mechanical ventilation in severe and critical COVID-19 patients

Zeng J, Qi XX, Cai WW, Pan YP, Xie Y

Observational Study

- 9491** Vitamin D deficiency, functional status, and balance in older adults with osteoarthritis

Montemor CN, Fernandes MTP, Marquez AS, Poli-Frederico RC, da Silva RA, Fernandes KBP

- 9500** Psychological impact of the COVID-19 pandemic on Chinese population: An online survey

Shah T, Shah Z, Yasmeen N, Ma ZR

- 9509** Outcomes of different minimally invasive surgical treatments for vertebral compression fractures: An observational study

Yeh KL, Wu SH, Liaw CK, Hou SM, Wu SS

META-ANALYSIS

- 9520** Glycated albumin as a biomarker for diagnosis of diabetes mellitus: A systematic review and meta-analysis

Xiong JY, Wang JM, Zhao XL, Yang C, Jiang XS, Chen YM, Chen CQ, Li ZY

CASE REPORT

- 9535** Rapid response to radiotherapy in unresectable tracheal adenoid cystic carcinoma: A case report

Wu Q, Xu F

- 9542** Clinical observation of pediatric-type follicular lymphomas in adult: Two case reports

Liu Y, Xing H, Liu YP

- 9549** Malignant adenomyoepithelioma of the breast: Two case reports and review of the literature

Zhai DY, Zhen TT, Zhang XL, Luo J, Shi HJ, Shi YW, Shao N

- 9557** Validation of diagnostic strategies of autoimmune atrophic gastritis: A case report

Sun WJ, Ma Q, Liang RZ, Ran YM, Zhang L, Xiao J, Peng YM, Zhan B

- 9564** Characteristics of primary giant cell tumor in soft tissue on magnetic resonance imaging: A case report

Kang JY, Zhang K, Liu AL, Wang HL, Zhang LN, Liu WV

- 9571** Acute esophageal necrosis as a complication of diabetic ketoacidosis: A case report
Moss K, Mahmood T, Spaziani R
- 9577** Simultaneous embolization of a spontaneous porto-systemic shunt and intrahepatic arterioportal fistula: A case report
Liu GF, Wang XZ, Luo XF
- 9584** Ureteroscopic holmium laser to transect the greater omentum to remove an abdominal drain: Four case reports
Liu HM, Luo GH, Yang XF, Chu ZG, Ye T, Su ZY, Kai L, Yang XS, Wang Z
- 9592** Forearm compartment syndrome due to acquired hemophilia that required massive blood transfusions after fasciotomy: A case report
Kameda T, Yokota T, Ejiri S, Konno SI
- 9598** Transforaminal endoscopic excision of bi-segmental non-communicating spinal extradural arachnoid cysts: A case report and literature review
Yun ZH, Zhang J, Wu JP, Yu T, Liu QY
- 9607** T-cell lymphoblastic lymphoma with extensive thrombi and cardiac thrombosis: A case report and review of literature
Ma YY, Zhang QC, Tan X, Zhang X, Zhang C
- 9617** Perfect pair, scopes unite – laparoscopic-assisted transumbilical gastroscopy for gallbladder-preserving polypectomy: A case report
Zheng Q, Zhang G, Yu XH, Zhao ZF, Lu L, Han J, Zhang JZ, Zhang JK, Xiong Y
- 9623** Bilateral hematoma after tubeless percutaneous nephrolithotomy for unilateral horseshoe kidney stones: A case report
Zhou C, Yan ZJ, Cheng Y, Jiang JH
- 9629** Atypical endometrial hyperplasia in a 35-year-old woman: A case report and literature review
Wu X, Luo J, Wu F, Li N, Tang AQ, Li A, Tang XL, Chen M
- 9635** Clinical features and literature review related to the material differences in thread rhinoplasty: Two case reports
Lee DW, Ryu H, Jang SH, Kim JH
- 9645** Concurrent tuberculous transverse myelitis and asymptomatic neurosyphilis: A case report
Gu LY, Tian J, Yan YP
- 9652** Diagnostic value of contrast-enhanced ultrasonography in mediastinal leiomyosarcoma mimicking aortic hematoma: A case report and review of literature
Xie XJ, Jiang TA, Zhao QY
- 9662** Misidentification of hepatic tuberculosis as cholangiocarcinoma: A case report
Li W, Tang YF, Yang XF, Huang XY

- 9670** Brunner's gland hyperplasia associated with lipomatous pseudohypertrophy of the pancreas presenting with gastrointestinal bleeding: A case report
Nguyen LC, Vu KT, Vo TTT, Trinh CH, Do TD, Pham NTV, Pham TV, Nguyen TT, Nguyen HC, Byeon JS
- 9680** Metachronous squamous cell carcinoma of pancreas and stomach in an elderly female patient: A case report
Kim JH, Kang CD, Lee K, Lim KH
- 9686** Iatrogenic giant pseudomeningocele of the cervical spine: A case report
Kim KW, Cho JH
- 9691** Traditional Chinese medicine for gait disturbance in adrenoleukodystrophy: A case report and review of literature
Kim H, Kim T, Cho W, Chang H, Chung WS

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Takeo Furuya, MD, PhD, Assistant Professor, Department of Orthopaedic Surgery, Chiba University Graduate School of Medicine, Chiba 2608670, Japan. furuya-takeo@chiba-u.jp

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for WJCC as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The WJCC's CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yan-Xia Xing; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lai Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Dennis A Bloomfield, Sandro Vento, Bao-Gan Peng

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

November 6, 2021

COPYRIGHT

© 2021 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Retrospective Study

Association of serum pepsinogen with degree of gastric mucosal atrophy in an asymptomatic population

Hao-Lei Cai, Yu-Ling Tong

ORCID number: Hao-Lei Cai 0000-0002-5669-1600; Yu-Ling Tong 0000-0002-3772-9327.

Author contributions: Cai HL performed the research and wrote the paper; Tong YL designed the research, contributed to the analysis, and supervised the report.

Supported by National Natural Science Foundation of China, No. 71804161 and No. 72074188.

Institutional review board statement: This study was reviewed and approved by the Human Research Ethics Committee of the Second Hospital Affiliated to Zhejiang University School of Medicine (No. 2015-079).

Informed consent statement: All subjects provided written informed consent prior to enrollment in this study.

Conflict-of-interest statement: The authors declare that they have no related financial relationships to disclose.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and

Hao-Lei Cai, Department of Surgery, The Second Affiliated Hospital of Zhejiang University School of Medicine, Hangzhou 310009, Zhejiang Province, China

Yu-Ling Tong, International Healthcare Center, The Second Affiliated Hospital of Zhejiang University School of Medicine, Hangzhou 310009, Zhejiang Province, China

Corresponding author: Yu-ling Tong, MD, Doctor, International Healthcare Center, The Second Affiliated Hospital of Zhejiang University School of Medicine, No.88 Jiefang Road, Hangzhou 310009, Zhejiang Province, China. tongyl0313@zju.edu.cn

Abstract

BACKGROUND

Atrophic gastritis is a precancerous lesion of the stomach. It has been reported that pepsinogen (PG) can reflect the morphology and function of the gastric mucosa, and it is therefore used as a marker for the early diagnosis of atrophic gastritis.

AIM

To evaluate the diagnostic value of serum PG for degree of gastric mucosal atrophy in asymptomatic Chinese upon physical examination.

METHODS

Medical data were collected from subjects who underwent transnasal gastroscopy between October 2016 and October 2018. For each study subject, serum PG levels and presence of *Helicobacter pylori* (*H. pylori*) infection were investigated. Pathology was evaluated using the Operative Link for Gastritis Assessment (OLGA) classification and Operative Link on Gastric Intestinal Metaplasia Assessment (OLGIM) systems. All statistical analyses were carried out using SPSS statistical software.

RESULTS

A total of 2256 subjects were enrolled and 1922 cases were finally included in the study. Based on the OLGA grading system, the levels of PGI were slightly decreased, while those of PGII were slightly increased. The PGI/PGII ratio (PGR) was reduced with increasing atrophy. The association between PG and OLGA grading was higher compared with that between PG and the OLGIM grading system. Compared with the OLGA-0 group, a statistically significant difference was observed in the mean age of OLGA-I, III, and IV groups ($P < 0.05$). In the *H.*

fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Specialty type: Gastroenterology and hepatology

Country/Territory of origin: China

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C, C
Grade D (Fair): D
Grade E (Poor): 0

Received: March 26, 2021

Peer-review started: March 26, 2021

First decision: April 28, 2021

Revised: May 6, 2021

Accepted: September 27, 2021

Article in press: September 27, 2021

Published online: November 6, 2021

P-Reviewer: Cho JH, Mohammadi M, Yücel O

S-Editor: Liu M

L-Editor: Wang TQ

P-Editor: Liu M



pylori-positive subjects, the PGR levels were notably lower in the OLGA-I, II, and III groups compared with the OLGA-0 group ($P < 0.05$). *H. pylori*-positive subjects exhibited significantly higher PGI and PGII serum levels and a significantly lower PGR compared with *H. pylori*-negative patients in different OLGA groups ($P < 0.05$).

CONCLUSION

Serum PG levels may represent a non-invasive screening marker for gastric mucosal atrophy in asymptomatic subjects.

Key Words: Pepsinogen; *Helicobacter pylori*; Operative Link for Gastritis Assessment; Atrophic gastritis; Gastric mucosal atrophy; Biomarker

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The current study evaluated the diagnostic value of serum pepsinogen (PG) as a screening marker for atrophic gastritis in asymptomatic healthy check-up populations in different regions of China. Serum PG levels were closely associated with Operative Link for Gastritis Assessment grading and could be used as an effective non-invasive screening tool for atrophic gastritis in asymptomatic subjects. In addition, better results could be obtained in *Helicobacter pylori*-positive individuals. Screening is more necessary in the elderly, and the application of the aforementioned screening tool may be beneficial for this population.

Citation: Cai HL, Tong YL. Association of serum pepsinogen with degree of gastric mucosal atrophy in an asymptomatic population. *World J Clin Cases* 2021; 9(31): 9431-9439

URL: <https://www.wjgnet.com/2307-8960/full/v9/i31/9431.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v9.i31.9431>

INTRODUCTION

In 2014, approximately 410400 new cases of gastric cancer (GC) were diagnosed in China, accounting for 10.79% of all cancer cases[1]. The prognosis of GC is closely associated with the times of diagnosis and treatment. The 5-year survival rate of advanced GC is less than 30%, while early GC, with a 5-year survival rate of over 90%, can be treated by endoscopy[2]. Therefore, improving the diagnostic rate of early GC is an efficient and feasible method to improve the survival rate of patients with GC.

GC follows the previously described Correa cascade of active gastritis-atrophic gastritis-intestinalization-intraepithelial neoplasia-GC, where atrophic gastritis is considered as the turning point and is referred to as a gastric precancerous lesion. Therefore, effective screening of subjects with gastric precancerous lesions and appropriate intervention can reduce the incidence of GC and increase the diagnostic rate of early GC.

In 2005, the International Atrophy Study Group proposed the Operative Link for Gastritis Assessment (OLGA) grading and staging system for chronic gastritis[3]. This staging system represents the extent and degree of gastric mucosal atrophy to link the histopathology of chronic gastritis to the risk of GC. In terms of pathological diagnosis, the consistency of gastric mucosal atrophy diagnosis among different physicians is low, while that of intestinal metaplasia is high. Therefore, in 2010, the standard Operative Link on Gastric Intestinal Metaplasia Assessment (OLGIM) staging system put forth the proposal to replace the term "atrophy" with "intestinal metaplasia"[4]. Several studies have now confirmed that the OLGA/OLGIM staging system, reflecting the severity of atrophic gastritis and the risk of GC, facilitates the identification of patients with a high risk of developing GC (OLGA/OLGIM stages III and IV), thus promoting the early diagnosis and prevention of the disease[5,6].

Endoscopy is an invasive diagnostic approach that requires a large number of samples, and its diagnostic accuracy is affected by the practicing pathologists themselves. Therefore, endoscopy is not considered as an ideal method in clinical practice, particularly for the physical examination of healthy individuals. Never-

theless, screening for atrophy and risk of GC in healthy individuals using non-invasive approaches is of great importance.

Serum levels of pepsinogen (PG) can reflect the morphology and function of the gastric mucosa. PGI reflects the function of gastric acid secretion of the gastric gland. Therefore, it has been reported that the levels of PGI are reduced when gastric mucosal glands are atrophied. Additionally, the PGII serum levels are mainly associated with the extent of lesions in the gastric mucosa of the gastric fundus. Increased PGII serum levels are associated with gastric fundus atrophy, intestinal metaplasia, or pseudo-pyloric metaplasia and dysplasia. Several studies have demonstrated that the PGI/PGII ratio (PGR) was associated with progressive gastric mucosal atrophy and GC[7,8].

The current study aimed to evaluate the diagnostic value of the serum PG levels and the presence of *Helicobacter pylori* (*H. pylori*) infection as a screening tool for atrophic gastritis in asymptomatic Chinese who attend standard health check-up in China.

MATERIALS AND METHODS

Study population

A total of nine health management centers from different areas of China participated in the study, which was conducted between October 2016 and October 2018. The centers were the No. 924 Hospital of the People's Liberation Army of China (southern China), the Second Affiliated Hospital of Zhejiang University College of Medicine, the Zhongshan Affiliated Hospital of Xiamen University, the Kunshan Hospital of Traditional Chinese Medicine (all in eastern China), the Sichuan Provincial People's Hospital, the Southwest Affiliated Hospital of the Third Military Medical University (both in southwest China), the Jilin City People's Hospital (northeast China), the PLA General Hospital, and the Jingzhou Hospital of Traditional Chinese Medicine (both in central-northern China). More specifically, subjects between 20-80 years of age, without gastrointestinal symptoms, and who presented for asymptomatic health check-up that included transnasal endoscopy were enrolled in the present study.

Participants with one of the following symptoms or clinical findings were excluded from the present study: (1) Subjects with contraindications to transnasal gastroscopy; (2) Previous history of definitive benign or malignant diseases of the upper gastrointestinal tract, including peptic ulcers, gastric polyps, esophageal cancer, and GC, or history of surgery; (3) Treatment with acid suppressants within the past month; (4) Serious organic diseases, including heart, liver, and kidney diseases; (5) Severe mental illness precluding an ability to cooperate; or (6) Current pregnancy or lactation.

Determination of serum PG levels, *H. pylori* test, and endoscopy

All participants received serum PG testing, *H. pylori* testing, and transnasal endoscopy in a single-day hospital visit. Approximately 5 mL of fasting blood was collected from each subject. Serum PG levels were measured by particle-based chemiluminescence immunoassay using the PGI and PGII kits (Abbott Laboratories Inc., Chicago, IL, United States). *H. pylori* infection was evaluated using a ¹³C-urea breath test (Shenzhen Zhonghe Headway Bio-Sci & Tech Co., Ltd., Shenzhen, China) or serum *H. pylori* antibodies (MP Biomedicals, Santa Ana, CA, United States) in combination with pathological *H. pylori* findings. Individuals with a positive result for any of the aforementioned three tests were categorized into a *H. pylori*-positive group, while those with a negative result for all tests were categorized into a *H. pylori*-negative group. Endoscopy was performed by experienced endoscopists who were blinded to the *H. pylori* and PG test results. Pathological samples were obtained from the gastric body, angulus, and antrum according to the Chinese Consensus on Screening and Endoscopic Diagnosis and Treatment of Early GC (2014 version)[9]. Additional biopsies were performed at the sites where lesions were detected.

Statistical analysis

The data are expressed as the mean \pm SD for normally distributed data and as the median \pm interquartile range for non-normally distributed data. Statistical analyses were performed using SPSS statistical software (version 20; IBM Corp., Armonk, NY, United States). Age, PGI and PGII levels, and PGR were tested for normal distribution by the Kolmogorov-Smirnov test. After confirming a normal distribution, ANOVA with *post hoc* Scheffe's test was used for further assessment. Bonferroni's correction and Pearson's chi-square test were used to evaluate the differences between *H. pylori*-negative and *H. pylori*-positive patients. Receiver operating characteristic (ROC) curve

analysis was used to estimate the cutoff values for PGI and PGR. *P* values less than 0.05 were considered statistically significant.

RESULTS

Among the total 2256 subjects included in the current study, 14 were diagnosed with GC (0.6%), including six cases of early GC. In addition, one subject was diagnosed with pharyngeal cancer. There were 326 cases (15.9%) of atrophic gastritis and 391 cases (19.0%) of intestinal metaplasia, while the total *H. pylori* infection rate was 41.7%. A total of 1922 subjects, with an average age of 52.3 ± 9.8 years and male to female ratio of 1.2:1 (1065/857), underwent serological, gastroscopic, and pathological examinations (Table 1).

Based on OLGA grading, the PGI serum levels were slightly decreased, while those of PGII were slightly increased. In addition, the PGR was reduced in conjunction with increasing degrees of atrophy. Compared with the OLGA-0 group, the PGII levels were slightly increased and PGR was slightly decreased in the OLGA-I to IV groups ($P < 0.05$). Additionally, compared with the OLGA-I and II groups, PGR levels in the OLGA-III and IV groups were significantly reduced ($P < 0.05$). A statistically significant difference was also observed in the PGR between the OLGA-II and OLGA-III/IV groups ($P < 0.05$). A notable difference in PGR was also identified between the GC group and both OLGA-0 and I groups ($P < 0.05$; Table 2).

Furthermore, as shown in Table 2, compared with the OLGIM-0 group, the PGI levels were significantly lower in the OLGIM-II, III, and IV groups ($P < 0.05$), and PGR was reduced in the OLGIM-II and III groups ($P < 0.05$). The PGI serum levels were significantly lower in the OLGIM-III/IV groups, and PGR was also markedly decreased in the OLGIM-II/III groups compared with the OLGIM-I group ($P < 0.05$). Finally, compared with the OLGIM-II group, the OLGIM-III group exhibited significantly reduced PGR ($P < 0.05$; Table 3).

The aforementioned results indicated that the PG serum levels were more relevant to OLGA grade compared with the OLGIM grade. Therefore, the OLGA grading system was applied for the subsequent analyses.

Rugge *et al.*'s [10] prospective study confirmed that the high-risk stage (defined as stage III or IV by the OLGA classification) is closely associated with high risk of GC. Accordingly, OLGA I and II groups were then combined in a low-risk group and compared with the OLGA-0 group. The best cutoff for low risk was estimated at PGI 73.14 ng/mL [area under the curve (AUC) = 0.585, sensitivity = 62.1%, specificity = 53.8%] and PGR 11.54 (AUC = 0.611, sensitivity = 43.2%, specificity = 77.7%) (Figure 1). The OLGA III and IV groups were combined in a high-risk group and compared with the remaining three groups; the best cutoff for high risk was estimated at PGI 64.00 ng/mL (AUC = 0.631, sensitivity = 67.2%, specificity = 61.2%) and PGR 9.11 (AUC = 0.740, sensitivity = 53.0%, specificity = 91.8%) (Figure 2).

Compared with the OLGA-0 group, the mean ages in the OLGA-I, III, and IV groups were significantly higher ($P < 0.05$) and increased with an increasing OLGA grade. In addition, a statistically significant difference was observed in the *H. pylori* positivity rate between the OLGA-I/II/III/IV groups and the OLGA-0 group ($P < 0.05$; Table 4).

Further analyses were performed with the *H. pylori*-negative or positive groups. In the OLGA-0 group, *H. pylori* positive subjects exhibited significantly higher PGI and PGII serum levels and a significantly lower PGR compared with *H. pylori*-negative patients ($P < 0.05$). The same trend was observed among *H. pylori*-positive/*H. pylori*-negative subjects in the OLGA-I group ($P < 0.05$). Additionally, *H. pylori*-positive subjects in the OLGA-II group had significantly elevated PGI and PGII levels compared with the *H. pylori*-negative subjects ($P < 0.05$). In the OLGA-III group, significantly increased serum levels of PGII were observed in the *H. pylori*-positive group compared with the *H. pylori*-negative group, while the ratio PGI/PGII was notably reduced. By contrast, no statistically significant differences were obtained in the PG levels between the *H. pylori*-positive/*H. pylori*-negative subjects in the OLGA-IV and GC groups ($P > 0.05$).

In the *H. pylori*-positive population, compared with the OLGA-0 group, both PGI levels and PGR were significantly lower in the OLGA-I and III groups ($P < 0.05$), and PGR was also notably reduced lower in the OLGA-II group ($P < 0.05$). In the *H. pylori*-negative population, the serum levels of PGI and PGR were markedly decreased in the OLGA-I/II ($P < 0.05$) and GC groups ($P < 0.05$; Table 5).

Table 1 Baseline characteristics of the 1922 subjects involved in the study, *n* (%)

Age	52.3 ± 9.8
Gender (male)	1065 (55.41)
<i>Helicobacter pylori</i>	757(39.39)
Gastric cancer	10 (0.52)

Table 2 Serum pepsinogen levels in different OLGA groups

OLGA grade	<i>n</i>	PGI (ng/mL)	PGII (ng/mL)	PGR
0	1590	90.4 ± 82.2	8.3 ± 8.6	10.1 ± 9.3
I	185	68.7 ± 71.2 ¹	7.8 ± 8.5	8.1 ± 5.6 ¹
II	88	73.7 ± 80.7 ¹	9.8 ± 8.8	7.5 ± 4.6 ¹
III	43	60.8 ± 51.1 ¹	11.7 ± 12.8 ^{1,2}	6.7 ± 4.4 ^{1,2,3}
IV	6	52.1 ± 37.3 ¹	15.5 ± 9.9	4.1 ± 3.7 ^{1,2,3}

¹*P* < 0.05, compared with the OLGA-0 group.²*P* < 0.05, compared with the OLGA-I group.³*P* < 0.05, compared with the OLGA-II group.

OLGA: Operative Link for Gastritis Assessment; PG: Pepsinogen; PGR: Pepsinogen I/pepsinogen II ratio.

Table 3 Serum pepsinogen levels in different OLGIM groups

OLGIM grade	<i>n</i>	PGI (ng/mL)	PGII (ng/mL)	PGR
0	1522	91.1 ± 83.8	8.5 ± 8.9	9.9 ± 8.8
I	252	79.2 ± 80.7 ¹	8.1 ± 8.4	9.0 ± 8.3
II	89	66.4 ± 76.5 ¹	8.8 ± 7.6	7.3 ± 6.3 ^{1,2,3}
III	41	56.3 ± 45.1 ^{1,2}	9.5 ± 6.7 ²	6.3 ± 4.1 ^{1,2,3}
IV	8	60.8 ± 26.4 ^{1,2}	8.6 ± 11.7	6.7 ± 18.8

¹*P* < 0.05, compared with the OLGIM-0 group.²*P* < 0.05, compared with the OLGIM-I group.³*P* < 0.05, compared with the OLGIM-II group.

OLGA: Operative Link for Gastritis Assessment; PG: Pepsinogen; PGR: Pepsinogen I/pepsinogen II ratio.

Table 4 Age, sex, and presence of *Helicobacter pylori* infection in different Operative Link for Gastritis Assessment groups, *n* (%)

OLGA grade	0	I	II	III	IV
Sex (male)	866 (54.5)	109 (58.9)	57 (64.8)	25 (58.1)	3 (50.0)
Age	51.9 ± 9.9	54.99 ± 9.1 ¹	53.49 ± 8.8	55.42 ± 9.9 ¹	56.33 ± 8.2 ¹
<i>Helicobacter pylori</i> -positive	583 (36.7)	86 (46.5) ¹	51 (60.0) ¹	27 (62.8) ¹	5 (83.3) ¹

¹*P* < 0.05, compared with the OLGA-0 group.

OLGA: Operative Link for Gastritis Assessment.

DISCUSSION


Atrophic gastritis is considered as one of the main precursor lesions in the Correa cascade, which may eventually result in GC. A study reported that the odds of developing GC 5 years after the diagnosis of atrophic gastritis, enterocolitis, mild to moderate atypical hyperplasia, and severe atypical hyperplasia were 0.1%, 0.25%,

Table 5 Serum pepsinogen levels in the OLGA groups after stratification according to the presence of *Helicobacter pylori* infection

	OLGA-0		I	II		III		IV		
	<i>Helicobacter pylori</i> -(n = 1007)	<i>Helicobacter pylori</i> + (583)	<i>Helicobacter pylori</i> - (n = 99)	<i>Helicobacter pylori</i> + (n = 86)	<i>Helicobacter pylori</i> - (n = 37)	<i>Helicobacter pylori</i> + (n = 51)	<i>Helicobacter pylori</i> - (n = 16)	<i>Helicobacter pylori</i> + (n = 27)	<i>Helicobacter pylori</i> -(n = 1)	<i>Helicobacter pylori</i> + (n = 5)
PGI (ng/mL)	82.18 ± 78.64	101.70 ± 89.01 ¹	56.71 ± 66.10 ²	73.05 ± 78.20 ^{1,2}	55.40 ± 54.18 ²	92.12 ± 73.51 ¹	60.00 ± 47.55	60.8 ± 54.30 ²	34.3	63.50 ± 43.88 ²
PGII (ng/mL)	7.35 ± 6.61	10.80 ± 11.36 ¹	6.60 ± 6.31	10.05 ± 10.23 ¹	7.10 ± 4.88	13.00 ± 10.49 ¹	7.00 ± 6.13	13.5 ± 10.64 ¹	3.8	16.40 ± 8.70
PGR	10.68 ± 8.76	8.82 ± 9.89 ¹	8.77 ± 5.07 ²	7.24 ± 6.28 ^{1,2}	7.94 ± 3.54 ²	7.18 ± 5.51 ²	8.32 ± 3.25	4.50 ± 3.45 ^{1,2}	9.026316	3.90 ± 2.65 ²

¹P < 0.05, within the same pathological subgroup, compared with the *Helicobacter pylori*- group.²P < 0.05, within the same *Helicobacter pylori* subgroup, compared with the OLGA-0 group.

OLGA: Operative Link for Gastritis Assessment; PG: Pepsinogen; PGR: Pepsinogen I/pepsinogen II ratio.

**Figure 1 Operating characteristics curve analysis of pepsinogen I and pepsinogen I/pepsinogen II ratio for the low-risk group.** The Operative Link for Gastritis Assessment (OLGA) I and II groups were combined as the low-risk group and compared with the OLGA-0 group. PG: Pepsinogen; PGR: Pepsinogen I/pepsinogen II ratio; OLGA: Operative Link for Gastritis Assessment.

0.6%, and 6%, respectively[1].

The current study found that approximately 15.9% of the asymptomatic individuals undergoing standard physical examination exhibited atrophic gastritis and 19.0% exhibited enterocolitis. Among these patients, 2.5% suffered from severe atrophic gastritis (OLGA grade III or IV). These patients have a higher risk of progression to GC after 5 years. Therefore, effective screening of such individuals in the asymptomatic population *via* endoscopic surveillance interventions are important for reducing the incidence of GC and improving the diagnostic rate of early GC.

Serum PG levels reflect the morphological and functional status of the gastric mucosa. It has been reported that the gradual decrease of PGR is associated with the progression of gastric mucosal atrophy and GC[2]. A meta-analysis of 31 studies, including 1520 patients with GC and 2265 with atrophic gastritis, suggested that serum PG levels could be considered as a potent, non-invasive, population-based screening tool for the diagnosis of GC and atrophic gastritis[11]. Additionally, a study by Zoalfaghari *et al*[12] indicated that the serum PGI levels and PGR could be potential serological markers for the diagnosis of atrophic gastritis, with a high sensitivity and specificity. The present study also demonstrated that PGI levels and PGR were significantly decreased with increasing atrophy, with PGR showing more significant diagnostic value. The association between PG levels and OLGA grade was more pronounced compared with the OLGA grading system, suggesting that PG could be more closely associated with atrophy compared with intestinal metaplasia. Furthermore, ROC curve analysis showed that serum PGI levels and PGR could be used to predict atrophy and intestinal metaplasia of the gastric mucosa in asymptomatic health check-up subjects in China.




Figure 2 Operating characteristics curve analysis of pepsinogen I and pepsinogen I/pepsinogen II ratio for the high-risk group. The Operative Link for Gastritis Assessment (OLGA) III and IV groups were combined as the high-risk group and compared with OLGA 0, I, and II groups. PG: Pepsinogen; PGR: Pepsinogen I/pepsinogen II ratio; OLGA: Operative Link for Gastritis Assessment.

Several regional studies have demonstrated that the *H. pylori* infection rate is approximately 40%-55% in China. The overall *H. pylori* infection rate in the study population was 39.39%. The results showed that the PGI and PGII serum levels were significantly increased, and PGR was markedly decreased in the *H. pylori*-infected group. *H. pylori* infection also affected the levels of PG in different pathological subgroups[13]. Al-Ezzy *et al*[14] reported that *H. pylori* infection can affect the expression of the *Fas* gene through inflammatory factors, thus regulating PG concentration.

Furthermore, the results of the current study also demonstrated that patients with high-grade lesions were older and had a higher rate of *H. pylori* infection. A positive association has been reported between the severity of gastric atrophy or intestinal metaplasia and the age of the subjects[15,16]. Additionally, a study showed that the prevalence of high-risk OLGA stage was increased with increasing age in *H. pylori*-negative subjects[17]. A Korean study demonstrated that advanced age, long-term smoking, and *H. pylori* infection were independent risk factors associated with advanced stages determined by the OLGA staging system[18]. The transition from *H. pylori* infection into GC is a multistep process, including the progression of chronic gastritis into precancerous lesions and ultimately to GC[19]. The presence or absence of *H. pylori* infection allows a preliminary prediction of whether a patient is at high risk by OLGA staging. Therefore, it has been suggested that routine endoscopy may not be necessary for all patients by OLGA staging or those at a low-risk by OLGIM stage, since *H. pylori* eradication can effectively prevent the development of GC. Effectively eradicating *H. pylori* before the age of 40 years could reduce the number of routine endoscopies in individuals requiring GC surveillance and substantially reduce the related healthcare burden.

The current study has some limitations in. First, the overall sample size in the study was not large enough; in particular, the number of patients with OLGA stage III/IV and GC was insufficient, which could cause statistical bias. Therefore, further studies with increased sample size are needed to verify the results of the present study. Second, there could be a selection bias in this study, since all enrolled subjects voluntarily underwent physical examination and gastroscopy, suggesting that their economic status was moderate.

CONCLUSION

The present study demonstrated that serum PG levels were closely associated with OLGA classification. The results suggested that PG levels could be used as an effective non-invasive screening tool for atrophic gastritis in the asymptomatic population undergoing standard physical examination, while the diagnostic value of PG levels could be more potent in patients with *H. pylori* infection. This screening tool could be valuable and represent an important screening method for the elderly population.

ARTICLE HIGHLIGHTS

Research background

Atrophic gastritis is a precancerous lesion of the stomach. Pepsinogen (PG) has been reported to reflect the morphology and function of the gastric mucosa.

Research motivation

PG can be used for non-invasive screening of atrophic gastritis and even gastric cancer (GC). Effective screening of subjects with gastric precancerous lesions and appropriate intervention can reduce the incidence of GC and increase the diagnostic rate of early GC.

Research objectives

The main objective of this research was to evaluate the diagnostic value of serum PG in the degree of gastric mucosal atrophy in an asymptomatic Chinese population undergoing standard physical examination.

Research methods

The study subjects underwent transnasal gastroscopy, and serum PG levels and the presence of *Helicobacter pylori* (*H. pylori*) infection were investigated to assess the diagnostic accuracy of PG for evaluating the degree of gastric mucosal atrophy. Pathology was evaluated using the Operative Link for Gastritis Assessment (OLGA) classification and Operative Link on Gastric Intestinal Metaplasia Assessment (OLGIM) systems. ANOVA with *post hoc* Scheffe's test was used for further assessment of the differences in age, PGI and PGII levels, and PGI/PGII ratio (PGR). Bonferroni's correction and Pearson's chi-square test were used to evaluate the differences between *H. pylori*-negative and *H. pylori*-positive patients. Receiver operating characteristic curve analysis was used to estimate the cutoff values for PGI, PGII, and PGR.

Research results

The association between PG and OLGA grade was higher compared with that between PG and the OLGIM grading system. Based on the OLGA grading system, the levels of PGI were slightly decreased, while those of PGII were slightly increased. PGR was reduced with increasing atrophy ($P < 0.05$). A slightly increasing trend was observed in the mean age of different OLGA groups. *H. pylori*-positive subjects exhibited significantly higher PGI and PGII serum levels and a significantly lower PGR compared with *H. pylori*-negative patients in the different OLGA groups ($P < 0.05$).

Research conclusions

Serum levels of PG are closely associated with OLGA stage and could be used as an effective non-invasive screening tool for evaluating the degree of gastric mucosal atrophy in asymptomatic subjects.

Research perspectives

Future studies could focus on the cutoff values of PG for the diagnosis of precancerous lesions or early gastric cancer in different regions of China.

REFERENCES

- 1 Yang L, Zheng R, Wang N, Yuan Y, Liu S, Li H, Zhang S, Zeng H, Chen W. Incidence and mortality of stomach cancer in China, 2014. *Chin J Cancer Res* 2018; **30**: 291-298 [PMID: 30046223 DOI: 10.21147/j.issn.1000-9604.2018.03.01]
- 2 Fang WL, Huang KH, Chen MH, Liu CA, Hung YP, Chao Y, Tai LC, Lo SS, Li AF, Wu CW, Shyr YM. Comparative study of the 7th and 8th AJCC editions for gastric cancer patients after curative surgery. *PLoS One* 2017; **12**: e0187626 [PMID: 29131840 DOI: 10.1371/journal.pone.0187626]
- 3 Rugge M, Genta RM. Staging and grading of chronic gastritis. *Hum Pathol* 2005; **36**: 228-233 [PMID: 15791566 DOI: 10.1016/j.humpath.2004.12.008]
- 4 Capelle LG, de Vries AC, Haringsma J, Ter Borg F, de Vries RA, Bruno MJ, van Dekken H, Meijer J, van Grieken NC, Kuipers EJ. The staging of gastritis with the OLGA system by using intestinal metaplasia as an accurate alternative for atrophic gastritis. *Gastrointest Endosc* 2010; **71**: 1150-1158 [PMID: 20381801 DOI: 10.1016/j.gie.2009.12.029]
- 5 Malfertheiner P, Megraud F, O'Morain CA, Atherton J, Axon AT, Bazzoli F, Gensini GF, Gisbert JP, Graham DY, Rokkas T, El-Omar EM, Kuipers EJ; European Helicobacter Study Group.

- Management of *Helicobacter pylori* infection--the Maastricht IV/ Florence Consensus Report. *Gut* 2012; **61**: 646-664 [PMID: [22491499](#) DOI: [10.1136/gutjnl-2012-302084](#)]
- 6 **Sugano K**, Tack J, Kuipers EJ, Graham DY, El-Omar EM, Miura S, Haruma K, Asaka M, Uemura N, Malfertheiner P; faculty members of Kyoto Global Consensus Conference. Kyoto global consensus report on *Helicobacter pylori* gastritis. *Gut* 2015; **64**: 1353-1367 [PMID: [26187502](#) DOI: [10.1136/gutjnl-2015-309252](#)]
 - 7 **Miki K**. Gastric cancer screening by combined assay for serum anti-*Helicobacter pylori* IgG antibody and serum pepsinogen levels - "ABC method". *Proc Jpn Acad Ser B Phys Biol Sci* 2011; **87**: 405-414 [PMID: [21785258](#) DOI: [10.2183/pjab.87.405](#)]
 - 8 **Yamaguchi Y**, Nagata Y, Hiratsuka R, Kawase Y, Tominaga T, Takeuchi S, Sakagami S, Ishida S. Gastric Cancer Screening by Combined Assay for Serum Anti-*Helicobacter pylori* IgG Antibody and Serum Pepsinogen Levels--The ABC Method. *Digestion* 2016; **93**: 13-18 [PMID: [26789514](#) DOI: [10.1159/000441742](#)]
 - 9 **Zou WB**, Yang F, Li ZS. [How to improve the diagnosis rate of early gastric cancer in China]. *Zhejiang Da Xue Xue Bao Yi Xue Ban* 2015; **44**: 9-14 [PMID: [25851969](#)]
 - 10 **Rugge M**, Meggio A, Pravadelli C, Barbareschi M, Fassan M, Gentilini M, Zorzi M, Pretis G, Graham DY, Genta RM. Gastritis staging in the endoscopic follow-up for the secondary prevention of gastric cancer: a 5-year prospective study of 1755 patients. *Gut* 2019; **68**: 11-17 [PMID: [29306868](#) DOI: [10.1136/gutjnl-2017-314600](#)]
 - 11 **Tu H**, Sun L, Dong X, Gong Y, Xu Q, Jing J, Bostick RM, Wu X, Yuan Y. A Serological Biopsy Using Five Stomach-Specific Circulating Biomarkers for Gastric Cancer Risk Assessment: A Multi-Phase Study. *Am J Gastroenterol* 2017; **112**: 704-715 [PMID: [28323271](#) DOI: [10.1038/ajg.2017.55](#)]
 - 12 **Zoalfaghari A**, Aletaha N, Roushan N, Taslimi R, Foroutan H, Faridnia B. Accuracy of pepsinogens for early diagnosis of atrophic gastritis and gastric cancer in Iranian population. *Med J Islam Repub Iran* 2014; **28**: 150 [PMID: [25695008](#)]
 - 13 **De Re V**, Orzes E, Canzonieri V, Maiero S, Fornasari M, Alessandrini L, Cervo S, Steffan A, Zanette G, Mazzon C, De Paoli P, Cannizzaro R. Pepsinogens to Distinguish Patients With Gastric Intestinal Metaplasia and *Helicobacter pylori* Infection Among Populations at Risk for Gastric Cancer. *Clin Transl Gastroenterol* 2016; **7**: e183 [PMID: [27441820](#) DOI: [10.1038/ctg.2016.42](#)]
 - 14 **Al-Ezzy AI**. Immunomodulatory Effect of *H. Pylori CagA* Genotype and Gastric Hormones On Gastric Versus Inflammatory Cells Fas Gene Expression in Iraqi Patients with Gastroduodenal Disorders. *Open Access Maced J Med Sci* 2016; **4**: 364-373 [PMID: [27703557](#) DOI: [10.3889/oamjms.2016.032](#)]
 - 15 **Graham DY**, Nurgalieva ZZ, El-Zimaity HM, Opekun AR, Campos A, Guerrero L, Chavez A, Cardenas V. Noninvasive vs histologic detection of gastric atrophy in a Hispanic population in North America. *Clin Gastroenterol Hepatol* 2006; **4**: 306-314 [PMID: [16527693](#) DOI: [10.1016/j.cgh.2005.11.003](#)]
 - 16 **Rokkas T**, Rokka A, Portincasa P. A systematic review and meta-analysis of the role of *Helicobacter pylori* eradication in preventing gastric cancer. *Ann Gastroenterol* 2017; **30**: 414-423 [PMID: [28655977](#) DOI: [10.20524/aog.2017.0144](#)]
 - 17 **Nam JH**, Choi IJ, Kook MC, Lee JY, Cho SJ, Nam SY, Kim CG. OLGA and OLGIM stage distribution according to age and *Helicobacter pylori* status in the Korean population. *Helicobacter* 2014; **19**: 81-89 [PMID: [24617667](#) DOI: [10.1111/hel.12112](#)]
 - 18 **Kim Y**, Jun JK, Choi KS, Lee HY, Park EC. Overview of the National Cancer screening programme and the cancer screening status in Korea. *Asian Pac J Cancer Prev* 2011; **12**: 725-730 [PMID: [21627372](#)]
 - 19 **Bornschein J**, Kandulski A, Selgrad M, Malfertheiner P. From gastric inflammation to gastric cancer. *Dig Dis* 2010; **28**: 609-614 [PMID: [21088411](#) DOI: [10.1159/000320061](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

