

World Journal of *Clinical Cases*

World J Clin Cases 2021 November 26; 9(33): 10052-10391



Contents

Thrice Monthly Volume 9 Number 33 November 26, 2021

REVIEW

- 10052** Effects of alcohol consumption on viral hepatitis B and C

Xu HQ, Wang CG, Zhou Q, Gao YH

MINIREVIEWS

- 10064** Effects of anti-diabetic drugs on sarcopenia: Best treatment options for elderly patients with type 2 diabetes mellitus and sarcopenia

Ma XY, Chen FQ

ORIGINAL ARTICLE

Retrospective Cohort Study

- 10075** Utility of cooling patches to prevent hand-foot syndrome caused by pegylated liposomal doxorubicin in breast cancer patients

Zheng YF, Fu X, Wang XX, Sun XJ, He XD

Retrospective Study

- 10088** Clinicopathological features of small T1 colorectal cancers

Takashina Y, Kudo SE, Ichimasa K, Kouyama Y, Mochizuki K, Akimoto Y, Maeda Y, Mori Y, Misawa M, Ogata N, Kudo T, Hisayuki T, Hayashi T, Wakamura K, Sawada N, Baba T, Ishida F, Yokoyama K, Daita M, Nemoto T, Miyachi H

- 10098** Comparison of dental pulp periodontal therapy and conventional simple periodontal therapy as treatment modalities for severe periodontitis

Li L, Chen HJ, Lian Y, Wang T

- 10106** Tripartite intensive intervention for prevention of rebleeding in elderly patients with hypertensive cerebral hemorrhage

Li CX, Li L, Zhang JF, Zhang QH, Jin XH, Cai GJ

- 10116** Clinical and electroencephalogram characteristics and treatment outcomes in children with benign epilepsy and centrotemporal spikes

Chen RH, Li BF, Wen JH, Zhong CL, Ji MM

- 10126** Endoscopic ultrasonography diagnosis of gastric glomus tumors

Bai B, Mao CS, Li Z, Kuang SL

- 10134** Learning curves of robot-assisted pedicle screw fixations based on the cumulative sum test

Yu J, Zhang Q, Fan MX, Han XG, Liu B, Tian W

- 10143** Value of GRACE and SYNTAX scores for predicting the prognosis of patients with non-ST elevation acute coronary syndrome

Wang XF, Zhao M, Liu F, Sun GR

- 10151** Effectiveness of enhanced recovery after surgery in the perioperative management of patients with bone surgery in China

Zhao LY, Liu XT, Zhao ZL, Gu R, Ni XM, Deng R, Li XY, Gao MJ, Zhu WN

Clinical Trials Study

- 10161** Association between plasma dipeptidyl peptidase-4 levels and cognitive function in perinatal pregnant women with gestational diabetes mellitus

Sana SRGL, Li EY, Deng XJ, Guo L

- 10172** Paricalcitol in hemodialysis patients with secondary hyperparathyroidism and its potential benefits

Chen X, Zhao F, Pan WJ, Di JM, Xie WN, Yuan L, Liu Z

Observational Study

- 10180** Did the severe acute respiratory syndrome-coronavirus 2 pandemic cause an endemic *Clostridium difficile* infection?

Cojocariu C, Girleanu I, Trifan A, Olteanu A, Muzica CM, Huiban L, Chiriac S, Singeap AM, Cuciureanu T, Sfarti C, Stanciu C

- 10189** Effect of nursing intervention based on Maslow's hierarchy of needs in patients with coronary heart disease interventional surgery

Xu JX, Wu LX, Jiang W, Fan GH

- 10198** Impacts of statin and metformin on neuropathy in patients with type 2 diabetes mellitus: Korean Health Insurance data

Min HK, Kim SH, Choi JH, Choi K, Kim HR, Lee SH

META-ANALYSIS

- 10208** Is endoscopic retrograde appendicitis therapy a better modality for acute uncomplicated appendicitis? A systematic review and meta-analysis

Wang Y, Sun CY, Liu J, Chen Y, Bhan C, Tuason JPW, Misra S, Huang YT, Ma SD, Cheng XY, Zhou Q, Gu WC, Wu DD, Chen X

- 10222** Prognostic value of ground glass opacity on computed tomography in pathological stage I pulmonary adenocarcinoma: A meta-analysis

Pan XL, Liao ZL, Yao H, Yan WJ, Wen DY, Wang Y, Li ZL

CASE REPORT

- 10233** Atrial fibrillation and concomitant left subclavian, axillary and brachial artery embolism after fiberoptic bronchoscopy: A case report

Yang CL, Zhou R, Jin ZX, Chen M, Zi BL, Li P, Zhou KH

- 10238** Streptococcal toxic shock syndrome after hemorrhoidectomy: A case report

Lee CY, Lee YJ, Chen CC, Kuo LJ

- 10244** Subsequent placenta accreta after previous mifepristone-induced abortion: A case report

Zhao P, Zhao Y, He J, Bai XX, Chen J

- 10249** Autosomal dominant tubulointerstitial kidney disease with a novel heterozygous missense mutation in the uromodulin gene: A case report
Zhang LL, Lin JR, Zhu TT, Liu Q, Zhang DM, Gan LW, Li Y, Ou ST
- 10257** Novel KDM6A mutation in a Chinese infant with Kabuki syndrome: A case report
Guo HX, Li BW, Hu M, Si SY, Feng K
- 10265** Pancreatic cancer with synchronous liver and colon metastases: A case report
Dong YM, Sun HN, Sun DC, Deng MH, Peng YG, Zhu YY
- 10273** Veno-venous-extracorporeal membrane oxygenation treatment for severe capillary leakage syndrome: A case report
Nong WX, Lv QJ, Lu YS
- 10279** Anticoagulant treatment for pulmonary embolism in patient with cerebral hemorrhage secondary to mechanical thrombectomy: A case report
Chen XT, Zhang Q, Zhou CQ, Han YF, Cao QQ
- 10286** Complete restoration of congenital conductive hearing loss by staged surgery: A case report
Yoo JS, Lee CM, Yang YN, Lee EJ
- 10293** Blastic plasmacytoid dendritic cell neoplasm with skin and bone marrow involvement: Report of three cases
Guo JH, Zhang HW, Wang L, Bai W, Wang JF
- 10300** Extracranial multiorgan metastasis from primary glioblastoma: A case report
Luan XZ, Wang HR, Xiang W, Li SJ, He H, Chen LG, Wang JM, Zhou J
- 10308** Transverse myelitis after infection with varicella zoster virus in patient with normal immunity: A case report
Yun D, Cho SY, Ju W, Seo EH
- 10315** Duodenal ulcer caused by coil wiggle after digital subtraction angiography-guided embolization: A case report
Xu S, Yang SX, Xue ZX, Xu CL, Cai ZZ, Xu CZ
- 10323** Crab lice infestation in unilateral eyelashes and adjacent eyelids: A case report
Tang W, Li QQ
- 10328** Local random flaps for cervical circumferential defect or tracheoesophageal fistula reconstruction after failed gastric pull-up: Two case reports
Zhang Y, Liu Y, Sun Y, Xu M, Wang XL
- 10337** Incurable and refractory spinal cystic echinococcosis: A case report
Zhang T, Ma LH, Liu H, Li SK
- 10345** Individualized treatment of breast cancer with chronic renal failure: A case report and review of literature
Cai JH, Zheng JH, Lin XQ, Lin WX, Zou J, Chen YK, Li ZY, Chen YX

- 10355** Persistent fibrinogen deficiency after snake bite: A case report
Xu MH, Li J, Han L, Chen C
- 10362** Successful prolonged cardiopulmonary resuscitation after intraoperative cardiac arrest due to povidone-iodine allergy: A case report
Xiang BB, Yao YT, Jiao SL
- 10369** Clinical algorithm for preventing missed diagnoses of occult cervical spine instability after acute trauma: A case report
Zhu C, Yang HL, Im GH, Liu LM, Zhou CG, Song YM
- 10374** Carbon ion radiotherapy for synchronous choroidal melanoma and lung cancer: A case report
Zhang YS, Hu TC, Ye YC, Han JH, Li XJ, Zhang YH, Chen WZ, Chai HY, Pan X, Wang X, Yang YL
- 10382** Heart failure as an adverse effect of infliximab for Crohn's disease: A case report and review of the literature
Grillo TG, Almeida LR, Beraldo RF, Marcondes MB, Queiróz DAR, da Silva DL, Quera R, Baima JP, Saad-Hossne R, Sasaki LY

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Jian-Wu Zhao, PhD, Chief Physician, Professor, Department of Orthopedics, Jilin University Second Hospital, Changchun 130000, Jilin Province, China. jianwu@jlu.edu.cn

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for WJCC as 1.337; IF without journal self cites: 1.301; 5-year IF: 1.742; Journal Citation Indicator: 0.33; Ranking: 119 among 169 journals in medicine, general and internal; and Quartile category: Q3. The WJCC's CiteScore for 2020 is 0.8 and Scopus CiteScore rank 2020: General Medicine is 493/793.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Ji-Hong Lin; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lai Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Dennis A Bloomfield, Sandro Vento, Bao-Gan Peng

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

November 26, 2021

COPYRIGHT

© 2021 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Is endoscopic retrograde appendicitis therapy a better modality for acute uncomplicated appendicitis? A systematic review and meta-analysis

Ying Wang, Chen-Yu Sun, Jie Liu, Yue Chen, Chandur Bhan, John Pocholo Whitaker Tuason, Sudha Misra, Yu-Ting Huang, Shao-Di Ma, Xing-Yu Cheng, Qin Zhou, Wen-Chao Gu, Dan-Dan Wu, Xia Chen

ORCID number: Ying Wang 0000-0002-8983-1307; Chen-Yu Sun 0000-0003-3812-3164; Jie Liu 0000-0001-6079-7566; Yue Chen 0000-0002-2502-9518; Chandur Bhan 0000-0002-2741-9798; John Pocholo Whitaker Tuason 0000-7689-0987-2349; Sudha Misra 0000-0002-5218-2753; Yu-Ting Huang 0000-0001-9986-5124; Shao-Di Ma 0000-0003-1930-3936; Xing-Yu Cheng 0000-0001-8803-4261; Qin Zhou 0000-0003-4177-6289; Wen-Chao Gu 0000-0002-1505-9887; Dan-Dan Wu 0000-0003-4171-9751; Xia Chen 0000-0003-1479-9802.

Author contributions: Wu DD, Chen X, Wang Y, Liu J and Sun CY designed the research study; Wang Y, Sun CY and Chen Y selected and collected the data; Wang Y, Sun CY and Liu J analyzed the data; Bhan C, Tuason JPW, Misra S, Huang YT, Ma SD, Cheng XY, Zhou Q and Gu WC provided critical opinions and revised the manuscript; Wang Y and Sun CY wrote the manuscript; Wang Y, Sun CY and Liu J contributed equally to this work and should be considered as co-first authors; all authors approved the final manuscript.

Conflict-of-interest statement:

All authors have no conflict(s) of interest to declare in relation to this manuscript.

Ying Wang, Dan-Dan Wu, Department of Endoscopy Center, The First Affiliated Hospital of USTC, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei 230001, Anhui Province, China

Chen-Yu Sun, Chandur Bhan, John Pocholo Whitaker Tuason, Sudha Misra, Internal Medicine, AMITA Health Saint Joseph Hospital Chicago, Chicago, IL 60657, United States

Jie Liu, Department of Gastroenterology, The First Affiliated Hospital of USTC, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei 230001, Anhui Province, China

Yue Chen, Xing-Yu Cheng, Department of Clinical Medicine, School of the First Clinical Medicine, Anhui Medical University, Hefei 230032, Anhui Province, China

Yu-Ting Huang, University of Maryland Medical Center Midtown Campus, Baltimore, MD 21201, United States

Shao-Di Ma, Department of Epidemiology and Health Statistics, School of Public Health Anhui Medical University, Hefei 230032, Anhui Province, China

Qin Zhou, Department of Radiation Oncology, Mayo Clinic, Rochester, MN 55905, United States

Wen-Chao Gu, Department of Diagnostic Radiology and Nuclear Medicine, Gunma University Graduate School of Medicine, Maebashi 371-8511, Japan

Xia Chen, Department of Nursing, The First Affiliated Hospital of USTC, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei 230001, Anhui Province, China

Corresponding author: Wu Dan-Dan, MSN, RN, Associate Chief Nurse, Department of Endoscopy Center, The First Affiliated Hospital of USTC, Division of Life Sciences and Medicine, University of Science and Technology of China, No. 17 Lujiang Road, Hefei 230001, Anhui Province, China. 16013255@qq.com

PRISMA 2009 Checklist statement:

The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised in accordance with this checklist.

Country/Territory of origin: China

Specialty type: Gastroenterology and hepatology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Received: June 5, 2021

Peer-review started: June 5, 2021

First decision: June 27, 2021

Revised: August 1, 2021

Accepted: October 14, 2021

Article in press: October 14, 2021

Published online: November 26, 2021

P-Reviewer: Hosseini MS, Maslennikov R

S-Editor: Wang LL

L-Editor: A

P-Editor: Wang LL

**Abstract****BACKGROUND**

Previous studies had shown endoscopic retrograde appendicitis therapy (ERAT) is an effective treatment for acute appendicitis. However, different studies reported conflicting outcomes regarding the effectiveness of ERAT in comparison with laparoscopic appendectomy (LA).

AIM

To compare the effectiveness of ERAT with LA.

METHODS

Randomized controlled trials (RCTs) and retrospective studies of ERAT for acute uncomplicated appendicitis were searched in PubMed, Cochrane Library, Web of Science, Embase database, China National Knowledge Infrastructure (CNKI), the WanFang Database, and Chinese Scientific Journals Database (VIP) from the establishment date to March 1 2021. Heterogeneity was assessed using the I-squared statistic. Pooled odds ratios (OR), weighted mean difference (WMD), and standard mean difference (SMD), with 95% confidence intervals (CI) were calculated through either fixed-effects or random-effects model. Sensitivity analysis was also performed. Publication bias was tested by Egger's test, and Begg's test. The quality of included RCT were evaluated by the Jadad scale, while Newcastle-Ottawa scale is adopted for assessing the methodological quality of case-control studies. All statistical analysis was performed using Stata 15.1 statistical software. All statistical analysis was performed using Stata 15.1 statistical software. This study is registered with PROSPERO, CRD42021243955.

RESULTS

After screening, 10 RCTs and 2 case-control studies were included in the current systematic review. Firstly, the length of hospitalizations [WMD = -1.15, 95%CI: -1.99, -0.31; $P = 0.007$] was shorter than LA group. Secondly, the level of postoperative CRP [WMD = -10.06, 95%CI: (-17.39, -2.73); $P = 0.007$], TNF- α [WMD = -7.70, 95%CI: (-8.47, -6.93); $P < 0.001$], and IL-6 Levels [WMD = -9.78, 95%CI: (-10.69, -8.88); $P < 0.001$; $P < 0.001$] in ERAT group was significantly lower than LA group. Thirdly, ERAT group had a lower incidence of intestinal obstruction than LA group. [OR = 0.19, 95%CI: (0.05, 0.79); $P = 0.020$]. Moreover, the quality of 10 RCTs were low with 0-3 Jadad scores, while the methodological quality of two case-control studies were fair with a score of 2 (each).

CONCLUSION

Compared with LA, ERAT reduces operation time, the level of postoperative inflammation, and results in fewer complications and shorter recovery time, with preserving the appendix and its immune and biological functions.

Key Words: Endoscopic retrograde appendicitis therapy; Acute appendicitis; Meta analysis; Laparoscopic appendectomy; Randomized controlled study

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Acute appendicitis is one of the common surgical emergencies all over the world, with a mean cost of about \$9000 per procedure. It is recognized that the conventional treatment of acute appendicitis was laparoscopic appendectomy (LA), while an increasing number of surgical complications, include bleeding, adhesive intestinal obstruction, infection of the incision, and intestinal fistula, have been reported. Therefore, we conducted a meta-analysis to compare the effectiveness of endoscopic retrograde appendicitis therapy (ERAT) with standard treatment. After screening, 12 studies were included in the current systematic review and we found that, compared with LA, ERAT reduces operation time, the level of postoperative inflammation, and results in fewer complications and shorter recovery time, with preserving the appendix and its immune and biological functions.

Citation: Wang Y, Sun CY, Liu J, Chen Y, Bhan C, Tuason JPW, Misra S, Huang YT, Ma SD, Cheng XY, Zhou Q, Gu WC, Wu DD, Chen X. Is endoscopic retrograde appendicitis therapy a better modality for acute uncomplicated appendicitis? A systematic review and meta-analysis. *World J Clin Cases* 2021; 9(33): 10208-10221

URL: <https://www.wjgnet.com/2307-8960/full/v9/i33/10208.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v9.i33.10208>

INTRODUCTION

Acute appendicitis is one of the common surgical emergencies all over the world, with a mean cost of about \$9000 per procedure[1,2]. Appendicitis is one of the most frequent specific underlying causes in patients presenting to emergency departments with abdominal pain[3,4]. The majority (approximately 70%-80%) of acute appendicitis cases are of uncomplicated nature[5,6]. It is reported that the incidence of appendicitis is rising, which is about 1 per 1,000 in the America[7,8]. At present, the etiology of acute appendicitis is still unknown. Common etiological factors, including luminal obstruction from appendiceal fecalith, stool, lymphoid hyperplasia, and neoplasm result in about half of the cases, with stool and appendiceal fecalith as more common causes[9].

LA is currently widely applied for the treatment of acute appendicitis. Although patients could benefit from LA with a decreased wound infection rate, shorter hospital stay, and better diagnostic power[10], some complications can not be ignored. Liang TJ *et al*[11] investigated 864 patients who developed acute appendicitis recurrence in a median follow-up of 6.5 years. The authors found that 258 patients were performed LA, which accounted for about 30%. What's more, an increasing number of surgical complications after LA, including bleeding, adhesive intestinal obstruction, infection of the incision, appendiceal remnants, and intestinal fistula[12].

In 2012, Liu *et al*[13] proposed a new endoscopic minimally invasive treatment for appendicitis, namely Endoscopic retrograde appendicitis therapy (ERAT). After preoperative bowel preparation, the appendix was intubated through the colonoscopy with a transparent cap at the head end, and the diagnosis of appendicitis was confirmed by angiography under X-ray monitoring. It can also relieve the obstruction of the appendix lumen, drain the pus, and flush the lumen to control the inflammation. It also allows the placement of drainage tube into the lumen to ensure the smooth drainage through the appendiceal orifice, reduce the risk of recurrence of appendicitis caused by obstruction.

Previous studies had shown ERAT as an effective treatment for acute appendicitis complicated with local perforation and/or periappendiceal abscess[14]. However, different studies reported conflicting outcomes regarding the effectiveness of ERAT in comparison with LA. Therefore, we conducted a meta-analysis to compare the effectiveness of ERAT with LA for adults.

MATERIALS AND METHODS

Preferred reporting items for systematic reviews and meta-analyses

The Preferred Reporting Items declared by the Systematic Review and Meta-Analysis (PRISMA)[15] was utilized in the performance of this study. The databases including PubMed, Cochrane Library, Web of Science, Embase database, China National Knowledge Infrastructure (CNKI), the WanFang Database, and Chinese Scientific Journals Database (VIP), were searched by using the searching terms including acute appendicitis (acute uncomplicated appendicitis) and endoscopic retrograde appendicitis therapy [endoscopic retrograde appendiceal radiography (ERAR), endoscopic appendiceal irrigation (EAI), and endoscopic appendiceal stent placement (ERSP)]. By taking the retrieval in PubMed as an example, the concrete retrieval strategies are as follows: (acute appendicitis [Mesh Terms] OR acute appendicitis [Title/Abstract] OR acute uncomplicated appendicitis [Mesh Terms] OR acute uncomplicated appendicitis [Title/Abstract]) AND (endoscopic retrograde appendicitis therapy [Mesh Terms] OR endoscopic retrograde appendicitis therapy [Title/Abstract] OR endoscopic retrograde appendiceal radiography [Mesh Terms] OR endoscopic appendiceal irrigation [Title/Abstract] OR endoscopic appendiceal stent placement [Title/Abstract]).

The retrieval time of each database is from the establishment of the database to March 1, 2021. The reference of related literatures and reviews were also retrieved manually to ensure that there was no omission, and the prospective study of ERAT on acute appendicitis published in the literatures are statistically analyzed. The protocol of this systematic review and meta-analysis has already prospectively registered in the PROSPERO (International Prospective Register of Systematic Reviews) database (reference no. CRD42021243955).

Study selection

Studies that met the following criteria were considered to be eligible for inclusion: (1) Study design: Randomized controlled trials, retrospective studies, and prospective studies; (2) Patients: The subjects were clinically diagnosed as acute uncomplicated appendicitis patients; (3) Outcomes: Literatures should provide accurate comprehensive statistical indicators: Sample Size, length of hospitalizations, operation time, recovery time, length of hospitalization, risk of complications; (4) Intervention and control: Intervention was endoscopic retrograde appendicitis therapy, while control group receiving LA; and (5) Articles published in English or Chinese. Exclusion criteria: (1) Duplicate publications; (2) Studies without sufficient data; and (3) Case reports, meta-analysis and reviews, study without English abstract and studies only with abstract were also excluded.

Literature quality evaluation and data extraction

Literature screened by two reviewers independently according to the inclusion and exclusion criteria mentioned above. Any disagreements were resolved through discussion with a third reviewer to reach a consensus. The following data were extracted: first author's name, the time of publication, the type of appendicitis, the participants of the experimental and control group, interventions, and outcomes (the bed rest time, time interval of body temperature returning to normal range, and time interval of white blood cell count returning to normal range, *et al*). Included RCT studies were evaluated by the Jadad scale regarding quality and methodology, where a higher score (total score of seven) suggests more rigorousness of a trial's methodological design[16]. For both case-control and cohort studies, Newcastle-Ottawa scale [17] is adopted for assessing the methodological quality, which provides a comprehensive score system with eight items.

Statistical analysis

Heterogeneity test was performed with Stata 15.0 statistical software (Stata Corp., College Station, TX). The bed rest time, body temperature return to normal time and white blood cells return to normal time were combined by standard mean difference (SMD) with 95%CI, while duration of operation, length of hospitalizations, and levels of inflammatory factors were combined by weighted mean difference (WMD) with 95%CI. Q-test and I²-test were used to analyze the heterogeneity of the studies included in this meta-analysis. If $P > 0.100$ and $I^2 < 50\%$, it was considered that there was small heterogeneity among the studies, and fixed effect model was chosen; otherwise, random effect model was used to merge SMD with 95%CI[18]. The pooled relative risk (RR) with 95%CI: Was performed to analyze the risk of complications. Data of the outcomes were recorded for this meta-analysis when three or more trials reported the same outcome. Sensitivity analyses were performed to investigate the robustness of this meta-analysis. Meanwhile, the risk of publication bias was evaluated by Egger's test, Begg's test, and funnel plots[19]. If the heterogeneity shown $P < 0.100$ and $I^2 > 50\%$, considered that there was large heterogeneity among the studies. Egger's test was assessed by using Stata 15.0.

RESULTS

From the 1,013 relevant records initially identified, 696 remained after excluding duplicates. Then 143 articles were excluded after subsequent scanning of the titles and abstracts. Full texts of the 161 records remained were scrutinized, and 12 studies[20-31] that met the inclusion criteria were selected in systematic review, while 8 studies[21-24,26,28,30,31] were included in meta analysis. The flow of selecting included studies was shown in Figure 1. The 12 included articles with 970 subjects were published between 2016 and 2020 and included 2 case-control[27,31] studies, and 10 RCTs. More detailed characteristics were summarized in Table 1. The Jadad scores of 10 included studies were 0-3 scores. Meanwhile, the methodological quality of two case-control

Table 1 Detailed characteristics of included studies in this meta analysis


Ref.	Studies types	Patients age	Treatment		Sample size		Disease	Outcomes
			Experiment	Control	Experiment	Control		
Kang <i>et al</i> [20], 2020	RCT	1 to 13 years old	Modified ERAT	Antibiotics treatment	36	47	Acute uncomplicated appendicitis in children	Length of hospital stay
Deng <i>et al</i> [21], 2018	RCT	18-62 years old	ERAT	Laparoscopic appendectomy	20	20	Acute appendicitis	Duration of operation, Bed rest time; time interval of body temperature returning to normal range; time interval of white blood cells count returning to normal time range, complication
Huang <i>et al</i> [22], 2020	RCT	18-65 years old	ERAT	Laparoscopic appendectomy	78	119	Acute appendicitis	Duration of operation, bed rest time, complication
Lin <i>et al</i> [23], 2016	RCT	18-70 years old	ERAT	Laparoscopic appendectomy/antibiotics treatment	44	45/36	Simple appendicitis	Length of hospital stay, bed rest time, time interval of body temperature returning to normal range, inflammatory factors, complication
Ma <i>et al</i> [24], 2020	RCT	19-74 years old	ERAT	Laparoscopic appendectomy	20	20	Non-complex appendicitis	Duration of operation, length of hospital stay, time interval of body temperature returning to normal range, inflammatory factors, complication
Wang <i>et al</i> [25], 2017	RCT	3 to 13 years old	ERAT	Laparoscopic appendectomy	42	42	Acute uncomplicated appendicitis in children	Duration of operation, length of hospital stay, bed rest time, time interval of body temperature returning to normal range, complication
Pan <i>et al</i> [26], 2018	RCT	19-62 years old	ERAT	Laparoscopic appendectomy	35	36	Acute appendicitis	Duration of operation, length of hospital stay, bed rest time, inflammatory factors
Shen <i>et al</i> [27], 2020	Case-control	NA	ERAT combined with antibiotics treatment	Antibiotics treatment	42	57	Acute appendicitis	Length of hospital stay
Ye <i>et al</i> [28], 2016	RCT	18-70 years old	ERAT	Laparoscopic appendectomy	57	57	Non-perforated acute appendicitis	Length of hospital stay, bed rest time, inflammatory factors, complication
Zhu <i>et al</i> [29], 2018	RCT	NA	ERAT	Antibiotics treatment	17	24	Atypical acute appendicitis	Complication
Yang <i>et al</i> [30], 2016	RCT	20-60 years old	ERAT	Laparoscopic appendectomy	35	35	Acute uncomplicated appendicitis	Duration of operation, bed rest time, length of hospital stay, time interval of body temperature returning to normal range
Li <i>et al</i> [31], 2016	Case-control	14-73 years old	ERAT	Laparoscopic appendectomy	21	20	Uncomplicated acute appendicitis	Duration of operation, length of hospital stay, bed rest time, time interval of body temperature returning to normal range, time interval of white blood cells count returning to normal time range, complication

ERAT: Endoscopic retrograde appendicitis therapy.

studies[27,31] were fair, with a score of 2 (each). The Jadad score of included studies were shown in Table 2 and Newcastle-Ottawa scale score was shown in Supplementary Table 1.

Table 2 Detailed quality assessment of included studies using modified Jadad score

Ref.	Randomization	Concealment of allocation	Double blinding	Description of withdrawals and dropouts	Total score
Kang <i>et al</i> [1], 2018	2	0	0	1	3
Deng <i>et al</i> [2], 2018	0	0	0	1	1
Huang <i>et al</i> [3], 2020	2	0	0	0	0
Lin <i>et al</i> [4], 2016	0	0	0	0	0
Ma <i>et al</i> [5], 2020	1	0	0	0	1
Wang <i>et al</i> [6], 2017	2	0	0	0	2
Pan <i>et al</i> [7], 2018	2	0	0	0	2
Wu <i>et al</i> [9], 2019	0	0	0	1	1
Ye <i>et al</i> [10], 2016	0	0	0	1	1
Zhang <i>et al</i> [11], 2017	1	0	0	0	1
Zhu <i>et al</i> [12], 2018	2	0	0	1	3
Yang <i>et al</i> [13], 2016	2	0	0	0	2

**Figure 1** Flow diagram representing the selection of studies.**Bed rest time**

Eight records reported the bed rest time in ERAT group and LA group. The bed rest time in ERAT group was shorter than LA group [WMD = -3.68, 95%CI: (-4.78, -2.58); $P < 0.001$], with high heterogeneity [$Q = 736.21$, P heterogeneity < 0.001 , $I^2 = 99.0\%$]. Shown in [Figure 2](#).

Time interval of body temperature returning to normal range

The time interval of body temperature returning to normal range in ERAT group was shorter than LA group based on 6 included studies. [SMD = -0.43, 95%CI: (-1.58, 0.73); $P = 0.481$] with high heterogeneity [$Q = 113.64$, P heterogeneity < 0.001 , $I^2 = 95.6\%$]. Shown in [Figure 3](#).




Figure 2 Forest plot of bed rest time.




Figure 3 Forest plot of time interval of body temperature returning to normal range.

Time interval of white blood cell count returning to normal range

Based on 2 included studies, the time interval of leukocyte count returning to normal range in patients receiving ERAT group was shorter than that in LA group [SMD = -1.11, 95%CI: (-1.58, -0.63); $P < 0.001$] with low heterogeneity [$Q = 0.24$, P heterogeneity = 0.630, $I^2 = 0.00\%$]. See Figure 4.

Duration of operation

Seven studies reported the duration of ERAT in comparison to LA. There was no difference regarding duration of operation between ERAT group and LA group [WMD = -13.90, 95%CI: (-29.56, 1.76); $P = 0.08$] with high heterogeneity [$Q = 227.42$, P heterogeneity < 0.001, $I^2 = 97.4\%$]. Shown in Figure 5.

Length of hospitalizations

Based on 8 included studies, the length of hospitalizations in ERAT group was shorter than LA group. [WMD = -1.15, 95%CI: (-1.99, -0.31); $P = 0.007$] with high heterogeneity [$Q = 289.85$, P heterogeneity < 0.001, $I^2 = 97.6\%$]. Shown in Figure 6.

Levels of inflammatory factors

C-reactive protein (CRP): Based on 3 included studies[24,26,28], there was no difference of pre-operative CRP levels between ERAT group and LA group [WMD = -0.28, 95%CI: (-1.14, 0.58); $P = 0.53$] with high heterogeneity [$Q = 7.21$, P heterogeneity = 0.03, $I^2 = 72.0\%$]. However, the level of post-operative CRP in ERAT group was significantly lower than that in LA group. [WMD = -10.06, 95%CI: (-17.39, -2.73); $P = 0.007$] with high heterogeneity [$Q = 109.28$, P heterogeneity < 0.001, $I^2 = 98.0\%$]. Shown in Table 3.

Tumor necrosis factor- α (TNF- α): Based on 2 included studies[24,26], there was no difference of pre-operative levels of TNF- α between ERAT group and LA group [WMD = -0.21, 95%CI: (-1.32, 0.90); $P = 0.71$] with low heterogeneity [$Q = 0.17$, P heterogeneity = 0.68, $I^2 = 0.00\%$]. However, the level of TNF- α in ERAT group was significantly lower than LA group after operating. [WMD = -7.70, 95%CI: (-8.47, -6.93); $P < 0.001$] with high heterogeneity [$Q = 138.67$, P heterogeneity < 0.001, $I^2 = 99.0\%$]. Shown in Table 3.

Table 3 Pooled results of inflammatory factors and complications

Outcomes	Categories	Number of records	OR/WMD and 95%CI	P	Heterogeneity with groups (I^2)	P_{het} value
Inflammatory factors						
	C-reactive protein (pre)	3	-0.28, [-1.14, 0.58]	0.53	72%	0.03
	C-reactive protein (post)	3	-10.06, [-17.39, -2.73]	0.007	98.0%	< 0.001
	Tumor necrosis factor- α (pre)	2	-0.21, [-1.32, 0.90]	0.71	0.0%	0.68
	Tumor necrosis factor- α (post)	2	7.70, [-8.47, -6.93]	< 0.001	99.0%	< 0.001
	Interleukin 6 (pre)	3	-0.11, [-1.04, 0.82]	0.81	6.0%	0.34
	Interleukin 6 (post)	3	-9.78, [-10.69, -8.88]	< 0.001	99.0%	< 0.001
Complications						
	Intestinal obstruction	4	0.19, [0.05, 0.79]	0.020	0.0%	0.95
	Abdominal infection	2	0.10, [0.01, 0.83]	0.030	0.0%	0.44
	Urinary tract infection	3	0.27, [0.04, 1.65]	0.160	0.0%	0.97




Figure 4 Forest plot of time interval of white blood cell count returning to normal range.




Figure 5 Forest plot of duration of operation.

Interleukin 6 (IL-6): Based on 3 included studies[24,26,28], no difference of pre-operative levels of IL-6 was found between ERAT group and LA group [WMD = -0.11, 95%CI: (-1.04, 0.82); $P = 0.81$] with low heterogeneity [$Q = 2.13$, P heterogeneity = 0.34, $I^2 = 6.0\%$]. However, the level of IL-6 in ERAT group was significantly lower than LA group, post-operatively. [WMD = -9.78, 95%CI: (-10.69, -8.88); $P < 0.001$] with high heterogeneity [$Q = 163.52$, P heterogeneity < 0.001, $I^2 = 99.0\%$]. Shown in Table 3.

Complications

Intestinal obstruction: Four studies[22,24,28,31] reported the intestinal obstruction after operation. The pooled result shown that ERAT group had a lower incidence of intestinal obstruction than LA group. [OR = 0.19, 95%CI: (0.05, 0.79); $P = 0.020$] with low heterogeneity [$Q = 0.34$, P heterogeneity = 0.95, $I^2 = 0.00\%$]. Shown in Table 3.

Abdominal infection: Two studies[24,31] reported the abdominal infection after




Figure 6 Forest plot of length of hospitalizations.

operation. The pooled result found that ERAT group had a lower incidence of abdominal infection than LA group [OR = 0.10, 95%CI: (0.01, 0.83); $P = 0.350$] with low heterogeneity [$Q = 0.60$, P heterogeneity = 0.44, $I^2 = 0.00\%$]. Shown in Table 3.

Urinary tract infection (UTI): The pooled result of 3 studies[25,28,31] reporting post-operative UTI did not find statistically significant difference between ERAT group and LA group [OR = 0.27, 95%CI: (0.04, 1.65); $P = 0.160$] with low heterogeneity [$Q = 0.07$, P heterogeneity = 0.97, $I^2 = 0.00\%$]. Shown in Table 3.

Sensitivity analysis

Furtherly, sensitivity analysis was performed to investigate the robustness of this meta-analysis. The results of sensitivity analysis shown that one study had a significant influence on the result of duration of operation[26], one study had a significant influence on the result of time interval of body temperature returning to the normal range[23], one study had a significant influence on the result of CRP (post-operative)[26], no study had a significant influence on the result of TNF (pre-operative) and one study had a significant influence on the result of IL-6 (pre-operative) [28].

Bias analysis

No obvious publication bias was depicted by the funnel plot (Supplementary Figure 1) and result from Egger's test ($t = -0.06$, $P = 0.954$) and Begg's test ($Z = 0.30$, $P = 0.764$) indicated no evidence of publication bias with regard to the duration of the operation. All outcomes of bias analysis were shown in Table 4.

DISCUSSION

Acute appendicitis, as one of the common surgical diseases, is the most common causes of surgical acute abdomen[32]. The latest study reported that the morbidity of acute appendicitis is as high as 6% in the population[33]. It has been found that the appendix can secrete a variety of useful substances and hormones (such as digestive enzymes, hormones that promote intestinal peristalsis, hormones related to growth), and play immune function to resist various diseases[34]. In addition, as the appendix contains a variety of intestinal microorganisms, it plays a key role in maintaining the balance of intestinal flora[35]. At present, the treatment for acute non-complex appendicitis includes surgery and conservative antibiotic treatment[36]. In order to preserve the potentially important function of the appendix, a retrograde endoscopic appendicitis treatment for acute simple appendicitis was first proposed in 2012. ERAT has the advantages of convenient operation, small trauma, and rapid relief of pain after the pressure of the appendix cavity is lifted[37]. In order to explore the safety of ERAT and provide more evidence for clinical treatment, this meta-analysis was conducted to investigate postoperative complications, length of hospitalizations, operation time, postoperative bed rest time, and indicators of recovery. The results showed that ERAT had shorter time intervals of white blood cell count returning to normal range, length of hospitalizations, and bed rest time. Meanwhile, the incidence of complications is lower, and the postoperative recovery time is faster compared with LA.

Table 4 Publication bias of outcomes by Egger's test and Begg's test

	Egger's test		Begg's test	
	<i>t</i>	<i>P</i>	<i>Z</i>	<i>P</i>
Time interval of body temperature returning to normal rangetime	1.17	0.306	0.75	0.452
Time interval of White white blood cells count returning to normal timerange	-	-	0.00	1.00
Duration of operation	-0.9	0.409	1.2	0.230
Length of hospitalizations (<i>vs</i> LA)	-0.48	0.648	0.37	0.711
Length of hospitalizations (<i>vs</i> Anti)	-1.72	0.336	0.00	1.00
CRP (pre-operative)	2.23	0.268	1.04	0.296
CRP (post-operative)	-0.19	0.878	0.00	1.00
TNF- α (pre-operative)	-	-	0.00	1.00
TNF- α (post-operative)	-	-	0.00	1.00
IL-6 (pre-operative)	-1.27	0.425	0.00	1.00
IL-6 (post-operative)	-7.43	0.085	1.04	0.296
Intestinal obstruction	2.03	0.179	1.70	0.089
Abdominal infection	-	-	0.00	1.00
Urinary tract infection	11.87	0.053	0.00	1.00
Bed rest time	-3.1	0.021	1.11	0.266

LA: Laparoscopic appendectomy.

In 2008, Mason *et al*[38] proposed that about 70% of patients with acute appendicitis do not need appendectomy and can be treated conservatively. Recently, Prechal *et al* [39] pointed out in a meta-analysis that appendectomy is more effective than antibiotic treatment in the treatment of acute uncomplicated appendicitis, and that the incidence of complications of the two treatment schemes is almost the same. Although ERAT emerges recently as a relatively new modality of treatment, it shows unique advantages. The latest research reported by Liu *et al*[18], the abdominal pain of 32 acute uncomplicated appendicitis patients resolved immediately after ERAT operation, and the clinical success rate was 97%. Colonoscopic irrigation, as a type of ERAT, was performed on 10 patients with acute appendicitis by Feng Jia *et al*[40]. Follow-up results found that there was no tenderness in the abdomen on physical examination, and no fever and other symptoms after operation. Notably, during the follow-up period of 1-8 mo, no complications occurred, and 9 cases had no recurrence of appendicitis. Chen *et al*[41] performed ERAT on 101 patients with acute appendicitis, the results showed that the success rate of appendiceal intubation was 96% (97/101), the success rate of treatment was 97.9% (94/96). Meanwhile, the operation time, the temperature recovery time, the white blood cell recovery time, and the abdominal pain relief time was shorter than the control group. What is more, no postoperative complications were detected. In addition, regarding the complication after ERAT, Li Yingchao *et al*[31] compared ERAT with LA and the results showed that perforation occurred in 1 case (5%) in ERAT group, and complications occurred in 3 cases (15%) in LA group. After more than half a year of follow-up, 2 cases in ERAT group were highly suspected of "chronic appendicitis" (recurrence rate 2/20, 10%), while no recurrence of appendicitis in LA group was reported, however, during a follow-up period of at least six months after surgery, 10 cases in LA group had postoperative diarrhea and constipation. Conversely, the results from Deng Ganlin *et al*[21] showed that the incidence of postoperative complications of the ERAT group was lower than that of the LA group, but the difference was not statistically significant ($P > 0.05$). Ma Zhuangfu *et al*[24] found that 1 sary intestinal obstruction occurred in ERAT group, while 6 sary intestinal obstructions occurred in LA group. Notably, our study shown that ERAT group had a lower incidence of intestinal obstruction than LA group based on 7 included studies. Lin *et al*[23] found that no patients with UTI and abdominal infection after ERAT, while 2 patients with UTI and 1 patient with abdominal infection were discovered in LA group, while this comprehensive meta-

analysis demonstrated that there was no difference between ERAT group and control group regarding abdominal infection and UTI.

The serum inflammatory factors of the patients between ERAT and control group were analyzed by Pan Hongwei[26], and the results showed the serum levels of hypersensitive CRP, IL-6, and TNF- α between ERAT group and LA group were significantly decreased after operation compared with those before operation, and the ERAT group was lower than the control group; The serum levels of hypersensitive CRP, IL-6, and TNF- α in the two groups were significantly decreased after operation compared with those before operation, and the ERAT group was lower than the control group ($P < 0.05$). CRP is an acute response protein secreted by the liver, and is also an essential inflammatory medium[42] to measure the intensity of response to trauma. IL-1 β , TNF- α , and IL-6 are common pro-inflammatory factors, and their secretion is increased in both acute and chronic inflammation, jointly promoting multiple pathological injury processes such as tissue destruction and edema formation [43,44]. IL-6 is also a typical pro-inflammatory factor, produced by activated T cells and fibroblasts, and can cooperatively activate inflammation-related signals with TNF- α to induce cascade reaction[45] and induce the production of other pro-inflammatory factors[46]. It is a common anti-inflammatory factor and has the effect of reducing inflammatory cell overactivation[47]. Therefore, we conducted the pooled analysis of these markers which shown that there was no difference in pre-operative levels of TNF- α , IL-6, and CRP between ERAT group and LA group, while the level of TNF- α , IL-6, and CRP in ERAT group was significantly lower than LA group after operating. However, we acknowledge that the timing of post-ERAT measurement of inflammatory factors is various across included studies, which may be one of the sources of heterogeneity.

Appendectomy has long been the most important method for the treatment of acute appendicitis. Although LA has faster recovery, less pain, and less wound infection compared with open surgery[48,49], there is still a certain risk of postoperative complications, and it has been reported[50,51] that the negative resection rate of appendix is as high as 8%-15%. Based on our meta-analysis, it is found that ERAT has its own unique advantages of being faster, more effective, and safer, compared with LA.

Limitation

First, the high heterogeneity across included studies was found, which could be attributed to different severities of the patients enrolled in each study, different mean ages of each study, different operating experience of ERAT of gastroenterologists and endoscopists in each study, and different study designs. Second, as little study compared LA with antibiotics treatment as well as compared adults with children, it is difficult to perform a meta-analysis regarding these outcomes. Third, limited studies were reported in other areas outside China.

CONCLUSION

Compared with LA treatment, ERAT reduces operation time, and results in fewer complications and shorter recovery time, with preserving the appendix and its immune and biological functions. However, given that only a limited number of studies were reported and most were conducted in China, more original studies with high quality in multi-centers from different countries and areas are still needed to further explore this novel modality of treatment for appendectomy.

ARTICLE HIGHLIGHTS

Research background

Evidence from previous studies shown that endoscopic retrograde appendicitis therapy (ERAT) is an effective treatment for acute appendicitis.

Research motivation

However, different studies reported conflicting outcomes regarding the effectiveness of ERAT in comparison with laparoscopic appendectomy (LA).

Research objectives

This meta-analysis was conducted to compare the effectiveness of ERAT with LA.

Research methods

Randomized controlled trials and retrospective studies of ERAT for acute uncomplicated appendicitis were searched in PubMed, Cochrane Library, Web of Science, Embase database, China National Knowledge Infrastructure (CNKI), the WanFang Database, and Chinese Scientific Journals Database (VIP).

Research results

10 randomized controlled studies (RCTs) and 2 case-control studies were included in the current systematic review. Firstly, the length of hospitalizations [WMD = -1.15, 95%CI: (-1.99, -0.31); $P = 0.007$] was shorter than LA group. Secondly, the level of post-operative CRP [WMD = -10.06, 95%CI: (-17.39, -2.73); $P = 0.007$], TNF- α [WMD = -7.70, 95%CI: (-8.47, -6.93); $P < 0.001$], and IL-6 Levels [WMD = -9.78, 95%CI: (-10.69, -8.88); $P < 0.001$; $P < 0.001$] in ERAT group was significantly lower than LA group. Thirdly, ERAT group had a lower incidence of intestinal obstruction than LA group. [OR = 0.19, 95%CI: (0.05, 0.79); $P = 0.020$].

Research conclusions

Based on our meta-analysis, it is found that ERAT has its own unique advantages of being more effective, safer compared with LA.

Research perspectives

As little study compared LA with antibiotics treatment, future study should focus on comparing the effectiveness between LA and antibiotics treatment.

REFERENCES

- 1 **Alore EA**, Ward JL, Rob TS. Population-level outcomes of early versus delayed appendectomy for acute appendicitis using the American College of Surgeons National Surgical Quality Improvement Program. *J Surg Res* 2018; 229: 234-242 [PMID: 29936996 DOI: 10.1016/j.jss.2018.04.011]
- 2 **Rentea RM**, Peter SDS, Snyder CL. Pediatric appendicitis: state of the art review. *Pediatr Surg Int* 2017; 33: 269-283 [PMID: 27743024 DOI: 10.1007/s00383-016-3990-2]
- 3 **Fagerström A**, Paajanen P, Saarelainen H, Ahonen-Siirtola M, Ukkonen M, Miettinen P, Paajanen H. Non-specific abdominal pain remains as the most common reason for acute abdomen: 26-year retrospective audit in one emergency unit. *Scand J Gastroenterol* 2017; 52: 1072-1077 [PMID: 28657380 DOI: 10.1080/00365521.2017.1342140]
- 4 **Georgiou R**, Eaton S, Stanton MP, Pierro A, Hall NJ. Efficacy and Safety of Nonoperative Treatment for Acute Appendicitis: A Meta-analysis. *Pediatrics* 2017; 139 [PMID: 28213607 DOI: 10.1542/peds.2016-3003]
- 5 **Drake FT**, Mottey NE, Farrokhi ET, Florence MG, Johnson MG, Mock C, Steele SR, Thirlby RC, Flum DR. Time to appendectomy and risk of perforation in acute appendicitis. *JAMA Surg* 2014; 149: 837-844 [PMID: 24990687 DOI: 10.1001/jamasurg.2014.77]
- 6 **Körner H**, Söndena K, Söreide JA, Andersen E, Nysted A, Lende TH, Kjelleveid KH. Incidence of acute nonperforated and perforated appendicitis: age-specific and sex-specific analysis. *World J Surg* 1997; 21: 313-317 [PMID: 9015177 DOI: 10.1007/s002689900235]
- 7 **Buckius MT**, McGrath B, Monk J, Grim R, Bell T, Ahuja V. Changing epidemiology of acute appendicitis in the United States: study period 1993-2008. *J Surg Res* 2012; 175: 185-190 [PMID: 22099604 DOI: 10.1016/j.jss.2011.07.017]
- 8 **Anderson JE**, Bickler SW, Chang DC, Talamini MA. Examining a common disease with unknown etiology: trends in epidemiology and surgical management of appendicitis in California, 1995-2009. *World J Surg* 2012; 36: 2787-2794 [PMID: 22948195 DOI: 10.1007/s00268-012-1749-z]
- 9 **Sanders NL**, Bollinger RR, Lee R, Thomas S, Parker W. Appendectomy and Clostridium difficile colitis: relationships revealed by clinical observations and immunology. *World J Gastroenterol* 2013; 19: 5607-5614 [PMID: 24039352 DOI: 10.3748/wjg.v19.i34.5607]
- 10 **Moberg AC**, Ahlberg G, Leijonmarck CE, Montgomery A, Reiertsen O, Rosseland AR, Stoerksson R. Diagnostic laparoscopy in 1043 patients with suspected acute appendicitis. *Eur J Surg* 1998; 164: 833-40; discussion 841 [PMID: 9845129 DOI: 10.1080/110241598750005246]
- 11 **Liang TJ**, Liu SI, Tsai CY, Kang CH, Huang WC, Chang HT, Chen IS. Analysis of Recurrence Management in Patients Who Underwent Nonsurgical Treatment for Acute Appendicitis. *Medicine (Baltimore)* 2016; 95: e3159 [PMID: 27015200 DOI: 10.1097/MD.00000000000003159]
- 12 **Huang L**, Yin Y, Yang L, Wang C, Li Y, Zhou Z. Comparison of Antibiotic Therapy and Appendectomy for Acute Uncomplicated Appendicitis in Children: A Meta-analysis. *JAMA Pediatr* 2017; 171: 426-434 [PMID: 28346589 DOI: 10.1001/jamapediatrics.2017.0057]

- 13 **Liu BR**, Song JT, Han FY, Li H, Yin JB. Endoscopic retrograde appendicitis therapy: a pilot minimally invasive technique (with videos). *Gastrointest Endosc* 2012; **76**: 862-866 [PMID: 22840292 DOI: 10.1016/j.gie.2012.05.029]
- 14 **Gonzalez DO**, Deans KJ, Minneci PC. Role of non-operative management in pediatric appendicitis. *Semin Pediatr Surg* 2016; **25**: 204-207 [PMID: 27521709 DOI: 10.1053/j.sempedsurg.2016.05.002]
- 15 **Page MJ**, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021; **372**: n71 [PMID: 33782057 DOI: 10.1136/bmj.n71]
- 16 **Jadad AR**, Moore RA, Carroll D, Jenkinson C, Reynolds DJ, Gavaghan DJ, McQuay HJ. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials* 1996; **17**: 1-12 [PMID: 8721797 DOI: 10.1016/0197-2456(95)00134-4]
- 17 **The Newcastle-Ottawa Scale for Assessing the Quality of Nonrandomized Studies in MetaAnalyses**. [cited 20 February 2021]. Available from: http://www.ohri.ca/programs/clinical_epidemiology/oxford.htm
- 18 **Higgins JP**, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002; **21**: 1539-1558 [PMID: 12111919 DOI: 10.1002/sim.1186]
- 19 **Egger M**, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; **315**: 629-634 [PMID: 9310563 DOI: 10.1136/bmj.315.7109.629]
- 20 **Kang J**, Zhang W, Zeng L, Lin Y, Wu J, Zhang N, Xie X, Zhang Y, Liu X, Wang B, Yang R, Jiang X. The modified endoscopic retrograde appendicitis therapy vs antibiotic therapy alone for acute uncomplicated appendicitis in children. *Surg Endosc* 2020 [PMID: 33146811 DOI: 10.1007/s00464-020-08129-8]
- 21 **Deng GL**, Ceng ZX, Chen LF. Effect of colonoscopy in the treatment of acute appendicitis. *Hainan Yixue Zazhi* 2018; **11**: 1594-1596 [DOI: 10.3969/j.issn.1003-6350.2018.11.035]
- 22 **Huang ZL**, Huo ZH, Shu YM. Application of endoscopic retrograde appendicitis therapy in acute appendicitis. *Hainan Yixue Zazhi* 2020; **11**: 1028-1103 [DOI: 10.3969/j.issn.1005-6483.2020.11.009]
- 23 **Lin BM**, Sun XZ, Liu YY. Comparison on the therapeutic effect of different treatments for simple appendicitis complicated with diabetes mellitus. *Hainan Yixue Zazhi* 2016; **19**: 3157-3160 [DOI: 10.3969/j.issn.1003-6350.2016.19.021]
- 24 **Ma ZF**, Huang RW. Endoscopic retrograde appendicitis therapy in treatment of acute non-complex appendicitis. *Zhongguo Neijing Zazhi* 2020; **26**: 7-12 [DOI: 10.3969/j.issn.1007-1989.2020.07.002]
- 25 **Wang GF**. Efficacy of endoscopic retrograde appendicitis in children with acute non perforated appendicitis. *Yingxiang Yanjiu Yixue Yingyong Zazhi* 2017; **18**: 230-231 [DOI: 10.3969/j.issn.2096-3807.2017.18.148]
- 26 **Pan HW**, Weng JJ. Prevention value of endoscopic retrograde appendicitis treatment for postoperative with appendicitis. *Zhonghua Xiaohua Neijing Zazhi* 2018; **35**: 405-409 [DOI: 10.3760/cma.j.issn.1007-5232.2018.06.006]
- 27 **Shen WY**, Tang J, Wu T. Comparative study on the efficacy of conservative treatment and endoscopic retrograde appendicitis treatment for acute appendicitis. *Hainan Yixue Zazhi* 2020; **24**: 3208-3210 [DOI: 10.3969/j.issn.1003-6350.2020.24.023]
- 28 **Ye Y**, Sun XZ, Yang LM. Application of endoscopic retrograde appendicitis therapy in non-perforated acute appendicitis. *Zhongguo Linchuang Yanjiu Zazhi* 2016; **29**: 741-745 [DOI: 10.13429/j.cnki.cjcr.2016.06.006]
- 29 **Zhu FY**, Chen T, Fu Z. Diagnostic and therapeutic value of endoscopic retrograde appendicitis therapy for atypical acute appendicitis. *Zhonghua Xiaohua Neijing Zazhi* 2018; **35**: 571-575 [DOI: 10.3760/cma.j.issn.1007-5232.2018.08.009]
- 30 **Yang GJ**, Hu XF. The Value of Endoscopic Retrograde Appendicitis in the Treatment of Acute Uncomplicated Appendicitis. *Zhongguo Jixu Yixue Jiaoyu* 2016; **8**: 108-109 [DOI: 10.3969/j.issn.1674-9308.2016.17.068]
- 31 **Li YC**, Mi C, Li WZ. Effect and safety of Endoscopic retrograde appendicitis therapy in treating patients with uncomplicated acute appendicitis. *Zhongguo Neijing Zazhi* 2016; **22**: 11-17 [DOI: 10.3969/j.issn.1007-1989.2016.03.003]
- 32 **Bhangu A**, Søreide K, Di Saverio S, Assarsson JH, Drake FT. Acute appendicitis: modern understanding of pathogenesis, diagnosis, and management. *Lancet* 2015; **386**: 1278-1287 [PMID: 26460662 DOI: 10.1016/S0140-6736(15)00275-5]
- 33 **Snyder MJ**, Guthrie M, Cagle S. Acute Appendicitis: Efficient Diagnosis and Management. *Am Fam Physician* 2018; **98**: 25-33 [PMID: 30215950]
- 34 **Chen SH**, Yeong EK, Tang YB, Chen HC. Free and pedicled appendix transfer for various reconstructive procedures. *Ann Plast Surg* 2012; **69**: 602-606 [PMID: 23151608 DOI: 10.1097/SAP.0b013e31827475e2]
- 35 **Chen J**, Sali A, Vitetta L. The gallbladder and vermiform appendix influence the assemblage of intestinal microorganisms. *Future Microbiol* 2020; **15**: 541-555 [PMID: 32478580 DOI: 10.2217/fmb-2019-0325]
- 36 **Coccolini F**, Fugazzola P, Sartelli M, Cicuttin E, Sibilla MG, Leandro G, De' Angelis GL, Gaiani F, Di Mario F, Tomasoni M, Catena F, Ansaloni L. Conservative treatment of acute appendicitis. *Acta Biomed* 2018; **89**: 119-134 [PMID: 30561405 DOI: 10.23750/abm.v89i9-S.7905]
- 37 **LIU Bingrong**, SONG Jitao, MA Xiao. Introduction of endoscopic retrograde appendicitis treatment

- technology. *Zhonghua Xiaohua Beijing Zazhi* 2013; **30**: 468 [DOI: [10.3760/cma.j.issn.1007-5232.2013.08.015](https://doi.org/10.3760/cma.j.issn.1007-5232.2013.08.015)]
- 38 **Mason RJ.** Surgery for appendicitis: is it necessary? *Surg Infect (Larchmt)* 2008; **9**: 481-488 [PMID: [18687030](https://pubmed.ncbi.nlm.nih.gov/18687030/) DOI: [10.1089/sur.2007.079](https://doi.org/10.1089/sur.2007.079)]
 - 39 **Podda M,** Gerardi C, Cillara N, Fearnhead N, Gomes CA, Birindelli A, Mulliri A, Davies RJ, Di Saverio S. Antibiotic Treatment and Appendectomy for Uncomplicated Acute Appendicitis in Adults and Children: A Systematic Review and Meta-analysis. *Ann Surg* 2019; **270**: 1028-1040 [PMID: [30720508](https://pubmed.ncbi.nlm.nih.gov/30720508/) DOI: [10.1097/SLA.0000000000003225](https://doi.org/10.1097/SLA.0000000000003225)]
 - 40 **FENG Jia,** FENG Zitan, SUN Rong. Effect Observation of endoscopic intracavity douching in treatment of patients with acute appendicitis[J]. *Jiefangjun Yixue Zazhi* 2014; **26**: 46-47 [DOI: [10.3969/j.issn.2095-140X.2014.01.014](https://doi.org/10.3969/j.issn.2095-140X.2014.01.014)]
 - 41 **Chen Y,** Wang M, Chen H, Zhao L, Liu L, Wang X, Huang J, Fan Z. WITHDRAWN: Endoscopic intervention for acute appendicitis: retrospective study of 101 cases. *Gastrointest Endosc* 2019 [PMID: [31228430](https://pubmed.ncbi.nlm.nih.gov/31228430/) DOI: [10.1016/j.gie.2019.06.012](https://doi.org/10.1016/j.gie.2019.06.012)]
 - 42 **Trial J,** Cieslik KA, Entman ML. Phosphocholine-containing ligands direct CRP induction of M2 macrophage polarization independent of T cell polarization: Implication for chronic inflammatory states. *Immun Inflamm Dis* 2016; **4**: 274-288 [PMID: [27621811](https://pubmed.ncbi.nlm.nih.gov/27621811/) DOI: [10.1002/iid3.112](https://doi.org/10.1002/iid3.112)]
 - 43 **Broekman W,** Amatngalim GD, de Mooij-Eijk Y, Oostendorp J, Roelofs H, Taube C, Stolk J, Hiemstra PS. TNF- α and IL-1 β -activated human mesenchymal stromal cells increase airway epithelial wound healing *in vitro* via activation of the epidermal growth factor receptor. *Respir Res* 2016; **17**: 3 [PMID: [26753875](https://pubmed.ncbi.nlm.nih.gov/26753875/) DOI: [10.1186/s12931-015-0316-1](https://doi.org/10.1186/s12931-015-0316-1)]
 - 44 **Zhang P,** Wu X, Li G, He Q, Dai H, Ai C, Shi J. Tumor necrosis factor- α gene polymorphisms and susceptibility to ischemic heart disease: A systematic review and meta-analysis. *Medicine (Baltimore)* 2017; **96**: e6569 [PMID: [28383437](https://pubmed.ncbi.nlm.nih.gov/28383437/) DOI: [10.1097/MD.0000000000000659](https://doi.org/10.1097/MD.0000000000000659)]
 - 45 **Chae JW,** Ng T, Yeo HL, Shwe M, Gan YX, Ho HK, Chan A. Impact of TNF- α (rs1800629) and IL-6 (rs1800795) Polymorphisms on Cognitive Impairment in Asian Breast Cancer Patients. *PLoS One* 2016; **11**: e0164204 [PMID: [27701469](https://pubmed.ncbi.nlm.nih.gov/27701469/) DOI: [10.1371/journal.pone.0164204](https://doi.org/10.1371/journal.pone.0164204)]
 - 46 **Shah S,** Ma Y, Scherzer R, Huhn G, French AL, Plankey M, Peters MG, Grunfeld C, Tien PC. Association of HIV, hepatitis C virus and liver fibrosis severity with interleukin-6 and C-reactive protein levels. *AIDS* 2015; **29**: 1325-1333 [PMID: [25870985](https://pubmed.ncbi.nlm.nih.gov/25870985/) DOI: [10.1097/QAD.0000000000000654](https://doi.org/10.1097/QAD.0000000000000654)]
 - 47 **Abraham BP,** Ahmed T, Ali T. Inflammatory Bowel Disease: Pathophysiology and Current Therapeutic Approaches. *Handb Exp Pharmacol* 2017; **239**: 115-146 [PMID: [28233184](https://pubmed.ncbi.nlm.nih.gov/28233184/) DOI: [10.1007/164_2016_122](https://doi.org/10.1007/164_2016_122)]
 - 48 **Dhingra AK,** Chopra B, Dass R, Mittal SK. An update on Anti-inflammatory Compounds: A Review. *Antiinflamm Antiallergy Agents Med Chem* 2015; **14**: 81-97 [PMID: [25973652](https://pubmed.ncbi.nlm.nih.gov/25973652/) DOI: [10.2174/1871523014666150514102027](https://doi.org/10.2174/1871523014666150514102027)]
 - 49 **Jaschinski T,** Mosch CG, Eikermann M, Neugebauer EA, Sauerland S. Laparoscopic vs open surgery for suspected appendicitis. *Cochrane Database Syst Rev* 2018; **11**: CD001546 [PMID: [30484855](https://pubmed.ncbi.nlm.nih.gov/30484855/) DOI: [10.1002/14651858.CD001546.pub4](https://doi.org/10.1002/14651858.CD001546.pub4)]
 - 50 **Ceresoli M,** Tamini N, Gianotti L, Braga M, Nespoli L. Are endoscopic loop ties safe even in complicated acute appendicitis? *Int J Surg* 2019; **68**: 40-47 [PMID: [31226406](https://pubmed.ncbi.nlm.nih.gov/31226406/) DOI: [10.1016/j.ijsu.2019.06.011](https://doi.org/10.1016/j.ijsu.2019.06.011)]
 - 51 **Myers E,** Kavanagh DO, Ghous H, Evoy D, McDermott EW. The impact of evolving management strategies on negative appendectomy rate. *Colorectal Dis* 2010; **12**: 817-821 [PMID: [19508509](https://pubmed.ncbi.nlm.nih.gov/19508509/) DOI: [10.1111/j.1463-1318.2009.01910.x](https://doi.org/10.1111/j.1463-1318.2009.01910.x)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

