



## Risk factors for local recurrence of middle and lower rectal carcinoma after curative resection

Ze-Yu Wu, Jin Wan, Gang Zhao, Lin Peng, Jia-Lin Du, Yuan Yao, Quan-Fang Liu, Hua-Huan Lin

Ze-Yu Wu, Jin Wan, Gang Zhao, Lin Peng, Jia-Lin Du, Yuan Yao, Quan-Fang Liu, Department of General Surgery, Guangdong Provincial People's Hospital, Guangzhou 510080, Guangdong Province, China

Hua-Huan Lin, Department of Pathology, Guangdong Provincial People's Hospital, Guangzhou 510080, Guangdong Province, China

**Author contributions:** Wu ZY and Wan J designed the research; Wu ZY, Zhao G, Peng L, Du JL, Yao Y, Liu QF and Lin HH performed research; Wu ZY analyzed the data; Wu ZY and Wan J wrote the paper.

**Supported by** The WST Foundation of Guangdong Province, No. 2000112736580706003

**Correspondence to:** Dr. Ze-Yu Wu, Department of General Surgery, Guangdong Provincial People's Hospital, Guangzhou 510080, Guangdong Province, China. [ljhde@163.com](mailto:ljhde@163.com)

Telephone: +86-20-83827812-60821 Fax: +86-20-83827812

Received: April 13, 2008

Revised: June 16, 2008

Accepted: June 23, 2008

Published online: August 14, 2008

### Abstract

**AIM:** To explore the risk factors for local recurrence of middle and lower rectal carcinoma after curative resection.

**METHODS:** Specimens of middle and lower rectal carcinoma from 56 patients who received curative resection at the Department of General Surgery of Guangdong Provincial People's Hospital were studied. A large slice technique was used to detect mesorectal metastasis and evaluate circumferential resection margin status. The relations between clinicopathologic characteristics, mesorectal metastasis and circumferential resection margin status were identified in patients with local recurrence of middle and lower rectal carcinoma.

**RESULTS:** Local recurrence of middle and lower rectal carcinoma after curative resection occurred in 7 of the 56 patients (12.5%), and was significantly associated with family history ( $\chi^2 = 3.929$ ,  $P = 0.047$ ), high CEA level ( $\chi^2 = 4.964$ ,  $P = 0.026$ ), cancerous perforation ( $\chi^2 = 8.503$ ,  $P = 0.004$ ), tumor differentiation ( $\chi^2 = 9.315$ ,  $P = 0.009$ ) and vessel cancerous emboli ( $\chi^2 = 11.879$ ,  $P = 0.001$ ). In contrast, no significant correlation was found between local recurrence of rectal carcinoma and other variables such as age ( $\chi^2 = 0.506$ ,  $P = 0.477$ ), gender ( $\chi^2 = 0.102$ ,  $\chi^2 = 0.749$ ), tumor diameter ( $\chi^2 = 0.421$ ,  $P = 0.516$ ),

tumor infiltration ( $\chi^2 = 5.052$ ,  $P = 0.168$ ), depth of tumor invasion ( $\chi^2 = 4.588$ ,  $P = 0.101$ ), lymph node metastases ( $\chi^2 = 3.688$ ,  $P = 0.055$ ) and TNM staging system ( $\chi^2 = 3.765$ ,  $P = 0.152$ ). The local recurrence rate of middle and lower rectal carcinoma was 33.3% (4/12) in patients with positive circumferential resection margin and 6.8% (3/44) in those with negative circumferential resection margin. There was a significant difference between the two groups ( $\chi^2 = 6.061$ ,  $P = 0.014$ ). Local recurrence of rectal carcinoma occurred in 6 of 36 patients (16.7%) with mesorectal metastasis, and in 1 of 20 patients (5.0%) without mesorectal metastasis. However, there was no significant difference between the two groups ( $\chi^2 = 1.600$ ,  $P = 0.206$ ).

**CONCLUSION:** Family history, high CEA level, cancerous perforation, tumor differentiation, vessel cancerous emboli and circumferential resection margin status are the significant risk factors for local recurrence of middle and lower rectal carcinoma after curative resection. Local recurrence may be more frequent in patients with mesorectal metastasis than in patients without mesorectal metastasis.

© 2008 The WJG Press. All rights reserved.

**Key words:** Middle and lower rectal carcinoma; Local recurrence; Circumferential resection margin; Mesorectal metastasis

**Peer reviewer:** Yik-Hong Ho, Professor, Department of Surgery, School of Medicine, James Cook University, Townsville 4811, Australia

Wu ZY, Wan J, Zhao G, Peng L, Du JL, Yao Y, Liu QF, Lin HH. Risk factors for local recurrence of middle and lower rectal carcinoma after curative resection. *World J Gastroenterol* 2008; 14(30): 4805-4809 Available from: URL: <http://www.wjgnet.com/1007-9327/14/4805.asp> DOI: <http://dx.doi.org/10.3748/wjg.14.4805>

### INTRODUCTION

It is well known that local recurrence is of rectal carcinoma plays an important role in its prognosis<sup>[1-3]</sup>. However, local recurrence of rectal carcinoma occurs in about 4%-50% of patients even after radical resection of primary tumors and lymph nodes<sup>[4-8]</sup>. The risk factors

for local recurrence of rectal carcinoma remain unclear. Therefore, the aim of the current study was to explore the risk factors for local recurrence of middle and lower rectal carcinoma after curative resection. Specimens of middle and lower rectal carcinoma from 56 patients who underwent total mesorectal excision (TME) at the Department of General Surgery, Guangdong Provincial People's Hospital, from November, 2001 to July, 2003, were studied. A large slice technique was used to detect mesorectal metastasis and evaluate circumferential resection margin status. The relationship between mesorectal metastasis, local recurrence and circumferential resection margin status of rectal carcinoma was observed. The clinicopathologic characteristics of middle and lower rectal carcinoma were also evaluated.

## MATERIALS AND METHODS

### *Patients and specimens*

Specimens of middle and lower rectal carcinoma from 56 patients who underwent TME at the Department of General Surgery, Guangdong Provincial People's Hospital, from November, 2001 to July, 2003, were studied. There were 37 men and 19 women, ranging in age from 30 to 86 years, with a mean age of 60.5 years. None of these patients received preoperative chemotherapy or radiotherapy. Twenty-six patients had lower rectal carcinoma and 30 had middle rectal carcinoma. Tumors  $\geq 5$  cm and in  $< 5$  cm diameter were found in 18 and 38 patients, respectively. Low anterior resection was performed in 40 patients and abdominoperineal resection in 16 patients. TNM stages were as follows: stage I in 5 patients, stage II in 22 patients, and stage III in 29 patients. Poorly-differentiated carcinoma was observed in 14 patients, moderately-differentiated carcinoma in 37 patients, and well-differentiated carcinoma in 5 patients, respectively. A large slice technique was used to detect mesorectal metastasis and evaluate circumferential resection margin status. Two pathologists who were blinded to the clinicopathological data observed the specimens independently. If tumor cells were detected within 1 mm of circumferential margin, they were classified to have a positive circumferential resection margin as previously described<sup>[9-11]</sup>.

### *Statistical analysis*

Statistical analysis was performed by the Pearson  $\chi^2$  test to examine the association between local recurrence, circumferential resection margin status and mesorectal metastasis of rectal carcinoma. Clinicopathologic characteristics of the patients with middle and lower rectal carcinoma were also analyzed.  $P < 0.05$  was considered statistically significant.

## RESULTS

### *Correlation between local recurrence and clinicopathologic characteristics of patients with middle and lower rectal carcinoma*

Local recurrence of middle and lower rectal carcinoma

after curative resection was found in 7 of 56 patients (12.5%), which was significantly related with family history ( $\chi^2 = 3.929$ ,  $P = 0.047$ ), high CEA level ( $\chi^2 = 4.964$ ,  $P = 0.026$ ), cancerous perforation ( $\chi^2 = 8.503$ ,  $P = 0.004$ ), tumor differentiation ( $\chi^2 = 9.315$ ,  $P = 0.009$ ) and vessel cancerous emboli ( $\chi^2 = 11.879$ ,  $P = 0.001$ ). In contrast, no significant correlation was found between local recurrence and other variables such as age ( $\chi^2 = 0.506$ ,  $P = 0.477$ ) and gender ( $\chi^2 = 0.102$ ,  $P = 0.749$ ), tumor diameter ( $\chi^2 = 0.421$ ,  $P = 0.516$ ), tumor infiltration ( $\chi^2 = 5.052$ ,  $P = 0.168$ ), depth of tumor invasion ( $\chi^2 = 4.588$ ,  $P = 0.101$ ), lymph node metastases ( $\chi^2 = 3.688$ ,  $P = 0.055$ ) and TNM staging system ( $\chi^2 = 3.765$ ,  $P = 0.152$ ) (Table 1).

### *Correlation between circumferential resection margin status and local recurrence of middle and lower rectal carcinoma*

A positive circumferential resection margin of middle and lower rectal carcinoma was observed in 12 of 56 patients (21.4%). Local recurrence of middle and lower rectal carcinoma was found in 4 of 12 patients (33.3%) with a positive circumferential resection margin and in 3 of 44 patients (6.8%) with a negative circumferential resection margin. There was a significant difference between the two groups ( $\chi^2 = 6.061$ ,  $P = 0.014$ ) (Table 1).

### *Correlation between mesorectal metastasis and local recurrence of middle and lower rectal carcinoma*

Mesorectal metastasis of middle and lower rectal carcinoma was detected in 36 of 56 patients (64.3%). The local recurrence rate of mesorectal metastasis was 16.7% (6 of 36 patients) and 5.0% (1 of 20 patients), respectively. However, there was no significant difference between the two groups ( $\chi^2 = 1.600$ ,  $P = 0.206$ ) (Table 1).

## DISCUSSION

It is well known that middle and lower rectal carcinoma is one of the most common carcinomas in China. Local recurrence of middle and lower rectal carcinoma after curative resection has significant morbidity and mortality<sup>[4,12-15]</sup> and its recurrence rate varies from less than 4% to greater than 50%. Since TME was adopted as the standard treatment of patients with rectal carcinoma, a significant decrease in local recurrence and a trend to improve relative survival have been reported<sup>[16-18]</sup>. In our study, local recurrence of middle and lower rectal carcinoma occurred in 7 of 56 patients (12.5%) after TME, indicating that TME can significantly reduce the local recurrence rate of middle and lower rectal carcinomas.

The correlation between circumferential resection margin status and local recurrence of rectal carcinoma is still controversial<sup>[10,11,19-21]</sup>. Wibe *et al*<sup>[10]</sup> reported that a positive circumferential resection margin has a significant and major prognostic impact on the local recurrence rate of rectal carcinoma after TME. However, Luna-Perez *et al*<sup>[19]</sup> reported that circumferential resection margin involvement is not correlated significantly with

**Table 1** Local recurrence and circumferential resection margin status, mesorectal metastasis, and clinicopathologic characteristics of middle and lower rectal carcinoma

Clinicopathologic variable	Patients (n)	Local recurrence		$\chi^2$	P
		Positive (%)	Negative (%)		
Gender					
Male	37	5 (13.5)	32 (86.5)	0.102	0.749
Female	19	2 (10.5)	17 (89.5)		
Age					
< 60 yr	25	4 (16.0)	21 (84.0)	0.506	0.477
≥ 60 yr	31	3 (9.7)	28 (90.3)		
Family history					
Yes	21	5 (23.8)	16 (76.2)	3.929	0.047
No	35	2 (5.7)	33 (94.3)		
CEA level					
High	26	6 (23.1)	20 (76.9)	4.964	0.026
Normal	30	1 (3.3)	29 (96.7)		
Cancerous perforation					
Yes	3	2 (33.3)	1 (66.7)	8.503	0.004
No	53	5 (9.4)	48 (90.6)		
Superficial diameter					
< 5 cm	38	4 (10.5)	34 (89.5)	0.421	0.516
≥ 5 cm	18	3 (16.7)	15 (83.3)		
Diameter of infiltration					
1/4	8	0 (0)	8 (100)	5.052	0.168
1/2	16	1 (6.3)	15 (93.7)		
3/4	18	2 (11.1)	16 (88.9)		
4/4	14	4 (28.6)	10 (71.4)		
Depth of invasion					
T <sub>1</sub>	6	0 (0)	6 (100)	4.588	0.101
T <sub>2</sub>	23	1 (4.3)	22 (95.7)		
T <sub>3</sub>	27	6 (22.2)	21 (77.8)		
Histologic differentiation					
Well	5	0 (0)	5 (100)	9.315	0.009
Moderate	37	2 (5.4)	35 (94.6)		
Poorly	14	5 (35.7)	9 (64.3)		
Lymph node metastasis					
Positive	29	6 (20.7)	23 (79.3)	3.688	0.055
Negative	27	1 (3.7)	26 (96.3)		
Vessel cancerous emboli					
Positive	12	5 (41.7)	7 (58.3)	11.879	0.001
Negative	44	2 (4.5)	42 (95.5)		
Circumferential resection margin					
Positive	12	4 (33.3)	8 (66.7)	6.061	0.014
Negative	44	3 (6.8)	41 (93.2)		
Mesorectal metastasis					
Positive	36	6 (16.7)	30 (83.3)	1.6	0.206
Negative	20	1 (5.0)	19 (95.0)		
TNM staging					
I	5	0 (0)	5 (100)	3.765	0.152
II	22	1 (4.5)	21 (95.5)		
III	29	6 (20.7)	23 (79.3)		

local recurrence of rectal adenocarcinoma ( $P = 0.33$ ). Hall *et al*<sup>[11]</sup> reported that the local recurrence rate of rectal carcinoma with a positive circumferential resection margin is 15% and 11% in of those with a negative circumferential resection margin. The difference between the two groups was not significant. Our results demonstrate that circumferential resection margin involvement had a significant correlation with local recurrence of middle and low rectal carcinoma. Local recurrence was more frequently observed in rectal carcinomas with a positive circumferential resection margin (4 of 12 patients, 33.3%) than in those with a negative circumferential resection margin

(3 of 44 patients, 6.8%) ( $P = 0.014$ ), suggesting that circumferential resection margin status is an important predictor for the local recurrence of middle and low rectal carcinoma. It has been shown that residual mesorectal metastasis observed in rectal surgery may be the most important factor for local recurrence of rectal carcinoma<sup>[22,23]</sup>. In the current study, the local recurrence of rectal carcinoma was more frequent in patients with mesorectal metastasis than in those without mesorectal metastasis (16.7% *vs* 5.0%). However, there was no significant difference between the two groups ( $P = 0.206$ ). TME may be also a plausible explanation for the observation.

Sugihara *et al*<sup>[24]</sup> investigated the correlation between local recurrence and clinicopathologic characteristics of rectal carcinoma by multivariate analysis, and found that local recurrence of lower rectal cancer is significantly associated lymph node metastasis. It has been demonstrated that pathologic stages T and N are the significant predictors for the local recurrence of rectal carcinoma<sup>[25]</sup>. In the present study, local recurrence of poorly- and moderately- differentiated rectal carcinomas was found in 5 of 34 patients (35.7%) and in 2 of 37 patients (5.4%), respectively ( $P = 0.009$ ), while no local recurrence of well-differentiated rectal carcinoma was observed in any patients, suggesting that local recurrence of rectal carcinoma is significantly correlated with tumor differentiation. We also found that the local recurrence rate of rectal carcinoma was also correlated with the depth of tumor invasion. Local recurrence of T<sub>3</sub> and T<sub>2</sub> tumors was observed in 6 of 27 patients (22.2%) and in 1 of 23 patients (4.3%), respectively, while no local recurrence of T<sub>1</sub> tumors was observed ( $P = 0.101$ ). Local recurrence of rectal carcinoma developed in 6 (20.7%) of the 29 patients with lymph node metastasis and in 1 (3.7%) of 27 patients without lymph node metastasis ( $P = 0.055$ ). These observations may be explained by the fact that the number of patients in our study was comparatively small. Further study with a larger sample size is needed.

Park *et al*<sup>[26]</sup> reported that change in perioperative serum CEA is a useful prognostic predictor for the occurrence of stage III rectal cancer and the survival of such patients. Oh *et al*<sup>[27]</sup> reported that vascular invasion is significantly associated with local recurrence of rectal cancer. Our results also demonstrate that local recurrence of rectal carcinoma had a significant correlation with high CEA level ( $P = 0.026$ ) and vessel cancerous emboli ( $P = 0.001$ ). We also found that family history and cancerous perforation were significantly correlated with local recurrence of rectal carcinoma ( $P < 0.05$ ).

In conclusion, extensive mesorectal excision and postoperative adjuvant chemotherapy should be used in the treatment of middle and lower rectal carcinoma.

## COMMENTS

### Background

It is well known that local recurrence is the most important prognostic factor for rectal carcinoma. However, local recurrence of rectal carcinoma occurs in about 4%-50% of patients even after radical resection of primary tumors and lymph nodes. The risk factors for local recurrence of rectal carcinoma remain unclear.

### Research frontiers

Since total mesorectal excision (TME) was adopted as the standard treatment of rectal carcinoma, a significant decrease in its local recurrence and a trend to improve the relative survival of rectal carcinoma patients have been reported. However, the local recurrence rate of rectal carcinoma is still high. The correlation between circumferential resection margin status and local recurrence of rectal carcinoma is still controversial.

### Innovations and breakthroughs

The aim of the current study was to explore the risk factors for local recurrence of middle and lower rectal carcinoma after curative resection. A large slice technique was used to detect its mesorectal metastasis and evaluate its circumferential resection margin status. The relationship between mesorectal

metastasis and circumferential resection margin status with local recurrence was identified. The clinicopathological characteristics of rectal carcinoma were described.

### Applications

Family history, high CEA level, cancerous perforation, tumor differentiation, vessel cancerous emboli and circumferential resection margin status are the significant risk factors of local recurrence of middle and lower rectal carcinoma after curative resection. Local recurrence of middle and lower rectal carcinoma may be more frequent observed in patients with mesorectal metastasis than in those without mesorectal metastasis.

### Terminology

Local recurrence of middle and lower rectal carcinoma was defined as any recurrence diagnosed or suspected in the pelvis (tumor bed, pelvic nodes, anastomosis, drain site, or perineum).

### Peer review

The authors explored the risk factors for local recurrence of middle and lower rectal carcinoma after curative resection, demonstrating that family history, high CEA level, cancerous perforation, tumor differentiation, vessel cancerous emboli and circumferential resection margin status are the significant risk factors for the local recurrence of middle and lower rectal carcinoma. The study was well designed. The data provided in the paper are interesting and valuable.

## REFERENCES

- 1 Noda K, Umekita N, Tanaka S, Ohkubo T, Inoue S, Kitamura M. [A clinical study of therapy for local recurrent rectal cancer] *Gan To Kagaku Ryoho* 2006; **33**: 1830-1833
- 2 Bedrosian I, Giasco G, Pederson L, Rodriguez-Bigas MA, Feig B, Hunt KK, Ellis L, Curley SA, Vauthey JN, Delclos M, Crane CH, Janjan N, Skibber JM. Outcome after curative resection for locally recurrent rectal cancer. *Dis Colon Rectum* 2006; **49**: 175-182
- 3 Moriya Y, Akasu T, Fujita S, Yamamoto S. Total pelvic exenteration with distal sacrectomy for fixed recurrent rectal cancer. *Surg Oncol Clin N Am* 2005; **14**: 225-238
- 4 Radice E, Dozois RR. Locally recurrent rectal cancer. *Dig Surg* 2001; **18**: 355-362
- 5 Piso P, Dahlke MH, Mirena P, Schmidt U, Aselmann H, Schlitt HJ, Raab R, Klempnauer J. Total mesorectal excision for middle and lower rectal cancer: a single institution experience with 337 consecutive patients. *J Surg Oncol* 2004; **86**: 115-121
- 6 Rengan R, Paty PB, Wong WD, Guillem JG, Weiser M, Temple L, Saltz L, Minsky BD. Ten-year results of preoperative radiation followed by sphincter preservation for rectal cancer: increased local failure rate in nonresponders. *Clin Colorectal Cancer* 2006; **5**: 413-421
- 7 Hohenberger W, Merkel S, Matzel K, Bittorf B, Papadopoulos T, Gohl J. The influence of abdomino-perineal (intersphincteric) resection of lower third rectal carcinoma on the rates of sphincter preservation and locoregional recurrence. *Colorectal Dis* 2006; **8**: 23-33
- 8 Lezoche E, Guerrieri M, De Sanctis A, Campagnacci R, Baldarelli M, Lezoche G, Paganini AM. Long-term results of laparoscopic versus open colorectal resections for cancer in 235 patients with a minimum follow-up of 5 years. *Surg Endosc* 2006; **20**: 546-553
- 9 Hermanek P, Junginger T. The circumferential resection margin in rectal carcinoma surgery. *Tech Coloproctol* 2005; **9**: 193-199; discussion 199-200
- 10 Wibe A, Rendedal PR, Svensson E, Norstein J, Eide TJ, Myrvold HE, Soreide O. Prognostic significance of the circumferential resection margin following total mesorectal excision for rectal cancer. *Br J Surg* 2002; **89**: 327-334
- 11 Hall NR, Finan PJ, al-Jaberi T, Tsang CS, Brown SR, Dixon MF, Quirke P. Circumferential margin involvement after mesorectal excision of rectal cancer with curative intent. Predictor of survival but not local recurrence? *Dis Colon Rectum* 1998; **41**: 979-983
- 12 Okaro AC, Worthington T, Stebbing JF, Broughton M, Caffarey S, Marks CG. Curative resection for low rectal adenocarcinoma: abdomino-perineal vs anterior resection.

- Colorectal Dis* 2006; **8**: 645-649
- 13 **Chiappa A**, Biffi R, Bertani E, Zbar AP, Pace U, Crotti C, Biella F, Viale G, Orecchia R, Pruneri G, Poldi D, Andreoni B. Surgical outcomes after total mesorectal excision for rectal cancer. *J Surg Oncol* 2006; **94**: 182-193; discussion 181
  - 14 **Martling AL**, Holm T, Rutqvist LE, Moran BJ, Heald RJ, Cedemark B. Effect of a surgical training programme on outcome of rectal cancer in the County of Stockholm. Stockholm Colorectal Cancer Study Group, Basingstoke Bowel Cancer Research Project. *Lancet* 2000; **356**: 93-96
  - 15 **Temple WJ**, Saettler EB. Locally recurrent rectal cancer: role of composite resection of extensive pelvic tumors with strategies for minimizing risk of recurrence. *J Surg Oncol* 2000; **73**: 47-58
  - 16 **Laurent C**, Nobili S, Rullier A, Vendrely V, Saric J, Rullier E. Efforts to improve local control in rectal cancer compromise survival by the potential morbidity of optimal mesorectal excision. *J Am Coll Surg* 2006; **203**: 684-691
  - 17 **Visser O**, Bakx R, Zoetmulder FA, Levering CC, Meijer S, Slors JF, van Lanschot JJ. The influence of total mesorectal excision on local recurrence and survival in rectal cancer patients: a population-based study in Greater Amsterdam. *J Surg Oncol* 2007; **95**: 447-454
  - 18 **Bernardshaw SV**, Ovrebø K, Eide GE, Skarstein A, Rokke O. Treatment of rectal cancer: reduction of local recurrence after the introduction of TME-experience from one University Hospital. *Dig Surg* 2006; **23**: 51-59
  - 19 **Luna-Perez P**, Bustos-Cholico E, Alvarado I, Maffuz A, Rodriguez-Ramirez S, Gutierrez de la Barrera M, Labastida S. Prognostic significance of circumferential margin involvement in rectal adenocarcinoma treated with preoperative chemoradiotherapy and low anterior resection. *J Surg Oncol* 2005; **90**: 20-25
  - 20 **Nagtegaal ID**, Marijnen CA, Kranenburg EK, van de Velde CJ, van Krieken JH. Circumferential margin involvement is still an important predictor of local recurrence in rectal carcinoma: not one millimeter but two millimeters is the limit. *Am J Surg Pathol* 2002; **26**: 350-357
  - 21 **Laurent C**, Nobili S, Rullier A, Vendrely V, Saric J, Rullier E. Efforts to improve local control in rectal cancer compromise survival by the potential morbidity of optimal mesorectal excision. *J Am Coll Surg* 2006; **203**: 684-691
  - 22 **Heald RJ**, Moran BJ, Ryall RD, Sexton R, MacFarlane JK. Rectal cancer: the Basingstoke experience of total mesorectal excision, 1978-1997. *Arch Surg* 1998; **133**: 894-899
  - 23 **Wan J**, Wu ZY, Du JL, Yao Y, Wang ZD, Lin HH, Luo XL, Zhang W. [Mesorectal metastasis of middle and lower rectal cancer] *Zhonghua Waike Zazhi* 2006; **44**: 894-896
  - 24 **Sugihara K**, Kobayashi H, Kato T, Mori T, Mochizuki H, Kameoka S, Shirouzu K, Muto T. Indication and benefit of pelvic sidewall dissection for rectal cancer. *Dis Colon Rectum* 2006; **49**: 1663-1672
  - 25 **Das P**, Skibber JM, Rodriguez-Bigas MA, Feig BW, Chang GJ, Hoff PM, Eng C, Wolff RA, Janjan NA, Delclos ME, Krishnan S, Levy LB, Ellis LM, Crane CH. Clinical and pathologic predictors of locoregional recurrence, distant metastasis, and overall survival in patients treated with chemoradiation and mesorectal excision for rectal cancer. *Am J Clin Oncol* 2006; **29**: 219-224
  - 26 **Park YA**, Lee KY, Kim NK, Baik SH, Sohn SK, Cho CW. Prognostic effect of perioperative change of serum carcinoembryonic antigen level: a useful tool for detection of systemic recurrence in rectal cancer. *Ann Surg Oncol* 2006; **13**: 645-650
  - 27 **Oh YT**, Kim MJ, Lim JS, Kim JH, Lee KY, Kim NK, Kim WH, Kim KW. Assessment of the prognostic factors for a local recurrence of rectal cancer: the utility of preoperative MR imaging. *Korean J Radiol* 2005; **6**: 8-16

S- Editor Li DL L- Editor Wang XL E- Editor Yin DH