



Cisapride in treatment of functional dyspepsia in 71 patients

Chang-Hua Wang, Ai-Xiu Ding, Xing-Jin Lu

Chang-Hua Wang, Ai-Xiu Ding, Xing-Jin Lu, Changzhou Guanghua Hospital, Changzhou 213001, Jiangsu Province, China

Author contributions: All authors contributed equally to the work.

Original title: *China National Journal of New Gastroenterology* (1995-1997) renamed *World Journal of Gastroenterology* (1998-).

Received: November 3, 1995

Revised: January 21, 1996

Accepted: May 19, 1996

Published online: September 15, 1996

Abstract

AIM: To observe the efficacy and safety of cisapride in treatment of functional dyspepsia.

METHODS: 71 patients (M 32, F 39; Age 41.3 ± 10.7 years) of functional dyspepsia were treated with cisapride in a dose of 5 mg

tid po for 4 wk.

RESULTS: The results showed that cisapride could significantly improve the symptoms including early satiety. Abdominal distention, epigastric pain, nausea, vomiting and anoxia. The pronounced response rate and the total response rate were 67.6% and 91.5%, respectively. The side effects were mild.

CONCLUSION: The above results indicated that the cisapride was safe and effective in treatment of functional dyspepsia.

Key words: Cisapride; Functional dyspepsia

© The Author(s) 1996. Published by Baishideng Publishing Group Inc. All rights reserved.

Wang CH, Ding AX, Lu XJ. Cisapride in treatment of functional dyspepsia in 71 patients. *World J Gastroenterol* 1996; 2(Suppl1): 210 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v2/iSuppl1/210.htm> DOI: <http://dx.doi.org/10.3748/wjg.v2.iSuppl1.210>

E- Editor: Liu WX



Published by **Baishideng Publishing Group Inc**
8226 Regency Drive, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>
<http://www.wjgnet.com>

