

World Journal of *Gastroenterology*

World J Gastroenterol 2017 September 7; 23(33): 6009-6196



**EDITORIAL**

- 6009** Helminths as an alternative therapy for intestinal diseases

Sipahi AM, Baptista DM

REVIEW

- 6016** Dextran sodium sulfate colitis murine model: An indispensable tool for advancing our understanding of inflammatory bowel diseases pathogenesis

Eichele DD, Kharbanda KK

- 6030** Autoimmune hepatitis: Standard treatment and systematic review of alternative treatments

Terziroli Beretta-Piccoli B, Mieli-Vergani G, Vergani D

MINIREVIEWS

- 6049** Colorectal cancer, screening and primary care: A mini literature review

Hadjipetrou A, Anyfantakis D, Galanakis CG, Kastanakis M, Kastanakis S

- 6059** Behavioral gastroenterology: An emerging system and new frontier of action

Jia L, Jiang SM, Liu J

ORIGINAL ARTICLE**Basic Study**

- 6065** Pharmacological evaluation of NSAID-induced gastropathy as a "Translatable" model of referred visceral hypersensitivity

Hummel M, Knappenberger T, Reilly M, Whiteside GT

- 6077** High yield reproducible rat model recapitulating human Barrett's carcinogenesis

Matsui D, Omstead AN, Kosovec JE, Komatsu Y, Lloyd EJ, Raphael H, Kelly RJ, Zaidi AH, Jobe BA

- 6088** Changes in expression of inhibitory substances in the intramural neurons of the stomach following streptozotocin- induced diabetes in the pig

Bulc M, Palus K, Zielonka L, Gajęcka M, Calka J

- 6100** HOX transcript antisense intergenic RNA represses E-cadherin expression by binding to EZH2 in gastric cancer

Chen WM, Chen WD, Jiang XM, Jia XF, Wang HM, Zhang QJ, Shu YQ, Zhao HB

- 6111 Ca^{2+} /calmodulin-dependent protein kinase II regulates colon cancer proliferation and migration *via* ERK1/2 and p38 pathways

Chen W, An P, Quan XJ, Zhang J, Zhou ZY, Zou LP, Luo HS

- 6119 Aberrant DNA-PKcs and ERGIC1 expression may be involved in initiation of gastric cancer

Wang FR, Wei YC, Han ZJ, He WT, Guan XY, Chen H, Li YM

Retrospective Cohort Study

- 6128 Real world treatment patterns of gastrointestinal neuroendocrine tumors: A claims database analysis

Benson III AB, Broder MS, Cai B, Chang E, Neary MP, Papayan E

Retrospective Study

- 6137 Patients with inflammatory bowel disease have increased risk of autoimmune and inflammatory diseases

Halling ML, Kjeldsen J, Knudsen T, Nielsen J, Koch-Hansen L

- 6147 Suspicious brush cytology is an indication for liver transplantation evaluation in primary sclerosing cholangitis

Boyd S, Vannas M, Jokelainen K, Isoniemi H, Mäkilä MA, Arola J

- 6155 Management of gastric mucosa-associated lymphoid tissue lymphoma in patients with extra copies of the *MALT1* gene

Iwamuro M, Takenaka R, Nakagawa M, Moritou Y, Saito S, Hori S, Inaba T, Kawai Y, Toyokawa T, Tanaka T, Yoshino T, Okada H

- 6164 Ketogenic diet poses a significant effect on imbalanced gut microbiota in infants with refractory epilepsy

Xie G, Zhou Q, Qiu CZ, Dai WK, Wang HP, Li YH, Liao JX, Lu XG, Lin SF, Ye JH, Ma ZY, Wang WJ

Observational Study

- 6172 Definition of colorectal anastomotic leakage: A consensus survey among Dutch and Chinese colorectal surgeons

van Rooijen SJ, Jongen ACHM, Wu ZQ, Ji JF, Slooter GD, Roumen RMH, Bouvy ND

CASE REPORT

- 6181 How to treat intestinal obstruction due to malignant recurrence after Whipple's resection for pancreatic head cancer: Description of 2 new endoscopic techniques

Mouradides C, Taha A, Borbath I, Deprez PH, Moreels TG

- 6187 Arteriportal shunt incidental to treatment with oxaliplatin that mimics recurrent gastric cancer

Kim HB, Park SG

LETTERS TO THE EDITOR

- 6194** Comment on "Effect of biofilm formation by clinical isolates of *Helicobacter pylori* on the efflux-mediated resistance to commonly used antibiotics"

Kazakos EI, Dorrell N, Polyzos SA, Deretzi G, Kountouras J

ABOUT COVER

Editorial Board Member of World Journal of Gastroenterology, Mostafa Sira, MD, Associate Professor of Pediatric Hepatology, Gastroenterology and Nutrition, National Liver Institute, Menofiya University, Menofiya 32511, Egypt

AIMS AND SCOPE

World Journal of Gastroenterology (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7th, 14th, 21st, and 28th each month. The *WJG* Editorial Board consists of 1375 experts in gastroenterology and hepatology from 68 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

INDEXING/ABSTRACTING

World Journal of Gastroenterology (*WJG*) is now indexed in Current Contents[®]/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch[®]), Journal Citation Reports[®], Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2017 edition of Journal Citation Reports[®] cites the 2016 impact factor for *WJG* as 3.365 (5-year impact factor: 3.176), ranking *WJG* as 29th among 79 journals in gastroenterology and hepatology (quartile in category Q2).

FLYLEAF

I-IX Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: Xiang Li
Responsible Electronic Editor: Yan Huang
Proofing Editor-in-Chief: Lian-Sheng Ma

Responsible Science Editor: Li-Juan Wei
Proofing Editorial Office Director: Jin-Lei Wang

NAME OF JOURNAL
World Journal of Gastroenterology

ISSN
ISSN 1007-9327 (print)
ISSN 2219-2840 (online)

LAUNCH DATE
October 1, 1995

FREQUENCY
Weekly

EDITORS-IN-CHIEF
Damian Garcia-Olmo, MD, PhD, Doctor, Professor, Surgeon, Department of Surgery, Universidad Autonoma de Madrid; Department of General Surgery, Fundacion Jimenez Diaz University Hospital, Madrid 28040, Spain

Stephen C Strom, PhD, Professor, Department of Laboratory Medicine, Division of Pathology, Karolinska Institutet, Stockholm 141-86, Sweden

Andrzej S Tarnawski, MD, PhD, DSc (Med), Professor of Medicine, Chief Gastroenterology, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach,

CA 90822, United States

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE
Jin-Lei Wang, Director
Yuan Qi, Vice Director
Ze-Mao Gong, Vice Director
World Journal of Gastroenterology
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>

<http://www.wjgnet.com>

PUBLICATION DATE
September 7, 2017

COPYRIGHT
© 2017 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.f6publishing.com>

Basic Study

Ca²⁺/calmodulin-dependent protein kinase II regulates colon cancer proliferation and migration *via* ERK1/2 and p38 pathways

Wei Chen, Ping An, Xiao-Jing Quan, Jun Zhang, Zhong-Yin Zhou, Li-Ping Zou, He-Sheng Luo

Wei Chen, Ping An, Xiao-Jing Quan, Jun Zhang, Zhong-Yin Zhou, He-Sheng Luo, Department of Gastroenterology, Renmin Hospital of Wuhan University, Wuhan 430060, Hubei Province, China

Li-Ping Zou, Department of Education, Renmin Hospital of Wuhan University, Wuhan 430060, Hubei Province, China

Ping An, Division of Gastroenterology and Hepatology, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA 02115, United States

Author contributions: Chen W and An P contributed equally to this work; An P designed the research; Chen W, Quan XJ and Zhang J performed the research; Zhou ZY and Zou LP analyzed the data; Luo HS and An P wrote the paper.

Supported by the National Natural Science Foundation of China, No. 81302131.

Institutional review board statement: This study was approved by the Institutional Review Board of Renmin Hospital of Wuhan University.

Conflict-of-interest statement: The authors have no relevant conflicts of interest to disclose.

Data sharing statement: Data can be obtained from the corresponding author.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Correspondence to: He-Sheng Luo, Professor, Department of

Gastroenterology, Renmin Hospital of Wuhan University, 238 Jiefang Road, Wuhan 430060, Hubei Province, China. xhnhk@163.com
Telephone: +86-27-88041911
Fax: +86-27-88041911

Received: February 6, 2017

Peer-review started: February 8, 2017

First decision: May 12, 2017

Revised: June 10, 2017

Accepted: July 4, 2017

Article in press: July 4, 2017

Published online: September 7, 2017

Abstract

AIM

To investigate the role of calmodulin-dependent protein kinase II (CaMK II) in colon cancer growth, migration and invasion.

METHODS

CaMK II expression in colon cancer and paracancerous tissues was evaluated *via* immunohistochemistry. Transcriptional and posttranscriptional levels of CaMK II in tissue samples and MMP2, MMP9 and TIMP-1 expression in the human colon cancer cell line HCT116 were assessed by qRT-PCR and western blot. Cell proliferation was detected with the MTT assay. Cancer cell migration and invasion were investigated with the Transwell culture system and wound-healing assay.

RESULTS

We first demonstrated that CaMK II was over-expressed in human colon cancers and was associated with cancer differentiation. In the human colon cancer cell line HCT116, the CaMKII-specific inhibitor KN93, but not its inactive analogue KN92, decreased cancer cell proliferation. Furthermore, KN93 also significantly

prohibited HCT116 cell migration and invasion. The specific inhibition of ERK1/2 or p38 decreased the proliferation and migration of colon cancer cells.

CONCLUSION

Our findings highlight CaMK II as a potential critical mediator in human colon tumor development and metastasis.

Key words: Ca²⁺/calmodulin-dependent protein kinase II; Colon cancer; Proliferation; Migration

© **The Author(s) 2017.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: In the present study, we demonstrated that calmodulin-dependent protein kinase II (CaMK II) was over-expressed in human colon cancers and was associated with cancer differentiation. We investigated the role of CaMKII in human colon cancer proliferation and migration. The results revealed that in the human colon cancer cell line HCT116, the CaMK II-specific inhibitor KN93, but not its inactive analogue KN92, decreased cancer cell proliferation. KN93 also significantly prohibited colon cancer cell migration and invasion. Additionally, we found that ERK1/2 and p38 were the targets of CaMK II regulation. These findings highlight CaMK II as a potential critical mediator in human colon tumor development and metastasis.

Chen W, An P, Quan XJ, Zhang J, Zhou ZY, Zou LP, Luo HS. Ca²⁺/calmodulin-dependent protein kinase II regulates colon cancer proliferation and migration via ERK1/2 and p38 pathways. *World J Gastroenterol* 2017; 23(33): 6111-6118 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v23/i33/6111.htm> DOI: <http://dx.doi.org/10.3748/wjg.v23.i33.6111>

INTRODUCTION

Colorectal cancer (CRC) is the third most common cancer and the fourth leading cause of cancer death globally. Researchers estimate that roughly 1.4 million new cases of CRC will be diagnosed and 694000 deaths will result from CRC per year^[1]. Despite improvements in multimodal anticancer strategies, the prognosis of advanced CRC is still poor, with 5-year survival rates for stage III and stage IV colon cancer at 65.4% and 12.8%, respectively^[2]. A detailed understanding of the biological processes that modulate the progression and metastasis of CRC may provide new molecular pathogenesis targets for CRC diagnosis and benefit anti-tumor therapy.

The calcium ion (Ca²⁺) is a ubiquitous intracellular signal responsible for a broad range of cellular events, such as cell growth, cytoskeletal organization, regulation of synaptic transmission, and Ca²⁺ homeostasis^[3]. The Ca²⁺/calmodulin (CaM)-dependent protein kinases (CaMK I, CaMK II and CaMKIV) are

multifunctional serine/threonine kinases whose activity is activated upon Ca²⁺/CaM binding^[4]. The upstream CaMK kinases (CaMKKs) phosphorylate a critical Thr 200 in the activation-loop to activate CaMKIV, whereas CaMKII is fully activated by the autophosphorylation of its own Thr 286^[5]. Studies have revealed that the calcium signal is involved in cancer cell proliferation, invasion, tumor growth and metastasis^[6-9]. However, fewer studies have focused on CaMKII, one of the most important sensors and regulators of the Ca²⁺ signal, in digestive cancers, especially in CRC.

In the present study, we first demonstrated that CaMKII was over-expressed in human colon cancers and was associated with cancer differentiation. We further sought to investigate the role of CaMKII in human colon cancer proliferation and migration. The results revealed that in the human colon cancer cell line HCT116, the CaMKII-specific inhibitor KN93, but not its inactive analogue KN92, decreased cancer cell proliferation. KN93 also significantly prohibits HCT116 cell migration and invasion. Additionally, we determined that CaMKII inhibition decreased the phosphorylation of ERK1/2 and p38. The specific inhibition of ERK1/2 or p38 decreased the proliferation and migration of colon cancer cells. These findings highlight CaMKII as a potential critical mediator in human colon tumor development and metastasis.

MATERIALS AND METHODS

Reagents

Antibodies against CaMKIIα, ERK1/2, phospho (p)-ERK1/2, p38, p-p38, MMP2, MMP9, TIMP-1 and GAPDH were purchased from Santa Cruz Biotechnology (Santa Cruz, CA, United States). KN-93 (2-[N-(2-Hydroxyethyl)]-N-(4-methoxybenzene-sulfonyl)] amino-N-(4-chlorocinnamyl)-N-methylbenzylamine), KN-92 (2-[N-(4-Methoxybenzenesulfonyl)] amino-N-(4-chlorocinnamyl)-N-methylbenzylamine), PD98059 and SB203580 were obtained from Calbiochem (La Jolla, CA, United States). Other chemicals of the highest purity were purchased from Sigma (St. Louis, MO, United States).

Tissue samples

Colon tissue samples were obtained from the Department of Gastrointestinal Surgery at Renmin Hospital of Wuhan University in accordance with local ethics committees. All samples were obtained from the primary colon sites of pretreatment cases of paracancerous lesions (*n* = 5), well-differentiated colon cancer tissues (*n* = 6) and poorly differentiated colon cancer tissues (*n* = 6). All samples were analyzed and assessed by two histological specialists working blindly.

Cell culture and treatment

The human colon cancer cell line HCT116 was obtained from the Type Culture Collection of the Chinese Academy of Sciences (Shanghai, China) and cultured

in Dulbecco's modified Eagle's medium (DMEM) (Gibco, Rockville, MD, United States) containing 10% heat-inactivated fetal bovine serum (FBS) (Gibco) and 1% penicillin/streptomycin (Life Technologies, Rockville, MD, United States) at 37 °C in a humidified atmosphere with 5% CO₂. For conditioned media, 3 × 10⁵ HCT116 cells were seeded in 6-well plates and then stimulated for the indicated amount of time.

Immunostaining

Immunohistochemistry (IHC) for CaMKII was performed on paraffin-embedded tissues from paracancerous tissues, well-differentiated colon cancers and poorly differentiated colon cancers. Specimens were fixed in a 4% formaldehyde solution for 12-48 h and embedded in paraffin. IHC was performed on 3.5 µm tissue sections on slides. Heat-induced epitope retrieval was carried out. For immunofluorescent staining for CaMKII, all slides were blocked with serum solution and incubated with the primary antibody overnight at 4 °C. After rinsing again, the slides were incubated for 45 min at room temperature with secondary antibodies. Visualization was achieved with the DAB (3,3'-diaminobenzidine) detection kit, and the cell nuclei were counterstained using hematoxylin. After staining, five fields (at a magnification of × 100) were randomly selected in each slide. Staining was quantified with Image J software and evaluated by two investigators following the "blinded" principle.

MTT assay

Cell proliferation was measured with the MTT assay (ATCC, Manassas, VA, United States). A total of 5 × 10³ cells were seeded at the well bottoms of 96-well culture plates and treated with TGFβ1 or conditioned medium from fibroblasts or colon cancer cells in triplicate. After treatment, MTT solution [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazoliumbromide] was added to the culture medium (0.5 mmol/L), and plates were incubated for 2 h at 37 °C with 5% CO₂. Detergent solution was then added to solubilize formazan crystals. Finally, the optical density was determined at 540 nm with a Benchmark Plus microplate reader (Bio-Rad, Hercules, CA, United States).

Western blot analysis

Protein from unstained paraffin-embedded samples was deparaffinized and rehydrated *via* successive washes in xylene/ethanol and water. After incubation in extraction buffer at 100 °C for 20 min followed by a 2 h incubation at 80 °C, proteins were obtained by centrifugation. Whole cell lysates were prepared with RIPA lysis buffer containing protease and phosphatase inhibitor cocktail. Protein concentrations of cell lysates were determined with the bicinchoninic acid protein assay. A total of 30-40 µg of lysate were loaded onto a 10% SDS-PAGE gel and subjected to gel electrophoresis. Resolved proteins were transferred to a polyvinylidene fluoride membrane. Membranes

were then blocked in 5% non-fat dry milk in Tris-buffered saline with Tween (TBS-T) for 1 h and then incubated with antibodies to GAPDH, CaMKIIα (Cell Signaling Technology), ERK1/2, phospho-ERK1/2, p38, p-p38, MMP2, MMP9 and TIMP-1 at 4 °C overnight. Membranes were washed with TBS-T and then incubated with horseradish peroxidase-conjugated secondary antibody for 1 h at room temperature. The blotting bands were visualized with enhanced chemiluminescence. GAPDH was used as a loading control.

Quantitative real-time PCR

Cells were seeded at 10⁶ cells per 10 cm dish and were allowed to grow to 80% confluency in complete media. Cells were removed with 0.025% ethylenediaminetetraacetic acid and centrifuged for 5 min at 1100 rpm. Cell pellets were resuspended in 1 mL of TRIzol (Life Technologies, Carlsbad, CA, United States), and RNA was extracted according to the manufacturer's protocol. Total RNA in unstained paraffin-embedded samples was extracted with the RNeasy FFPE Kit (Qiagen, Hilden, Germany). The SuperScript First-Strand Kit (Life Technologies) was used to synthesize cDNA from 5 µg of total RNA. Quantitative PCR was established with the RT2 SYBR Green Fluor FAST Mastermix (Qiagen, Venlo, the Netherlands) and run on a Bio-Rad CFX96 Real-Time PCR Detection System (Bio-Rad). Data were analyzed with Bio-Rad CFX Manager 3.0 software and are shown as relative fold-changes.

For all PCR reactions, GAPDH was used as an endogenous control, and CT values were normalized to levels of GAPDH expression. Primers were designed with Integrated DNA Technologies PrimerQuest software (Coralville, IA, United States). Sequences used to analyze RNA expression include the following: CaMKII, Forward: 5'-GAG AGC ACC AAC ACC ACC ATC G-3' and Reverse: 5'-AGG CTG ACT CGT CGC CCA TCA GG-3'; MMP2, Forward: 5'-TCT CCT GAC ATT GAC CTT GGC-3' and Reverse: 5'-CAA GGT GCT GGC TGA GTA GAT C-3'; MMP9, Forward: 5'-TTG ACA GCG ACA AGA AGT GG-3' and Reverse: 5'-GCC ATT CAC GTC GTC CTT AT-3'; TIMP-1, Forward: 5'-CTT CTG GCA TCC TGT TGT TG-3' and Reverse: 5'-GGT ATA AGG TGG TCT GGT TG-3'; GAPDH, Forward: 5'-ACA GTC CAT GCC ATC ACT GCC-3' and Reverse: 5'-GCC TGC TTC ACC ACC TTC TTG-3'. The results are averaged from three independent experiments.

Migration and invasion assay

A 24-well Transwell plate (Fisher Scientific, Hampton, NH, United States) was used to measure migratory and invasive ability *in vitro*. A total of 2 × 10⁵ HCT-116 cells were plated on the bottom of each well. For the migration assay, 1 × 10⁵ colon cancer cells were plated in the top chamber with a non-coated membrane. For the invasion assay, 2 × 10⁵ HCT-116 cells were seeded in the top chamber coated with Matrigel (BD Biosciences, San Jose, CA, United States). In both




Figure 1 CaMKII over-expressed in colon cancer. A: Immunohistochemistry staining of CaMKII in paracancerous tissue, well-differentiated colon cancer and poorly differentiated colon cancer samples. Scale bar, 100 μ m; B: qRT-PCR of CaMKII mRNA in paraffin-embedded paracancerous tissue, well-differentiated colon cancer and poorly differentiated colon cancer samples; C: Western blot of CaMKII protein in paraffin-embedded paracancerous tissue, well-differentiated colon cancer and poorly differentiated colon cancer samples. All data are presented as the mean \pm standard error of at least three independent experiments. ^a $P < 0.05$, ^b $P < 0.01$ vs paracancerous tissue.

assays, cells were cultured in DMEM containing 10% FBS with the indicated treatment. After incubation at 37 °C for 24 h, migrated cells were fixed and stained with the Diff-Quik stain and counted in four random fields. The experiments were performed in triplicate wells, and each experiment was performed at least two or three times as indicated.

Wound-healing assay

A wound-healing assay was performed with 6-well plates. HCT116 cells were seeded at confluency 24 h before the experiment. The cell layers were carefully scratched with 200-mL sterile pipette tips and washed twice with fresh medium. The cells were photographed under a light microscope at $\times 200$ magnification 24 h later.

Statistical analysis

All experiments were performed in triplicate. Data are expressed as the mean \pm SD. Differences between

groups were calculated *via* one-way analysis of variance. Differences were considered significant at $P < 0.05$.

RESULTS

CaMKII is over-expressed in colon cancers

The expression of CaMKII was evaluated by IHC with a monoclonal CaMKII antibody on sections obtained from formalin-fixed, paraffin-embedded samples of paracancerous tissue, well-differentiated colon cancer samples and poorly differentiated colon cancer samples. As expected, CaMKII was significantly overexpressed in the poorly differentiated colon cancer samples and well-differentiated cancer samples compared to the paracancerous tissues (Figure 1A). A quantification assay suggested that the expression of CaMKII was highest in poorly differentiated colon cancer tissues. We further evaluated CaMKII protein and mRNA levels in cancer and paracancerous




Figure 2 CaMKII is required for colon cancer proliferation and migration. Human colon cancer cells HCT116 were treated with or without KN93 or KN92 at 10 μ mol/L for 24 h. A: MTT analysis for cell proliferation; B: Western blot of MMP2, MMP9 and TIMP-1; C: qRT-PCR assay of MMP2, MMP9 and TIMP-1; D: Immigration and invasion analysis by T transwell assay; E: Wound-healing test. All data are presented as the mean \pm standard error SE of at least three independent experiments. ^a $P < 0.05$, ^b $P < 0.01$ vs control group.




Figure 3 CaMKII regulates colon cancer cells proliferation and migration via ERK1/2- and p38-dependent pathways. Human colon cancer cells HCT116 were treated with or without PD98059 (50 μ mol/L) or SB203580 (10 μ mol/L) for 24 h. A: MTT assay for cell proliferation; B: Western blot analysis of ERK1/2, phosphor-ERK1/2, p38 and phosphor-p38; C: Immigration and invasion analysis by T transwell assay. All data are presented as the mean \pm standard error SE of at least three independent experiments. ^a $P < 0.05$, vs control group.

tissues. The results were similar to the findings of the above studies (Figure 1B and C). We deduced that CaMKII may be a potential regulator in colon cancer development.

CaMKII is required for colon cancer proliferation and migration

CaMKII has been shown to drive cancer growth and metastasis. To ascertain whether CaMKII regulates

colon cancer proliferation and migration, the CaMKII specific inhibitor KN93 was selected. Human colon cancer HCT116 cells were treated with KN93, and its inactive analogue, KN92, was used as an additional control. We found that KN93 inhibited colon cancer cell proliferation (Figure 2A). Importantly, in migration and invasion experiments, the motility of HCT116 cells was significantly attenuated (Figure 2B and C). Furthermore, CaMKII inhibition decreased MMP2 and MMP9 expression at both transcriptional and posttranscriptional levels. The opposite results were detected for TIMP-1 expression (Figure 2D and E). Thus, we concluded that CaMKII is required for colon cancer cell proliferation and migration.

CaMKII regulates colon cancer cell proliferation and migration via ERK1/2 and p38-dependent pathways

We further investigated the molecular mechanism of CaMKII regulation in human colon cancer cells. As shown in Figure 3, KN93, but not KN92, decreased the phosphorylation of ERK1/2 and p38, which are involved in colon cancer development. Pretreatment with specific inhibitors of ERK1/2 or p38 (PD98059, SB203580) decreased HCT116 cell proliferation, migration and invasion. These data demonstrated that the effects of CaMKII on human colon cancer are ERK1/2- and p38-dependent.

Taken together, these results indicate that CaMKII is an important regulator of human colon cancer growth and migration.

DISCUSSION

It is widely accepted that intracellular Ca^{2+} signaling pathways regulate various biological events, including cell proliferation, motility, activation and differentiation, during cancer development^[10,11]. In the present study, we demonstrated that CaMKII is over-expressed in colon cancers and highly expressed in poorly differentiated colon tumors. We highlight the role of CaMKII in colon cancer growth and metastasis.

CaMKII inhibitors can inhibit CaMKII activity by interacting with the Ca^{2+} /CaM-binding site or interfering with its catalytic activities. Previous studies revealed that CaMKII-specific inhibitors such as KN-62, KN-93, and autocamtide 2-related inhibitory peptide inhibit CaMKII-dependent processes in tumor and normal cells, causing cell cycle arrest, cellular apoptosis, or the inhibition of cell proliferation^[12-14].

Some studies have implicated CaMKII as an important player in cancer cell proliferation. CaMKII inhibition decreased ovarian cancer cell motility and decreased tumor growth and metastasis^[15]. The *in vivo* administration of KN-93 to mice xenografted with human osteosarcoma cells significantly decreased intratibial and subcutaneous tumor growth^[16]. In CRC, we demonstrated that CaMKII inhibition strongly

inhibited human colon cancer cell proliferation (Figure 2A).

Recent studies support the important role of CaMKII in cancer invasion and metastasis. The up-regulation of CaMKII α was found in primary osteosarcoma tissues from patients and in aggressive osteosarcoma cell lines^[17]. CaMKII α depletion decreased motility and invasion, whereas CaMKII α over-expression increased the tumorigenic properties of osteosarcoma cells *in vitro*. Our current results revealed the important role of CaMKII in colon cancer migration and invasion. KN93 dramatically decreased the expression of MMP2 and MMP9, increased TIMP-1 levels and prohibited the invasive motility of colon cancer cells (Figure 2). CaMKII is a critical regulator of colon cancer progression.

The MAPK pathway is one of the important downstream targets in CaMKII regulation^[4]. CaMKII directly or indirectly up-regulated multiple signaling pathways, such as ERK1/2, AKT1 and β -catenin, and is involved in regulating the survival and proliferation of non-small cell lung cancer cells^[6]. We further investigated whether ERK1/2 and p38 were involved in CaMKII pathways. Our data showed that the inhibition of CaMKII by KN93 significantly decreased ERK1/2 and p38 phosphorylation (Figure 3). The specific inhibition of ERK1/2 or p38 decreased the proliferation, migration and invasion of HCT116 cells. Therefore, CaMKII mediates colon cancer growth and migration via ERK1/2- and p38-dependent pathways.

Taken together, the current study provided evidence for CaMKII as a pivotal regulator of colon cancer cell proliferation and migration. CaMKII is a potentially important target in CRC treatment.

COMMENTS

Background

Ca^{2+} /calmodulin (CaM)-dependent protein kinase II (CaMKII) is one of the most important sensors and regulators of the Ca^{2+} signal which is involved in cancer cell proliferation, invasion, tumor growth and metastasis. Despite the critical role of CaMKII, fewer studies focus on it in digestive cancers, especially in colorectal cancer.

Research frontiers

Recently, studies have revealed CaMKII is an effective controller in ovarian cancer and osteosarcoma cell motility, cell growth and metastasis.

Innovations and breakthroughs

In the present study, the authors demonstrated that CaMKII was over-expressed in human colon cancers and was associated with cancer differentiation. Meanwhile, in the human colon cancer cell line HCT116, the CaMKII-specific inhibitor KN93, but not its inactive analogue KN92, decreased cancer cell proliferation. CaMKII inhibition also significantly prohibited colon cancer cell migration and invasion. Additionally, ERK1/2 and p38 were the targets of CaMKII regulation.

Applications

These findings highlight CaMKII as a potential critical mediator in human colon tumor development and metastasis. Its relationship to colon cancer

differentiation indicates CaMKII a potential target for clinical diagnosis and prognosis.

Terminology

CaMKII is a ubiquitous serine/threonine protein kinase. Autophosphorylation of CaMKII at threonine 286 switches the kinase from Ca^{2+} -dependent to Ca^{2+} -independent activity and leads to sustained CaMKII auto-activation. More importantly, this activation is thought to phosphorylate numerous different proteins and regulate diverse signaling pathways.

Peer-review

This is a brief study concerning the role of Ca^{2+} /calmodulin-dependent protein kinase II involved in invasion and migration of the colon cancer. The study is interesting and topical. The experiments were well designed. The manuscript is clear and structured very well.

REFERENCES

- 1 Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015; **136**: E359-E386 [PMID: 25220842 DOI: 10.1002/ijc.29210]
- 2 Lan YT, Yang SH, Chang SC, Liang WY, Li AF, Wang HS, Jiang JK, Chen WS, Lin TC, Lin JK. Analysis of the seventh edition of American Joint Committee on colon cancer staging. *Int J Colorectal Dis* 2012; **27**: 657-663 [PMID: 22146786 DOI: 10.1007/s00384-011-1366-6]
- 3 Leung CS, Yeung TL, Yip KP, Pradeep S, Balasubramanian L, Liu J, Wong KK, Mangala LS, Armaiz-Pena GN, Lopez-Berestein G, Sood AK, Birrer MJ, Mok SC. Calcium-dependent FAK/CREB/TNNC1 signalling mediates the effect of stromal MFAP5 on ovarian cancer metastatic potential. *Nat Commun* 2014; **5**: 5092 [PMID: 25277212 DOI: 10.1038/ncomms6092]
- 4 Wang YY, Zhao R, Zhe H. The emerging role of CaMKII in cancer. *Oncotarget* 2015; **6**: 11725-11734 [PMID: 25961153 DOI: 10.18632/oncotarget.3955]
- 5 Hook SS, Means AR. Ca^{2+} /CaM-dependent kinases: from activation to function. *Annu Rev Pharmacol Toxicol* 2001; **41**: 471-505 [PMID: 11264466 DOI: 10.1146/annurev.pharmtox.41.1.471]
- 6 Chai S, Qian Y, Tang J, Liang Z, Zhang M, Si J, Li X, Huang W, Xu R, Wang K. RETRACTED: Ca^{2+} /calmodulin-dependent protein kinase II γ , a critical mediator of the NF- κ B network, is a novel therapeutic target in non-small cell lung cancer. *Cancer Lett* 2014; **344**: 119-128 [PMID: 24189456 DOI: 10.1016/j.canlet.2013.10.022]
- 7 Britschgi A, Bill A, Brinkhaus H, Rothwell C, Clay I, Duss S, Rebhan M, Raman P, Guy CT, Wetzel K, George E, Popa MO, Lilley S, Choudhury H, Gosling M, Wang L, Fitzgerald S, Borawski J, Baffoe J, Labow M, Gaither LA, Bentires-Alj M. Calcium-activated chloride channel ANO1 promotes breast cancer progression by activating EGFR and CAMK signaling. *Proc Natl Acad Sci USA* 2013; **110**: E1026-E1034 [PMID: 23431153 DOI: 10.1073/pnas.1217072110]
- 8 Wang T, Guo S, Liu Z, Wu L, Li M, Yang J, Chen R, Liu X, Xu H, Cai S, Chen H, Li W, Xu S, Wang L, Hu Z, Zhuang Q, Wang L, Wu K, Liu J, Ye Z, Ji JY, Wang C, Chen K. CAMK2N1 inhibits prostate cancer progression through androgen receptor-dependent signaling. *Oncotarget* 2014; **5**: 10293-10306 [PMID: 25296973 DOI: 10.18632/oncotarget.2511]
- 9 Wang C, Li N, Liu X, Zheng Y, Cao X. A novel endogenous human CaMKII inhibitory protein suppresses tumor growth by inducing cell cycle arrest via p27 stabilization. *J Biol Chem* 2008; **283**: 11565-11574 [PMID: 18305109 DOI: 10.1074/jbc.M800436200]
- 10 Zayzafoon M. Calcium/calmodulin signaling controls osteoblast growth and differentiation. *J Cell Biochem* 2006; **97**: 56-70 [PMID: 16229015 DOI: 10.1002/jcb.20675]
- 11 Bourguignon LY, Gilad E, Brightman A, Diedrich F, Singleton P. Hyaluronan-CD44 interaction with leukemia-associated RhoGEF and epidermal growth factor receptor promotes Rho/Ras co-activation, phospholipase C epsilon- Ca^{2+} signaling, and cytoskeleton modification in head and neck squamous cell carcinoma cells. *J Biol Chem* 2006; **281**: 14026-14040 [PMID: 16565089 DOI: 10.1074/jbc.M507734200]
- 12 An P, Tian Y, Chen M, Luo H. Ca^{2+} /calmodulin-dependent protein kinase II mediates transforming growth factor- β -induced hepatic stellate cells proliferation but not in collagen $\alpha 1(\text{I})$ production. *Hepatol Res* 2012; **42**: 806-818 [PMID: 22414022 DOI: 10.1111/j.1872-034X.2012.00983.x]
- 13 Eleftheriadis T, Pissas G, Karioti A, Antoniadis G, Goulinopoulos S, Liakopoulos V, Mamara A, Speletas M, Koukoulis G, Stefanidis I. Uric acid induces caspase-1 activation, IL-1 β secretion and P2X7 receptor dependent proliferation in primary human lymphocytes. *Hippokratia* 2013; **17**: 141-145 [PMID: 24376319]
- 14 An P, Zhu JY, Yang Y, Lv P, Tian YH, Chen MK, Luo HS. KN-93, a specific inhibitor of CaMKII inhibits human hepatic stellate cell proliferation in vitro. *World J Gastroenterol* 2007; **13**: 1445-1448 [PMID: 17457979 DOI: 10.3748/wjg.v13.i9.1445]
- 15 Ma S, Yang Y, Wang C, Hui N, Gu L, Zhong H, Cai Z, Wang Q, Zhang Q, Li N, Cao X. Endogenous human CaMKII inhibitory protein suppresses tumor growth by inducing cell cycle arrest and apoptosis through down-regulation of the phosphatidylinositol 3-kinase/Akt/HDM2 pathway. *J Biol Chem* 2009; **284**: 24773-24782 [PMID: 19586909 DOI: 10.1074/jbc.M109.028621]
- 16 Yuan K, Chung LW, Siegal GP, Zayzafoon M. α -CaMKII controls the growth of human osteosarcoma by regulating cell cycle progression. *Lab Invest* 2007; **87**: 938-950 [PMID: 17632540 DOI: 10.1038/labinvest.3700658]
- 17 Daft PG, Yuan K, Warram JM, Klein MJ, Siegal GP, Zayzafoon M. α -CaMKII plays a critical role in determining the aggressive behavior of human osteosarcoma. *Mol Cancer Res* 2013; **11**: 349-359 [PMID: 23364534 DOI: 10.1158/1541-7786.MCR-12-0572]

P- Reviewer: Pan HC, Perse M S- Editor: Ma YJ

L- Editor: Filipodia E- Editor: Xu XR





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>



ISSN 1007-9327



9 771007 932045