

World Journal of *Gastroenterology*

World J Gastroenterol 2018 March 21; 24(11): 1181-1284



**REVIEW**

- 1181 Functional macrophages and gastrointestinal disorders

Liu YH, Ding Y, Gao CC, Li LS, Wang YX, Xu JD

ORIGINAL ARTICLE**Basic Study**

- 1196 *NOD2*- and disease-specific gene expression profiles of peripheral blood mononuclear cells from Crohn's disease patients

Schäffler H, Rohde M, Rohde S, Huth A, Gittel N, Hollborn H, Koczan D, Glass A, Lamprecht G, Jaster R

- 1206 Three-microRNA signature identified by bioinformatics analysis predicts prognosis of gastric cancer patients

Zhang C, Zhang CD, Ma MH, Dai DQ

Retrospective Cohort Study

- 1216 Differing profiles of people diagnosed with acute and chronic hepatitis B virus infection in British Columbia, Canada

Binka M, Butt ZA, Wong S, Chong M, Buxton JA, Chapinal N, Yu A, Alvarez M, Darvishian M, Wong J, McGowan G, Torban M, Gilbert M, Tyndall M, Krajden M, Janjua NZ

- 1228 Role of relevant immune-modulators and cytokines in hepatocellular carcinoma and premalignant hepatic lesions

Zekri AN, El Deeb S, Bahnassy AA, Badr AM, Abdellateif MS, Esmat G, Salama H, Mohanad M, El-dien AE, Rabah S, Abd Elkader A

Retrospective Study

- 1239 Serum autotaxin levels are correlated with hepatic fibrosis and ballooning in patients with non-alcoholic fatty liver disease

Fujimori N, Umemura T, Kimura T, Tanaka N, Sugiura A, Yamazaki T, Joshita S, Komatsu M, Usami Y, Sano K, Igarashi K, Matsumoto A, Tanaka E

- 1250 Epidemiological features of chronic hepatitis C infection caused by remunerated blood donors: A nearly 27-year period survey

Tan YW, Tao Y, Liu LG, Ye Y, Zhou XB, Chen L, He C

Clinical Trials Study

- 1259 Low-FODMAP vs regular rye bread in irritable bowel syndrome: Randomized SmartPill® study

Pirkola L, Laatikainen R, Lopenon J, Hongisto SM, Hillilä M, Nuora A, Yang B, Linderborg KM, Freese R

Prospective Study

- 1269** Fatty liver in hepatitis C patients post-sustained virological response with direct-acting antivirals

Noureddin M, Wong MM, Todo T, Lu SC, Sanyal AJ, Mena EA

- 1278** Low-pressure pneumoperitoneum with abdominal wall lift in laparoscopic total mesorectal excision for rectal cancer: Initial experience

Xia PT, Yusofu M, Han HF, Hu CX, Hu SY, Yu WB, Liu SZ

ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Chang Moo Kang, MD, PhD, Associate Professor, Division of Hepatobiliary and Pancreatic Surgery, Department of Surgery, Yonsei University, Pancreaticobiliary Cancer Clinic, Yonsei Cancer Center, Severance Hospital, Seoul 120-752, South Korea

AIMS AND SCOPE

World Journal of Gastroenterology (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7th, 14th, 21st, and 28th each month. The *WJG* Editorial Board consists of 642 experts in gastroenterology and hepatology from 59 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

INDEXING/ABSTRACTING

World Journal of Gastroenterology (*WJG*) is now indexed in Current Contents[®]/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch[®]), Journal Citation Reports[®], Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2017 edition of Journal Citation Reports[®] cites the 2016 impact factor for *WJG* as 3.365 (5-year impact factor: 3.176), ranking *WJG* as 29th among 79 journals in gastroenterology and hepatology (quartile in category Q2).

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: Xiang Li
Responsible Electronic Editor: Yan Huang
Proofing Editor-in-Chief: Lian-Sheng Ma

Responsible Science Editor: Ze-Mao Gong
Proofing Editorial Office Director: Jin-Lei Wang

NAME OF JOURNAL
World Journal of Gastroenterology

ISSN
ISSN 1007-9327 (print)
ISSN 2219-2840 (online)

LAUNCH DATE
October 1, 1995

FREQUENCY
Weekly

EDITORS-IN-CHIEF
Damian Garcia-Olmo, MD, PhD, Doctor, Professor, Surgeon, Department of Surgery, Universidad Autonoma de Madrid; Department of General Surgery, Fundacion Jimenez Diaz University Hospital, Madrid 28040, Spain

Stephen C Strom, PhD, Professor, Department of Laboratory Medicine, Division of Pathology, Karolinska Institutet, Stockholm 141-86, Sweden

Andrzej S Tarnawski, MD, PhD, DSc (Med), Professor of Medicine, Chief Gastroenterology, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach,

CA 90822, United States

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE
Ze-Mao Gong, Director
World Journal of Gastroenterology
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
March 21, 2018

COPYRIGHT
© 2018 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.f6publishing.com>

Retrospective Cohort Study

Role of relevant immune-modulators and cytokines in hepatocellular carcinoma and premalignant hepatic lesions

Abdel-Rahman N Zekri, Somaya El Deeb, Abeer A Bahnassy, Abeer M Badr, Mona S Abdellateif, Gamal Esmat, Hosny Salama, Marwa Mohanad, Ahmed Esam El-dien, Shimaa Rabah, Assmaa Abd Elkader

Abdel-Rahman N Zekri, Molecular Virology and Immunology Unit, Department of Cancer Biology, National Cancer Institute, Cairo University, Cairo 11976, Egypt

Somaya El Deeb, Abeer M Badr, Ahmed Esam El-dien, Shimaa Rabah, Assmaa Abd Elkader, Department of Zoology, Faculty of Science, Cairo University, Giza 12613, Egypt

Abeer A Bahnassy, Department of Pathology, National Cancer Institute, Cairo University, Cairo 11976, Egypt

Mona S Abdellateif, Medical Biochemistry and Molecular Biology, Department of Cancer Biology, National Cancer Institute, Cairo University, Cairo 11976, Egypt

Gamal Esmat, Hosny Salama, Department of Hepatology and Tropical Medicine, Faculty of Medicine, Cairo University, Cairo 11441, Egypt

Marwa Mohanad, Department of Biochemistry, Misr University for Science and Technology, 6th October 12945, Giza Governorate, Egypt

ORCID number: Abdel-Rahman N Zekri (0000-0003-3939-0416); Somaya El Deeb (0000-0002-1826-1503); Abeer A Bahnassy (0000-0001-5454-6576); Abeer M Badr (0000-0003-3503-6639); Mona S Abdellateif (0000-0002-5510-4435); Gamal Esmat (0000-0001-8614-1629); Hosny Salama (0000-0003-0670-3770); Marwa Mohanad (0000-0002-3827-9647); Ahmed Esam El-dien (0000-0002-7357-3235); Shimaa Rabah (0000-0003-0477-8672); Assmaa Abd Elkader (0000-0001-7945-1610).

Author contributions: Zekri AN was the main supervisor, put forth the idea and planned of work, and performed data interpretation; El Deeb S supervised the biochemical work; Bahnassy AA supervised the flow cytometry work and revised the paper; Badr AM revised the paper; Abdellateif MS analyzed the data and wrote the paper; Esmat G and Salama H performed the medical care and the follow-up of patients; Mohanad M analyzed the data; El-dien AE performed the immunophenotyping; Rabah

S and Abd Elkader A performed the ELISA work.

Institutional review board statement: The study was reviewed and approved for publication by our Institutional Reviewer.

Informed consent statement: All study participants or their legal guardian provided informed written consent about personal and medical data collection prior to study enrolment.

Conflict-of-interest statement: All the authors have no conflict of interest related to the manuscript.

Data sharing statement: The original anonymized dataset is available on request from the corresponding author at ncizekri@yahoo.com.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Abdel-Rahman N Zekri, MSc, PhD, Professor, Molecular Virology and Immunology Unit, Department of Cancer Biology, National Cancer Institute, Cairo University, Kasr Al-Aini Street, Fom El-Khaleeg, Cairo 11976, Egypt. ncizekri@yahoo.com
Telephone: +2-10-01413521
Fax: +2-202-23644720

Received: November 5, 2017

Peer-review started: November 6, 2017

First decision: November 27, 2017

Revised: December 24, 2017

Accepted: January 16, 2018

Article in press: January 16, 2018
Published online: March 21, 2018

Group Inc. All rights reserved.

Abstract

AIM

To assess the levels of different immune modulators in patients with hepatocellular carcinoma (HCC), in relation to other hepatic diseases.

METHODS

Eighty-eight patients were included in the current study and represented patients with HCC (20), liver cirrhosis (28) and chronic hepatitis (CH; 25), and normal controls (NC; 15). Peripheral blood was isolated for immunophenotyping of active myeloid dendritic cells (mDCs; CD1c and CD40), mature inactive myeloid cells (CD1c and HLA), active plasmacytoid cells (pDCs; CD303 and CD40), mature inactive pDCs (CD30 and HLA), active natural killer (NK) cells (CD56 and CD161), active NK cells (CD56 and CD314) and inactive NK cells (CD56 and CD158) was done by flow cytometry. Serum levels of interleukin (IL)-2, IL-10, IL-12, IL-1 β , interferon (IFN)- α , IFN- γ and tumor necrosis factor (TNF)- α R2 were assessed by ELISA.

RESULTS

Active mDCs (CD1c+/CD40+) and inactive mDCs (CD1c+/HLA+) were significantly decreased in HCC patients in relation to NC ($P < 0.001$). CD40+ expression on active pDCs was decreased in HCC patients ($P < 0.001$), and its level was not significantly changed among other groups. Inactive pDCs (CD303+/HLA+), inactive NKs (CD56+/CD158+) and active NKs (CD56+/CD161+) were not statistically changed among the four groups studied; however, the latter was increased in CH ($P < 0.05$). NKG2D was statistically decreased in HCC, CH and cirrhosis ($P < 0.001$), and it was not expressed in 63% (12/20) of HCC patients. There was significant decrease of IL-2, IFN- α and IFN- γ ($P < 0.001$), and a significant increase in IL-10, IL-1 β , and TNF- α R2 ($P < 0.01$, $P < 0.001$ and $P < 0.001$; respectively) in HCC patients. There was inverted correlation between IL-12 and IL-1 β in HCC ($r = -0.565$, $P < 0.01$), with a strong correlation between pDCs (CD303+/CD40+) and NKs (CD56+/CD161+; $r = 0.512$, $P < 0.05$) as well as inactive mDCs (CD1c+/HLA+) and inactive NK cells (CD56+/CD158+; $r = 0.945$, $P < 0.001$).

CONCLUSION

NKG2D, CD40, IL-2 and IL-10 are important modulators in the development and progression of HCC.

Key words: Hepatocellular carcinoma; Hepatitis C virus; NKG2D; CD40; Interleukin-2; Interleukin-10; Myeloid dendritic cells; Plasmacytoid cells; Natural killer cell; Cytokines

© The Author(s) 2018. Published by Baishideng Publishing

Core tip: We assessed the levels of different immune modulatory cytokines and innate immune cells as natural killer (NK) cells and dendritic cells (DCs) in patients with disease progression of hepatocarcinogenesis. Our results showed significant down-regulation in active mDCs and pDCs expressing CD40 as well as NK cells expressing NKG2D. The expression of NKG2D on NKs was not expressed in 63% of hepatocellular carcinoma (HCC) patients. Also, there was significant decrease of interleukin (IL)-2, interferon- α and interferon- γ , and a significant increase in IL-10, IL-1 β , and TNF- α R2 in HCC patients. These factors could be implicated in the pathogenesis of HCC, and represent attractive targets for therapy in chronic hepatitis C virus hepatitis and HCC.

Zekri AN, El Deeb S, Bahnassy AA, Badr AM, Abdellateif MS, Esmat G, Salama H, Mohanad M, El-dien AE, Rabah S, Abd Elkader A. Role of relevant immune-modulators and cytokines in hepatocellular carcinoma and premalignant hepatic lesions. *World J Gastroenterol* 2018; 24(11): 1228-1238 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v24/i11/1228.htm> DOI: <http://dx.doi.org/10.3748/wjg.v24.i11.1228>

INTRODUCTION

Hepatocellular carcinoma (HCC) is a major health problem worldwide. It is the sixth most common cancer and the second leading cause of cancer-related death in the world^[1]. In Egypt, HCC ranks the first among cancers in males (33.6%), and the 2nd in females after breast cancer (13.5%)^[2]. This high incidence is attributed to the high prevalence of hepatitis C virus (HCV) infection, especially of genotype IV, in Egypt^[3]. Despite advances in treatment modalities, sorafenib is still the only treatment approved by the Food and Drug Administration (FDA) for HCC; however, it extends the overall survival by 3 mo only. Hence, it is crucial to understand the underlying biological and immunological changes in HCC and to develop new treatment modalities based on these data^[4,5].

The body's immune defense mechanism(s) plays an important role in the inhibition or progression of cancer. However, tumors can escape immune surveillance by producing a local and/or systemic immune-suppressive environment^[6]. Although the underlying molecular mechanisms are not yet fully clear, recent studies show that the immune response in cancer patients is usually down-regulated by immunosuppressive cells, mainly T regulatory cells (Tregs) and myeloid-derived suppressor cells (MDSCs), which suppress the immune system and promote immunologic tolerance^[7]. These inhibitory cells accumulate during advanced cancer stages. They are involved in chronic inflammation and tumor progression and they can inhibit many immune cells

(e.g., CD4+, CD8+, natural killer (NK) cells)^[8]. They also secrete many immune-suppressive cytokines, such as interleukin (IL)-6, IL-10, and transforming growth factor (TGF)- β , creating a tolerogenic and suppressive environment^[9].

The NK cells represent 25%-30% of the human liver lymphocytes, compared to 10%-20% of the total peripheral blood lymphocytes^[10]. They are the main effector cells of innate immunity, which play an important role in tumor surveillance. The NK cells achieve their functions through the release of cytolytic mediators, such as perforin and granzymes, or the induction of apoptosis *via* the expression of tumor necrosis factor (TNF) ligands and interferon (IFN)- γ ^[11]. Activation of NK cells is strictly regulated by a balance between activating and inhibitory signals, whereas the inhibitory signals are mainly induced by receptors for MHC class I molecules (KIRs, CD94/NKG2A). The activating signals are mainly achieved through NKG2D, a C-type lectin-like receptor which is expressed on NK cells, $\gamma\delta$ T cells and CD8+ T cells^[12,13]. NKG2D is responsible for detection and elimination of transformed cells *via* binding to MICA (MHC class I-related chain), MICB and the UL16-binding proteins^[14]. Activation of NKG2D provides unique costimulation to antigen-specific CD8 T cells that is non-redundant to CD28 costimulation^[15].

Dendritic cells (DCs) are the most professional antigen-presenting cells which respond rapidly to microenvironment signals. Upon maturation by tumor antigen, cross-priming to T and B lymphocytes occurs to produce antitumor adaptive immune responses^[16]. There are two subsets of DCs: the myeloid dendritic cells (mDCs), which are characterized by their ability to produce IL-12; and, the plasmacytoid cells (pDCs), which are responsible for the production of over 95% of type-I IFNs in response to viral infection^[17]. On activation, the CD8+ cytotoxic T cells (CTLs) eliminate tumor cells by the production of IFN- γ , whereas CD4+ T helper cells stimulate B cells to support the cytotoxic and humoral immune response^[18].

IL-1 β is a regulatory cytokine produced by tumor and immune cells, such as MDSCs. It also induces COX-2 expression, which prevents maturation and activation of antigen presenting cells at the tumor site^[19].

In the current study, we assessed the role(s) of different immune regulatory cells and cytokines in the development and progression of chronic liver disease (chronic active hepatitis, cirrhosis and HCC) compared to the normal healthy volunteers. We believe that this could allow for better understanding of different pathways implicated in the development and progression of hepatocarcinogenesis, and could also help in designing new immune-therapeutic drugs.

MATERIALS AND METHODS

Patients

The current study included 88 patients who attended

the medical oncology clinics of the National Cancer Institute (NCI), Cairo University during the period from 2014 to 2016. Patients were divided into four groups: HCC (20 patients - G1), liver cirrhosis (28 patients - G2), chronic hepatitis (CH; 25 patients - G3) and normal healthy volunteers as a control group (NC; 15 persons, matched for age and sex - G4). The Ethical Committee of the NCI, Egypt approved the study protocol and an informed consent was obtained from all participants before enrollment in the study.

Characterization of immune cells by flow cytometry

Peripheral blood samples were isolated from patients and healthy subjects for immunophenotyping of active mDCs (CD1c and CD40), mature inactive myeloid cells (CD1c and HLA), active pDCs (CD303 and CD40), mature inactive pDCs (CD303 and HLA), active NK cells (CD56 and CD161), active NK cells (CD56F CD314) and inactive NK cells (CD56 and CD158) using the specific monoclonal antibodies according to manufacturer's protocols (all monoclonal antibodies were purchased from Invitrogen, eBioscience, San Diego, CA, United States). Samples were then analyzed by flow cytometry (FACSCaliber; Becton and Dickinson, Franklin Lakes, NJ, United States).

Detection of cytokine levels by ELISA

Serum levels of IL-2, IL-10, IL-12, IL-1 β , IFN- α , IFN- γ and TNF- α R2 were measured by the ELISA technique, according to manufacturer's instructions (Invitrogen, eBioscience), using ELISA reader (Sunrise; Tecan, Mannedorf, Switzerland).

Statistical analysis

Data were expressed as mean \pm SE of mean for continuous variables. Comparison between cytokine levels and immune cells were performed using one-way analysis of variance followed by Tukeys' post hoc test. Pearson's correlation was used to assess the strength of correlation between different variables and a two-tailed *P*-value was determined. The *P*-value was considered significant at < 0.05 . All statistical analyses were performed using SPSS, version 22 (IBM SPSS, Armonk, NY, United States).

RESULTS

Patients' characteristics

The mean age of the patients included in the current study was 57.1 ± 4.76 for HCC, patients, 51.29 ± 9.03 for liver cirrhosis patients, 49.64 ± 7.2 for CH patients and 42.9 ± 10.2 for NC (Table 1).

Flow cytometry data

There was a significant decrease in active mDCs (CD1c+/CD40+) and inactive mDCs (CD1c+/HLA+) in HCC patients compared to the NC group ($P < 0.001$), as well as in active mDCs (CD1c+/CD40+) in cirrhotic patients compared to CH patients ($P = 0.003$; Table

Table 1 Patient characteristics

	Normal	Chronic hepatitis	Cirrhosis	HCC
Sex				
Male	11 (73.3)	9 (36.0)	19 (67.9)	19 (95.0)
Female	4 (26.7)	16 (64.0)	9 (32.1)	1 (5.0)
Age, mean \pm SD	42.9 \pm 10.2	49.64 \pm 7.2	51.29 \pm 9.03	57.1 \pm 4.76
HCV	+ve	+ve	+ve	+ve
Total	15	25	28	20

Data are presented as *n* (%), unless otherwise specified. HCC: Hepatocellular carcinoma; HCV: Hepatitis C virus.




Figure 1 Heatmap of the differential levels of (A) active myeloid dendritic cells, (B) inactive myeloid dendritic cells, (C) active plasmacytoid cells, and (D) inactive plasmacytoid cells in the four groups studied. HCC: Hepatocellular carcinoma.

2). Also, the expression level of CD40+ on active pDCs (CD303+) was significantly decreased in HCC compared to the NC group ($P < 0.001$). However, there was no significant difference from the other groups (cirrhosis and CH). Meanwhile, the level of inactive pDCs (CD303+/HLA+) did not differ significantly between the four groups studied (Table 2, Figure 1).

The level of the nonactive NK cells (CD56+/CD158+) did not differ significantly among groups. However, the active NK cells (CD56+/CD161+) showed a significant increase in the CH group ($P < 0.05$), whereas the level of active NK cells (CD56+/CD314+) was statistically

decreased in HCC, CH and cirrhotic patients compared to the NC group ($P < 0.001$), indicating an important role of these cells in the pathogenesis of HCC (Table 2). The expression of all cell types in each patient showed that active NK cells (CD56+/CD314+) were not expressed in 63% (12/20) of HCC patients. However, 2 patients showed unexplainable high variability (Figure 2).

We also found a significant decrease in the active NK cells (CD56+/CD314+; $P < 0.05$) and active pDCs (CD303+, CD40+; $P < 0.05$) compared to the inactive NK cells (CD56+/CD158+) and inactive pDCs (CD303+/HLA+) respectively in HCC. To the contrary,

Table 2 Differential counts of nature killer cells and dendritic cells among the studied groups

	Normal	Chronic hepatitis	Cirrhosis	HCC	P value
Active mDCs (CD1C+/CD40+)	652.4 ± 15.9 ^{A1}	154.1 ± 40.9 ^B	25.5 ± 4.3 ^C	82.8 ± 28.7 ^{BC}	< 0.001
Inactive mature mDCs (CD1c+/HLA+)	627.1 ± 14.8 ^A	134.4 ± 37.1 ^B	54.7 ± 12.3 ^B	121.8 ± 44.8 ^B	< 0.001
Active pDCs (CD303+/CD40+)	782.2 ± 89.6 ^A	237.8 ± 56.2 ^B	164.1 ± 45.9 ^B	251 ± 55.9 ^B	< 0.001
Inactive mature pDCs (CD303+/HLA+)	727.5 ± 18.7 ^A	318.5 ± 61.4 ^A	385.7 ± 233.3 ^A	339.2 ± 67.5 ^A	0.339
Active NK cells (CD56+/CD161+)	535 ± 83.4 ^{AB}	1166.3 ± 426.3 ^B	399.8 ± 139.8 ^{AB}	145.5 ± 30.2 ^A	< 0.05
Inactive NK cells (CD56+/CD158+)	710.3 ± 127.9 ^A	656.3 ± 235.7 ^A	462.6 ± 105.3 ^A	456.7 ± 218.1 ^A	0.708
Active NK cells (CD56+/CD314+)	585.5 ± 35.9 ^A	188.5 ± 69.4 ^B	123.5 ± 47.6 ^B	196 ± 110.8 ^B	< 0.001

Data are presented as mean ± SEM. ¹Groups having the same letters in the same variable are statistically insignificant. HCC: Hepatocellular carcinoma; mDCs: Myeloid dendritic cells; NK: Natural killer; pDCs: Plasmacytoid cells.




Figure 2 Heatmap of the differential levels of (A) active natural killer cells (CD56+/CD314+), (B) inactive natural killer cells (CD56+/CD158+), and (C) natural killer cells (CD56+/CD161+) in the four groups studied. HCC: Hepatocellular carcinoma.

the NK cells (CD56+/CD161+) and the mDCs did not differ significantly between groups (Figure 3).

Serum levels of cytokines

There was a significant decrease in serum levels of IL-2, IFN- α and IFN- γ in HCC patients compared to the NC group ($P < 0.001$), and a significant increase in serum levels of IL-10, IL-1 β and TNF- α R2 in HCC

compared to the NC group ($P < 0.01$, $P < 0.001$ and $P < 0.001$ respectively). However, there was no statistically significant change in the serum level of IL-12 ($P = 0.393$) among the four groups studied (Table 3, Figure 4).

Significant correlation was also reported in HCC patients between (1) serum level of IL-12 and IL-1 β ($r = -0.565$, $P < 0.01$), (2) expression of CD40 on

Table 3 Levels of the studied cytokines among different studied groups

	Normal	Chronic hepatitis	Cirrhosis	HCC	P value
IL-2	22.13 ± 2.21 ^{A1}	16.28 ± 2.12 ^B	11.36 ± 0.95 ^C	10.24 ± 0.64 ^C	< 0.001
IL-10	74.81 ± 1.34 ^A	98.46 ± 4.01 ^{AB}	104.11 ± 9.35 ^B	132.1 ± 14.26 ^C	< 0.01
IL-12	0.06 ± 0.01 ^A	0.09 ± 0.01 ^A	0.15 ± 0.06 ^A	0.08 ± 0.01 ^A	0.393
IL-1β	3.46 ± 0.15 ^A	6.48 ± 0.58 ^A	6.71 ± 2.39 ^A	21.38 ± 4.88 ^B	< 0.001
IFN-α	31.20 ± 0.67 ^A	32.28 ± 1.44 ^A	21.31 ± 1.01 ^B	18.49 ± 1.37 ^B	< 0.001
IFN-γ	20.47 ± 0.49 ^A	29.75 ± 0.68 ^B	17.51 ± 1.18 ^C	10.46 ± 0.56 ^D	< 0.001
TNF-αR2	6.03 ± 0.07 ^A	13.25 ± 1.73 ^A	10.09 ± 0.97 ^A	47.42 ± 6.74 ^B	< 0.001

Data are presented as mean ± SEM. ¹Groups having the same letters in the same variable are statistically insignificant. HCC: Hepatocellular carcinoma; IFN: Interferon; IL: Interleukin; TNF: Tumor necrosis factor.




Figure 3 Balance between active and inactive natural killer cells, plasmacytoid cells, and myeloid dendritic cells in hepatocellular carcinoma patients in relation to the normal group. ^a $P < 0.05$. HCC: Hepatocellular carcinoma; mDCs: Myeloid dendritic cells; NK: Natural killer; pDCs: Plasmacytoid cells.

pDCs (CD303+, CD40+) and active NK cells (CD56+/CD161+; $r = 0.512$, $P < 0.05$), and (3) inactive mDCs (CD1c+/HLA+) and inactive NK cells (CD56+/CD158+; $r = 0.945$, $P < 0.001$) (Figure 5A-C).

Patients with chronic active hepatitis showed strong significant correlation between active NK cells (CD56+/CD161+) and both IL-2 and IL-12 ($r = 0.549$, $P = 0.004$ and $r = 0.660$, $P < 0.001$ respectively). Strong correlations were also reported between the expression of CD314 on NK cells and IL-2 ($r = 0.548$, $P < 0.01$) and active mDCs CD1c+/CD40+ ($r = 0.577$, $P = 0.003$). The level of IL-10 was strongly correlated with (1) increased inactive NK cells (CD56+/CD158+; $r = 0.604$, $P = 0.001$) and (2) inactive mDCs CD1c+/HLA+ ($r = 0.588$, $P = 0.002$; Table 4).

DISCUSSION

Identification of inflammatory mediators which promote

or prevent the progression of cancer depends on the tumor microenvironment. Thus, this study was designed to assess the levels of different immune modulators and cytokines in patients with HCC, nonmalignant hepatic diseases (e.g., CH and liver cirrhosis) compared to a normal group. This would enable us to study different pathways that help in identifying immune cells or biomarkers that may be implicated in hepatocellular carcinogenesis. This may also explain why liver disease progresses more rapidly in some patients than in others and provides additional therapeutic modalities for viral infections and HCC (immunotherapy).

One of the most powerful antigen presenting cells is the DC, which plays an important role in antitumor immune response. Our data revealed that active and inactive mDCs were significantly decreased in HCC patients compared to the NC group ($P < 0.001$); however, the level of active mDCs (CD1c+/CD40+) was significantly decreased in cirrhosis compared to CH

Table 4 Correlation between dendritic cell and nature killer cell count and the levels of studied cytokines in chronic active hepatitis

	IL-2	IL-12	IL-10	Active mDCs (CD1c+/CD40+)
Active NK cells (CD56+/CD161+)	$r = 0.549^b$ $P = 0.004$	$r = 0.660^b$ $P < 0.001$		
Active NK cells (CD56+/CD314+)	$r = 0.548^a$ $P = 0.005$			$r = 0.577^b$ $P = 0.003$
Inactive NK cells (CD56+/CD158+)			$r = 0.604^b$ $P = 0.001$	
Inactive mDCs (CD1c+/HLA+)			$r = 0.588^b$ $P = 0.002$	

^a $P < 0.05$ (2-tailed); ^b $P < 0.01$ (2-tailed). IL: Interleukin; mDCs: Myeloid dendritic cells; NK: Natural killer.

patients ($P = 0.003$). These data are consistent with Obermajer *et al.*^[20], who reported that the inhibition of DC maturation in HCC may prove to be a critical feature of tumor escape, as well as with another two recent studies which reported numerical and functional defects in the peripheral DCs in hepatitis B- and C-associated HCC patients^[21,22].

The pDC (CD303+, CD40+) represents the major cell responsible for antiviral immunity, and the main source of IFN- α in the body. Our data revealed that the expression level of CD40+ on active pDCs was significantly decreased in HCC patients compared to NC ($P < 0.001$). Meanwhile, inactive pDCs (CD303+/HLA+) did not change significantly among the four groups studied. This was also confirmed by the significant decrease of serum IFN- α , and the increase of IL-10 in HCC patients compared to the NC group ($P < 0.001$ and $P < 0.01$ respectively). This is in concordance with previous reports showing that IL-10 inhibits IFN- α production and promotes apoptosis of human pDCs^[23,24]. Moreover, Gonzalez-Carmona *et al.*^[25] demonstrated that cancer patients had low CD40 expression on DCs or CD40L on T cells, which is associated with an impaired immune response, and Shurin *et al.*^[26] reported that the tumor-derived IL-10 inhibits CD40 expression on DCs and DC precursors and suppresses their maturation and function.

NK cells play an important role in controlling viral hepatitis, liver fibrosis and carcinogenesis. Their functions are modulated by different classes of receptors present on its surface^[27]. Thus, it is essential to identify the roles of these cells in different stages of HCC development. Among these, we investigated the role of the killer cell immunoglobulin-like receptors (KIR; CD158), NKR-P1A (CD161) and NKG2D (CD314) in HCV CH patients, and in liver cirrhosis and its progression to HCC.

Our results revealed that the expression of the inhibitory KIR (CD158+) receptors and the activating NKR-P1A (CD161) on NK cells was not statistically changed between HCC and NC ($P = 0.827$ and $P = 0.788$ respectively). Thus, these two pathways may not be involved in the pathogenesis of HCC. However, the latter were statistically increased in CH ($P < 0.05$). This result was supported by our finding of a strong

statistical correlation between the expression of CD161 on NK cells and IL-12 ($r = 0.77$, $P < 0.001$) in patients with HCV chronic active hepatitis. This illustrates the role of NK cells (CD56+/CD161+) in viral hepatitis.

On the other hand, the level of active NK cells expressing NKG2D were statistically decreased in HCC, CH and cirrhosis in relation to the NC group ($P < 0.001$), which could indicate the implication of these cells in the progression of HCC. This was clarified by representing its expression in each patient, it was not expressed in 63% (12/20) of HCC patients, emphasizing its important role in HCC development. These data were consistent with Lanier *et al.*^[28]. Yoon *et al.*^[29] reported that direct interaction between human NK cells and HCV-infected hepatoma cells down-regulates NK cell NKG2D expression and effector function with diminished IFN- γ production. Another study carried out by Kamiya *et al.*^[30] demonstrated that NKG2D was an important mediator of antiHCC activity. On the contrary, Oliviero *et al.*^[31] found a decreased percentage of NK cells expressing the inhibitory receptor KIR3DL1 and a concomitant increase in the proportion of NKG2D⁺ NK cells in Italian patients with chronic HCV. This could be explained by racial and environmental differences with different viral genotypes, and also according to the clinical stage.

Our results regarding DCs and NK cells were supported by assessment of the cytokine serum levels in the four groups studied. We found that there was a significant increase in IL-1 β ($P < 0.001$) in HCC patients compared to the NC group. However, there was no statistically significant change in the serum level of IL-12 ($P = 0.393$) among the four groups studied. Furthermore, we had a significant increase in the serum level of IL-10 in the cirrhosis and HCC groups compared to the NC group. This is consistent with Accapezzato *et al.*^[32]. IL-10 is important for allowing liver NK cells to maintain immune-tolerant states^[33]. Its significant increase in HCC patients is responsible for the abnormal tolerant NK cells that cannot eliminate infected or transformed cells, which leads to immune evasion.

In the current study, there was a significant decrease of the serum levels of IFN- γ in chronic HCV hepatitis, cirrhosis and HCC patients in comparison to NC ($P < 0.001$). This is in agreement with Li *et al.*^[34],




Figure 4 Different levels of serum cytokines in the four groups studied. A: IL-2; B: IL-10; C: IL-12; D: IL-1B; E: IFN- α ; F: IFN- γ ; G: TNF- α R2. IFN: Interferon; IL: Interleukin; TNF: Tumor necrosis factor.




Figure 5 Correlation between immune cells and cytokine levels in hepatocellular carcinoma patients. IL: Interleukin; NK: Natural killer.

who found that NK cells cultured with cancer-associated fibroblasts from HCC (H-CAFs) down-regulate NKG2D and NKp46 and decrease expression of granzyme B, perforin and $\text{INF-}\gamma$. Later, another study showed that $\text{TGF-}\beta$ and IL-10 in hepatic carcinoma patients induce the expression of microRNA (miR)-146a, which causes reduced $\text{INF-}\gamma$ production and cytotoxicity, resulting in a poorer prognosis^[35]. Meanwhile, Zhang *et al.*^[36] found that MDSCs selectively suppressed the $\text{INF-}\gamma$ production deriving from NKT cells through membrane-bound $\text{TGF-}\beta$.

IL-2 plays a critical role in activation of the immune system, which could be used to eradicate cancer. It has been demonstrated as a monotherapy, and was approved for metastatic renal cell carcinoma and metastatic melanoma by the FDA^[37]. Our results showed a significant increase in $\text{TNF-}\alpha\text{R2}$ ($P < 0.001$) and significant decrease in IL-2 ($P < 0.001$) in HCC patients in relation to other groups, consistent with our previous results^[38,39] that concluded the possible use of serum TNFR-II , IL-2Ra and IL-8 as combined biomarkers in HCV-infected patients at high risk of developing HCC.

Another important finding in the current study was that HCC patients had a significant decrease of active NK cells ($\text{CD56+}/\text{CD314+}$; $P < 0.05$) and active pDCs (CD303+ , CD40+ ; $P < 0.05$) in comparison to inactive NK cells ($\text{CD56+}/\text{CD158+}$) and inactive pDCs ($\text{CD303+}/\text{HLA+}$) respectively. NK cells expressing

$\text{CD56+}/\text{CD161+}$ and mDCs were not statistically affected. This revealed the importance of these two subsets of cells [active NK cells ($\text{CD56+}/\text{CD314+}$), and active pDCs (CD303+ , CD40+)] in the pathogenesis of HCC, and allows for further research around the state of this imbalance that emerges in HCC patients.

We conclude that there are immunological changes occurring in HCC patients which could be possible candidate(s) for future immunotherapy for these patients. Among these are the NK cells expressing NKG2D and pDCs expressing CD40, IL-2 and IL-10. These factors could be implicated in the pathogenesis of HCC, and provide an attractive target for therapeutics in chronic HCV hepatitis and liver cancer.

ARTICLE HIGHLIGHTS

Research background

Hepatocellular carcinoma (HCC) is a major health problem worldwide and mainly in Egypt due to the high prevalence of hepatitis C virus (HCV) infection, especially of genotype IV. It ranks the first among cancers in males (33.6%), and the 2nd in females after breast cancer. Sorafenib is still the only treatment approved by the Food and Drug Administration for HCC; however, it extends the overall survival by 3 mo only. Hence, it is crucial to understand the underlying biological and immunological changes in HCC Egyptian patients, and to develop new treatment modalities based on these data.

Research motivation

The immune system plays an important role in suppression of cancer. However, tumors can escape immune surveillance by producing an immune-suppressive

environment, for which the underlying mechanisms are not fully clear. Recent studies show that the immune response in HCC patients is usually down-regulated by immunosuppressive cells (myeloid-derived suppressor cells and T regulatory cells) that are involved in chronic inflammation and tumor progression. Also, these inhibitory cells secrete many immune-suppressive cytokines, such as interleukin (IL)-6, IL-10 and transforming growth factor- β , creating a tolerogenic and suppressive environment^[9].

Research objectives

The objective of this study was to assess the levels of different immune modulators and cytokines that may play a role in the pathogenesis of HCC and other hepatic diseases (e.g., chronic hepatitis and liver cirrhosis) compared to a normal group. This would help in identifying additional therapeutic modalities for viral infections and HCC in the context of immunotherapy.

Research methods

This retrospective cohort study included 88 patients who attended the medical oncology clinics of the National Cancer Institute, Cairo University during the period from 2014 to 2016. Patients were divided into the HCC group (20 patients-G1), liver cirrhosis group (28 patients-G2), chronic hepatitis group (CH; 25 patients-G3) and normal healthy volunteers as a control group (NC; 15 persons). The immune system of the patients was assessed through immunophenotyping of CD1c and CD40, CD1c and HLA, CD303 and CD40, CD303 and HLA, CD56 and CD161, CD56 and CD314, and CD56 and CD158 by flow cytometry. On the other hand, serum levels of IL-2, IL-10, IL-12, IL-1 β , interferon (IFN)- α , IFN- γ and tumor necrosis factor (TNF)- α R2 were measured by the ELISA technique, according to the manufacturer's instructions. Data were expressed as mean \pm SE of mean, and statistical comparison between cytokine levels and immune cells were performed.

Research results

There was a significant decrease in active and inactive mDCs in HCC patients compared to the NC group, as well as active mDCs (CD1C+/CD40+) in cirrhotic patients compared to CH patients. The expression level of CD40+ on active pDCs (CD303+) was significantly decreased in HCC compared to the NC. However, there was no significant difference with the other groups (cirrhosis and CH). Meanwhile, the level of inactive pDCs (CD303+/HLA+) and inactive NK cells (CD56+/CD158+) did not differ significantly between the four groups studied.

The active NK cells (CD56+/CD161+) showed a significant increase in the CH group, whereas the level of active NK cells (CD56+/CD314+) was statistically decreased in HCC, CH and cirrhotic patients compared to the NC group, indicating an important role of these cells in the pathogenesis of HCC. The individual expression of each cell type in patients showed that active NK cells (CD56+/CD314+) were not expressed in 63% (12/20) of HCC patients.

There was a significant decrease in serum levels of IL-2, IFN- α and IFN- γ in HCC patients compared to the NC group, and a significant increase in serum levels of IL-10, IL-1 β and TNF- α R2 in HCC compared to the NC group. However, there was no statistically significant change in the serum level of IL-12 among the four groups studied.

Research conclusions

We conclude that there are immunological changes occurring in HCC patients in relation to other liver diseases. The related immunological factors are NKG2D expressed on NK cells, and pDCs expressing CD40, IL-2 and IL-10. These factors could be implicated in the pathogenesis of HCC, and represent attractive targets for therapeutics in chronic HCV hepatitis and liver cancer.

Research perspectives

NKG2D, CD40, IL-2 and IL-10 could be a possible candidate for future immunotherapy for HCC patients. However, further studies are recommended regarding correlation of these factors and clinicopathological features as well as the overall survival of the patients.

REFERENCES

1 Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo

- M, Parkin DM, Forman D, Bray F. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015; **136**: E359-E386 [PMID: 25220842 DOI: 10.1002/ijc.29210]
- 2 Ibrahim AS, Khaled HM, Mikhail NN, Baraka H, Kamel H. Cancer incidence in Egypt: results of the national population-based cancer registry program. *J Cancer Epidemiol* 2014; **2014**: 437971 [PMID: 25328522 DOI: 10.1155/2014/437971]
- 3 Zekri AR, Hafez MM, Bahnassy AA, Hassan ZK, Mansour T, Kamal MM, Khaled HM. Genetic profile of Egyptian hepatocellular-carcinoma associated with hepatitis C virus Genotype 4 by 15 K cDNA microarray: preliminary study. *BMC Res Notes* 2008; **1**: 106 [PMID: 18959789 DOI: 10.1186/1756-0500-1-106]
- 4 Knudsen ES, Gopal P, Singal AG. The changing landscape of hepatocellular carcinoma: etiology, genetics, and therapy. *Am J Pathol* 2014; **184**: 574-583 [PMID: 24388934 DOI: 10.1016/j.ajpath.2013.10.028]
- 5 Llovet JM, Villanueva A, Lachenmayer A, Finn RS. Advances in targeted therapies for hepatocellular carcinoma in the genomic era. *Nat Rev Clin Oncol* 2015; **12**: 408-424 [PMID: 26054909 DOI: 10.1038/nrclinonc.2015.103]
- 6 Ostrand-Rosenberg S. Tolerance and immune suppression in the tumor microenvironment. *Cell Immunol* 2016; **299**: 23-29 [PMID: 26435343 DOI: 10.1016/j.cellimm.2015.09.011]
- 7 Gabrilovich DI, Ostrand-Rosenberg S, Bronte V. Coordinated regulation of myeloid cells by tumours. *Nat Rev Immunol* 2012; **12**: 253-268 [PMID: 22437938 DOI: 10.1038/nri3175]
- 8 Shrihari TG. Dual role of inflammatory mediators in cancer. *Ecancermedicalscience* 2017; **11**: 721 [PMID: 28275390 DOI: 10.3332/ecancer.2017.721]
- 9 Costa A, Scholer-Dahirel A, Mechta-Grigoriou F. The role of reactive oxygen species and metabolism on cancer cells and their microenvironment. *Semin Cancer Biol* 2014; **25**: 23-32 [PMID: 24406211 DOI: 10.1016/j.semcancer.2013.12.007]
- 10 Zeromski J, Mozer-Lisewska I, Kaczmarek M, Kowala-Piaskowska A, Sikora J. NK cells prevalence, subsets and function in viral hepatitis C. *Arch Immunol Ther Exp (Warsz)* 2011; **59**: 449-455 [PMID: 21972016 DOI: 10.1007/s00005-011-0145-y]
- 11 Zhong Z, Sanchez-Lopez E, Karin M. Autophagy, Inflammation, and Immunity: A Troika Governing Cancer and Its Treatment. *Cell* 2016; **166**: 288-298 [PMID: 27419869 DOI: 10.1016/j.cell.2016.05.051]
- 12 Lanier LL. NK cell recognition. *Annu Rev Immunol* 2005; **23**: 225-274 [PMID: 15771571 DOI: 10.1146/annurev.immunol.23.021704.115526]
- 13 Dai Z, Turtle CJ, Booth GC, Riddell SR, Gooley TA, Stevens AM, Spies T, Groh V. Normally occurring NKG2D+CD4+ T cells are immunosuppressive and inversely correlated with disease activity in juvenile-onset lupus. *J Exp Med* 2009; **206**: 793-805 [PMID: 19289577 DOI: 10.1084/jem.20081648]
- 14 Guerra N, Tan YX, Joncker NT, Choy A, Gallardo F, Xiong N, Knoblaugh S, Cado D, Greenberg NM, Raulet DH. NKG2D-deficient mice are defective in tumor surveillance in models of spontaneous malignancy. *Immunity* 2008; **28**: 571-580 [PMID: 18394936 DOI: 10.1016/j.immuni.2008.02.016]
- 15 Wu J, Groh V, Spies T. T cell antigen receptor engagement and specificity in the recognition of stress-inducible MHC class I-related chains by human epithelial gamma delta T cells. *J Immunol* 2002; **169**: 1236-1240 [PMID: 12133944 DOI: 10.4049/jimmunol.169.3.1236]
- 16 Burkholder B, Huang RY, Burgess R, Luo S, Jones VS, Zhang W, Lv ZQ, Gao CY, Wang BL, Zhang YM, Huang RP. Tumor-induced perturbations of cytokines and immune cell networks. *Biochim Biophys Acta* 2014; **1845**: 182-201 [PMID: 24440852 DOI: 10.1016/j.bbcan.2014.01.004]
- 17 Tel J, van der Leun AM, Figdor CG, Torensma R, de Vries IJ. Harnessing human plasmacytoid dendritic cells as professional APCs. *Cancer Immunol Immunother* 2012; **61**: 1279-1288 [PMID: 22294456 DOI: 10.1007/s00262-012-1210-z]
- 18 Korniluk A, Koper O, Kemona H, Dymicka-Piekarska V. From

- inflammation to cancer. *Ir J Med Sci* 2017; **186**: 57-62 [PMID: 27156054 DOI: 10.1007/s11845-016-1464-0]
- 19 **Kanterman J**, Sade-Feldman M, Baniyash M. New insights into chronic inflammation-induced immunosuppression. *Semin Cancer Biol* 2012; **22**: 307-318 [PMID: 22387003 DOI: 10.1016/j.semcancer.2012.02.008]
 - 20 **Obermajer N**, Muthuswamy R, Lesnock J, Edwards RP, Kalinski P. Positive feedback between PGE2 and COX2 redirects the differentiation of human dendritic cells toward stable myeloid-derived suppressor cells. *Blood* 2011; **118**: 5498-5505 [PMID: 21972293 DOI: 10.1182/blood-2011-07-365825]
 - 21 **Nakamoto Y**, Guidotti LG, Kuhlen CV, Fowler P, Chisari FV. Immune pathogenesis of hepatocellular carcinoma. *J Exp Med* 1998; **188**: 341-350 [PMID: 9670046 DOI: 10.1084/jem.188.2.341]
 - 22 **Kakumu S**, Ito S, Ishikawa T, Mita Y, Tagaya T, Fukuzawa Y, Yoshioka K. Decreased function of peripheral blood dendritic cells in patients with hepatocellular carcinoma with hepatitis B and C virus infection. *J Gastroenterol Hepatol* 2000; **15**: 431-436 [PMID: 10824889 DOI: 10.1046/j.1440-1746.2000.02161.x]
 - 23 **Dolganiuc A**, Chang S, Kodys K, Mandrekar P, Bakis G, Cormier M, Szabo G. Hepatitis C virus (HCV) core protein-induced, monocyte-mediated mechanisms of reduced IFN- α and plasmacytoid dendritic cell loss in chronic HCV infection. *J Immunol* 2006; **177**: 6758-6768 [PMID: 17082589 DOI: 10.4049/jimmunol.177.10.6758]
 - 24 **Duramad O**, Fearon KL, Chan JH, Kanzler H, Marshall JD, Coffman RL, Barrat FJ. IL-10 regulates plasmacytoid dendritic cell response to CpG-containing immunostimulatory sequences. *Blood* 2003; **102**: 4487-4492 [PMID: 12946990 DOI: 10.1182/blood-2003-07-2465]
 - 25 **Gonzalez-Carmona MA**, Lukacs-Kornek V, Timmerman A, Shabani S, Kornek M, Vogt A, Yildiz Y, Sievers E, Schmidt-Wolf IG, Caselmann WH, Sauerbruch T, Schmitz V. CD40ligand-expressing dendritic cells induce regression of hepatocellular carcinoma by activating innate and acquired immunity in vivo. *Hepatology* 2008; **48**: 157-168 [PMID: 18537185 DOI: 10.1002/hep.22296]
 - 26 **Shurin MR**, Yurkovetsky ZR, Tourkova IL, Balkir L, Shurin GV. Inhibition of CD40 expression and CD40-mediated dendritic cell function by tumor-derived IL-10. *Int J Cancer* 2002; **101**: 61-68 [PMID: 12209589 DOI: 10.1002/ijc.10576]
 - 27 **Tian Z**, Chen Y, Gao B. Natural killer cells in liver disease. *Hepatology* 2013; **57**: 1654-1662 [PMID: 23111952 DOI: 10.1002/hep.26115]
 - 28 **Lanier LL**. NKG2D Receptor and Its Ligands in Host Defense. *Cancer Immunol Res* 2015; **3**: 575-582 [PMID: 26041808 DOI: 10.1158/2326-6066.CIR-15-0098]
 - 29 **Yoon JC**, Lim JB, Park JH, Lee JM. Cell-to-cell contact with hepatitis C virus-infected cells reduces functional capacity of natural killer cells. *J Virol* 2011; **85**: 12557-12569 [PMID: 21937646 DOI: 10.1128/JVI.00838-11]
 - 30 **Kamiya T**, Chang YH, Campana D. Expanded and Activated Natural Killer Cells for Immunotherapy of Hepatocellular Carcinoma. *Cancer Immunol Res* 2016; **4**: 574-581 [PMID: 27197065 DOI: 10.1158/2326-6066.CIR-15-0229]
 - 31 **Oliviero B**, Varchetta S, Paudice E, Michelone G, Zaramella M, Mavilio D, De Filippi F, Bruno S, Mondelli MU. Natural killer cell functional dichotomy in chronic hepatitis B and chronic hepatitis C virus infections. *Gastroenterology* 2009; **137**: 1151-1160, 1160.e1-1160.e7 [PMID: 19470388 DOI: 10.1053/j.gastro.2009.05.047]
 - 32 **Accapezzato D**, Francavilla V, Paroli M, Casciaro M, Chircu LV, Cividini A, Abrignani S, Mondelli MU, Barnaba V. Hepatic expansion of a virus-specific regulatory CD8(+) T cell population in chronic hepatitis C virus infection. *J Clin Invest* 2004; **113**: 963-972 [PMID: 15057302 DOI: 10.1172/JCI200420515]
 - 33 **Wu Y**, Tian Z, Wei H. Developmental and Functional Control of Natural Killer Cells by Cytokines. *Front Immunol* 2017; **8**: 930 [PMID: 28824650 DOI: 10.3389/fimmu.2017.00930]
 - 34 **Li T**, Yang Y, Hua X, Wang G, Liu W, Jia C, Tai Y, Zhang Q, Chen G. Hepatocellular carcinoma-associated fibroblasts trigger NK cell dysfunction via PGE2 and IDO. *Cancer Lett* 2012; **318**: 154-161 [PMID: 22182446 DOI: 10.1016/j.canlet.2011.12.020]
 - 35 **Xu D**, Han Q, Hou Z, Zhang C, Zhang J. miR-146a negatively regulates NK cell functions via STAT1 signaling. *Cell Mol Immunol* 2017; **14**: 712-720 [PMID: 26996068 DOI: 10.1038/cmi.2015.113]
 - 36 **Zhang H**, Li Z, Wang L, Tian G, Tian J, Yang Z, Cao G, Zhou H, Zhao L, Wu Z, Yin Z. Critical Role of Myeloid-Derived Suppressor Cells in Tumor-Induced Liver Immune Suppression through Inhibition of NKT Cell Function. *Front Immunol* 2017; **8**: 129 [PMID: 28243237 DOI: 10.3389/fimmu.2017.00129]
 - 37 **Jiang T**, Zhou C, Ren S. Role of IL-2 in cancer immunotherapy. *Oncoimmunology* 2016; **5**: e1163462 [PMID: 27471638 DOI: 10.1080/2162402X.2016.1163462]
 - 38 **Zekri AR**, Youssef AS, Bakr YM, Gabr RM, El-Rouby MN, Hammad I, Ahmed EA, Marzouk HA, Nabil MM, Hamed HA, Aly YH, Zachariah KS, Esmat G. Serum biomarkers for early detection of hepatocellular carcinoma associated with HCV infection in Egyptian patients. *Asian Pac J Cancer Prev* 2015; **16**: 1281-1287 [PMID: 25735368 DOI: 10.7314/APJCP.2015.16.3.1281]
 - 39 **Zekri AR**, Alam El-Din HM, Bahnassy AA, Zayed NA, Mohamed WS, El-Masry SH, Gouda SK, Esmat G. Serum levels of soluble Fas, soluble tumor necrosis factor-receptor II, interleukin-2 receptor and interleukin-8 as early predictors of hepatocellular carcinoma in Egyptian patients with hepatitis C virus genotype-4. *Comp Hepatol* 2010; **9**: 1 [PMID: 20051112 DOI: 10.1186/1476-5926-9-1]

P- Reviewer: Liu XL **S- Editor:** Wang JL **L- Editor:** Filipodia
E- Editor: Huang Y





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>



ISSN 1007-9327

