

World Journal of *Gastroenterology*

World J Gastroenterol 2021 January 21; 27(3): 233-304



EDITORIAL

- 233 Screening colonoscopy: The present and the future
Hayman CV, Vyas D

ORIGINAL ARTICLE

Basic Study

- 240 Circular RNA AKT3 governs malignant behaviors of esophageal cancer cells by sponging miR-17-5p
Zang HL, Ji FJ, Ju HY, Tian XF

Retrospective Study

- 255 Serum vitamin D and vitamin-D-binding protein levels in children with chronic hepatitis B
Huang CZ, Zhang J, Zhang L, Yu CH, Mo Y, Mo LY
- 267 Trends in the management of anorectal melanoma: A multi-institutional retrospective study and review of the world literature
Bleicher J, Cohan JN, Huang LC, Peche W, Pickron TB, Scaife CL, Bowles TL, Hyngstrom JR, Asare EA

Observational Study

- 281 Comparative study on artificial intelligence systems for detecting early esophageal squamous cell carcinoma between narrow-band and white-light imaging
Li B, Cai SL, Tan WM, Li JC, Yalikong A, Feng XS, Yu HH, Lu PX, Feng Z, Yao LQ, Zhou PH, Yan B, Zhong YS

CASE REPORT

- 294 Peritoneal dissemination of pancreatic cancer caused by endoscopic ultrasound-guided fine needle aspiration: A case report and literature review
Kojima H, Kitago M, Iwasaki E, Masugi Y, Matsusaka Y, Yagi H, Abe Y, Hasegawa Y, Hori S, Tanaka M, Nakano Y, Takemura Y, Fukuhara S, Ohara Y, Sakamoto M, Okuda S, Kitagawa Y

ABOUT COVER

Michael Linnebacher, PhD, Academic Fellow, Research Fellow, Research Scientist, Senior Researcher, Senior Scientist, Molecular Oncology and Immunotherapy, Department of General, Visceral, Vascular and Transplantation Surgery, University Medical Center Rostock, Schillingallee 69, Rostock 18057, Germany. michael.linnebacher@med.uni-rostock.de

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (WJG, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, and Scopus. The 2020 edition of Journal Citation Report® cites the 2019 impact factor (IF) for WJG as 3.665; IF without journal self cites: 3.534; 5-year IF: 4.048; Ranking: 35 among 88 journals in gastroenterology and hepatology; and Quartile category: Q2. The WJG's CiteScore for 2019 is 7.1 and Scopus CiteScore rank 2019: Gastroenterology is 17/137.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Yun-Jie Ma*; Production Department Director: *Xiang Li*; Editorial Office Director: *Ze-Mao Gong*.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski, Subrata Ghosh

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

January 21, 2021

COPYRIGHT

© 2021 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Trends in the management of anorectal melanoma: A multi-institutional retrospective study and review of the world literature

Josh Bleicher, Jessica N Cohan, Lyen C Huang, William Peche, T Bartley Pickron, Courtney L Scaife, Tawnya L Bowles, John R Hyngstrom, Elliot A Asare

ORCID number: Josh Bleicher 0000-0001-6137-4426; Jessica N Cohan 0000-0002-5461-4716; Lyen C Huang 0000-0002-8605-2631; William Peche 0000-0001-9770-2137; T Bartley Pickron 0000-0002-0880-1959; Courtney L Scaife 0000-0003-2497-4961; Tawnya L Bowles 0000-0002-1883-1510; John R Hyngstrom 0000-0002-4833-3838; Elliot A Asare 0000-0002-0176-3865.

Author contributions: Bleicher J, Bowles TL, Hyngstrom JR and Asare EA designed the study and created the methodology; Bleicher J and Asare EA performed data analysis; Bleicher J created the first draft of the manuscript; Asare EA supervised the research; all authors participated in data acquisition and provided edits and helped with subsequent writing.

Institutional review board statement: The Institutional Review Boards of the University of Utah and Intermountain Health Care approved this study.

Conflict-of-interest statement: I confirm that I have no financial disclosures to declare. As the corresponding author for this manuscript, I also declare that none of my co-authors have financial disclosures to declare.

Josh Bleicher, Jessica N Cohan, Lyen C Huang, William Peche, T Bartley Pickron, Courtney L Scaife, Tawnya L Bowles, John R Hyngstrom, Elliot A Asare, Department of Surgery, Huntsman Cancer Institute at the University of Utah, Salt Lake City, UT 84114, United States

William Peche, John R Hyngstrom, Department of Surgery, George E Wahlen Department of Veterans Affairs Medical Center, Salt Lake City, UT 84114, United States

Tawnya L Bowles, Elliot A Asare, Department of Surgery, Intermountain Medical Center, Murray, UT 84107, United States

Corresponding author: Elliot A Asare, MD, MS, Assistant Professor, Surgical Oncologist, Department of Surgery, Huntsman Cancer Institute at the University of Utah, 1950 Circle of Hope Drive, Suite N6345, Salt Lake City, UT 84114, United States. elliott.asare@hci.utah.edu

Abstract

BACKGROUND

Anorectal melanoma (ARM) is a rare disease with a poor prognosis. Evidence on optimal treatment is limited and surgical management varies widely. We hypothesized that the frequency of abdominoperineal resection used as primary treatment of ARM has decreased over the past several decades.

AIM

To update our understanding of outcomes for patients with ARM and analyze management trends around the world.

METHODS

This is a multi-institutional, retrospective study of patients treated for ARM at 7 hospitals. Hospitals included both large, academic, tertiary care centers and smaller, general community hospitals. Using prospectively maintained institutional tumor registries, we identified 24 patients diagnosed with ARM between January 2000 and May 2019. We analyzed factors prognostic for recurrence and survival. We then used Cox regression to measure overall survival (OS) and melanoma-specific survival. We also performed a literature review to assess trends in surgical management and outcomes.

RESULTS

Of the 24 patients diagnosed with ARM, 12 (50.0%) had local, 8 (33.3%) regional,

Data sharing statement: Because of the small number of participants involved in this cohort and the inability to obtain informed consent, it is difficult to ensure patient anonymity in our dataset and this will not be made available. Statistical code is available from the first author at

josh.bleicher@hsc.utah.edu.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Specialty type: Gastroenterology and hepatology

Country/Territory of origin: United States

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B, B, B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

Received: December 1, 2020

Peer-review started: December 1, 2020

First decision: December 21, 2020

Revised: December 28, 2020

Accepted: January 7, 2021

Article in press: January 7, 2021

Published online: January 21, 2021

P-Reviewer: Gong N, Taher M

S-Editor: Fan JR

L-Editor: A

P-Editor: Li JH

and 4 (16.7%) distant disease at diagnosis. Median time to recurrence was 10.4 mo [interquartile range (IQR) 7.5-17.2] with only 2 patients (9.3%) not developing recurrence following surgical resection. Median OS was 18.8 mo (IQR 13.5-33.9). One patient is still alive without recurrence at 21.4 mo from diagnosis; no other patient survived 5 years. Primary surgical management with abdominoperineal resection (APR) *vs* wide excision (WE) did not lead to differences in OS [hazard ratio = 1.4 (95%CI: 0.3-6.8)]. Review of the literature revealed geographic differences in surgical management of ARM, with increased use of WE in the United States and Europe over time and more frequent use of APR in Asia and India. There was no significant improvement in survival over time.

CONCLUSION

There is wide variation in the management of ARM and survival outcomes remain poor regardless of approach. Surgical management should aim to minimize morbidity.

Key Words: Melanoma; Anorectal melanoma; Literature review; Melanoma surgery; Surgical oncology; Colorectal surgery

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This is a retrospective study to evaluate current trends in management of anorectal melanoma (ARM). On review of 24 patients from 7 hospitals in Utah, we found that ARM is a highly lethal disease with overall survival of 18.8 mo (interquartile range 13.5-33.9) and no 5-year survivors. Only 2 patients underwent abdominoperineal resection (APR) as primary surgical management. Review of the literature demonstrated wide variation in surgical management of ARM over time and around the world. Whether APR or wide excision was used, outcomes remained poor. With this data, we recommend that surgical management should aim to minimize morbidity.

Citation: Bleicher J, Cohan JN, Huang LC, Peche W, Pickron TB, Scaife CL, Bowles TL, Hyngstrom JR, Asare EA. Trends in the management of anorectal melanoma: A multi-institutional retrospective study and review of the world literature. *World J Gastroenterol* 2021; 27(3): 267-280

URL: <https://www.wjgnet.com/1007-9327/full/v27/i3/267.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v27.i3.267>

INTRODUCTION

Anorectal melanoma (ARM) is a rare malignancy with a poor prognosis. The estimated annual incidence in the United States is less than 5 cases per 10 million^[1]. Overall 5-year survival is between 10% and 20%^[2]. This low survival is due to the late diagnosis of most tumors and aggressive biology of ARM^[3]. Most tumors are first recognized from symptoms such as bleeding, obstruction, pain, or changes in bowel habits^[4-6]. When these tumors are recognized, they are often misdiagnosed as hemorrhoids or other benign anorectal pathology^[7].

National Clinical Cancer Network (NCCN) guidelines on melanoma do not currently include recommendations for treatment of ARM^[8]. Without guidelines, and due to the rare nature of the tumor, treatment is highly variable. Controversy exists over optimal primary surgical therapy. Some advocate abdominoperineal resection (APR) for initial treatment, while others report similar oncologic outcomes with wide excision (WE) alone^[9,10]. As outcomes are universally poor, many providers recommend the less invasive and lower morbidity WE as primary treatment^[11]. Optimal primary nodal management strategy is also unknown. Non-surgical therapy is even more varied. Radiotherapy, chemotherapy, and targeted therapies (including interferon, checkpoint-inhibitors, anti-BRAF therapy, and tyrosine kinase inhibitors) have all been used alone or in various combinations^[12-17]. No clear treatment strategy has emerged as the gold standard for treatment of this rare but aggressive disease.



Given the lack of guidelines and variability in reported practice patterns, we analyzed outcomes from a multi-institutional cohort of patients with ARM. We also provide an updated review of the literature to compare outcomes from across the decades and around the world. This review allows for analysis of overall trends to help guide treatment decisions for patients with ARM.

MATERIALS AND METHODS

Study design

We retrospectively reviewed patients diagnosed with ARM between January 1, 2000 and January 1, 2019. This allowed for at least 12 mo of follow-up for all patients. Patients were identified using international classification of diseases-9/10 codes in prospectively maintained institutional tumor registries at 7 centers near Salt Lake City, Utah. These centers included the University of Utah Huntsman Cancer Institute and 6 hospitals affiliated with Intermountain Health Care. All names were linked across institutions to ensure only unique patients were included in the study.

Data Collection

We abstracted data from the electronic medical record and institutional tumor registries. Manual chart review was performed for all records to verify data and obtain additional information. Data abstracted includes patient demographics, primary tumor characteristics, treatment details, and cancer-related outcomes. Both adjuvant therapy and therapy at time of relapse were recorded. Specific chemotherapy and immunotherapy agents were noted. Vital status was available for all but one patient.

Extent of disease was categorized into local, regional, or distant depending on whether disease was confined to the anorectum, involved regional lymph nodes, or other organs^[3]. Extent of disease classification was based on clinical documentation. The extent of primary surgical therapy was also determined by clinical documentation. The Institutional Review Boards of the University of Utah and Intermountain Health Care approved this study.

Statistical analyses

Statistical analyses were performed using Stata Version 15.1 (Stata Corp, College Station, TX, United States). We analyzed patient demographics, initial tumor characteristics, and treatment details using descriptive statistics. We calculated median time to recurrence, melanoma-specific survival (MSS), and overall survival (OS) for the cohort and determined MSS and OS at 2, 3, and 5-year intervals. Patients with unknown survival outcomes were excluded from OS analysis and patients with unknown cause of death were excluded from MSS analysis. We graphically evaluated these outcomes using the Kaplan-Meier method. Time-zero for all time-to-event outcomes was the date of diagnosis. Recurrence was defined as re-appearance of disease on physical exam or radiographically in patients who had been initially rendered free of disease after initial treatment. Patients were determined to be free of disease following initial therapy based on intention to treat, as described in clinical documentation.

Cox regression was used to assess for any factors associated with survival. Results were considered statistically significant if the two-sided $P < 0.05$. Analysis of outcomes associated with different surgical and non-surgical treatment options was performed in a similar fashion. Multivariable analysis was not performed because of the small sample size of this cohort.

Review of the literature

We performed a literature review using the PubMed database. The 2009 PRISMA checklist was used to ensure transparent reporting of search and review methodology^[18]. The search term used was "ARM." Search results did not differ significantly when "anal melanoma" or "rectal melanoma" were considered separately. All English language articles were included. Articles were excluded if they did not describe outcomes of a unique cohort of at least 10 patients. When multiple articles described overlapping patient cohorts, the most recent and inclusive article was used. Studies describing patient outcomes from national databases in the United States were excluded, as these patients are often represented in other institutional studies. National database studies from other countries were included when other cohorts from these countries did not exist. All titles and abstracts were reviewed for

inclusion.

Once this review was complete, all full-length articles were reviewed. Outcomes of interest were surgical management of patients, median OS, and 5-year OS. No summary of outcomes was performed because of the significant heterogeneity among the various studies.

RESULTS

Twenty-four patients met inclusion criteria. Two-thirds of patients were female, with median age of 65.5 [interquartile range (IQR) 54-76] (Table 1). Patients were from five different states (UT, ID, WY, NV, CO) and approximately 20% of patients were from rural communities. Of 13 patients with information on Breslow depth, 7 (53.8%) were > 5 mm. There were 9 (37.5%) patients whose melanoma exhibited ulceration (Table 1). Half of the patients had advanced disease at diagnosis; 8 with nodal disease and 4 with distant metastases.

Fifteen patients (62.5%) underwent WE at diagnosis and 2 patients (8.3%) underwent APR (Table 2). Seven patients (29.2%) received biopsy alone, including 2/4 patients with distant disease at diagnosis. The primary operation took place at a median of 27 d after diagnosis (IQR 0-47). Sentinel lymph node biopsy (SLNB) was performed in 6 (25%) patients. Of those with local or nodal disease, nearly half of patients received surgery alone as primary management. The remainder of patients received systemic therapy of some form following surgical resection. There was wide variation in adjuvant treatment. Some patients received chemotherapy, radiation, interferon, or checkpoint-inhibitor therapy alone; others received these in various combinations (Table 2).

Of 21 patients with complete follow-up data, only 2 (9.5%) remained free of disease after resection. One of these patients died of metastatic colon adenocarcinoma 13.9 mo after ARM diagnosis. The other patient was alive at last follow-up with no evidence of disease 21.4 mo from diagnosis. Excluding the 4 patients with distant disease at diagnosis, 3 (20.0%) of the remaining 15 patients were never free of disease and the remaining 12 (80.0%) recurred after initial treatment. One patient who was never free of disease underwent APR as salvage therapy. Median time to recurrence was 10.4 mo (IQR 7.5-17.2). At the time of recurrence, 4 patients (33.3%) opted not to pursue further therapy given their age, comorbidities, and/or overall prognosis. Of those who received further treatment ($n = 8$), only 1 patient had surgery (repeat WE and bilateral inguinal lymph node dissection). Use of systemic therapy was highly variable. Individual patient treatments and outcomes are shown in Supplementary Table 1.

Survival status was known for all but 1 patient. Of these 23 patients, 1 (4.3%) was alive at last follow-up (21.4 mo from diagnosis). Fourteen (60.9%) died from ARM. The cause of death was unknown for 7 (30.4%) patients. Median OS was 18.8 mo (IQR 13.5-33.9) with 0 survivors at 5 years (Figure 1A). Two-year OS was 21.0% (95%CI: 6.8%-40.3%) and 3-year OS was 10.4% (95%CI: 1.8%-27.9%). Median MSS was 19.5 mo (IQR 14.8-35.1) (Figure 1B). Two-year MSS was 29.1% (95%CI: 9.1%-53.0%) and 3-year MSS was 14.6% (95%CI: 2.4%-37.0%). Excluding patients with distant disease at diagnosis, median OS was 19.9 mo (IQR 16.0-35.1) with 2-year OS of 25.5% (95%CI: 8.2%-47.3%) and 3-year OS of 12.7% (95%CI: 2.2%-33.0%). Median MSS was 19.9 mo (IQR 16.4-39.8) with 2-year MSS of 36.4% (95%CI: 11.2%-62.7%) and 3-year MSS of 18.2% (95%CI: 2.9%-44.2%).

Age, sex, rural location, mitoses, ulceration, and Breslow depth were not prognostic of OS. Patients with distant disease at diagnosis had higher risk of mortality than patients with local disease [hazard ratio (HR) = 14.6 (95%CI: 2.5-86.7)] or nodal disease [HR = 14.4 (95%CI: 2.2-92.1)]. No differences in OS were noted for patients who underwent APR *vs* WE as their primary operation [HR = 1.4 (95%CI: 0.3-6.8)]. There was no significant difference in OS between patients who underwent nodal surgery [SLNB or completion lymph node dissection (CLND)] and those who did not [HR = 0.4 (95%CI: 0.1-1.1)]. Exclusion of patients with distant disease at diagnosis did not alter these results. No individual adjuvant therapy (immunotherapy, radiation, or chemotherapy) demonstrated a benefit over another therapy for patients with local or nodal disease treated with surgery as initial treatment.

Review of the literature

This search revealed 360 unique articles, of which 33 were included for review (Figure 2). Cohorts differed across studies, with some including all patients diagnosed with ARM and others limited to only patients with local or nodal disease or patients

Table 1 Demographics and primary tumor characteristics for cohort with anorectal melanoma (*n* = 24)

Classification		<i>n</i>
Age, years (median, IQR)		65.5 (54-76)
	≤ 50	3 (12.5)
	51-60	7 (29.2)
	61-70	6 (25.0)
	71-80	6 (25.0)
	> 80	2 (8.3)
Sex	M	8 (33.3)
	F	16 (66.7)
Race	White	23 (95.8)
	Latino	1 (4.2)
Rural	Yes	5 (20.8)
	No	19 (79.2)
Breslow depth (mm)	≤ 5	6 (25.0)
	> 5	7 (29.2)
	Unknown	11 (45.8)
Ulceration	Present	9 (37.5)
	Absent	2 (8.3)
	Unknown	13 (54.2)
Mitoses	> 1	7 (29.2)
	Unknown	17 (70.8)
Stage	I	12 (50.0)
	II	8 (33.3)
	III	4 (16.7)

IQR: Interquartile range.

treated with curative intent. Twenty-five studies reported median OS (Table 3). Median OS ranged from 7-49.5 mo and 21 (84%) studies had median OS < 25 mo. There was wide variation in the type of surgical management across studies. At some centers, all patients received WE while other centers treated all patients with APR^[10,19,20].

Eight studies achieved a 5-year OS rate of 20%. Three of these studies included patients diagnosed before 1980, with 1 study including patients from the 1930s^[10,21,22]. The other five study cohorts spanned into the 2000s. Surgical management of patients was mixed in this subset of studies. In a study of 54 patients with ARM treated at MD Anderson Cancer Center (MDACC) from 1989-2008, all patients with local disease underwent WE followed by radiation therapy and a 5-year OS of 30% was reported^[10]. In another study from South Korea, authors described 12 patients who underwent APR and 7 who underwent WE with significantly improved OS with APR compared to WE^[9]. In the remainder of studies, 3 studies reported no significant differences between APR and WE and 3 did not report results of this comparison^[7,21-24]. No other dominant themes in surgical or non-surgical treatment were noted across these studies with superior survival outcomes.

Across all studies, the number of APRs was similar to WEs. In total, 427 patients had APR and 436 underwent WE. Studies from the same institution at different time points showed a trend towards performing fewer APRs with time. In two studies from Memorial Sloan Kettering Cancer Center (MSKCC) looking at cohorts from 1950-1977 and 1984-2003, 73.3% of patients underwent APR in the older cohort compared to 41.3% in the more recent cohort^[25,26]. At MDACC, Ross *et al*^[27] reported APR in 53.8% of patients from 1952-1988 while Kelly *et al*^[10] reported exclusive treatment with WE

Table 2 Treatment details for cohort with anorectal melanoma (*n* = 24)

	Treatment	
Primary operation, <i>n</i>	WE	15 (62.5)
	APR	2 (8.3)
	Biopsy alone	7 (29.2)
Primary nodal operation, <i>n</i>	SLNB	6 (25.0)
	CLND	3 (12.5)
	None	15 (62.5)
Adjuvant therapy, <i>n</i> ¹	Chemotherapy alone	0 (0)
	Radiation alone	1 (4.2)
	Interferon alone	3 (12.5)
	Checkpoint Inhibitor	3 (12.5)
	Combination chemotherapy/radiation	2 (8.3)
	Combination chemotherapy/immunotherapy	3 (12.5)
	Combination radiation/immunotherapy	1 (4.2)
	None	11 (45.8)
Surgery at recurrence ²	APR	1 (9.1)
	WE	1 (9.1)
Non-operative therapies at recurrence ²	Chemotherapy	1 (9.1)
	Radiation	0 (0)
	Interferon	0 (0)
	Checkpoint Inhibitor	2 (18.2)
	Combination chemotherapy/radiation	4 (26.7)
	Combination chemotherapy/immunotherapy	2 (13.3)
	Combination radiation/immunotherapy	2 (13.3)

¹Excluding patients with distant disease at diagnosis (*n* = 4).

²Percentages calculated from total patients with recurrence who underwent treatment (*n* = 8).

APR: Abdominoperineal resection; WE: Wide excision; SLNB: Sentinel lymph node biopsy; CLND: Completion lymph node dissection.

between 1989-2008, as noted previously^[10,27].

Geographic variation in surgical management exists. In United States cohorts, 45.7% (132/289) of surgical patients underwent APR, down to 24.3% over the past 40 years (35/144). European cohorts were similar with 45.1% of patients undergoing APR (123/273). Asian (China, South Korea, Japan, and Taiwan) and Indian cohorts had higher rates of APR with 69.9% (200/286) and 79.7% (51/64) of patients receiving APR as primary surgical therapy respectively. Das *et al*^[19] and Ranjith *et al*^[20] report cohorts from India with 100% of patients undergoing APR^[19,20].

DISCUSSION

Dr. George Pack wrote in 1967, “cures are possible although they do not occur with encouraging frequency^[19]”. This study confirms the dismal prognosis associated with ARM. Only 1 patient from our study cohort was alive at last follow up, and there were no 5-year survivors. Only 6/33 studies reviewed reported a 5-year OS > 20%, and some of these studies included only patients with local disease. Most studies reported median OS of less than 2 years, and many less than 1 year. There is no compelling evidence from this review that a significant improvement in survival has been made for patients with ARM since 1967.

This study also demonstrates the wide variation in surgical treatment for ARM, both

Table 3 Studies included in literature review and select outcomes

Ref.	Date published	Location	Dates included	Sample size	Median age	M/F (n)	APR/WE (n)	Median OS	APR vs WE (% survival to 5 yr)	Overall 5 yr survival (%)
Pack <i>et al</i> ^[40]	1967	Pack Medical Foundation, United States	1930-1965	20	53.5 ¹	5/15	11/3	-	-	5
Abbas <i>et al</i> ^[41]	1980	Roswell Park Memorial Institute, United States	1930-1979	20	61.7	4/16	11/7	18.8 ¹	20.1 mo vs 8.5 mo ³	7
Ward <i>et al</i> ^[42]	1986	St. Mark's Hospital, United Kingdom	1932-1982	21	-	12/9	9/6	8.8 ¹	-	0
Thibault <i>et al</i> ^[21]	1996	Mayo Clinic, United States	1939-1993	50	63 ¹	15/35	26/10	26 ^{1,2}	18 vs 19	22
Wanebo <i>et al</i> ^[25]	1990	Memorial Sloan Kettering Cancer Center, United States	1950-1977	36	60	15/21	22/8	14	16/7 mo vs 21.5 mo ^{3,4}	8
Ross <i>et al</i> ^[27]	1990	MD Anderson Cancer Center, United States	1952-1988	32	-	-	14/12	18.6	19.5 mo vs 18.9 mo ³	3
Dodds <i>et al</i> ^[43]	2019	Melanoma Institute Australia, Australia	1958-2016	43	61	21/22	20/15	24	-	16
Siegal <i>et al</i> ^[44]	1983	Sheba Medical Center, Israel	1960-1981	30	64 ¹	13/17	15/9	10.5 ¹	-	7
Roumen <i>et al</i> ^[45]	1996	Eindhoven Cancer Registry, Netherlands	1960-1995	63	66	27/36	21/18	-	-	6
Nilsson <i>et al</i> ^[46]	2010	Swedish National Cancer Registry, Sweden	1960-1999	251	73	101/150	66/86	11.2	7 vs 15, P = 0.08	
Podnos <i>et al</i> ^[47]	2006	City of Hope National Medical Center, United States	1973-2001	126	69.2 ¹	39/87	-	15	-	19
Slingluff <i>et al</i> ^[48]	1990	Duke University Medical Center, United States	1974-1988	24	64 ¹	7/17	13/8	18	18 vs 12	8
Belli <i>et al</i> ^[49]	2008	National Institute of Cancer, Italy	1975-2006	40	63	19/21	13/18	17	18.5 vs 18.5, P = 0.97	-
Che <i>et al</i> ^[22]	2011	Peking Union Medical College, China	1975-2008	56	-	22/34	36/20	21	24.6 vs 9.9, P = 0.65	20
Pessaux <i>et al</i> ^[50]	2004	Institut Gustave Roussy, France	1977-2002	30	58.1	7/23	9/21	17	16 vs 33	17
Ramakrishnan <i>et al</i> ^[51]	2008	Cancer Institute (WIA), India	1980-2004	63	-	34/29	3/8	9.5 ²	-	5
Yeh <i>et al</i> ^[26]	2006	Memorial Sloan Kettering Cancer Center, United States	1984-2003	46	59	18/28	19/27	39 ⁵	32 vs 35, P = 0.66 ⁵	-
Homsy <i>et al</i> ^[52]	2007	H. Lee Moffitt Cancer Center, United States	1987-2004	12	67	3/9	5/6	-	-	-
Bullard <i>et al</i> ^[53]	2003	University of Minnesota, United States	1988-2002	15	65 ¹	6/9	4/11	18 ⁶	14 mo vs 19 mo ³	-
Kelly <i>et al</i> ^[10]	2010	MD Anderson Cancer Center, United States	1989-2008	54	61	19/35	0/54	29	-	30
Das <i>et al</i> ^[19]	2003	Tata Memorial Hospital, India	1990-2001	72	49 ¹	20/52	24/0	13 ^{2,5}	-	8 ²
Hicks <i>et al</i> ^[28]	2014	John Hopkins Hospital, United States	1991-2012	18	64	10/8	7/11	15.5	11.5 mo vs 13.5 mo, P = 0.75 ³	-
Yen <i>et al</i> ^[54]	2013	Chang Gung Memorial Hospital, China	1993-2011	22	58.4 ¹	8/14	12/8	-	0 vs 28.6, P = 0.06	9
Zhang <i>et al</i> ^[7]	2010	First Affiliated Hospital of Guangxi Medical University, China	1995-2007	54	54	21/33	39/15	25	30 vs 16, P = 0.28	26

Aytac <i>et al</i> ^[55]	2010	Uludag University, Turkey	1997-2004	14	58	8/6	11/3	7.5	-	0
Ishizone <i>et al</i> ^[32]	2008	Shinshu University Hospital, Japan	1997-2006	79	65.8 ¹	34/45	63/14	22	-	29
Belbaraka <i>et al</i> ^[56]	2012	National Institute of Oncology, Morocco	1998-2007	17	58 ¹	12/5	7/3	8	-	-
Choi <i>et al</i> ^[9]	2010	Samsung Medical Center, South Korea	1999-2008	19	61	8/11	12/7	45.9	50 vs 0	32
Miguel <i>et al</i> ^[23]	2015	IPOFG, Portugal	2000-2011	10	70.5	2/8	5/1	9.3	-	20
Ranjith <i>et al</i> ^[20]	2018	Regional Cancer Center Thiruvananthapuram, India	2001-2013	31	56	12/19	9/0	9	-	0
Ren <i>et al</i> ^[24]	2018	Fudan University Shanghai Cancer Center, China	2005-2017	60	61	18/42	38/22	-	-	33.3 ⁵
Nusrath <i>et al</i> ^[57]	2018	Basavatakrakam Indo American Cancer Hospital, India	2010-2015	30	50	15/15	15/5	13	13 vs 36, <i>P</i> = 0.48	-
Sahu <i>et al</i> ^[58]	2017	Tata Memorial Hospital, India	2013-2015	37	54	25/12	-	7	-	-

¹Mean reported instead of median.

²Only cases treated with curative intent included in analysis.

³Median overall survival (OS) [abdominoperineal resection (APR) *vs* wide excision].

⁴APR OS grouped as lymph node negative/lymph node positive.

⁵Melanoma specific survival.

⁶Mean OS of deceased patients only.

APR: Abdominoperineal resection; WE: Wide excision; mo: Month; OS: Overall survival.

within and between medical centers. Geographic variation also exists, with United States and European centers more likely to perform WE and Asian and Indian centers more likely to perform APR. This finding was true in our cohort, with few patients undergoing APR. While low sample size limits analysis, there was no difference in survival outcomes between patients undergoing WE *vs* APR. Multiple other groups have demonstrated similar or better outcomes with WE compared to APR^[10,28]. This same conclusion has been reached using larger cohorts from Surveillance, Epidemiology, and End Results and National Cancer Database (NCDB)^[1,11,12]. A prior systematic review concluded that while APR may reduce local recurrence, there is no improvement in OS or recurrence-free survival compared to WE^[29].

WE allows for avoidance of a colostomy and significantly reduced morbidity compared to APR. A study of 49 patients undergoing WE demonstrated the safety of this procedure; 3 patients had minor infections requiring antibiotics and 1 patient required a second operation for postoperative bleeding. No other complications from surgery occurred^[10]. While most studies of APR for ARM have not reported complication rates, APR for other indications is known to be associated with significant morbidities. Perineal wound complications occur in up to 40% of patients and 50% of patients develop genitourinary and/or sexual dysfunction postoperatively^[30]. No studies currently exist in ARM that compare quality of life between WE and APR^[31]. Some centers continue to routinely perform APR for ARM patients; however, we did not find evidence in this review to support this practice^[9,19,20,32].

Nodal management also differs widely. In our cohort, there were no significant differences in survival outcomes between patients who underwent initial nodal surgery (SLNB or CLND) and those who did not. Nearly 2/3 of patients did not receive any nodal staging or treatment. Older studies hypothesized that the benefit of APR was largely secondary to the mesorectal lymphadenectomy performed with this procedure^[25]. However, Yeh *et al*^[26] found that the presence of lymph node metastases had no prognostic significance on survival in 19 patients who underwent APR at MSKCC^[26]. Many patients in this cohort received local surgery alone and did not receive additional therapy until the time of recurrence. Some of these patients likely had unidentified nodal disease at the time of initial surgery. If patients had undergone SLNB and were found to have positive nodal disease, adjuvant systemic therapy could have been initiated sooner. The impact this may have had on survival is unknown. This review did not find studies with large enough patient numbers to make




Figure 1 Overall mortality and melanoma-specific mortality in cohort of patients with anorectal melanoma. A: Overall mortality; B: Melanoma-specific mortality.

conclusions regarding the benefits of nodal surgery.

Use of immune checkpoint inhibitors and targeted therapies is also controversial and evidence is lacking to help with decision making. While checkpoint inhibitors, tyrosine kinase inhibitors, and BRAF/MEK inhibitors have significantly improved outcomes for cutaneous melanoma over the past decade, their role in treatment of ARM remains unknown^[33-36]. Results of immune checkpoint inhibitor therapy for ARM are limited and show mixed outcomes. Tokuhara *et al*^[13] reported a case of a 67-year-old male with ARM who had no oncologic progression of disease for 17 mo after initiation of anti-programmed death 1 (PD-1) therapy^[13]. Conversely, Faure *et al*^[37] reported a case of a 77 year-old male with ARM who progressed rapidly on anti-PD-1 therapy^[37]. Higher level evidence of the effectiveness of immune checkpoint inhibitor therapy in treating ARM is lacking^[13,38]. While immune checkpoint inhibitor therapy has helped individual patients with ARM, the efficacy of this treatment in most ARMs has been questioned as most ARMs do not exhibit 1-PD-ligand expression and few have tumor-infiltrating lymphocytes^[25]. Evidence for other targeted therapies is similarly poor^[2]. The genomic profiles of ARMs differ from cutaneous melanomas, with very low BRAF expression and few NRAS and KIT mutations^[39]. ARM likely has different drivers of metastases with fewer targetable mutations. Although a rare disease, clinical trials are necessary to determine what therapies are most useful for ARM.

This study is limited by its retrospective nature and small cohort size. ARM is an extremely rare disease and only 24 cases were identified over a 20-year period. Additionally, the lack of a synoptic report for this disease has resulted in many

"Anorectal melanoma"
PubMed database. Last review February 21, 2020




Figure 2 Flow diagram of selection of articles for literature review. ARM: Anorectal melanoma.

missing pertinent variables which would have strengthened this study.

CONCLUSION

ARM is a highly lethal disease. Over the past 50 years, outcomes have remained largely unchanged. Without good evidence to drive treatment decisions, surgical and non-surgical management remains highly variable across the United States and the world. Even within our own cohort, management differed between patients. Review of the literature was also unable to resolve many questions on ARM. There does not appear to be survival benefit of APR over WE. With no clear advantage to APR, surgical management should aim to minimize morbidity. Many other questions on ARM management remain unanswered. Improving the quality of data on ARM is necessary. A consensus meeting of experts aimed at the identification of pertinent variables to collect would be a good first step. Additionally, clinical trials to assess the role of sentinel lymph node biopsy, targeted therapies, radiation therapy, and treatment sequencing are needed.

ARTICLE HIGHLIGHTS

Research background

Anorectal melanoma (ARM) is a rare disease with poor outcomes. 5-year survival remains < 20%.

Research motivation

Optimal surgical management of ARM remains unknown. Abdominoperineal resection (APR) and wide excision (WE) are both used and no gold standard for primary tumor management currently exists. Understanding trends in management and outcomes is critical to determining appropriate surgical management.

Research objectives

We aimed to update our understanding of treatment outcomes for patients with ARM

and analyze trends across countries and time.

Research methods

We performed a retrospective study of patients who were diagnosed with ARM at 7 hospitals in the Salt Lake City, UT region. We analyzed factors prognostic for recurrence and survival. We also performed a review of the literature to assess regional and temporal trends in ARM management.

Research results

We identified 24 patients diagnosed with ARM between 2000-01 and 2019-05. 12 (50.0%) had local, 8 (33.3%) regional, and 4 (16.7%) distant disease at diagnosis. Only 2 patients who had surgical resection of their primary tumor with curative intent failed to recur. Median time to recurrence was 10.4 mo [interquartile range (IQR) 7.5–17.2] and median overall survival was 18.8 mo (IQR 13.5–33.9). No patients survived to 5 years. No survival differences were noted for patients managed with WE *vs* APR. Review of the literature demonstrated regional trends in surgical management of ARM, with WE favored in the United States and Europe and APR used more frequently in Asia.

Research conclusions

ARM remains a highly lethal disease regardless of surgical treatment. Patients who undergo WE and APR have poor outcomes. No convincing evidence exists to favor APR over WE. Despite this, APR continues to be used for primary surgical management, although with decreasing frequency in the United States and Europe in recent years. We feel that surgical management should aim to minimize morbidity. WE should be favored over APR for primary surgical treatment.

Research perspectives

Further research should focus on better risk stratification and the role of targeted therapies, radiation therapy, and treatment sequencing. Improving non-surgical therapies will be critical to improving survival for patients with ARM.

ACKNOWLEDGEMENTS

We thank Emily Z. Keung, MD, MD Anderson Cancer Center, Houston, TX, United States.

REFERENCES

- 1 **Chen H**, Cai Y, Liu Y, He J, Hu Y, Xiao Q, Hu W, Ding K. Incidence, Surgical Treatment, and Prognosis of Anorectal Melanoma From 1973 to 2011: A Population-Based SEER Analysis. *Medicine (Baltimore)* 2016; **95**: e2770 [PMID: 26886623 DOI: 10.1097/MD.0000000000002770]
- 2 **Paolino G**, Didona D, Macri G, Calvieri S, Mercuri SR. Anorectal Melanoma. In: Scott JF, Gerstenblith MR, editors. *Noncutaneous Melanoma* [Internet]. *Brisbane (AU): Codon Publications* 2018; **6**: 83-98 [PMID: 29874013 DOI: 10.15586/codon.noncutaneousemelanoma.2018.ch6]
- 3 **Patrick RJ**, Fenske NA, Messina JL. Primary mucosal melanoma. *J Am Acad Dermatol* 2007; **56**: 828-834 [PMID: 17349716 DOI: 10.1016/j.jaad.2006.06.017]
- 4 **Mihajlovic M**, Vlajkovic S, Jovanovic P, Stefanovic V. Primary mucosal melanomas: a comprehensive review. *Int J Clin Exp Pathol* 2012; **5**: 739-753 [PMID: 23071856]
- 5 **van Schaik PM**, Ernst MF, Meijer HA, Bosscha K. Melanoma of the rectum: a rare entity. *World J Gastroenterol* 2008; **14**: 1633-1635 [PMID: 18330962 DOI: 10.3748/wjg.14.1633]
- 6 **Khan M**, Bucher N, Elhassan A, Barbaryan A, Ali AM, Hussain N, Mirzakhimov AE. Primary anorectal melanoma. *Case Rep Oncol* 2014; **7**: 164-170 [PMID: 24748866 DOI: 10.1159/000360814]
- 7 **Zhang S**, Gao F, Wan D. Effect of misdiagnosis on the prognosis of anorectal malignant melanoma. *J Cancer Res Clin Oncol* 2010; **136**: 1401-1405 [PMID: 20130908 DOI: 10.1007/s00432-010-0793-z]
- 8 **National Comprehensive Cancer Network**. Cutaneous Melanoma: Clinical Practice Guidelines in Oncology. 2019
- 9 **Choi BM**, Kim HR, Yun HR, Choi SH, Cho YB, Kim HC, Yun SH, Lee WY, Chun HK. Treatment outcomes of anorectal melanoma. *J Korean Soc Coloproctol* 2011; **27**: 27-30 [PMID: 21431094 DOI: 10.3393/jksc.2011.27.1.27]
- 10 **Kelly P**, Zagars GK, Cormier JN, Ross MI, Guadagnolo BA. Sphincter-sparing local excision and hypofractionated radiation therapy for anorectal melanoma: a 20-year experience. *Cancer* 2011; **117**: 4747-4755 [PMID: 21446049 DOI: 10.1002/cncr.26088]

- 11 **Ciarrocchi A**, Pietroletti R, Carlei F, Amicucci G. Extensive surgery and lymphadenectomy do not improve survival in primary melanoma of the anorectum: results from analysis of a large database (SEER). *Colorectal Dis* 2017; **19**: 158-164 [PMID: [27317493](#) DOI: [10.1111/codi.13412](#)]
- 12 **Taylor JP**, Stem M, Yu D, Chen SY, Fang SH, Gearhart SL, Safar B, Efron JE. Treatment Strategies and Survival Trends for Anorectal Melanoma: Is it Time for a Change? *World J Surg* 2019; **43**: 1809-1819 [PMID: [30830243](#) DOI: [10.1007/s00268-019-04960-w](#)]
- 13 **Tokuhara K**, Nakatani K, Tanimura H, Yoshioka K, Kiyohara T, Kon M. A first reported case of metastatic anorectal amelanotic melanoma with a marked response to anti-PD-1 antibody nivolumab: A case report. *Int J Surg Case Rep* 2017; **31**: 188-192 [PMID: [28171845](#) DOI: [10.1016/j.ijscr.2017.01.028](#)]
- 14 **Knowles J**, Lynch AC, Warriar SK, Henderson M, Heriot AG. A case series of anal melanoma including the results of treatment with imatinib in selected patients. *Colorectal Dis* 2016; **18**: 877-882 [PMID: [26546509](#) DOI: [10.1111/codi.13209](#)]
- 15 **Shoushtari AN**, Bluth MJ, Goldman DA, Bitas C, Lefkowitz RA, Postow MA, Munhoz RR, Buchar G, Hester RH, Romero JA, Fitzpatrick LJ, Weiser MR, Panageas KS, Wolchok JD, Chapman PB, Carvajal RD. Clinical features and response to systemic therapy in a historical cohort of advanced or unresectable mucosal melanoma. *Melanoma Res* 2017; **27**: 57-64 [PMID: [27792058](#) DOI: [10.1097/CMR.0000000000000306](#)]
- 16 **Kim KB**, Sanguino AM, Hodges C, Papadopoulos NE, Eton O, Camacho LH, Broemeling LD, Johnson MM, Ballo MT, Ross MI, Gershenwald JE, Lee JE, Mansfield PF, Prieto VG, Bedikian AY. Biochemotherapy in patients with metastatic anorectal mucosal melanoma. *Cancer* 2004; **100**: 1478-1483 [PMID: [15042682](#) DOI: [10.1002/cncr.20113](#)]
- 17 **Lian B**, Si L, Cui C, Chi Z, Sheng X, Mao L, Li S, Kong Y, Tang B, Guo J. Phase II randomized trial comparing high-dose IFN- α 2b with temozolomide plus cisplatin as systemic adjuvant therapy for resected mucosal melanoma. *Clin Cancer Res* 2013; **19**: 4488-4498 [PMID: [23833309](#) DOI: [10.1158/1078-0432.CCR-13-0739](#)]
- 18 PRISMA. Systematic Review and Meta-analysis Checklist. 2009
- 19 **Das G**, Gupta S, Shukla PJ, Jagannath P. Anorectal melanoma: a large clinicopathologic study from India. *Int Surg* 2003; **88**: 21-24 [PMID: [12731727](#)]
- 20 **Ranjith S**, Muralee M, Sajeed A, Arun PM, Cherian K, Nair CK, Augustine P, Ahamed I. Anorectal melanoma: experience from a tertiary cancer care centre in South India. *Ann R Coll Surg Engl* 2018; **100**: 185-189 [PMID: [29046101](#) DOI: [10.1308/rcsann.2017.0184](#)]
- 21 **Thibault C**, Sagar P, Nivatvongs S, Ilstrup DM, Wolff BG. Anorectal melanoma--an incurable disease? *Dis Colon Rectum* 1997; **40**: 661-668 [PMID: [9194459](#) DOI: [10.1007/BF02140894](#)]
- 22 **Che X**, Zhao DB, Wu YK, Wang CF, Cai JQ, Shao YF, Zhao P. Anorectal malignant melanomas: retrospective experience with surgical management. *World J Gastroenterol* 2011; **17**: 534-539 [PMID: [21274385](#) DOI: [10.3748/wjg.v17.i4.534](#)]
- 23 **Miguel I**, Freire J, Passos MJ, Moreira A. Anorectal malignant melanoma: retrospective analysis of management and outcome in a single Portuguese Institution. *Med Oncol* 2015; **32**: 445 [PMID: [25502089](#) DOI: [10.1007/s12032-014-0445-2](#)]
- 24 **Ren M**, Lu Y, Lv J, Shen X, Kong J, Dai B, Kong Y. Prognostic factors in primary anorectal melanoma: a clinicopathological study of 60 cases in China. *Hum Pathol* 2018; **79**: 77-85 [PMID: [29763716](#) DOI: [10.1016/j.humpath.2018.05.004](#)]
- 25 **Wanebo HJ**, Woodruff JM, Farr GH, Quan SH. Anorectal melanoma. *Cancer* 1981; **47**: 1891-1900 [PMID: [6164474](#) DOI: [10.1002/1097-0142\(19810401\)47:7<1891::aid-cncr282047030>3.0.co;2-k](#)]
- 26 **Yeh JJ**, Shia J, Hwu WJ, Busam KJ, Paty PB, Guillem JG, Coit DG, Wong WD, Weiser MR. The role of abdominoperineal resection as surgical therapy for anorectal melanoma. *Ann Surg* 2006; **244**: 1012-1017 [PMID: [17122627](#) DOI: [10.1097/01.sla.0000225114.56565.f9](#)]
- 27 **Ross M**, Pezzi C, Pezzi T, Meurer D, Hickey R, Balch C. Patterns of failure in anorectal melanoma. A guide to surgical therapy. *Arch Surg* 1990; **125**: 313-316 [PMID: [2306178](#) DOI: [10.1001/archsurg.1990.01410150035007](#)]
- 28 **Hicks CW**, Pappou EP, Magruder JT, Gazer B, Fang S, Wick EC, Gearhart SL, Ahuja N, Efron JE. Clinicopathologic Presentation and Natural History of Anorectal Melanoma: A Case Series of 18 Patients. *JAMA Surg* 2014; **149**: 608-611 [PMID: [24848283](#) DOI: [10.1001/jamasurg.2013.4643](#)]
- 29 **Matsuda A**, Miyashita M, Matsumoto S, Takahashi G, Matsutani T, Yamada T, Kishi T, Uchida E. Abdominoperineal resection provides better local control but equivalent overall survival to local excision of anorectal malignant melanoma: a systematic review. *Ann Surg* 2015; **261**: 670-677 [PMID: [25119122](#) DOI: [10.1097/SLA.0000000000000862](#)]
- 30 **Peterson CY**, Garcia-Aguilar J. Abdominoperineal resection for rectal cancer. In: Fischer's Mastery of Surgery. 2019: 1841-1852
- 31 **Smith HG**, Glen J, Turnbull N, Peach H, Board R, Payne M, Gore M, Nugent K, Smith MJF. Less is more: A systematic review and meta-analysis of the outcomes of radical vs conservative primary resection in anorectal melanoma. *Eur J Cancer* 2020; **135**: 113-120 [PMID: [32563895](#) DOI: [10.1016/j.ejca.2020.04.041](#)]
- 32 **Ishizone S**, Koide N, Karasawa F, Akita N, Muranaka F, Uhara H, Miyagawa S. Surgical treatment for anorectal malignant melanoma: report of five cases and review of 79 Japanese cases. *Int J Colorectal Dis* 2008; **23**: 1257-1262 [PMID: [18633625](#) DOI: [10.1007/s00384-008-0529-6](#)]
- 33 **Freeman M**, Laks S. Surveillance imaging for metastasis in high-risk melanoma: importance in individualized patient care and survivorship. *Melanoma Manag* 2019; **6**: MMT12 [PMID: [31236204](#)]

- DOI: [10.2217/mmt-2019-0003](https://doi.org/10.2217/mmt-2019-0003)]
- 34 **Joseph RW**, Elassaiss-Schaap J, Kefford R, Hwu WJ, Wolchok JD, Joshua AM, Ribas A, Hodi FS, Hamid O, Robert C, Daud A, Dronca R, Hersey P, Weber JS, Patnaik A, de Alwis DP, Perrone A, Zhang J, Kang SP, Ebbinghaus S, Anderson KM, Gangadhar TC. Baseline Tumor Size Is an Independent Prognostic Factor for Overall Survival in Patients with Melanoma Treated with Pembrolizumab. *Clin Cancer Res* 2018; **24**: 4960-4967 [PMID: [29685882](https://pubmed.ncbi.nlm.nih.gov/29685882/) DOI: [10.1158/1078-0432.CCR-17-2386](https://doi.org/10.1158/1078-0432.CCR-17-2386)]
 - 35 **Long GV**, Flaherty KT, Stroyakovskiy D, Gogas H, Levchenko E, de Braud F, Larkin J, Garbe C, Jouary T, Hauschild A, Chiarion-Sileni V, Lebbe C, Mandalà M, Millward M, Arance A, Bondarenko I, Haanen JBAG, Hansson J, Utikal J, Ferraresi V, Mohr P, Probachai V, Schadendorf D, Nathan P, Robert C, Ribas A, Davies MA, Lane SR, Legos JJ, Mookerjee B, Grob JJ. Dabrafenib plus trametinib vs dabrafenib monotherapy in patients with metastatic BRAF V600E/K-mutant melanoma: long-term survival and safety analysis of a phase 3 study. *Ann Oncol* 2017; **28**: 1631-1639 [PMID: [28475671](https://pubmed.ncbi.nlm.nih.gov/28475671/) DOI: [10.1093/annonc/mdx176](https://doi.org/10.1093/annonc/mdx176)]
 - 36 **Kaufman H**, Amatruda T, Nemunaitis JJ, Chesney JA, Delman KA, Spitler LE, Collichio FA, Ross MI, Zhang Y, Shilkut M, Andtbacka RHI. Tumor size and clinical outcomes in melanoma patients (MEL pts) treated with talimogene laherparepvec (T-VEC). *J Clin Oncol* 2015; **33**: 9074 [DOI: [10.1200/jco.2015.33.15_suppl.9074](https://doi.org/10.1200/jco.2015.33.15_suppl.9074)]
 - 37 **Faure M**, Rochigneux P, Olive D, Taix S, Brenot-Rossi I, Gilibert M. Hyperprogressive Disease in Anorectal Melanoma Treated by PD-1 Inhibitors. *Front Immunol* 2018; **9**: 797 [PMID: [29725330](https://pubmed.ncbi.nlm.nih.gov/29725330/) DOI: [10.3389/fimmu.2018.00797](https://doi.org/10.3389/fimmu.2018.00797)]
 - 38 **Cai YT**, Cao LC, Zhu CF, Zhao F, Tian BX, Guo SY. Multiple synchronous anorectal melanomas with different colors: A case report. *World J Clin Cases* 2019; **7**: 1337-1343 [PMID: [31236398](https://pubmed.ncbi.nlm.nih.gov/31236398/) DOI: [10.12998/wjcc.v7.i11.1337](https://doi.org/10.12998/wjcc.v7.i11.1337)]
 - 39 **Yang HM**, Hsiao SJ, Schaeffer DF, Lai C, Remotti HE, Horst D, Mansukhani MM, Horst BA. Identification of recurrent mutational events in anorectal melanoma. *Mod Pathol* 2017; **30**: 286-296 [PMID: [27739435](https://pubmed.ncbi.nlm.nih.gov/27739435/) DOI: [10.1038/modpathol.2016.179](https://doi.org/10.1038/modpathol.2016.179)]
 - 40 **Pack GT**, Oropeza R. A comparative study of melanoma and epidermoid carcinoma of the anal canal: A review of 20 melanomas and 29 epidermoid carcinomas (1930 to 1965). *Dis Colon Rectum* 1967; **10**: 161-176 [PMID: [6026797](https://pubmed.ncbi.nlm.nih.gov/6026797/) DOI: [10.1007/BF02617173](https://doi.org/10.1007/BF02617173)]
 - 41 **Abbas JS**, Karakousis CP, Holyoke ED. Anorectal melanoma: clinical features, recurrence and patient survival. *Int Surg* 1980; **65**: 423-426 [PMID: [7451062](https://pubmed.ncbi.nlm.nih.gov/7451062/)]
 - 42 **Ward MW**, Romano G, Nicholls RJ. The surgical treatment of anorectal malignant melanoma. *Br J Surg* 1986; **73**: 68-69 [PMID: [3947881](https://pubmed.ncbi.nlm.nih.gov/3947881/) DOI: [10.1002/bjs.1800730127](https://doi.org/10.1002/bjs.1800730127)]
 - 43 **Dodds TJ**, Wilmott JS, Jackett LA, Lo SN, Long GV, Thompson JF, Scolyer RA. Primary anorectal melanoma: clinical, immunohistology and DNA analysis of 43 cases. *Pathology* 2019; **51**: 39-45 [PMID: [30497801](https://pubmed.ncbi.nlm.nih.gov/30497801/) DOI: [10.1016/j.pathol.2018.09.060](https://doi.org/10.1016/j.pathol.2018.09.060)]
 - 44 **Siegal B**, Cohen D, Jacob ET. Surgical treatment of anorectal melanomas. *Am J Surg* 1983; **146**: 336-338 [PMID: [6614323](https://pubmed.ncbi.nlm.nih.gov/6614323/) DOI: [10.1016/0002-9610\(83\)90410-5](https://doi.org/10.1016/0002-9610(83)90410-5)]
 - 45 **Roumen RM**. Anorectal melanoma in The Netherlands: a report of 63 patients. *Eur J Surg Oncol* 1996; **22**: 598-601 [PMID: [9005147](https://pubmed.ncbi.nlm.nih.gov/9005147/) DOI: [10.1016/S0748-7983\(96\)92346-X](https://doi.org/10.1016/S0748-7983(96)92346-X)]
 - 46 **Nilsson PJ**, Ragnarsson-Olding BK. Importance of clear resection margins in anorectal malignant melanoma. *Br J Surg* 2010; **97**: 98-103 [PMID: [20013935](https://pubmed.ncbi.nlm.nih.gov/20013935/) DOI: [10.1002/bjs.6784](https://doi.org/10.1002/bjs.6784)]
 - 47 **Podnos YD**, Tsai NC, Smith D, Ellenhorn JD. Factors affecting survival in patients with anal melanoma. *Am Surg* 2006; **72**: 917-920 [PMID: [17058735](https://pubmed.ncbi.nlm.nih.gov/17058735/)]
 - 48 **Slingluff CL Jr**, Vollmer RT, Seigler HF. Anorectal melanoma: clinical characteristics and results of surgical management in twenty-four patients. *Surgery* 1990; **107**: 1-9 [PMID: [2296748](https://pubmed.ncbi.nlm.nih.gov/2296748/)]
 - 49 **Belli F**, Gallino GF, Lo Vullo S, Mariani L, Poiasina E, Leo E. Melanoma of the anorectal region: the experience of the National Cancer Institute of Milano. *Eur J Surg Oncol* 2009; **35**: 757-762 [PMID: [18602790](https://pubmed.ncbi.nlm.nih.gov/18602790/) DOI: [10.1016/j.ejso.2008.05.001](https://doi.org/10.1016/j.ejso.2008.05.001)]
 - 50 **Pessaix P**, Pocard M, Elias D, Duvillard P, Avril MF, Zimmerman P, Lasser P. Surgical management of primary anorectal melanoma. *Br J Surg* 2004; **91**: 1183-1187 [PMID: [15449271](https://pubmed.ncbi.nlm.nih.gov/15449271/) DOI: [10.1002/bjs.4592](https://doi.org/10.1002/bjs.4592)]
 - 51 **Ramakrishnan AS**, Mahajan V, Kannan R. Optimizing local control in anorectal melanoma. *Indian J Cancer* 2008; **45**: 13-19 [PMID: [18453735](https://pubmed.ncbi.nlm.nih.gov/18453735/) DOI: [10.4103/0019-509x.40641](https://doi.org/10.4103/0019-509x.40641)]
 - 52 **Homs J**, Garrett C. Melanoma of the anal canal: a case series. *Dis Colon Rectum* 2007; **50**: 1004-1010 [PMID: [17468984](https://pubmed.ncbi.nlm.nih.gov/17468984/) DOI: [10.1007/s10350-007-0242-5](https://doi.org/10.1007/s10350-007-0242-5)]
 - 53 **Bullard KM**, Tuttle TM, Rothenberger DA, Madoff RD, Baxter NN, Finne CO, Spencer MP. Surgical therapy for anorectal melanoma. *J Am Coll Surg* 2003; **196**: 206-211 [PMID: [12595048](https://pubmed.ncbi.nlm.nih.gov/12595048/) DOI: [10.1016/S1072-7515\(02\)01538-7](https://doi.org/10.1016/S1072-7515(02)01538-7)]
 - 54 **Yen CI**, Chen HH, Chiang SF, Yeh CY, Chen JS, Hsieh PS, Chiang JM, Tsai WS, Tang R, Changchien CR, Wang JY. Anorectal melanoma: review of 22 consecutive cases. *Hepatogastroenterology* 2013; **60**: 89-93 [PMID: [22829553](https://pubmed.ncbi.nlm.nih.gov/22829553/) DOI: [10.5754/hge12453](https://doi.org/10.5754/hge12453)]
 - 55 **Aytac B**, Adim SB, Yerci O, Yilmazlar T. Anorectal malignant melanomas: experience of Uludag University. *Kaohsiung J Med Sci* 2010; **26**: 658-662 [PMID: [21186014](https://pubmed.ncbi.nlm.nih.gov/21186014/) DOI: [10.1016/S1607-551X\(10\)70100-5](https://doi.org/10.1016/S1607-551X(10)70100-5)]
 - 56 **Belbaraka R**, Elharroudi T, Ismaili N, Fetohi M, Tijami F, Jalil A, Errihani H. Management of anorectal melanoma: report of 17 cases and literature review. *J Gastrointest Cancer* 2012; **43**: 31-35 [PMID: [20886311](https://pubmed.ncbi.nlm.nih.gov/20886311/) DOI: [10.1007/s12029-010-9216-2](https://doi.org/10.1007/s12029-010-9216-2)]

- 57 **Nusrath S**, Thammineedi SR, Patnaik SC, Raju KVVN, Pawar S, Goel V, Chavali RN, Murthy S. Anorectal Malignant Melanoma-Defining the Optimal Surgical Treatment and Prognostic Factors. *Indian J Surg Oncol* 2018; **9**: 519-523 [PMID: [30538382](#) DOI: [10.1007/s13193-018-0791-1](#)]
- 58 **Sahu A**, Ramaswamy A, Singhal N, Doshi V, Mirani J, Desouza A, Banavali S, Saklani A, Ostwal V. Metastatic anorectal melanomas - An exploratory retrospective analysis on the benefits of systemic therapy vs best supportive care in a resource-limited setting from India. *South Asian J Cancer* 2017; **6**: 147-150 [PMID: [29404289](#) DOI: [10.4103/sajc.sajc_276_16](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

