

World Journal of *Gastroenterology*

World J Gastroenterol 2023 June 14; 29(22): 3385-3573



REVIEW

- 3385 Overview of current detection methods and microRNA potential in *Clostridioides difficile* infection screening
Bocchetti M, Ferraro MG, Melisi F, Grisolia P, Scrima M, Cossu AM, Yau TO
- 3400 Epidemiology of small intestinal bacterial overgrowth
Efremova I, Maslennikov R, Poluektova E, Vasilieva E, Zharikov Y, Suslov A, Letyagina Y, Kozlov E, Levshina A, Ivashkin V

ORIGINAL ARTICLE

Basic Study

- 3422 Mechanism of annexin A1/N-formylpeptide receptor regulation of macrophage function to inhibit hepatic stellate cell activation through Wnt/ β -catenin pathway
Fan JH, Luo N, Liu GF, Xu XF, Li SQ, Lv XP
- 3440 Study of the roles of caspase-3 and nuclear factor kappa B in myenteric neurons in a P2X7 receptor knockout mouse model of ulcerative colitis
Magalhães HIR, Machado FA, Souza RF, Caetano MAF, Figliuolo VR, Coutinho-Silva R, Castelucci P
- 3469 Comparative analysis of the gut microbiota of wild adult rats from nine district areas in Hainan, China
Niu LN, Zhang GN, Xuan DD, Lin C, Lu Z, Cao PP, Chen SW, Zhang Y, Cui XJ, Hu SK

Case Control Study

- 3482 Plasma exosomal hsa_circ_0079439 as a novel biomarker for early detection of gastric cancer
Li X, Lin YL, Shao JK, Wu XJ, Li X, Yao H, Shi FL, Li LS, Zhang WG, Chang ZY, Chai NL, Wang YL, Linghu EQ

Retrospective Cohort Study

- 3497 Per-oral endoscopic myotomy is safe and effective for pediatric patients with achalasia: A long-term follow-up study
Bi YW, Lei X, Ru N, Li LS, Wang NJ, Zhang B, Yao Y, Linghu EQ, Chai NL

Retrospective Study

- 3508 Efficacy and safety of modified tetracycline dosing in a quadruple therapy for *Helicobacter pylori*: A retrospective single center study
Sun YC, Zhu MJ, Chen XQ, Yue L, Zhao YR, Wang XJ, Kim JJ, Du Q, Hu WL

Observational Study

- 3519 Pre-transjugular-intrahepatic-portosystemic-shunt measurement of hepatic venous pressure gradient and its clinical application: A comparison study
Wang XX, Yin XC, Gu LH, Guo HW, Cheng Y, Liu Y, Xiao JQ, Wang Y, Zhang W, Zou XP, Wang L, Zhang M, Zhu-Ge YZ, Zhang F

- 3534** Difference and clinical value of metabolites in plasma and feces of patients with alcohol-related liver cirrhosis

Xu YF, Hao YX, Ma L, Zhang MH, Niu XX, Li Y, Zhang YY, Liu TT, Han M, Yuan XX, Wan G, Xing HC

Prospective Study

- 3548** Prospective study comparing hepatic steatosis assessment by magnetic resonance imaging and four ultrasound methods in 105 successive patients

Collin R, Magnin B, Gaillard C, Nicolas C, Abergel A, Buchard B

SCIENTOMETRICS

- 3561** Research trends on artificial intelligence and endoscopy in digestive diseases: A bibliometric analysis from 1990 to 2022

Du RC, Ouyang YB, Hu Y

ABOUT COVER

Editorial Board Member of *World Journal of Gastroenterology*, Wolfgang Kreisel, MD, Emeritus Professor, Department of Medicine II, Gastroenterology, Hepatology, Endocrinology, and Infectious Diseases, Faculty of Medicine, Medical Center-University of Freiburg, Hugstetter Str. 55, Freiburg 79106, Germany. Wolfgang.Kreisel@uniklinik-freiburg.de

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (WJG, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports, Index Medicus, MEDLINE, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJG as 5.374; IF without journal self cites: 5.187; 5-year IF: 5.715; Journal Citation Indicator: 0.84; Ranking: 31 among 93 journals in gastroenterology and hepatology; and Quartile category: Q2. The WJG's CiteScore for 2021 is 8.1 and Scopus CiteScore rank 2021: Gastroenterology is 18/149.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Hua-Ge Yin; Production Department Director: Xu Guo; Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

June 14, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Basic Study

Mechanism of annexin A1/N-formylpeptide receptor regulation of macrophage function to inhibit hepatic stellate cell activation through Wnt/ β -catenin pathway

Jun-Hua Fan, Na Luo, Geng-Feng Liu, Xiao-Fang Xu, Shi-Quan Li, Xiao-Ping Lv

Specialty type: Gastroenterology and hepatology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Dauey K, Kazakhstan; Mohamed GA, Egypt; Nakade Y, Japan

Received: February 15, 2023

Peer-review started: February 15, 2023

First decision: March 28, 2023

Revised: April 10, 2023

Accepted: May 11, 2023

Article in press: May 11, 2023

Published online: June 14, 2023



Jun-Hua Fan, Na Luo, Geng-Feng Liu, Xiao-Fang Xu, Shi-Quan Li, Xiao-Ping Lv, Department of Gastroenterology, The First Affiliated Hospital of Guangxi Medical University, Nanning 530021, Guangxi Zhuang Autonomous Region, China

Corresponding author: Jun-Hua Fan, Doctor, MD, PhD, Associate Professor, Doctor, Department of Gastroenterology, The First Affiliated Hospital of Guangxi Medical University, No. 6 Shuangyong Road, Nanning 530021, Guangxi Zhuang Autonomous Region, China. fanjunhuaxiaoshe@hotmail.com

Abstract

BACKGROUND

Hepatic fibrosis is a common pathological process of chronic liver diseases with various causes, which can progress to cirrhosis.

AIM

To evaluate the effect and mechanism of action annexin (Anx)A1 in liver fibrosis and how this could be targeted therapeutically.

METHODS

CCl₄ (20%) and active N-terminal peptide of AnxA1 (Ac2-26) and N-formylpeptide receptor antagonist N-Boc-Phe-Leu-Phe-Leu-Phe (Boc2) were injected intraperitoneally to induce liver fibrosis in eight wild-type mice/Anxa1 knockout mice, and to detect expression of inflammatory factors, collagen deposition, and the role of the Wnt/ β -catenin pathway in hepatic fibrosis.

RESULTS

Compared with the control group, AnxA1, transforming growth factor (TGF)- β 1, interleukin (IL)-1 β and IL-6 expression in the liver of mice with hepatic fibrosis induced by CCl₄ was significantly increased, which promoted collagen deposition and expression of α -smooth muscle actin (α -SMA), collagen type I and connective tissue growth factor (CTGF), and increased progressively with time. CCl₄ induced an increase in TGF- β 1, IL-1 β and IL-6 in liver tissue of AnxA1 knockout mice, and the degree of liver inflammation and fibrosis and expression of α -SMA, collagen I and CTGF were significantly increased compared with in wild-type mice. After treatment with Ac2-26, expression of liver inflammatory factors, degree of collagen deposition and expression of α -SMA, collagen I and CTGF were decreased.

sed compared with before treatment. Boc2 inhibited the anti-inflammatory and antifibrotic effects of Ac2-26. AnxA1 downregulated expression of the Wnt/ β -catenin pathway in CCl₄-induced hepatic fibrosis. *In vitro*, lipopolysaccharide (LPS) induced hepatocyte and hepatic stellate cell (HSC) expression of AnxA1. Ac2-26 inhibited LPS-induced RAW264.7 cell activation and HSC proliferation, decreased expression of α -SMA, collagen I and CTGF in HSCs, and inhibited expression of the Wnt/ β -catenin pathway after HSC activation. These therapeutic effects were inhibited by Boc2.

CONCLUSION

AnxA1 inhibited liver fibrosis in mice, and its mechanism may be related to inhibition of HSC Wnt/ β -catenin pathway activation by targeting formylpeptide receptors to regulate macrophage function.

Key Words: Annexin A1; Active N-terminal peptide of annexin A1; Macrophage; Hepatic stellate cell; Wnt/ β -catenin; Liver fibrosis

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Hepatic stellate cells (HSCs) are the main receiver of inflammatory signals and the main promoter of fibrosis. Kupffer cells are activated by damage-associated molecular patterns or lipopolysaccharide, and other pathogen-associated molecular patterns, producing and releasing cytokines to activate HSCs, causing secretion and deposition of extracellular matrix and liver fibrosis. Annexin (Anx)A1 is a glucocorticoid regulatory protein that enhances efferocytosis in macrophages, inhibits activation of phospholipase A2, downregulates expression of inflammatory arachidonic acid, and negatively regulates inflammatory cells and factors. Whether AnxA1 is involved in the formation of liver fibrosis and its mechanism of action were the focus of this study.

Citation: Fan JH, Luo N, Liu GF, Xu XF, Li SQ, Lv XP. Mechanism of annexin A1/N-formylpeptide receptor regulation of macrophage function to inhibit hepatic stellate cell activation through Wnt/ β -catenin pathway. *World J Gastroenterol* 2023; 29(22): 3422-3439

URL: <https://www.wjgnet.com/1007-9327/full/v29/i22/3422.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v29.i22.3422>

INTRODUCTION

Various chronic liver diseases cause inflammation and necrosis of liver cells, and abnormal hyperplasia of fibrous connective tissue, leading to fibrosis, which is the early reversible stage of cirrhosis and one of the most important threats to human health worldwide[1,2]. When liver parenchyma is damaged, extracellular matrix (ECM) synthesis increases due to liver repair. When the cause is not removed, the continuously damaged liver leads to the failure of timely degradation of a large amount of synthetic ECM and excessive deposition in the liver tissue, which eventually develops into fibrosis[1-4]. The process involves a variety of fibrosis-promoting cells, including vascular smooth muscle cells, portal fibroblasts, and hepatic stellate cells (HSCs), and the activation and proliferation of HSCs is the main cell source of ECM[5-10]. Liver injury leads to necrosis or apoptosis of liver cells, and immune cells, mainly Kupffer cells are activated. Transforming growth factor (TGF)- β 1, interleukin (IL)-1 β , IL-6 and other cytokines are released, which promote the activation and proliferation of HSCs and secretion of inflammatory and fibrogenic factors. The activated HSCs express α -smooth muscle actin (α -SMA) and cytokines induce HSCs to differentiate into myofibroblasts, which secrete ECM components and cause liver fibrosis[5-11]. These changes may become an important target for the treatment of liver fibrosis.

Annexin (Anx)A1 is a member of the structurally related calcium-dependent phospholipid binding protein superfamily, which is widely distributed in all human tissues and cells, and involved in cell proliferation, differentiation, signal transduction, apoptosis, immunity and other physiological processes[12]. The proteins of the AnxA1 superfamily are homologous in structure, and each member has its unique N-terminal structure, which is related to its different functions and physiological effects [12]. After hydrolysis of full-length AnxA1, the N terminus is disconnected, and 26 amino acids in the N segment form a peptide chain fragment N-terminal peptide of AnxA1 (Ac2-26)[13]. AnxA1 and Ac2-26 both activate and bind to a specific G-protein-coupled receptor known as N-formylpeptide receptor (FPR)2[13], and Ac2-26 binds to FPR1[14]. AnxA1 inhibits expression of α -SMA and collagen I genes in TGF- β 1-induced renal cortical fibroblasts[15] and regulates cigarette-smoke-induced pulmonary fibrosis

[16]. However, whether AnxA1 regulates cell proliferation and collagen deposition in liver tissue remains unclear. The purpose of this study was to investigate the effects of AnxA1 on CCl₄-induced inflammation, proliferation and collagen deposition in liver injury.

The Wnt/ β -catenin signaling pathway is highly evolutionarily conserved and widespread in multicellular organisms. It is essential for cell growth, differentiation, ontogeny, genetic stability, apoptosis, self-renewal of stem cells, and homeostatic maintenance of adult tissues[17,18]. In normal liver tissue, the Wnt/ β -catenin signaling pathway is almost inactive, but its abnormal activation plays an important role in the physiological and pathological processes of the liver. Increasing evidence has shown that abnormal expression of the Wnt/ β -catenin signaling pathway is involved in the incidence of liver tumors, and the occurrence of liver inflammation, cirrhosis, and focal nodular hyperplasia[19]. AnxA1 has been reported as an important regulator of Wnt/ β -catenin signaling[20], but whether AnxA1 contributes to the formation of liver fibrosis was the focus of this study.

MATERIALS AND METHODS

Animal treatment

Six-to-eight-week-old C57BL/6 mice (weighing 18-25 g; SLC, Guangxi, China) and C57BL/6JGpt Anxa1-KO mice (B6/JGptAnxa1em4Cd9634/Gpt, Strain NO. T016864; weighing 18-25 g; SLC, Jiangsu, China) were maintained in climate-controlled rooms under a 12-h light-dark cycle. All experiments were conducted in accordance with the institutional guidelines of Guangxi Medical University for the care and use of laboratory animals.

Thirty-two wild-type mice and 32 *Anxa1*^{-/-} mice were each randomized into four subgroups of eight: Control, CCl₄-induced liver fibrosis, liver fibrosis + Ac2-26, and liver fibrosis + Ac2-26 + FPR antagonist N-Boc-Phe-Leu-Phe-Leu-Phe (Boc-2). Mice in the control group received olive oil for intraperitoneal injection, while mice in the other three subgroups received 20% CCl₄ (500 mL; Shanghai Acme Biochemical Co., Ltd, China), and CCl₄ + Ac2-26 1 mg/kg (500 mg; MedChemExpress, CAT: HY-P1098A/CS-0093919) with or without N-Boc2 1 mg/kg (500 mL; MedChemExpress, CAT: HY-103473A) for intraperitoneal injection twice weekly. Eight mice were used for each time point. At 4 and 8 wk after injection, animals were anesthetized with pentobarbital sodium (50 mg/kg intravenous injection), and blood samples were collected from the inferior vena cava. The livers were removed, and a small portion of liver tissue was fixed with 4% formaldehyde solution, and the remaining liver tissue was immediately frozen and stored in liquid nitrogen for RNA and protein extraction.

Histopathological examination of liver

Hematoxylin-eosin (HE) staining and modified Masson three-color staining (Modified Masson's Trichrome Staining Kit; Beijing Solarbio Science & Technology,) were performed after liver tissue fixation, dehydration, paraffin embedding, sectioning, dewaxing and hydration (bar: 50 mm).

In situ expression of AnxA1 in liver tissue determined by immunohistochemistry

The sections were dewaxed and hydrated according to AnxA1 monoclonal antibody instructions and SP-9000 Histostain™ - SP Kits (6 mL, Beijing Zhongshan Jinqiao Biotechnology, cat: K197712D/K196915F) requirements, antigenic alkaline repair, endogenous peroxidase blocking, goat serum blocking, and AnxA1 monoclonal antibody dilution 1: 1000 (Abcam, United Kingdom, cat: ab214486) were incubated overnight at 4 °C, the biotin-labeled goat anti-mouse/rabbit immunoglobulin G (IgG) (Invitrogen, United States, cat: TJ26211) was incubated for 15 min, and the horseradish-labeled Streptomyces vitellin was incubated for 15 min. Diaminobenzidine staining, counterstaining, dehydration, transparency, sealing, and microscopic examination. For each sample, five high-power fields (bar: 100 mm) were randomly selected, each containing an average of 400 cells, and the number of positive cells was counted for each field.

Preparation of HSCs, hepatocytes and Kupffer cells

JS1 HSCs and AML1 hepatocytes were separately cultivated in Williams' E medium containing 10% calf serum, 2 mmol/L L-glutamate and antibiotics (100 U/mL penicillin G and 100 μ g/mL streptomycin). RAW246.7 cells were cultured in RPMI 1640 medium containing 10% calf serum and antibiotics. Cells were plated at (2 \times 10⁶)-(3 \times 10⁶)/mL and preincubated for 24 h at 37 °C in a humidified atmosphere containing 5% CO₂. The purity and viability of cells ranged from 87% to 95%. RAW246.7 cells HSCs and hepatocytes were incubated with 100 ng/mL lipopolysaccharide (LPS) (Sigma, St. Louis, MO, United States), LPS + 100 ng/mL Ac2-26, or LPS + Ac2-26 + 100 ng/mL Boc2 for 1, 3, 6, 12 and 24 h to activate expression of AnxA1, α -SMA, TGF- β 1, connective tissue growth factor (CTGF), collagen I, IL-1 β and IL-6. Ac2-26 or Ac2-26 + Boc2 dissolved in medium was added to the incubation medium 1 h prior to the addition of LPS.

Quantitative real-time polymerase chain reaction

Total RNA (1 µg) was extracted from liver or primary cells using TRIzol reagent (Takara Bio, Beijing, China) and reverse transcribed using a ReverTra Ace qPCR RT kit (Takara Bio). Target mRNA expression was quantified using quantitative real-time polymerase chain reaction (qPCR) as described previously[21]. TaqMan Gene Expression Assays for AnxA1 (forward primer 5'-AGCAGATCAAGGC-CGCGTA-3', reverse primer 5'-CATGGCACCACGGAGTTCA-3'), IL-6 (forward primer 5'-AGGATAC-CACTCCCAACAGACC-3', reverse primer 5'-GCACAACCTCTTTCTCATTTCAC-3'), TGF-β1 (forward primer 5'-TACGGCAGTGGCTGAACCA-3', reverse primer 5'-CGGTTTCATGTCATG-GATGGTG-3'), IL-1β (forward primer 5'-CACTACAGGCTCCGAGATGAACAAC-3', reverse primer 5'-TGTCGTTGCTTGGTTCTCCTTGTA-3'), α-SMA (forward primer 5'-CTGACAGAGGCACCACTGAA-3', reverse primer 5'-CATCTCCAGAGTCCAGCACA-3'), collagen I (forward primer 5'-AAGAGGCGA-GAGGTTTCC-3', reverse primer 5'-AGAACCATCAGCACCTTTGG-3'), CTGF (forward primer 5'-CAAAGCAGCAAATACCA-3', reverse primer 5'-GGCCAAATGTGTCTTCCAGT-3'), and GAPDH (forward primer 5'-TGTGTCCGTCGTGGATCTGA-3', reverse primer 5'-TTGCTGTTGAAGTCG-CAGGAG-3') were purchased from Takara Bio.

Western blotting

Protein samples (30-50 µg) were prepared from livers and cells and separated by SDS-PAGE, followed by transfer to polyvinylidene difluoride membranes as reported previously[21]. Blots were incubated with antibodies (1:1000) against AnxA1 (Abcam, cat: ab214486), α-SMA (50 µL, Affinity Biosciences, Jiangsu, cat: AF1032), β-catenin (50 mL, MedChemExpress, cat: 51067-2-AP) and GAPDH (Abcam, cat: ab181602), followed by incubation with a peroxidase-conjugated goat anti-mouse/rabbit IgG(H + L) (1:5000; Invitrogen, cat: TJ26211). Proteins were visualized using fluorescence scanner imaging system (Odyssey; Bio-Rad, Hercules, CA, United States).

Statistical analysis

Results are expressed as mean ± SE ($n = 8$), unless otherwise indicated. Statistical analyses were performed using Kruskal-Wallis one-way analysis of variance. $P < 0.05$ was considered significant.



RESULTS

Effects of AnxA1 on liver appearance

The liver of the control mice was red with bright color and smooth surface. In the CCl₄-induced wild-type mice group, the liver tissue became light in color, rough on the surface, dull around the edge, and hard in texture, and nodules or even cellulose-like changes appeared over time. The lesions in the *AnxA1*^{-/-} group were more severe than in the wild-type mice. After treatment with Ac2-26, the appearance of the liver in the CCl₄-induced mice was significantly improved with dark red color, and the surface roughness and nodulation were significantly reduced, the effects of Ac2-26 were inhibited by Boc2 (Figure 1A). Compared with the control group, AnxA1 expression was significantly increased in the CCl₄ model group, and was higher at 8 wk than 4 wk, suggesting that AnxA1 was related to the development of liver fibrosis. AnxA1 expression was higher in liver tissues of the CCl₄ model group after Ac2-26 treatment, while Boc2 inhibited AnxA1 expression (Figures 1B and C), and the difference was significant ($P < 0.05$). Western blotting and qPCR results were consistent with immunohistochemical results (Figure 1D).

AnxA1 inhibits CCl₄-induced liver injury and inflammatory cell infiltration in mice

HE staining showed that the structure of hepatic lobules in the control group was clear, and there was no vacuole formation or obvious inflammatory cell infiltration. In the CCl₄-induced wild model group, HE staining for 4 wk showed that the hepatocytes were swollen and enlarged, with scattered degeneration and necrosis, as well as infiltration of monocytes and lymphocyte cells. Degeneration and necrosis of hepatocytes were observed at 8 wk. In the *AnxA1*^{-/-} model group with fibrosis induced by CCl₄, liver tissue destruction was accompanied by a large number of inflammatory cell infiltration and steatosis. At 4 wk, there was massive necrosis of liver cells, massive infiltration of inflammatory cells, and disorder of hepatic lobular structure. At 8 wk, there was obvious structural destruction that became more severe over time. The infiltration of liver inflammatory cells and destruction of liver lobular structure in the *AnxA1*^{-/-} mice were more obvious than those in the wild-type mice. After treatment with Ac2-26, the damage to the hepatic lobular structure and inflammatory cell infiltration in the CCl₄-induced mice was reduced compared with that before treatment. AnxA1 mimics the peptide Ac2-26, which largely controls inflammation and is protective against hepatic fibrosis. In the model group treated with Ac2-26 and Boc2, Boc2 almost completely eliminated the effects of Ac2-26 at 4 wk (Figure 2A), suggesting that AnxA1 inhibited CCl₄-induced liver tissue injury and reduced inflammatory cell infiltration. AnxA1 works by targeting mainly FPR receptors.







Figure 1 Annexin A1 inhibited CCl₄-induced liver injury. A: Gross view of liver tissue. The liver of mice in the control group was red with bright color and smooth surface. In the CCl₄-induced model group, the liver was light in color, rough and grainy on the surface. Liver tissue nodules appeared at 8 wk, and fibrosis appeared at 8 wk in *AnxA1*^{-/-} mice. After Ac2-26 treatment, the liver was mostly dark red, and the surface roughness was reduced. The effect of Ac2-26 was reversed by Boc2, and the liver color became lighter, the surface was grainy, the edge was blunt, and the texture was hard; B: Annexin (Anx)A1 was expressed in liver tissue. AnxA1 was stained brown. Compared with the control group, the liver tissue in mice treated with CCl₄ showed obvious AnxA1 staining. More liver tissue was stained at 8 wk than at 4 wk. The *AnxA1*^{-/-} model groups were hardly stained. More tissue staining was observed in the CCl₄-induced model group after Ac2-26 treatment, while staining was almost absent after Boc2 intervention; C: *AnxA1* mRNA expression; D: Protein expression of AnxA1, α-smooth muscle actin, β-catenin, and GAPDH (*n* = 8). Results are presented as ratios of target mRNA or protein normalized to internal GAPDH. ^a*P* < 0.05 vs control; ^b*P* < 0.01 vs CCl₄; ^c*P* < 0.05 vs CCl₄ + Ac2-26; ^d*P* < 0.05 vs intra-group. Ac2-26: Active N-terminal peptide of annexin A1; AnxA1: Annexin A1; Boc2: N-formyl peptide receptor antagonist N-Boc-Phe-Leu-Phe-Leu-Phe; α-SMA: α-smooth muscle actin.


AnxA1 inhibits CCl₄-induced collagen deposition in mouse liver tissue


Collagen deposition is an important indicator of the degree of liver fibrosis, and can be visualized by Masson staining. Staining clearly showed that there was no collagen staining in the control group of wild-type and *AnxA1*^{-/-} mice. In the CCl₄-induced wild model group, blue collagen fibers proliferated in the portal area at 4 wk, but the degree of collagen fibrosis was mild. Similarly, the proliferation of blue-stained collagen fibers in the portal area was observed in the wild model group at 8 wk, but the collagen fibers were further increased, widened and extended to the surrounding hepatic lobular space than at 4 wk, suggesting that the liver fibers were gradually aggravated with the extension of modeling time. Compared with wild model mice, CCl₄ induced more significant collagen deposition in the liver tissue of *AnxA1*^{-/-} mice. At 4 wk, blue collagen fibers proliferated obviously in hepatic portal area and began to extend to liver parenchyma, and collagen fibers began to segment liver parenchyma. At 8 wk, a large amount of collagen surrounded the liver cells to form pseudolobules, which seriously damaged the normal structure of the liver tissue, indicating formation of fibrosis. The collagen deposition in the liver of *AnxA1*^{-/-} mice was significantly aggravated by CCl₄, and the degree of liver fibrosis was even more severe. The collagen deposition and wrapping in the liver of mice in the CCl₄-induced mice was significantly weakened after Ac2-26 treatment, and the severity of liver fibrosis was relieved. In the model group, the damage to the liver tissue structure and collagen deposition after treatment with Ac2-26 and Boc2 were similar to those before treatment (Figure 3A).

A



B





DOI: 10.3748/wjg.v29.i22.3422 Copyright ©The Author(s) 2023.

Figure 2 Annexin A1 inhibits CCl₄-induced hepatic inflammatory cytokines release. A: Annexin A1 (AnxA1) inhibited CCl₄-induced liver injury and inflammatory cell infiltration in mice. Normal mouse liver lobule structure was clear, with no obvious vacuole formation or inflammatory cell infiltration. In the wild-type model group, hepatic lobules were severely damaged; a large number of hepatic nuclei were shrunken, hepatocytes were vacuolated, and inflammatory cells infiltrated. *Anxa1*^{-/-} mice had more severe vacuolation and extensive inflammatory cell infiltration than wild-type mice had. After treatment with active N-terminal peptide of AnxA1 (Ac2-26), the damage to hepatic lobules was reduced, hepatocyte vacuolation was reduced but necrosis was observed, clusters of regenerated hepatocytes were observed in some tissues, and inflammatory cell infiltration was also reduced. N-formylpeptide receptor antagonist N-Boc-Phe-Leu-Phe-Leu-Phe almost reversed the therapeutic effect of Ac2-26, but the inflammatory infiltration and damage to hepatic lobule structure were not significantly improved. Initial magnification: 200 ×; B: Effect of Ac2-26 on interleukin (IL)-1β mRNA expression; C: Effect of Ac2-26 on IL-6 mRNA expression; total RNA was extracted using TRIzol reagent and relative concentration of mRNA was quantified using quantitative real-time polymerase chain reaction (*n* = 8). Results are presented as ratios of target mRNA or protein normalized to internal GAPDH. ^a*P* < 0.05 and ^b*P* < 0.01 vs control; ^c*P* < 0.05 vs CCl₄; ^d*P* < 0.05 and ^e*P* < 0.01 vs CCl₄ + Ac2-26; ^f*P* < 0.05 and ^g*P* < 0.01 vs *Anxa1*^{-/-} + control; ^h*P* < 0.05 vs *Anxa1*^{-/-} + CCl₄; ⁱ*P* < 0.05 and ^j*P* < 0.01 vs *Anxa1*^{-/-} + CCl₄ + Ac2-26; ^k*P* < 0.05 vs intra-group; ^l*P* < 0.05 vs inter-group. Ac2-26: Active N-terminal peptide of AnxA1; AnxA1: Annexin A1; Boc2: N-formyl peptide receptor antagonist N-Boc-Phe-Leu-Phe-Leu-Phe; IL: Interleukin.

AnxA1 inhibits CCl₄-induced HSC activation in liver tissue

α-SMA is a marker of HSC activation and transformation into fibroblasts. HSC activation generates and secretes ECM, and type I collagen is the ECM marker of HSCs. In this study, qPCR and western blotting were used to detect the expression levels of α-SMA and collagen I in liver tissue, and to evaluate the activation status of HSCs. Expression of α-SMA and type I collagen in the liver of wild-type mice was significantly increased by CCl₄, while expression of α-SMA and collagen I in the liver of mice in the *Anxa1*^{-/-} group was higher and increased with time, which was in accordance with the progression of liver fibrosis in the CCl₄-induced mice. All of the above effects were inhibited by Ac2-26, suggesting that AnxA1 inhibited HSC activation and collagen deposition. In contrast, Boc2 reversed the therapeutic effect of Ac2-26 (Figures 3B and C), and the difference was significant (*P* < 0.05). Western blotting and qPCR results were consistent with immunohistochemical results (Figure 1D).


AnxA1 inhibits CCl₄-induced release of hepatic inflammatory cytokines

CCl₄ induces expression of various cytokines in liver tissues, among which IL-1β, IL-6 and TGF-β1 are important factors leading to the formation of fibrosis. Expression of IL-1β, IL-6 and TGF-β1 in the liver of the CCl₄-induced wild model group was increased simultaneously, and upregulation of inflammatory cytokines was more obvious and increased with time in the *Anxa1*^{-/-} model group, while the Ac2-26 significantly inhibited expression of IL-1β, IL-6 and TGF-β1. Boc2 inhibited the anti-inflammatory effect of Ac2-26 (Figures 2B, 2C and 3E), and the difference was significant (*P* < 0.05). This suggested that AnxA1 exerted its anti-inflammatory and antifibrotic effects by inhibiting expression of TGF-β1, IL-1β and IL-6 through FPR.


AnxA1 reduces the level of CTGF in the liver


CTGF mediates TGF-β1 promotion of ECM production and accumulation. CTGF regulates the growth of fibroblasts and epidermal cells, promoting cell adhesion, proliferation and ECM synthesis. Compared with the control group, the concentration of CTGF in the liver tissue of the CCl₄-induced wild model

A



B





DOI: 10.3748/wjg.v29.i22.3422 Copyright ©The Author(s) 2023.

Figure 3 Annexin A1 inhibits CCl₄ induced hepatic stellate cell activation in liver tissue. A: Annexin A1 (AnxA1) inhibited collagen deposition induced by CCl₄ in mouse liver tissue. Collagen was stained blue by Masson staining. The collagen content in the liver tissues in the control group was low, while collagen deposition in the liver of the CCl₄-induced model group was high. Collagen deposition in the liver of the Anxa1^{-/-} model group was significantly aggravated

compared with that in the wild-type model group. Active N-terminal peptide of AnxA1 (Ac2-26) significantly reduced collagen deposition in the CCl₄-induced model group, and the collagen deposition in the mice after the intervention of Ac2-26 + N-formylpeptide receptor antagonist N-Boc-Phe-Leu-Phe-Leu-Phe was similar to that in the CCl₄-induced model group. Initial magnification: 100 ×; B: Effect of Ac2-26 on α -smooth muscle actin mRNA expression; C: Effect of Ac2-26 on collagen I mRNA expression; D: Effect of Ac2-26 on connective tissue growth factor mRNA expression; E: Effect of Ac2-26 on transforming growth factor- β 1 mRNA expression; total RNA was extracted using TRIzol reagent and relative mRNA concentration was quantified using quantitative real-time polymerase chain reaction ($n = 8$). Results are presented as ratios of target mRNA or protein normalized to internal GAPDH. ^a $P < 0.05$ and ^b $P < 0.01$ vs control; ^c $P < 0.05$ vs CCl₄; ^d $P < 0.05$ vs CCl₄ + Ac2-26; ^e $P < 0.05$ and ^f $P < 0.01$ vs AnxA1^{-/-} + control; ^g $P < 0.05$ vs AnxA1^{-/-} + CCl₄; ^h $P < 0.05$ and ⁱ $P < 0.01$ vs AnxA1^{-/-} + CCl₄ + Ac2-26; ^j $P < 0.05$ vs intra-group; ^k $P < 0.05$ vs inter-group. Ac2-26: Active N-terminal peptide of AnxA1; AnxA1: Annexin A1; Boc2: N-formyl peptide receptor antagonist N-Boc-Phe-Leu-Phe-Leu-Phe; CTGF: Connective tissue growth factor; TGF- β 1: Transforming growth factor β 1.

group and AnxA1^{-/-} model group was higher, and paralleled the progression of liver fibrosis. The more severe the fibrosis, the higher the expression of CTGF, suggesting that CTGF was closely related to the occurrence and development of liver fibrosis. After Ac2-26 treatment, expression of CTGF in liver tissues of mice in the CCl₄-induced mice was significantly decreased. AnxA1 inhibited expression of CTGF in liver tissue and alleviated the occurrence of liver fibrosis, while Boc2 almost reversed the therapeutic effect of Ac2-26 (Figure 3D), and the difference was significant ($P < 0.05$).

AnxA1 inhibits activation of RAW264.7 cells and HSCs


LPS induced activation of RAW264.7 cells and the release of inflammatory cytokines. Compared with the control group, expression of TGF- β 1, IL-1 β and IL-6 in RAW264.7 cells was significantly induced by LPS, and reached a peak within 1 h, then gradually decreased to the normal level. After activation of RAW264.7 cells, AnxA1 was not significantly increased, but Ac2-26 inhibited TGF β 1, and IL-6 expression in RAW264.7 cells, while Boc2 blocked the inhibitory effect of Ac2-26 on RAW264.7 cells (Figures 4B-D), and the difference was significant ($P < 0.05$). LPS induced activation of HSCs, which expressed TGF β 1, α -SMA, CTGF, and collagen I simultaneously (Figures 4B and 5A-C). Expression of TGF- β 1 and CTGF began to increase at 1 h and peaked at 3 and 6 h, respectively. AnxA1 expression began to increase at 3 h and peaked at 12 h. Expression of α -SMA and collagen I began to increase significantly at 12 h and peaked at 24 h. Ac2-26 treatment significantly decreased LPS-induced expression of TGF- β 1, α -SMA, CTGF and collagen I, and Boc2 significantly inhibited this effect (Figures 4B and 5A-C and the difference was significant ($P < 0.05$). LPS induced activation of hepatocytes. Compared with the control group, AnxA1 was highly expressed after hepatocyte (HC) activation, which was time-dependent and increased from 4 to 24 h (Figure 4A). Ac2-26 significantly upregulated HSC AnxA1 expression, while Boc2 inhibited AnxA1 expression (Figure 4A), and the difference was significant ($P < 0.05$). The results of western blotting and qPCR were consistent (Figures 4E and F).


AnxA1 inhibits Wnt/ β -catenin activation in liver tissue

The Wnt/ β -catenin pathway plays an important role in the formation of liver fibrosis. We used western blotting to detect β -catenin expression, which was associated with inflammation and fibrosis. Compared with the low expression of β -catenin in control mice, expression of β -catenin was upregulated in the model group and higher in the AnxA1^{-/-} group, and expression was higher at 8 wk than at 4 wk. After Ac2-26 treatment, expression of β -catenin protein in the CCl₄-induced model group was significantly decreased compared with that before treatment, while Boc2 inhibited the effect of Ac2-26 (Figure 1D). The results of the animal experiments were verified by *in vitro* cell experiments. β -catenin was highly expressed in LPS-induced HSC activation, and increased progressively with time, starting at 3 h and reaching a peak at 24 h. Ac2-26 inhibited expression of β -catenin protein in LPS-induced HSCs, while Boc2 inhibition blocked this effect, β -catenin protein expression was maintained at a high level in LPS-induced HSCs (Figure 4F).

DISCUSSION

Hepatic fibrosis is a self-repair response of the liver to sustained injury. It is the abnormal structure and function of the liver caused by the excessive accumulation of ECM in the intercellular spaces caused by activation of HSCs and imbalance of synthesis and degradation of ECM. Chronic inflammation caused by sustained injury is the initiating factor of hepatic fibrosis[15]. When the pathogenic factors act on the body, a variety of injurious changes occur, but they also stimulate the body to produce defense, adaptation, compensation and other anti-injury responses. AnxA1 is a glucocorticoid regulatory protein that inhibits neutrophil apoptosis, enhances efferocytosis in macrophages, inhibits activation of phospholipase A2, downregulates expression of inflammatory arachidonic acid, and has a significant negative regulatory effect on several inflammatory cells and factors[22]. In this study, a mouse model of experimental liver fibrosis was established by intraperitoneal injection of 20%CCl₄, and treatment or intervention was performed by knockout of AnxA1 gene, intraperitoneal injection of AnxA1-simulated peptide Ac2-26 and AnxA1 receptor inhibitor Boc2, to observe the effects of AnxA1 on liver fibrosis and






DOI: 10.3748/wjg.v29.i22.3422 Copyright ©The Author(s) 2023.

Figure 4 Annexin A1 inhibits activation of RAW264.7 cells and hepatic stellate cells. A: Changes in annexin A1 (AnxA1) expression in hepatocyte and hepatic stellate cells; B: Transforming growth factor- β 1 mRNA expression; C: Interleukin (IL)-1 β mRNA was quantified using quantitative real-time polymerase chain reaction; D: Changes in IL-6 mRNA in RAW264.7; E: Immunoblotting assay for the detection of AnxA1 in hepatocytes; F: Immunoblotting assay for AnxA1, α -smooth muscle actin (α -SMA) and β -catenin, lysates were analyzed with AnxA1, α -SMA or β -catenin antibody. GAPDH was used as an internal control. ^a $P < 0.05$, ^b $P < 0.01$ and ^c $P < 0.001$ vs controls; ^d $P < 0.05$ and ^e $P < 0.01$ vs lipopolysaccharide; ^f $P < 0.05$, ^g $P < 0.01$ and ^h $P < 0.001$ vs lipopolysaccharide + Ac2-26 ($n = 8$). AnxA1: Annexin A1; α -SMA: α -smooth muscle actin; HSC: Hepatic stellate cell; LPS: Lipopolysaccharide; TGF- β 1: Transforming growth factor β 1; HCs: Hepatocyte; IL: Interleukin; Ac2-26: Active N-terminal peptide of AnxA1; Boc2: N-formyl peptide receptor antagonist N-Boc-Phe-Leu-Phe-Leu-Phe.

explore its main targets. We found that AnxA1 protein and mRNA levels in the liver fibrosis model group were higher than those in the normal control group, and were time-dependent. AnxA1 began to increase at 4 wk, peaked at 8 wk, and then began to decline, consistent with the severity of liver fibrosis in mice, suggesting that AnxA1 is associated with the onset of liver fibrosis. Further studies showed that AnxA1 inhibited fibrosis. Firstly, gross liver specimens revealed that the liver surface roughness and nodulation of mice in the *AnxA1*^{-/-} model group were more severe than those in the wild-type model group, while the lesions in the Ac2-26 group were significantly reduced after treatment, suggesting that Ac2-26 alleviated liver damage. Secondly, HE staining showed that compared with the wild-type model group, liver damage and inflammatory cell infiltration were further aggravated in the *AnxA1*^{-/-} model group, suggesting serious liver damage, which was significantly reduced after treatment with Ac2-26. In addition, Masson staining indicated collagen deposition in the portal area in the wild-type model group, CCl₄ induced more significant collagen deposition in the liver tissue of the *AnxA1*^{-/-} model group, with a wide area, especially at 8 wk, when large amounts of collagen surrounded the liver tissue to form pseudolobules rather than concentrating in the portal area. Collagen deposition in liver tissue of the *AnxA1*^{-/-} model group progressed significantly, and was significantly improved in the group treated with Ac2-26. AnxA1 inhibited liver inflammation and collagen deposition, and relieved liver fibrosis. AnxA1 may play an important role in the occurrence and development of liver fibrosis. While Boc2 blocked the anti-inflammatory and antifibrotic effects of Ac2-26, AnxA1 may have participated in the pathogenesis of liver fibrosis by targeting FPR. Chronic liver disease is progressive, in which inflammation and fibrosis form a vicious cycle, and eventually the excess deposition of ECM, especially type I collagen, makes the lesions difficult to reverse[19,20]. The results of our study showed that the collagen I concentration in liver tissues of mice in the CCl₄-induced model group was significantly increased, and collagen I content in the *AnxA1*^{-/-} model group was higher than that in the wild-type model group, and Boc2 reversed the therapeutic effect of Ac2-26 by reducing the collagen I concentration in liver tissues. High collagen I expression was maintained in the CCl₄-induced model group. It was verified that AnxA1 inhibited liver collagen deposition and relieved liver fibrosis through FPR. Therefore, AnxA1 has a protective effect on hepatic fibrosis.

There have been few studies on the effects of AnxA1 on liver fibrosis. The possible protective mechanism of AnxA1 on liver fibrosis is discussed. HSCs play an important role in the formation of liver fibrosis[19,20], and activation of HSCs is the main cause of deposition of fibrous collagen (types I, III and IV) and other stroma-like proteins in chronic liver disease[23,24], and α -SMA is an activation marker of HSCs. The activation and development of HSCs into fibroblasts is accompanied by increased expression of α -SMA[25], and the degree of HSC activation can be determined by detecting the expression level of α -SMA. In this study, the expression of α -SMA was detected by western blotting and qPCR. The expression of α -SMA in liver tissue of mice in the CCl₄-induced model group was higher than that in the control group, and the activation of HSCs induced by CCl₄ in *AnxA1*^{-/-} mice was higher



DOI: 10.3748/wjg.v29.i22.3422 Copyright ©The Author(s) 2023.

Figure 5 Annexin A1 inhibits hepatic stellate cell activation. A: Changes in α -smooth muscle actin expression in hepatic stellate cell; B: Connective tissue growth factor mRNA was quantified using quantitative real-time polymerase chain reaction; C: Collagen I mRNA expression. GAPDH was used as an internal control. ^a $P < 0.05$ and ^b $P < 0.01$ vs controls; ^c $P < 0.05$ vs lipopolysaccharide; ^d $P < 0.05$ and ^e $P < 0.01$ vs lipopolysaccharide + Ac2-26 ($n = 8$). AnxA1: Annexin A1; α -SMA: α -smooth muscle actin; HSC: Hepatic stellate cell; CTGF: Connective tissue growth factor; LPS: Lipopolysaccharide; Ac2-26: Active N-terminal peptide of AnxA1; Boc2: N-formyl peptide receptor antagonist N-Boc-Phe-Leu-Phe-Leu-Phe.

than that in wild-type mice. Supplementation of exogenous Ac2-26 significantly inhibited the expression of α -SMA, and Boc2 blocked the inhibitory effect of Ac2-26. It is suggested that Ac2-26 inhibits HSC activation, which may be one of the key factors in inhibiting liver fibrosis by AnxA1, in which FPR receptors are involved.

HSCs are the main receivers of inflammatory signals and the main promoters of liver fibrosis. Hepatocyte injury triggers an inflammatory response, releases various cytokines and inflammatory factors, activates HSCs, and causes secretion and deposition of ECM, leading to liver fibrosis[26,27]. Immune cells, especially Kupffer cells and macrophages in the liver, are key regulators of inflammation [28]. Kupffer cells are intrinsic macrophages in the liver, which play an important role in the elimination of and defense against bacteria and pathogenic substances from the gastrointestinal tract[29]. In the case

of liver injury, Kupffer cells are activated by damage-associated pathogenic patterns or LPS, viral DNA, and other pathogen-associated molecular patterns, producing and releasing TGF- β 1, IL-1 β and IL-6 to activate HSCs, causing the secretion and deposition of ECM and leading to liver fibrosis[30-32]. Our results showed that hepatic fibrosis was accompanied by inflammatory infiltration and secretion of inflammatory cytokines TGF- β 1, IL-1 β and IL-6, which increased with time, suggesting that hepatic inflammation is related to the severity of fibrosis. Exogenous Ac2-26 significantly reduced hepatic inflammatory cell infiltration and inflammatory factor expression in the CCl₄-induced model group, while alleviating the formation of liver fibrosis, and Boc2 inhibited the anti-inflammatory effect of Ac2-26. AnxA1 may reduce fibrosis by inhibiting the inflammatory response in the liver, and its mechanism may be related to inhibition of HSC activation by targeting FPR to regulate macrophage function.

TGF- β 1 is a key regulatory factor in the process of liver fibrosis[32], and CTGF, as its downstream factor, mediates TGF- β 1, promoting ECM production and accumulation[33]. HSCs are the main source of CTGF, and blocking CTGF can significantly inhibit HSC activation and proliferation, and reduce tissue inhibitors of metalloproteinase, matrix metalloproteinase (MMP)2, MMP9 and type I collagen. Our results showed that increased expression of CTGF in the liver tissue of mice in the CCl₄-induced model group was also inhibited by Ac2-26, suggesting that the inhibitory effect of AnxA1 on liver fibrosis is closely related to the expression of TGF- β 1 and its downstream factor CTGF.

In order to verify the results of the animal experiments and to better investigate the mechanism of AnxA1 in liver fibrosis, we conducted *in vitro* cell experiments. Ac2-26 significantly inhibited LPS-induced activation of RAW264.7 cells and downregulated TGF- β 1 and other inflammatory factors expressed in RAW264.7 cells after activation, and Boc2 reversed the above results. LPS induced HCs and HSCs to increase Anxa1 expression, but LPS induced RAW264.7 cells did not upregulate AnxA1 expression, and the activated HSCs that overexpressed SMA, collagen, and CTGF were inhibited by exogenous Ac2-26. AnxA1 mediated by FPR may affect the function of macrophages through HC paracrine or HSC autocrine effects, control inflammatory responses, inhibit HSC activation, and reduce ECM deposition by negative feedback, so as to prevent liver fibrosis formation.

Recent studies have shown that the Wnt/ β -catenin signaling pathway can promote liver fibrosis and play an important role in HSC proliferation, metastasis and ECM formation[34]. Inhibition of Wnt/ β -catenin signaling can inhibit the proliferation of HSCs and induce their apoptosis, thus alleviating CCl₄-induced liver fibrosis[34-36]. In addition, TGF- β 1 activates the Wnt/ β -catenin pathway by inhibiting the expression of Wnt/ β -catenin inhibitors[36]. Our results suggest that Ac2-26 downregulates the expression of β -catenin in the Wnt signaling pathway in mouse liver fibrosis, and inhibits the expression of activated HSC β -catenin. Combined with previous findings, the therapeutic effects of AnxA1 may alleviate the development of liver fibrosis by inhibiting the Wnt/ β -catenin pathway of HSCs.

CONCLUSION

AnxA1 may play an important role in the formation of liver fibrosis, including regulating macrophage function, reducing inflammation, inhibiting HSC activation, and improving collagen deposition. AnxA1 mediated by FPR receptors may regulate the activity of macrophages through HSC autocrine or paracrine functions, inhibit the Wnt/ β -catenin pathway, and inhibit HSC activity. Although its further mechanism of action has not been clearly elucidated, our results suggest that AnxA1 has potential activity against hepatic fibrosis and can block the occurrence of fibrosis at multiple targets, which may provide a new option for liver fibrosis treatment.

ARTICLE HIGHLIGHTS

Research background

Liver fibrosis is a pathological change caused by chronic liver injury, and is a key link in the progression of chronic liver diseases to cirrhosis or liver cancer. How to prevent and reverse liver fibrosis is currently a hot and difficult research topic, and fully understanding and elucidating the pathogenesis of liver fibrosis is the foundation of prevention and treatment. At present, there is still a lack of safe and effective treatment for liver cirrhosis clinically. It is of great significance to deeply study the pathogenesis of liver fiber and prevent and treat it in order to reduce the incidence rate and mortality of liver cirrhosis.

Research motivation

Studies have shown that liver fibrosis is closely related to inflammation in chronic liver disease, and annexin (Anx)A1 has strong anti-inflammatory effects. Therefore, this study focuses on the pathogenesis of liver fibrosis and investigates the role of AnxA1 in liver fibrosis. Provide valuable information for antifibrosis therapy.

Research objectives

To investigate molecular and cellular mechanisms of liver fibrosis and evaluate the effect of AnxA1 in hepatic fibrosis.

Research methods

Biomarkers as drug development tools, biomarker discovery and validation using a combination of *in vitro* and *in vivo* studies. A large and wide range of animal models and cell experiments can be used, and experiments can be repeated multiple times.

Research results

AnxA1/formylpeptide receptor (FPR) inhibits the activation of hepatic stellate cells by regulating macrophage function through the Wnt/ β -catenin pathway in the CCl₄-induced hepatic fibrosis mice.

Research conclusions

AnxA1 may involve in the pathogenesis of liver fibrosis and as a potential therapeutic target.

Research perspectives

This study investigates the mechanism of Anxa1 in liver fibrosis by using wild mice/Anxa1 knockout mice in combination with Anxa1 mimetic peptide and FPR receptor inhibitor to provide an experimental basis for the possible use of Anxa1 in the treatment of liver fibrosis.

FOOTNOTES

Author contributions: Fan JH, Luo N and Liu GF performed most of the experiments; Luo N, Xu XF and Li SQ analyzed the data; Fan JH and Lv XP designed the study and wrote the paper.

Supported by a Grant-in-Aid for Scientific Research from National Natural Science Foundation of China, No. 81860120 and 81860104; Guangxi Natural Science Foundation, No. 2017GXNSFBA198134, 2017GXNSFAA198299 and 2015GXNSFCA139024.

Institutional animal care and use committee statement: All experiments were conducted in accordance with the institutional guidelines of Guangxi Medical University for the care and use of laboratory animals.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: No additional data are available.

ARRIVE guidelines statement: The authors have read the ARRIVE guidelines, and the manuscript was prepared and revised according to the ARRIVE guidelines.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Jun-Hua Fan 0000-0002-7292-7484; Na Luo 0000-0001-7269-4476; Geng-Feng Liu 0000-0002-2414-7707; Xiao-Fang Xu 0000-0001-7986-1740; Shi-Quan Li 0000-0003-3744-2737; Xiao-Ping Lv 0000-0001-7144-9508.

S-Editor: Wang JJ

L-Editor: A

P-Editor: Wang JJ

REFERENCES

- Caligiuri A, Gentilini A, Pastore M, Gitto S, Marra F. Cellular and Molecular Mechanisms Underlying Liver Fibrosis Regression. *Cells* 2021; **10** [PMID: 34685739 DOI: 10.3390/cells10102759]
- Liu X, Mi X, Wang Z, Zhang M, Hou J, Jiang S, Wang Y, Chen C, Li W. Ginsenoside Rg3 promotes regression from hepatic fibrosis through reducing inflammation-mediated autophagy signaling pathway. *Cell Death Dis* 2020; **11**: 454 [PMID: 32532964 DOI: 10.1038/s41419-020-2597-7]
- Seki E, Brenner DA. Recent advancement of molecular mechanisms of liver fibrosis. *J Hepatobiliary Pancreat Sci* 2015;

- 22: 512-518 [PMID: [25869468](#) DOI: [10.1002/jhbp.245](#)]
- 4 **Ezhilarasan D**, Sokal E, Najimi M. Hepatic fibrosis: It is time to go with hepatic stellate cell-specific therapeutic targets. *Hepatobiliary Pancreat Dis Int* 2018; **17**: 192-197 [PMID: [29709350](#) DOI: [10.1016/j.hbpd.2018.04.003](#)]
- 5 **Koo JH**, Lee HJ, Kim W, Kim SG. Endoplasmic Reticulum Stress in Hepatic Stellate Cells Promotes Liver Fibrosis via PERK-Mediated Degradation of HNRNPA1 and Up-regulation of SMAD2. *Gastroenterology* 2016; **150**: 181-193.e8 [PMID: [26435271](#) DOI: [10.1053/j.gastro.2015.09.039](#)]
- 6 **Parola M**, Pinzani M. Liver fibrosis: Pathophysiology, pathogenetic targets and clinical issues. *Mol Aspects Med* 2019; **65**: 37-55 [PMID: [30213667](#) DOI: [10.1016/j.mam.2018.09.002](#)]
- 7 **Lan T**, Li C, Yang G, Sun Y, Zhuang L, Ou Y, Li H, Wang G, Kisseleva T, Brenner D, Guo J. Sphingosine kinase 1 promotes liver fibrosis by preventing miR-19b-3p-mediated inhibition of CCR2. *Hepatology* 2018; **68**: 1070-1086 [PMID: [29572892](#) DOI: [10.1002/hep.29885](#)]
- 8 **Cheng D**, Chai J, Wang H, Fu L, Peng S, Ni X. Hepatic macrophages: Key players in the development and progression of liver fibrosis. *Liver Int* 2021; **41**: 2279-2294 [PMID: [33966318](#) DOI: [10.1111/liv.14940](#)]
- 9 **Li P**, He K, Li J, Liu Z, Gong J. The role of Kupffer cells in hepatic diseases. *Mol Immunol* 2017; **85**: 222-229 [PMID: [28314211](#) DOI: [10.1016/j.molimm.2017.02.018](#)]
- 10 **Alegre F**, Pelegrin P, Feldstein AE. Inflammasomes in Liver Fibrosis. *Semin Liver Dis* 2017; **37**: 119-127 [PMID: [28564720](#) DOI: [10.1055/s-0037-1601350](#)]
- 11 **Gan C**, Cai Q, Tang C, Gao J. Inflammasomes and Pyroptosis of Liver Cells in Liver Fibrosis. *Front Immunol* 2022; **13**: 896473 [PMID: [35707547](#) DOI: [10.3389/fimmu.2022.896473](#)]
- 12 **Senchenkova EY**, Ansari J, Becker F, Vital SA, Al-Yafeai Z, Sparkenbaugh EM, Pawlinski R, Stokes KY, Carroll JL, Dragoi AM, Qin CX, Ritchie RH, Sun H, Cuellar-Saenz HH, Rubinstein MR, Han YW, Orr AW, Perretti M, Granger DN, Gavins FNE. Novel Role for the AnxA1-Fpr2/ALX Signaling Axis as a Key Regulator of Platelet Function to Promote Resolution of Inflammation. *Circulation* 2019; **140**: 319-335 [PMID: [31154815](#) DOI: [10.1161/CIRCULATIONAHA.118.039345](#)]
- 13 **Xu X**, Gao W, Li L, Hao J, Yang B, Wang T, Bai X, Li F, Ren H, Zhang M, Zhang L, Wang J, Wang D, Zhang J, Jiao L. Annexin A1 protects against cerebral ischemia-reperfusion injury by modulating microglia/macrophage polarization via FPR2/ALX-dependent AMPK-mTOR pathway. *J Neuroinflammation* 2021; **18**: 119 [PMID: [34022892](#) DOI: [10.1186/s12974-021-02174-3](#)]
- 14 **Vecchi L**, Alves Pereira Zóia M, Goss Santos T, de Oliveira Beserra A, Colaço Ramos CM, França Matias Colombo B, Paiva Maia YC, Piana de Andrade V, Teixeira Soares Mota S, Gonçalves de Araújo T, Van Petten de Vasconcelos Azevedo F, Soares FA, Oliani SM, Goulart LR. Inhibition of the AnxA1/FPR1 autocrine axis reduces MDA-MB-231 breast cancer cell growth and aggressiveness *in vitro* and *in vivo*. *Biochim Biophys Acta Mol Cell Res* 2018; **1865**: 1368-1382 [PMID: [29932988](#) DOI: [10.1016/j.bbamer.2018.06.010](#)]
- 15 **Neymeyer H**, Labes R, Reverte V, Saez F, Stroth T, Dathe C, Hohberger S, Zeisberg M, Müller GA, Salazar J, Bachmann S, Paliege A. Activation of annexin A1 signalling in renal fibroblasts exerts antifibrotic effects. *Acta Physiol (Oxf)* 2015; **215**: 144-158 [PMID: [26332853](#) DOI: [10.1111/apha.12586](#)]
- 16 **Lai T**, Li Y, Mai Z, Wen X, Lv Y, Xie Z, Lv Q, Chen M, Wu D, Wu B. Annexin A1 is elevated in patients with COPD and affects lung fibroblast function. *Int J Chron Obstruct Pulmon Dis* 2018; **13**: 473-486 [PMID: [29440885](#) DOI: [10.2147/COPD.S149766](#)]
- 17 **Vallée A**, Lecarpentier Y, Guillemin R, Vallée JN. Interactions between TGF- β 1, canonical WNT/ β -catenin pathway and PPAR γ in radiation-induced fibrosis. *Oncotarget* 2017; **8**: 90579-90604 [PMID: [29163854](#) DOI: [10.18632/oncotarget.21234](#)]
- 18 **Yu FJ**, Dong PH, Fan XF, Lin Z, Chen YP, Li J. Down-regulation of angiotensin II by shRNA reduces collagen synthesis in hepatic stellate cells. *Int J Mol Med* 2010; **25**: 801-806 [PMID: [20372825](#) DOI: [10.3892/ijmm.00000407](#)]
- 19 **Katoh M**. Multilayered prevention and treatment of chronic inflammation, organ fibrosis and cancer associated with canonical WNT/ β -catenin signaling activation (Review). *Int J Mol Med* 2018; **42**: 713-725 [PMID: [29786110](#) DOI: [10.3892/ijmm.2018.3689](#)]
- 20 **Zhang C**, Zhang Y, Zhang W, Tong H, Li S, Yan Y. WISP1 promotes bovine MDSC differentiation via recruitment of ANXA1 for the regulation of the TGF- β signalling pathway. *Mol Cell Biochem* 2020; **470**: 215-227 [PMID: [32458119](#) DOI: [10.1007/s11010-020-03763-1](#)]
- 21 **Fan JH**, Feng GG, Huang L, Tang GD, Jiang HX, Xu J. Naofen promotes TNF- α -mediated apoptosis of hepatocytes by activating caspase-3 in lipopolysaccharide-treated rats. *World J Gastroenterol* 2014; **20**: 4963-4971 [PMID: [24803807](#) DOI: [10.3748/wjg.v20.i17.4963](#)]
- 22 **Zharkova O**, Salamah MF, Babak MV, Rajan E, Lim LHK, Andrade F, Gil CD, Oliani SM, Moraes LA, Vaiyapuri S. Deletion of Annexin A1 in Mice Upregulates the Expression of Its Receptor, Fpr2/3, and Reactivity to the AnxA1 Mimetic Peptide in Platelets. *Int J Mol Sci* 2023; **24** [PMID: [36834844](#) DOI: [10.3390/ijms24043424](#)]
- 23 **Hall A**, Cotoi C, Luong TV, Watkins J, Bhathal P, Quaglia A. Collagen and elastic fibres in acute and chronic liver injury. *Sci Rep* 2021; **11**: 14569 [PMID: [34267266](#) DOI: [10.1038/s41598-021-93566-1](#)]
- 24 **Rong X**, Yang Y, Zhang G, Zhang H, Li C, Wang Y. Antler stem cells as a novel stem cell source for reducing liver fibrosis. *Cell Tissue Res* 2020; **379**: 195-206 [PMID: [31428875](#) DOI: [10.1007/s00441-019-03081-z](#)]
- 25 **Wang R**, Zhang D, Tang D, Sun K, Peng J, Zhu W, Yin S, Wu Y. Amygdalin inhibits TGF β 1-induced activation of hepatic stellate cells (HSCs) *in vitro* and CCl(4)-induced hepatic fibrosis in rats *in vivo*. *Int Immunopharmacol* 2021; **90**: 107151 [PMID: [33296784](#) DOI: [10.1016/j.intimp.2020.107151](#)]
- 26 **Delgado ME**, Cárdenas BI, Farran N, Fernandez M. Metabolic Reprogramming of Liver Fibrosis. *Cells* 2021; **10** [PMID: [34944111](#) DOI: [10.3390/cells10123604](#)]
- 27 **Papachristoforou E**, Ramachandran P. Macrophages as key regulators of liver health and disease. *Int Rev Cell Mol Biol* 2022; **368**: 143-212 [PMID: [35636927](#) DOI: [10.1016/bs.ircmb.2022.04.006](#)]
- 28 **Seo HY**, Lee SH, Han E, Hwang JS, Han S, Kim MK, Jang BK. Evogliptin Directly Inhibits Inflammatory and Fibrotic Signaling in Isolated Liver Cells. *Int J Mol Sci* 2022; **23** [PMID: [36232933](#) DOI: [10.3390/ijms231911636](#)]

- 29 **Wu J**, Zern MA. Hepatic stellate cells: a target for the treatment of liver fibrosis. *J Gastroenterol* 2000; **35**: 665-672 [PMID: [11023037](#) DOI: [10.1007/s005350070045](#)]
- 30 **Ohara M**, Ohnishi S, Hosono H, Yamamoto K, Yuyama K, Nakamura H, Fu Q, Maehara O, Suda G, Sakamoto N. Extracellular Vesicles from Amnion-Derived Mesenchymal Stem Cells Ameliorate Hepatic Inflammation and Fibrosis in Rats. *Stem Cells Int* 2018; **2018**: 3212643 [PMID: [30675167](#) DOI: [10.1155/2018/3212643](#)]
- 31 **Xiang DM**, Sun W, Ning BF, Zhou TF, Li XF, Zhong W, Cheng Z, Xia MY, Wang X, Deng X, Wang W, Li HY, Cui XL, Li SC, Wu B, Xie WF, Wang HY, Ding J. The HLF/IL-6/STAT3 feedforward circuit drives hepatic stellate cell activation to promote liver fibrosis. *Gut* 2018; **67**: 1704-1715 [PMID: [28754776](#) DOI: [10.1136/gutjnl-2016-313392](#)]
- 32 **Zigmond E**, Samia-Grinberg S, Pasmanik-Chor M, Brazowski E, Shibolet O, Halpern Z, Varol C. Infiltrating monocyte-derived macrophages and resident kupffer cells display different ontogeny and functions in acute liver injury. *J Immunol* 2014; **193**: 344-353 [PMID: [24890723](#) DOI: [10.4049/jimmunol.1400574](#)]
- 33 **Trampuž SR**, van Riet S, Nordling Å, Ingelman-Sundberg M. The Role of CTGF in Liver Fibrosis Induced in 3D Human Liver Spheroids. *Cells* 2023; **12** [PMID: [36672237](#) DOI: [10.3390/cells12020302](#)]
- 34 **Rong X**, Liu J, Yao X, Jiang T, Wang Y, Xie F. Human bone marrow mesenchymal stem cells-derived exosomes alleviate liver fibrosis through the Wnt/ β -catenin pathway. *Stem Cell Res Ther* 2019; **10**: 98 [PMID: [30885249](#) DOI: [10.1186/s13287-019-1204-2](#)]
- 35 **Lee IH**, Im E, Lee HJ, Sim DY, Lee JH, Jung JH, Park JE, Shim BS, Kim SH. Apoptotic and antihepatofibrotic effect of honokiol *via* activation of GSK3 β and suppression of Wnt/ β -catenin pathway in hepatic stellate cells. *Phytother Res* 2021; **35**: 452-462 [PMID: [32776713](#) DOI: [10.1002/ptr.6824](#)]
- 36 **El-Fadaly AA**, Afifi NA, El-Eraky W, Salama A, Abdelhameed MF, El-Rahman SSA, Ramadan A. Fisetin alleviates thioacetamide-induced hepatic fibrosis in rats by inhibiting Wnt/ β -catenin signaling pathway. *Immunopharmacol Immunotoxicol* 2022; **44**: 355-366 [PMID: [35255766](#) DOI: [10.1080/08923973.2022.2047198](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

