

# World Journal of *Gastroenterology*

*World J Gastroenterol* 2024 March 14; 30(10): 1261-1469



## EDITORIAL

- 1261 Bridging the gap: Unveiling the crisis of physical inactivity in inflammatory bowel diseases  
*Stafie R, Singeap AM, Rotaru A, Stanciu C, Trifan A*
- 1266 Double role of depression in gastric cancer: As a causative factor and as consequence  
*Christodoulidis G, Konstantinos-Elleftherios K, Marina-Nektaria K*
- 1270 Capsule endoscopy and panendoscopy: A journey to the future of gastrointestinal endoscopy  
*Rosa B, Cotter J*
- 1280 Vonoprazan-amoxicillin dual regimen with *Saccharomyces boulardii* as a rescue therapy for *Helicobacter pylori*: Current perspectives and implications  
*Dirjayanto VJ, Audrey J, Simadibrata DM*
- 1287 Women health and microbiota: Different aspects of well-being  
*Nannini G, Amedei A*
- 1291 Nomograms and prognosis for superficial esophageal squamous cell carcinoma  
*Lin HT, Abdelbaki A, Krishna SG*

## REVIEW

- 1295 Overview of the immunological mechanisms in hepatitis B virus reactivation: Implications for disease progression and management strategies  
*Ma H, Yan QZ, Ma JR, Li DF, Yang JL*
- 1313 Optimizing nutrition in hepatic cirrhosis: A comprehensive assessment and care approach  
*Mendez-Guerrero O, Carranza-Carrasco A, Chi-Cervera LA, Torre A, Navarro-Alvarez N*
- 1329 Optimizing prediction models for pancreatic fistula after pancreatectomy: Current status and future perspectives  
*Yang F, Windsor JA, Fu DL*

## ORIGINAL ARTICLE

## Retrospective Cohort Study

- 1346 Cumulative effects of excess high-normal alanine aminotransferase levels in relation to new-onset metabolic dysfunction-associated fatty liver disease in China  
*Chen JF, Wu ZQ, Liu HS, Yan S, Wang YX, Xing M, Song XQ, Ding SY*
- 1358 Time trends and outcomes of gastrostomy placement in a Swedish national cohort over two decades  
*Skogar ML, Sundbom M*

**Retrospective Study**

- 1368 Stage at diagnosis of colorectal cancer through diagnostic route: Who should be screened?

*Agatsuma N, Utsumi T, Nishikawa Y, Horimatsu T, Seta T, Yamashita Y, Tanaka Y, Inoue T, Nakanishi Y, Shimizu T, Ohno M, Fukushima A, Nakayama T, Seno H*

**Observational Study**

- 1377 Differential diagnosis of Crohn's disease and intestinal tuberculosis based on ATR-FTIR spectroscopy combined with machine learning

*Li YP, Lu TY, Huang FR, Zhang WM, Chen ZQ, Guang PW, Deng LY, Yang XH*

**Prospective Study**

- 1393 Establishment and validation of an adherence prediction system for lifestyle interventions in non-alcoholic fatty liver disease

*Zeng MH, Shi QY, Xu L, Mi YQ*

**Basic Study**

- 1405 Alkaline sphingomyelinase deficiency impairs intestinal mucosal barrier integrity and reduces antioxidant capacity in dextran sulfate sodium-induced colitis

*Tian Y, Li X, Wang X, Pei ST, Pan HX, Cheng YQ, Li YC, Cao WT, Petersen JDD, Zhang P*

- 1420 Preliminary exploration of animal models of congenital choledochal cysts

*Zhang SH, Zhang YB, Cai DT, Pan T, Chen K, Jin Y, Luo WJ, Huang ZW, Chen QJ, Gao ZG*

- 1431 Serotonin receptor 2B induces visceral hyperalgesia in rat model and patients with diarrhea-predominant irritable bowel syndrome

*Li ZY, Mao YQ, Hua Q, Sun YH, Wang HY, Ye XG, Hu JX, Wang YJ, Jiang M*

**META-ANALYSIS**

- 1450 Shear-wave elastography to predict hepatocellular carcinoma after hepatitis C virus eradication: A systematic review and meta-analysis

*Esposito G, Santini P, Galasso L, Mignini I, Ainora ME, Gasbarrini A, Zocco MA*

**LETTER TO THE EDITOR**

- 1461 Current considerations on intraductal papillary neoplasms of the bile duct and pancreatic duct

*Pavlidis ET, Galanis IN, Pavlidis TE*

- 1466 Are we ready to use new endoscopic scores for ulcerative colitis?

*Quera R, Núñez F P*

**ABOUT COVER**

Editorial Board Member of *World Journal of Gastroenterology*, Toru Mizuguchi, MD, PhD, Professor, Surgeon, Department of Nursing, Division of Surgical Science, Sapporo Medical University Postgraduate School of Health Science, Sapporo, Hokkaido 0608556, Japan. [tmizu@sapmed.ac.jp](mailto:tmizu@sapmed.ac.jp)

**AIMS AND SCOPE**

The primary aim of *World Journal of Gastroenterology* (WJG, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

**INDEXING/ABSTRACTING**

The WJG is now abstracted and indexed in Science Citation Index Expanded (SCIE), MEDLINE, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJG as 4.3; Quartile category: Q2. The WJG's CiteScore for 2021 is 8.3.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xiang Li*; Cover Editor: *Jia-Ru Fan*.

**NAME OF JOURNAL**

*World Journal of Gastroenterology*

**ISSN**

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

**LAUNCH DATE**

October 1, 1995

**FREQUENCY**

Weekly

**EDITORS-IN-CHIEF**

Andrzej S Tarnawski

**EXECUTIVE ASSOCIATE EDITORS-IN-CHIEF**

Xian-Jun Yu (Pancreatic Oncology), Jian-Gao Fan (Chronic Liver Disease), Hou-Bao Liu (Biliary Tract Disease)

**EDITORIAL BOARD MEMBERS**

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

**PUBLICATION DATE**

March 14, 2024

**COPYRIGHT**

© 2024 Baishideng Publishing Group Inc

**PUBLISHING PARTNER**

Shanghai Pancreatic Cancer Institute and Pancreatic Cancer Institute, Fudan University  
Biliary Tract Disease Institute, Fudan University

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**POLICY OF CO-AUTHORS**

<https://www.wjgnet.com/bpg/GerInfo/310>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

**PUBLISHING PARTNER's OFFICIAL WEBSITE**

<https://www.shca.org.cn>  
<https://www.zs-hospital.sh.cn>



## Double role of depression in gastric cancer: As a causative factor and as consequence

Grigorios Christodoulidis, Koumarelas Konstantinos-Eleftherios, Kouliou Marina-Nektaria

**Specialty type:** Gastroenterology and hepatology

**Provenance and peer review:** Invited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): 0  
Grade C (Good): 0  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Tan JK, Malaysia

**Received:** December 16, 2023

**Peer-review started:** December 16, 2023

**First decision:** January 4, 2024

**Revised:** January 13, 2024

**Accepted:** February 23, 2024

**Article in press:** February 23, 2024

**Published online:** March 14, 2024



**Grigorios Christodoulidis, Koumarelas Konstantinos-Eleftherios, Kouliou Marina-Nektaria,** Department of General Surgery, University Hospital of Larissa, Larissa 41110, Greece

**Corresponding author:** Grigorios Christodoulidis, MD, PhD, Surgeon, Department of General Surgery, University Hospital of Larissa, Mezourlo, Larissa 41110, Greece. [gregsurg@yahoo.gr](mailto:gregsurg@yahoo.gr)

### Abstract

In this editorial we comment on the article "Hotspots and frontiers of the relationship between gastric cancer and depression: A bibliometric study". Gastric cancer (GC) is a common malignancy in the digestive system with increased mortality and morbidity rates globally. Standard treatments, such as gastrectomy, negatively impact patients' quality of life and beyond the physical strain, GC patients face psychological challenges, including anxiety and depression. The prevalence of depression can be as high as 57%, among gastrointestinal cancer patients. Due to the advancements in treatment effectiveness and increased 5-year overall survival rates, attention has shifted to managing psychological effects. However, the significance of managing the depression doesn't lie solely in the need for a better psychological status. Depression leads to chronic stress activating the sympathetic nervous system and the hypothalamus-pituitary-adrenal axis, leading release of catecholamines inducing tumor proliferation, migration, and metastasis, contributing to GC progression. The dysregulation of neurotransmitters and the involvement of various signaling pathways underscore the complex interplay between depression and GC. Comprehensive strategies are required to address the psychological aspects of GC, including region-specific interventions and increased monitoring for depression. Understanding the intricate relationship between depression and GC progression is essential for developing effective therapeutic strategies and improving overall outcomes for patients facing this complex disease. In this Editorial we delve into double role of depression in the pathogenesis of GC and as a complication of it.

**Key Words:** Gastric cancer; Depression; Anxiety; Chronic stress; Pathogenesis of gastric cancer

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Gastric cancer (GC), a prevalent malignancy in the digestive system, poses a dual challenge with both physical and psychological implications. While standard treatments like gastrectomy impact patients' quality of life, the psychological burden, including anxiety and depression, cannot be overlooked. Depression, reaching prevalence rates of 57%, significantly influences cancer outcomes, affecting mental well-being, treatment adherence, and overall quality of life. Chronic stress and neurotransmitter dysregulation play a pivotal role in GC development, activating pathways that induce tumor progression. Understanding the intricate connection between depression and GC not only highlights the need for comprehensive psychological support but also unveils potential therapeutic targets. Addressing both the physical and psychological aspects of GC is essential for enhancing the overall well-being and outcomes of patients grappling with this complex disease.

**Citation:** Christodoulidis G, Konstantinos-Eleftherios K, Marina-Nektaria K. Double role of depression in gastric cancer: As a causative factor and as consequence. *World J Gastroenterol* 2024; 30(10): 1266-1269

**URL:** <https://www.wjgnet.com/1007-9327/full/v30/i10/1266.htm>

**DOI:** <https://dx.doi.org/10.3748/wjg.v30.i10.1266>

## INTRODUCTION

Gastric cancer (GC) stands as the most prevalent malignant tumor in the digestive system, holding the record for the third-highest mortality and fifth-highest morbidity rates among all cancers. Global statistics underscore the gravity of the situation, revealing an estimated 1 million new cases and 760000 deaths in 2020 alone[1-4]. The standard treatment for GC, gastrectomy, while common, has detrimental effects on patients' quality of life (QoL) and mental well-being. Total gastrectomy, an aspect of this treatment, triggers substantial weight loss, thereby impacting the nutritional status of individuals with the disease. GC alone, can cause disturbing and disabling nausea, vomiting, diarrhea having a significant impact on the patients' nutritional status[2,3,5]. Beyond the physical strain, patients diagnosed with GC confront some psychological challenges, including anxiety, depression, pain, and fatigue[1,3,5,6]. The prevalence of anxiety and depression can reach as high as 47.2% and 57% of patients with gastrointestinal cancer[6]. The last years, having an increased effectiveness of the treatment options and an increased 5-year overall survival, the attention shifts to managing the psychological effects accompanying the disease and the treatment. These challenges emphasize the urgent need for interventions aimed to enhance the overall QoL of these patients. Depression emerges as a pervasive issue among cancer patients, particularly affecting their mental well-being. Contributing factors include the dysregulation of miRNA expression, abnormalities in receptors, and structural changes in the brain[1,7]. Such emotional distress not only shapes the attitude of cancer patients but also influences treatment adherence, underscoring its critical role in determining overall QoL[1,6]. Psychological distress becomes a notable risk factor for treatment non-compliance, increasing the mortality rates. The repercussions of depression extend further, exerting a negative influence on the prognosis of GC and resulting in poor survival outcomes. Depression in the context of cancer, including GC, is linked to chronic psychological stress. Stress-associated neurotransmitters, particularly catecholamines, emerge as potential influencers of cancer progression[5,8]. Chronic stress, often manifesting as anxiety and depression, can trigger tumor development through pathways involving  $\beta$ 2-adrenergic receptors and epithelial-mesenchymal transition (EMT). Despite the acknowledgment of chronic stress and  $\beta$ 2-adrenergic receptors in tumor progression, the precise mechanisms of how EMT is regulated by  $\beta$ 2-AR remain elusive[4,8]. Consequently, there is a need for a deeper understanding of these mechanisms to guide more effective therapeutic strategies. Recognizing the gravity of depression's impact on cancer outcomes, proper treatment is deemed essential for cancer patients. This treatment aims not only to mitigate adverse effects but also to improve symptoms, ensuring the long-term efficacy of interventions for individuals grappling with the complexities of GC.

## ROLE OF DEPRESSION IN GASTRIC CANCER DEVELOPMENT

Under the influence of chronic stress, the sympathetic nervous system is activated as well as the hypothalamus-pituitary-adrenal axis, thereby triggering the release of neurotransmitters such as norepinephrine and epinephrine ( $P < 0.005$ )[5]. The increased expression of catecholamines within the tumor microenvironment has been revealed to induce the proliferation, migration, and metastasis of many tumors, such as breast, lung, and colon cancer. Catecholamines play a significant role in promoting EMT by utilizing signaling pathways like c-Jun[4,5,8]. Anxiety and depression can accelerate the onset and advancement of GC through multifaceted mechanisms (e.g., influencing reactive oxygen species-activated ABL1) and modulating the hypothalamic-pituitary-adrenal axis (e.g., FK506 binding protein 5 gene polymorphisms), thereby inducing disease deterioration and increasing the possibility of recurrence in GC patients[9,10].

Functioning as neurotransmitters, catecholamines can influence tumor characteristics, including phenotypic transformation, apoptosis, and drug resistance. The acquisition of a neuroendocrine phenotype in cancer cells strongly correlates with neoplasm metastasis, drug resistance, advanced cancer stage, and the increased expression of neuroendocrine markers-synaptophysin (SYP), CD44, and chromogranin A[5]. The binding of catecholamines to the beta-2 adrenergic receptor ( $\beta$ 2-AR) upregulates MACC1 expression, leading to neuroendocrine phenotypic transformation, GC invasion, and metastasis. In this process,  $\alpha$ -AR does not exhibit any discernible role. MACC1, an oncogene regulated by c-



Jun, controls c-Met transcriptional levels, enhancing EMT. The activation of the hepatocyte growth factor receptor (c-Met) orchestrates neuroendocrine features in advanced prostate cancer, assuming a parallel role in GC development. Reversal of these effects in mouse models and *in vitro* is achieved through treatment with  $\beta$ 2-AR antagonists or MACC1 silencing [5]. MACC1 also forms a complex with SYP, a marker of neuroendocrine phenotypic characteristics, utilizing the MACC1/SYP signaling pathway in the neuroendocrine phenotypic transformation triggered by catecholamine. Targeting  $\beta$ 2-AR mitigates depression-induced neuroendocrine phenotypic transformation and lung metastasis of GC, providing potential therapeutic targets for enhancing outcomes in GC patients with concurrent depression.  $\beta$ 2-AR stimulation may additionally induce EMT, migration, and invasion by ERK (Extracellular-signal-regulated kinase) phosphorylation [8]. Lu *et al* [8] observed that salbutamol, a  $\beta$ 2-AR agonist, heightened the expression of the mesenchymal marker N-cadherin and reduced the epithelial marker E-cadherin in transplanted tumor tissue, thereby inducing further EMT [8]. They also supported the idea that the  $\beta$ 2-AR agonist isoproterenol promotes EMT of GC cells through the STAT3-CD44 pathway, shedding light on the association of depression with GC [11]. The  $\beta$ 2-AR-HIF-1 $\alpha$ -Snail signaling pathway influences the EMT of GC cells, promoting the invasion and migration of GC [12]. Last but not least, Liu *et al* [4] observed that the activation of  $\beta$ 2-AR increases the expression of PlexinA1, activates JAK-STAT3 signaling, and further promotes EMT in human GC cells. Consequently, chronic stress is intricately linked with the pathogenesis of GC [4].

T helper (Th) cells modulate the stress response, oxidative stress, and neuroinflammation, potentially participating in the pathogenesis of anxiety, depression, and cognitive impairment. Th1 ( $P = 0.017$ ) and Th17 ( $P = 0.049$ ) levels were found to be elevated in patients with depression compared to those without depression [13].

When depression is quantified by the Patient Health Questionnaire-9 score, a positive correlation is observed with serum levels of epinephrine, noradrenaline, MACC1, as well as tumor-node-metastasis (TNM) stage, supporting the association of depression with GC pathogenesis [5].

## DEPRESSION AS A RESULT OF GASTRIC CANCER

GC patients face many psychological challenges, including anxiety, depression, pain, and fatigue, underscoring the need to prioritize their QoL. The prevalence of depression among these patients is often underestimated, despite its effects on prognosis and QoL [2]. Notably, depression has been linked to increased suicidal thoughts, anxiety, distress, and fatigue in cancer patients, with studies emphasizing in the need of increased monitoring for this condition [14]. The impact of depression is particularly significant among GC patients, while they are already at risk for malnutrition, lower body mass index, reduced physical activity, and social isolation, exacerbating their susceptibility to depression [2].

The comorbidities accompanying GC and its treatment introduces additional challenges, as gastrectomy, a commonly employed strategy for curative resection, profoundly affects patients' QoL and mental well-being. Total gastrectomy, in particular, results in substantial postoperative malnutrition, with patients experiencing significant weight loss within the first year of surgery [2].

Several studies highlight the prevalence of depression among GC patients, ranging from 4.0% to 68% with a mean of 37% (95% CI). The variability in prevalence underscores the need for comprehensive and region-specific approaches to address this psychological aspect of the disease. A study by Kouhestani *et al* [2] in 2022, drawing data from the National Health Service Sample Cohort, revealed a higher risk of new-onset depression in GC patients, particularly in females aged 60-69 living in metropolitan regions with high income [2].

Kwon *et al* [6] investigate the correlation between depression and stomach cancer further, emphasizing the stressful aspect of cancer diagnosis and therapy, which causes anxiety and depression in a considerable proportion of patients [6].

Patients with recurrent stomach cancer had greater levels of anxiety and sadness than newly diagnosed patients and healthy controls. Age above 60 years, diabetes, TNM stage at diagnosis, shorter duration to recurrence, and distant metastases at recurrence were all risk factors for anxiety. Age above 60 years, diabetes, tumor site upon diagnosis, and shorter time to recurrence were all risk factors for depression [6].

Liu investigates factors associated with anxiety and depression in GC patients, revealing that coping style, type D personality, and neutrophil-to-lymphocyte ratio contribute to preoperative anxiety and depression. Additionally, genetic factors, including polymorphisms in genes related to apoptosis, may play a role in susceptibility to GC and associated psychological distress [3,15].

However, Lou *et al* [15] observed that polymorphisms of BNIP3 and DAPK1 were associated with a protective effect against GC. These two genes are shown to also have a protective effect against depression [15].

## CONCLUSION

Considering the global significance of GC as the fifth most frequently diagnosed cancer, efforts to understand and address the psychological impact of the disease, particularly depression and anxiety, are essential. Efforts should involve a multidisciplinary approach, considering both the physical and mental well-being of patients to improve overall outcomes and QoL.

## FOOTNOTES

**Author contributions:** Christodoulidis G, Konstantinos-Eleftherios K and Marina-Nektaria K contributed to this paper; Christodoulidis G designed the overall concept and outline of the manuscript; Christodoulidis G, Konstantinos-Eleftherios K and Marina-Nektaria K contributed to the discussion and design of the manuscript; Christodoulidis G, Konstantinos-Eleftherios K and Marina-Nektaria K contributed to the writing, editing the manuscript, and review of literature.

**Conflict-of-interest statement:** All the authors report no relevant conflicts of interest for this article.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** Greece

**ORCID number:** Grigorios Christodoulidis 0000-0003-3413-0666; Koumarelas Konstantinos-Eleftherios 0000-0002-5614-4770; Kouliou Marina-Nektaria 0000-0002-2055-2297.

**S-Editor:** Li L

**L-Editor:** A

**P-Editor:** Zhao YQ

## REFERENCES

- Liu JY, Zheng JQ, Yin CL, Tang WP, Zhang JN. Hotspots and frontiers of the relationship between gastric cancer and depression: A bibliometric study. *World J Gastroenterol* 2023; **29**: 6076-6088 [PMID: 38130743 DOI: 10.3748/wjg.v29.i46.6076]
- Kouhestani M, Ahmadi Gharaei H, Fararouei M, Hosienpour Ghahremanloo H, Ghaiasvand R, Dianatinasab M. Global and regional geographical prevalence of depression in gastric cancer: a systematic review and meta-analysis. *BMJ Support Palliat Care* 2022; **12**: e526-e536 [PMID: 32434923 DOI: 10.1136/bmjspcare-2019-002050]
- Liu P, Wang Z. Postoperative anxiety and depression in surgical gastric cancer patients: their longitudinal change, risk factors, and correlation with survival. *Medicine (Baltimore)* 2022; **101** [PMID: 35356898 DOI: 10.1097/MD.00000000000028765]
- Liu Y, Hao Y, Zhao H, Zhang Y, Cheng D, Zhao L, Peng Y, Lu Y, Li Y. PlexinA1 activation induced by  $\beta$ 2-AR promotes epithelial-mesenchymal transition through JAK-STAT3 signaling in human gastric cancer cells. *J Cancer* 2022; **13**: 2258-2270 [PMID: 35517411 DOI: 10.7150/jca.70000]
- Pan C, Wu J, Zheng S, Sun H, Fang Y, Huang Z, Shi M, Liang L, Bin J, Liao Y, Chen J, Liao W. Depression accelerates gastric cancer invasion and metastasis by inducing a neuroendocrine phenotype via the catecholamine/ $\beta$ (2) -AR/MACC1 axis. *Cancer Commun (Lond)* 2021; **41**: 1049-1070 [PMID: 34288568 DOI: 10.1002/cac2.12198]
- Kwon S, Kim J, Kim T, Jeong W, Park EC. Association between gastric cancer and the risk of depression among South Korean adults. *BMC Psychiatry* 2022; **22**: 207 [PMID: 35313847 DOI: 10.1186/s12888-022-03847-w]
- Wang HQ, Wang ZZ, Chen NH. The receptor hypothesis and the pathogenesis of depression: Genetic bases and biological correlates. *Pharmacol Res* 2021; **167**: 105542 [PMID: 33711432 DOI: 10.1016/j.phrs.2021.105542]
- Lu Y, Zhang Y, Zhao H, Li Q, Liu Y, Zuo Y, Xu Q, Zuo H, Li Y. Chronic stress model simulated by salbutamol promotes tumorigenesis of gastric cancer cells through  $\beta$ 2-AR/ERK/EMT pathway. *J Cancer* 2022; **13**: 401-412 [PMID: 35069890 DOI: 10.7150/jca.65403]
- Huang T, Zhou F, Wang-Johanning F, Nan K, Wei Y. Depression accelerates the development of gastric cancer through reactive oxygen species-activated ABL1 (Review). *Oncol Rep* 2016; **36**: 2435-2443 [PMID: 27666407 DOI: 10.3892/or.2016.5127]
- Kang JI, Chung HC, Jeung HC, Kim SJ, An SK, Namkoong K. FKBP5 polymorphisms as vulnerability to anxiety and depression in patients with advanced gastric cancer: a controlled and prospective study. *Psychoneuroendocrinology* 2012; **37**: 1569-1576 [PMID: 22459275 DOI: 10.1016/j.psyneuen.2012.02.017]
- Lu YJ, Geng ZJ, Sun XY, Li YH, Fu XB, Zhao XY, Wei B. Isoprenaline induces epithelial-mesenchymal transition in gastric cancer cells. *Mol Cell Biochem* 2015; **408**: 1-13 [PMID: 26253173 DOI: 10.1007/s11010-015-2477-0]
- Shan T, Cui X, Li W, Lin W, Li Y, Chen X, Wu T. Novel regulatory program for norepinephrine-induced epithelial-mesenchymal transition in gastric adenocarcinoma cell lines. *Cancer Sci* 2014; **105**: 847-856 [PMID: 24815301 DOI: 10.1111/cas.12438]
- Zhou Y, Yu K. Th1, Th2, and Th17 cells and their corresponding cytokines are associated with anxiety, depression, and cognitive impairment in elderly gastric cancer patients. *Front Surg* 2022; **9**: 996680 [PMID: 36386524 DOI: 10.3389/fsurg.2022.996680]
- Walker J, Hansen CH, Martin P, Symeonides S, Ramessur R, Murray G, Sharpe M. Prevalence, associations, and adequacy of treatment of major depression in patients with cancer: a cross-sectional analysis of routinely collected clinical data. *Lancet Psychiatry* 2014; **1**: 343-350 [PMID: 26360998 DOI: 10.1016/S2215-0366(14)70313-X]
- Lou X, Hu D, Li Z, Teng Y, Lou Q, Huang S, Zou Y, Wang F. Associations of BNIP3 and DAPK1 gene polymorphisms with disease susceptibility, clinicopathologic features, anxiety, and depression in gastric cancer patients. *Int J Clin Exp Pathol* 2021; **14**: 633-645 [PMID: 34093949]





Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [office@baishideng.com](mailto:office@baishideng.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

