

World Journal of *Gastroenterology*

World J Gastroenterol 2024 March 14; 30(10): 1261-1469



EDITORIAL

- 1261 Bridging the gap: Unveiling the crisis of physical inactivity in inflammatory bowel diseases
Stafie R, Singeap AM, Rotaru A, Stanciu C, Trifan A
- 1266 Double role of depression in gastric cancer: As a causative factor and as consequence
Christodoulidis G, Konstantinos-Eleftherios K, Marina-Nektaria K
- 1270 Capsule endoscopy and panendoscopy: A journey to the future of gastrointestinal endoscopy
Rosa B, Cotter J
- 1280 Vonoprazan-amoxicillin dual regimen with *Saccharomyces boulardii* as a rescue therapy for *Helicobacter pylori*: Current perspectives and implications
Dirjayanto VJ, Audrey J, Simadibrata DM
- 1287 Women health and microbiota: Different aspects of well-being
Nannini G, Amedei A
- 1291 Nomograms and prognosis for superficial esophageal squamous cell carcinoma
Lin HT, Abdelbaki A, Krishna SG

REVIEW

- 1295 Overview of the immunological mechanisms in hepatitis B virus reactivation: Implications for disease progression and management strategies
Ma H, Yan QZ, Ma JR, Li DF, Yang JL
- 1313 Optimizing nutrition in hepatic cirrhosis: A comprehensive assessment and care approach
Mendez-Guerrero O, Carranza-Carrasco A, Chi-Cervera LA, Torre A, Navarro-Alvarez N
- 1329 Optimizing prediction models for pancreatic fistula after pancreatectomy: Current status and future perspectives
Yang F, Windsor JA, Fu DL

ORIGINAL ARTICLE

Retrospective Cohort Study

- 1346 Cumulative effects of excess high-normal alanine aminotransferase levels in relation to new-onset metabolic dysfunction-associated fatty liver disease in China
Chen JF, Wu ZQ, Liu HS, Yan S, Wang YX, Xing M, Song XQ, Ding SY
- 1358 Time trends and outcomes of gastrostomy placement in a Swedish national cohort over two decades
Skogar ML, Sundbom M

Retrospective Study

- 1368 Stage at diagnosis of colorectal cancer through diagnostic route: Who should be screened?

Agatsuma N, Utsumi T, Nishikawa Y, Horimatsu T, Seta T, Yamashita Y, Tanaka Y, Inoue T, Nakanishi Y, Shimizu T, Ohno M, Fukushima A, Nakayama T, Seno H

Observational Study

- 1377 Differential diagnosis of Crohn's disease and intestinal tuberculosis based on ATR-FTIR spectroscopy combined with machine learning

Li YP, Lu TY, Huang FR, Zhang WM, Chen ZQ, Guang PW, Deng LY, Yang XH

Prospective Study

- 1393 Establishment and validation of an adherence prediction system for lifestyle interventions in non-alcoholic fatty liver disease

Zeng MH, Shi QY, Xu L, Mi YQ

Basic Study

- 1405 Alkaline sphingomyelinase deficiency impairs intestinal mucosal barrier integrity and reduces antioxidant capacity in dextran sulfate sodium-induced colitis

Tian Y, Li X, Wang X, Pei ST, Pan HX, Cheng YQ, Li YC, Cao WT, Petersen JDD, Zhang P

- 1420 Preliminary exploration of animal models of congenital choledochal cysts

Zhang SH, Zhang YB, Cai DT, Pan T, Chen K, Jin Y, Luo WJ, Huang ZW, Chen QJ, Gao ZG

- 1431 Serotonin receptor 2B induces visceral hyperalgesia in rat model and patients with diarrhea-predominant irritable bowel syndrome

Li ZY, Mao YQ, Hua Q, Sun YH, Wang HY, Ye XG, Hu JX, Wang YJ, Jiang M

META-ANALYSIS

- 1450 Shear-wave elastography to predict hepatocellular carcinoma after hepatitis C virus eradication: A systematic review and meta-analysis

Esposito G, Santini P, Galasso L, Mignini I, Ainora ME, Gasbarrini A, Zocco MA

LETTER TO THE EDITOR

- 1461 Current considerations on intraductal papillary neoplasms of the bile duct and pancreatic duct

Pavlidis ET, Galanis IN, Pavlidis TE

- 1466 Are we ready to use new endoscopic scores for ulcerative colitis?

Quera R, Núñez F P

ABOUT COVER

Editorial Board Member of *World Journal of Gastroenterology*, Toru Mizuguchi, MD, PhD, Professor, Surgeon, Department of Nursing, Division of Surgical Science, Sapporo Medical University Postgraduate School of Health Science, Sapporo, Hokkaido 0608556, Japan. tmizu@sapmed.ac.jp

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (WJG, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now abstracted and indexed in Science Citation Index Expanded (SCIE), MEDLINE, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJG as 4.3; Quartile category: Q2. The WJG's CiteScore for 2021 is 8.3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xiang Li*; Cover Editor: *Jia-Ru Fan*.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski

EXECUTIVE ASSOCIATE EDITORS-IN-CHIEF

Xian-Jun Yu (Pancreatic Oncology), Jian-Gao Fan (Chronic Liver Disease), Hou-Bao Liu (Biliary Tract Disease)

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

March 14, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

PUBLISHING PARTNER

Shanghai Pancreatic Cancer Institute and Pancreatic Cancer Institute, Fudan University
Biliary Tract Disease Institute, Fudan University

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

POLICY OF CO-AUTHORS

<https://www.wjgnet.com/bpg/GerInfo/310>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

PUBLISHING PARTNER's OFFICIAL WEBSITE

<https://www.shca.org.cn>
<https://www.zs-hospital.sh.cn>



Retrospective Cohort Study

Cumulative effects of excess high-normal alanine aminotransferase levels in relation to new-onset metabolic dysfunction-associated fatty liver disease in China

Jing-Feng Chen, Zhuo-Qing Wu, Hao-Shuang Liu, Su Yan, You-Xiang Wang, Miao Xing, Xiao-Qin Song, Su-Ying Ding

Specialty type: Gastroenterology and hepatology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Gorrell MD, Australia

Received: November 7, 2023

Peer-review started: November 7, 2023

First decision: December 27, 2023

Revised: January 12, 2024

Accepted: February 18, 2024

Article in press: February 18, 2024

Published online: March 14, 2024



Jing-Feng Chen, Hao-Shuang Liu, Su Yan, Xiao-Qin Song, Su-Ying Ding, Health Management Center, The First Affiliated Hospital of Zhengzhou University, Zhengzhou 450052, Henan Province, China

Zhuo-Qing Wu, Institute of Systems Engineering, Dalian University of Technology, Dalian 116024, Liaoning Province, China

You-Xiang Wang, College of Public Health, Zhengzhou University, Zhengzhou 450001, Henan Province, China

Miao Xing, School of Basic Medicine and Forensic Medicine, Henan University of Science and Technology, Luoyang 471003, Henan Province, China

Corresponding author: Su-Ying Ding, MD, Director, Doctor, Health Management Center, The First Affiliated Hospital of Zhengzhou University, No. 1 Longhu Middle Ring Road, Jinshui District, Zhengzhou 450052, Henan Province, China. fccdingsy@zzu.edu.cn

Abstract

BACKGROUND

Within the normal range, elevated alanine aminotransferase (ALT) levels are associated with an increased risk of metabolic dysfunction-associated fatty liver disease (MAFLD).

AIM

To investigate the associations between repeated high-normal ALT measurements and the risk of new-onset MAFLD prospectively.

METHODS

A cohort of 3553 participants followed for four consecutive health examinations over 4 years was selected. The incidence rate, cumulative times, and equally and unequally weighted cumulative effects of excess high-normal ALT levels (ehALT) were measured. Cox proportional hazards regression was used to analyse the association between the cumulative effects of ehALT and the risk of new-onset MAFLD.

RESULTS

A total of 83.13% of participants with MAFLD had normal ALT levels. The incidence rate of MAFLD showed a linear increasing trend in the cumulative ehALT group. Compared with those in the low-normal ALT group, the multivariate adjusted hazard ratios of the equally and unequally weighted cumulative effects of ehALT were 1.651 [95% confidence interval (CI): 1.199-2.273] and 1.535 (95%CI: 1.119-2.106) in the third quartile and 1.616 (95%CI: 1.162-2.246) and 1.580 (95%CI: 1.155-2.162) in the fourth quartile, respectively.

CONCLUSION

Most participants with MAFLD had normal ALT levels. Long-term high-normal ALT levels were associated with a cumulative increased risk of new-onset MAFLD.

Key Words: Metabolic dysfunction-associated fatty liver disease; High-normal alanine aminotransferase level; Cumulative effect; Cox proportional hazards regression; Cohort study

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Limited evidence exists regarding the association between persistently elevated high-normal alanine transaminase (ALT) levels and the risk of new-onset metabolic dysfunction-associated fatty liver disease (MAFLD). This cohort study analysed 3553 participants followed for four consecutive health examinations between 2017 and 2020 and measured the cumulative effects of excess high-normal ALT (ehALT). Among the participants, the incidence rate of MAFLD showed a linear increasing trend for the cumulative ehALT group. The hazard ratios of new-onset MAFLD were significantly increased in the third and fourth quartiles of the equally and unequally weighted cumulative effects of ehALT. Among Chinese adults, long-term high-normal ALT levels were related to a cumulative increased risk of new-onset MAFLD.

Citation: Chen JF, Wu ZQ, Liu HS, Yan S, Wang YX, Xing M, Song XQ, Ding SY. Cumulative effects of excess high-normal alanine aminotransferase levels in relation to new-onset metabolic dysfunction-associated fatty liver disease in China. *World J Gastroenterol* 2024; 30(10): 1346-1357

URL: <https://www.wjgnet.com/1007-9327/full/v30/i10/1346.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v30.i10.1346>

INTRODUCTION

Metabolic dysfunction-associated fatty liver disease (MAFLD) is defined as the combination of nonalcoholic fatty liver disease (NAFLD) and metabolic dysfunction and includes overweight/obesity, type 2 diabetes, or other metabolic disorders, as indicated by liver biopsy or imaging examination or even blood biomarker tests suggesting the presence of fatty liver[1,2]. MAFLD has become a growing public health problem, affecting up to a third of the global population, and its burden has grown in parallel with rising rates of type 2 diabetes mellitus and obesity[3,4]. The prevalence of MAFLD is reportedly 25.0% among adults worldwide[5], 29.2% in China[6], and 33.9% in Korea after sex and age standardization [7], with an increasing incidence each year. MAFLD is a multisystemic disease beyond the liver that can increase the risk of heart failure, obstructive sleep apnoea, and malignancy and can result in an increase in cancer-related and cardiovascular disease mortality[8]. Given the high harm and enormous disease burden of MAFLD, a comprehensive analysis of risk factors is essential[9].

Alanine aminotransferase (ALT) has been recognized globally as a reliable indicator reflecting the degree of liver cell damage, such as the damage associated with NAFLD, chronic hepatitis, and cirrhosis[10]. Many studies have suggested that liver damage can occur in the presence of normal ALT levels[11]. Recently, a growing body of evidence has indicated that an ALT level that is within the normal range is an important biomarker for predicting NAFLD; additionally, nonalcoholic steatohepatitis (NASH) or advanced fibrosis is diagnosed in up to 37.5%-59% of patients with NAFLD who have normal ALT levels[12,13]. Our previous work also indicated that an ALT trajectory at a normal level is associated with the risk of new-onset MAFLD based on a cohort study[14]. Thus, we hypothesized that a specific ALT level, particularly a long-term high-normal ALT level, is associated with the risk of new-onset MAFLD.

Some evidence has suggested that the Youden index, a popular summary statistic for receiver-operating characteristic (ROC) curves, provides the optimal cut-off point for a biomarker to distinguish diseased and healthy individuals[15]. In a study of adolescents with obesity, the optimal ALT cut-off points for diagnosing NAFLD were 36 U/L for males and 33 U/L for females[16]. However, limited evidence exists on the determination of optimal ALT cut-off points for diagnosing MAFLD and on the associations between repeated high-normal ALT measurements and both the incidence of MAFLD and risk of new-onset MAFLD. In light of the public health burden of MAFLD in China, we investigated the association between repeated ALT levels that are high-normal and new-onset MAFLD using an ambispective cohort from a health examination population.

MATERIALS AND METHODS

Data sources and recruitment

This retrospective and prospective population-based cohort study was based on data from an ambispective cohort from a health examination population in Henan Province. All eligible participants were interviewed by uniformly trained medical staff to gather information about common chronic diseases and factors influencing health. Participants aged ≥ 18 years who underwent a health examination at the First Affiliated Hospital of Zhengzhou University over a period of 3 consecutive years from January 2017 to December 2019 were retrospectively selected. We identified a total of 7817 participants (4975 male and 2842 female individuals); 4521 had no diagnosis of MAFLD according to three consecutive health examinations and were followed up at their fourth health examination in 2020. During the follow-up period, 738 participants did not participate in the health examination for various reasons or had missing information on some of the studied factors; we also excluded 230 participants with viral hepatitis, alcoholic hepatitis, autoimmune hepatitis, severe cardiovascular and cerebrovascular diseases, or malignant tumours. Finally, 3553 eligible participants (1741 male and 1812 female individuals) were selected from the pool of 7817 participants.

Data collection

The data collected included physical measurements, laboratory test results, abdominal colour Doppler ultrasound results, and diagnostic criteria for MAFLD, as described below: (1) Physical measurements. Participants' body mass index (BMI), systolic blood pressure (SBP), diastolic blood pressure (DBP), and waist circumference (WC) were measured by clinicians using a uniform measurement instrument; (2) Laboratory tests. An automatic biochemical analyser was used to measure fasting plasma glucose (FPG), glycated haemoglobin (HbA1c), total cholesterol, triglyceride (TG), high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein cholesterol (LDL-C), serum uric acid (SUA), and ALT levels[15]; (3) Abdominal colour Doppler ultrasound. Ultrasound was used to determine the presence of diffuse echogenic changes in the liver; and (4) Diagnostic criteria for MAFLD. These findings included diffuse echogenic changes in the liver as revealed by abdominal colour Doppler ultrasonography and were accompanied by at least one of the following conditions: (1) Overweight/obese (BMI $> 23 \text{ kg/m}^2$); (2) Type 2 diabetes; and (3) Metabolic dysfunction, defined as the presence of at least two of the following conditions: (1) WC $\geq 90 \text{ cm}$ for men and WC $\geq 80 \text{ cm}$ for women; (2) Hypertension or use of blood pressure-lowering medication or SBP $\geq 130 \text{ mmHg}$ and/or DBP $\geq 85 \text{ mmHg}$; (3) FPG $\geq 5.6 \text{ mmol/L}$ or 2-h postprandial glucose $\geq 7.8 \text{ mmol/L}$ or HbA1c $\geq 5.7\%$; (4) TG $\geq 1.7 \text{ mmol/L}$ or use of lipid-lowering drugs; and (5) HDL-C $< 1.0 \text{ mmol/L}$ in men and HDL-C $< 1.3 \text{ mmol/L}$ in women or use of lipid-lowering medication[16].

Study design and measurements

In total, 7817 participants with three consecutive health examinations from 2017 to 2019 were analysed to determine the optimal ALT cut-off points for the diagnosis of MAFLD, with ALT within the normal range (0–40 U/L). The participants were divided into three groups: Those with a low-normal ALT (lALT) level, those with a high ALT (hALT) level, and those with an abnormal ALT (aALT) level. A follow-up cohort of 3553 participants who completed their fourth health examination in 2020 was subsequently analysed to calculate the cumulative effects of excess hALT (ehALT) and explore its association with the risk of new-onset MAFLD. In this study, the cumulative effects of ehALT were classified into the following three categories: (1) Cumulative number of ehALT occurrences; (2) Equally weighted cumulative effects of ehALT; and (3) Unequally weighted cumulative effects of ehALT[17]. Finally, a single ehALT occurrence (noncumulative ehALT) was also included as a control for the cumulative effects of ehALT. The study design is shown in Figure 1, and some terms are defined in Table 1.

Statistical analyses

The ROC curve with the maximum value of the Youden index (sensitivity + specificity–1) was used to determine the ALT cut-off points for the diagnosis of MAFLD using ALT in 7817 participants from 2017 to 2019. Normally distributed continuous data are presented as the means (SD); comparisons among groups were performed using one-way analysis of variance (ANOVA) along with pairwise comparisons, the least significant difference test for homogeneous variance, and Dunnett's T3 test for nonhomogeneous variance. Continuous data with a skewed distribution are presented as medians [interquartile ranges (IQRs)], and comparisons were performed using nonparametric tests. Categorical data are described as counts (percentages), and comparisons of rates were performed using Pearson's χ^2 test.

A total of 3553 participants who were followed up in 2020 were used to analyse the association between the cumulative effects of ehALT and new-onset MAFLD *via* a Cox proportional hazards regression model. MAFLD and follow-up time were regarded as dependent variables, and the cumulative number of ehALT occurrences (four groups: 0, 1, 2, 3 occurrences, with 0 occurrences regarded as the reference group), equally weighted cumulative effect, unequally weighted cumulative effect, and single ehALT occurrence [continuous variable, analysed with the per-SD increase after Z score standardization; discrete variable, five groups, a cumulative effect of 0 as the reference group; more than 0 was divided into four quartile groups (Q_1 , Q_2 , Q_3 , Q_4)] were the independent variables. Hazard ratios (HRs) and 95% confidence intervals (CIs) were calculated, and trend tests were also conducted. Furthermore, we used restricted cubic splines with five knots at the 5th, 35th, 65th, and 95th percentiles to flexibly model the associations of the equally or unequally weighted effects of ehALT with new-onset MAFLD and adopted variance analysis to verify whether there was a nonlinear correlation (12). All the data management and statistical analyses were performed using the statistical software R version 4.2.0 (The R Project for Statistical Computing, Vienna, Austria). $P < 0.05$ was considered to indicate statistical significance.

Table 1 Definitions of some specific terms

Term	Definition
lALT group	ALT \leq optimal ALT cut-off points (U/L)
hALT group	Optimal ALT cut-off point < ALT \leq 40 (U/L)
aALT group	ALT > 40 (U/L)
ehALT	ALT-optimal ALT cut-off point, if ehALT < 0, redefine ehALT = 0
Cumulative occurrences of ehALT	Sum of times that ehALT > 0 in 2017–2019, time = {0, 1, 2, 3}
Equally weighted cumulative effects of ehALT	Sum of ehALT levels with a weight of 1 in 2017–2019, <i>i.e.</i> , ehALT ₂₀₁₇ + ehALT ₂₀₁₈ + ehALT ₂₀₁₉
Unequally weighted cumulative effects of ehALT	Sum of ehALT levels with an increasing weight in 2017–2019, <i>i.e.</i> , 1 \times ehALT ₂₀₁₇ + 2 \times ehALT ₂₀₁₈ + 3 \times ehALT ₂₀₁₉
Single ehALT occurrence	ehALT ₂₀₁₉ along with ehALT ₂₀₁₇ = 0 and ehALT ₂₀₁₈ = 0

ALT: Alanine aminotransferase; lALT: Low-normal alanine aminotransferase; hALT: High-normal alanine aminotransferase; aALT: Abnormal alanine aminotransferase; ehALT: Excess high-normal alanine aminotransferase.




Figure 1 Schematic description of the study design. 7817 participants with three consecutive health examinations from 2017 to 2019 were analysed to determine the optimal alanine aminotransferase (ALT) cut-off points for the diagnosis of metabolic dysfunction-associated fatty liver disease (MAFLD). A follow-up cohort of 3553 participants who completed their fourth health examination in 2020 was subsequently analysed to calculate the cumulative effects of excess high-normal ALT and explore its association with the risk of new-onset MAFLD. ALT: Alanine aminotransferase; MAFLD: Metabolic dysfunction-associated fatty liver disease; lALT: Low-normal alanine aminotransferase; hALT: High-normal alanine aminotransferase; aALT: Abnormal alanine aminotransferase; ehALT: Excess high-normal alanine aminotransferase.

RESULTS

Prevalence of MAFLD in participants

The results of 7817 participants at baseline in 2017 showed that the prevalence of MAFLD was 34.27% (2679/7817), and 83.13% (36.36% + 46.77%) of participants with MAFLD had normal ALT levels (≤ 40 U/L) (Figure 2A). Analysis of differences based on nonparametric tests indicated that ALT levels were significantly greater in MAFLD patients than in healthy individuals ($P < 0.001$), with median (IQR) ALT levels of 24 (18, 35) U/L and 17 (13, 23) U/L (Figure 2B).




Figure 2 The prevalence of metabolic dysfunction-associated fatty liver disease in a large-scale, longitudinal population-based cohort. A: Distribution of alanine aminotransferase (ALT) levels in participants with metabolic dysfunction-associated fatty liver disease (MAFLD); B: Differential analysis of ALT levels between healthy participants and those with MAFLD. The prevalence of MAFLD was 34.27%, and 83.13% of participants with MAFLD had normal ALT levels. ALT levels were significantly greater in MAFLD patients than in healthy individuals. ALT: Alanine aminotransferase; MAFLD: Metabolic dysfunction-associated fatty liver disease.

The optimal ALT cut-off points for the diagnosis of MAFLD

The ROC curve was obtained for the annual and 3-year health examination data of the 7817 participants from 2017 to 2019. The optimal ALT cut-off points and corresponding sensitivity and specificity for diagnosing MAFLD were determined according to the maximum Youden index. The results showed that the optimal ALT cut-off points were 18.5 U/L in 2017, 18.5 U/L in 2018, 17.5 U/L in 2019, and 18.5 U/L in 2017-2019, as shown in Figure 3. Therefore, the optimal ALT cut-off point was 18.5 U/L based on ROC curve and Youden index, and the cut-off point for hALT was 18.6-40 U/L.

Baseline characteristics of follow-up participants

According to the definition of hALT (18.6-40 U/L), 3553 participants who were eligible for follow-up were included in the baseline analysis in 2017; their mean age was 48.39 (15.13) years, and 49% were male. The baseline characteristics of the 3553 participants according to hALT level are shown in Table 2. Compared with those in the IALT group, participants in the hALT and aALT groups were significantly more likely to have a higher DBP, BMI, SUA, and TG and lower HDL-C levels ($P < 0.001$). Compared with those in the hALT group, participants in the aALT group were more likely to be younger and to have lower FPG and HbA1c values.

Incidence rate of MAFLD under the cumulative effects of ehALT

Taking the date of the participants' health examinations in 2019 as the starting point of follow-up and the occurrence of MAFLD (no = 0, yes = 1) in the health examinations of participants in 2020 as the outcome, we calculated the incidence rate of MAFLD under the cumulative effects of ehALT in different groups and analysed the linear associations, as shown in Figure 4. For the cumulative number of ehALT occurrences of 0, 1, 2, and 3 times before the end of follow-up (by 2020), the incidence rates of MAFLD were 5.81%, 9.64%, 13.67%, and 18.36%, respectively. For the equally weighted cumulative effects of ehALT, the incidence rates of MAFLD in group 0 (reference), Q_1 , Q_2 , Q_3 , and Q_4 were 5.81%, 8.47%, 11.11%, 14.91%, and 17.62%, respectively. For the unequally weighted cumulative effects of ehALT, the incidence rates of MAFLD in group 0 (reference), Q_1 , Q_2 , Q_3 , and Q_4 were 5.81%, 9.85%, 10.67%, 13.47%, and 17.84%, respectively. Furthermore, $P < 0.001$ for both the χ^2 test and linear association test of the abovementioned cumulative effects, indicating that the incidence rates of MAFLD in the different groups were significantly different, with a linear increasing trend. For a single ehALT occurrence (control group), the incidence rates of MAFLD in group 0 (reference), Q_1 , Q_2 , Q_3 , and Q_4 were 5.81%, 4.11%, 3.92%, 7.27%, and 9.62%, respectively. The χ^2 test and linear association test showed $P = 0.682$ and $P = 0.434$, respectively, indicating that no significant difference existed in the incidence rate of MAFLD among the five groups and that no linear increasing trend was shown.

Cox regression analysis of the cumulative effects of ehALT and risk of new-onset MAFLD

Taking the cumulative effects of ehALT in different groups and confounders such as age, sex, and WC as independent variables and new-onset MAFLD (no = 0, yes = 1) in the health examinations of participants in 2020 as the outcome, we used a Cox proportional hazards regression model to analyse the association between the cumulative effects of ehALT in different groups and the risk of new-onset MAFLD. The univariate Cox proportional hazards regression model revealed that sex, WC, SBP, DBP, BMI, SUA, TG, LDL-C, FPG, and HbA1c were risk factors for new-onset MAFLD, with an HR > 1 ($P < 0.05$). HDL-C was a protective factor against new-onset MAFLD, with an HR < 1 ($P < 0.05$).

The results of a multivariate Cox proportional hazards regression model showed the following: (1) Cumulative number of ehALT occurrences: After adjustment for relevant confounding factors, the risk of new-onset MAFLD in the group of patients with 2 and 3 cumulative episodes of ehALT was 1.443 (95%CI: 1.050-1.982) and 1.551 (95%CI: 1.135-2.119), respectively, higher than that in the group with 0.82% ehALT. Furthermore, the trend test indicated that the risk of new-

Table 2 Baseline characteristics of 3553 participants on follow-up

Variables	Total (3553)	IALT group (2409)	hALT group (1046)	aALT group (98)	$F/\chi^2/H$	P value
Sex, n (%)					264.16	< 0.001
Male	1741 (49.00)	955 (39.64)	712 (68.07)	74 (75.51)		
Female	1812 (51.00)	1454 (60.36)	334 (31.93) ^a	24 (24.49) ^a		
Age (yr), mean (SD)	48.39 (15.13)	47.82 (15.39)	50.17 (14.54) ^a	43.51 (12.59) ^{a,b}	14.11	< 0.001
WC (cm), mean (SD)	85.81 (1.81)	85.79 (1.67)	85.85 (1.94)	85.85 (3.11)	0.44	0.643
SBP (mmHg), mean (SD)	120.33 (16.66)	119.18 (16.64)	122.93 (16.70) ^a	121.08 (13.41)	18.77	< 0.001
DBP (mmHg), mean (SD)	72.68 (10.27)	71.70 (10.03)	74.81 (10.60) ^a	74.07 (8.92) ^a	35.09	< 0.001
BMI (kg/m ²), mean (SD)	22.80 (2.41)	22.46 (2.39)	23.46 (2.32) ^a	23.94 (2.21) ^a	76.76	< 0.001
SUA (μmol/L), mean (SD)	290.53 (72.07)	280.17 (69.87)	311.33 (71.51) ^a	323.01 (74.16) ^a	82.00	< 0.001
TC (mmol/L), mean (SD)	4.53 (0.84)	4.51 (0.82)	4.58 (0.88) ^a	4.55 (0.84)	2.72	0.066
TG (mmol/L), median (IQR)	0.97 (0.72, 1.29)	0.93 (0.70, 1.26)	1.06 (0.79, 1.40) ^a	1.09 (0.82, 1.52) ^a	79.73	< 0.001
HDL-C (mmol/L), mean (SD)	1.47 (0.34)	1.50 (0.34)	1.42 (0.34) ^a	1.39 (0.39) ^a	18.51	< 0.001
LDL-C (mmol/L), mean (SD)	2.77 (0.75)	2.74 (0.73)	2.85 (0.78) ^a	2.83 (0.72)	7.76	< 0.001
FPG (mmol/L), median (IQR)	4.96 (4.68, 5.25)	4.95 (4.67, 5.21)	5.00 (4.71, 5.30) ^a	4.94 (4.65, 5.27) ^b	10.65	0.005
HbA1c (%), median (IQR)	5.70 (5.50, 5.70)	5.70 (5.50, 5.70)	5.70 (5.40, 5.70)	5.70 (5.40, 5.70) ^{a,b}	10.69	0.005

^a $P < 0.05$, compared with the low-normal alanine aminotransferase level group.^b $P < 0.05$, compared with the high-normal alanine aminotransferase level group.

IALT: Low-normal alanine aminotransferase; hALT: High-normal alanine aminotransferase; aALT: Abnormal alanine aminotransferase; WC: Waist circumference; SBP: Systolic blood pressure; DBP: Diastolic blood pressure; BMI: Body mass index; TC: Total cholesterol; FPG: Fasting plasma glucose; HbA1c: Glycated haemoglobin; TG: Triglycerides; HDL-C: High-density lipoprotein cholesterol; LDL-C: Low-density lipoprotein cholesterol; SUA: Serum uric acid; IQR: Interquartile range.

onset MAFLD showed an increasing trend; (2) Equal and unequally weighted cumulative effects of ehALT: After adjustment for the relevant confounding factors, the risk of new-onset MAFLD increased by 8.8% (95%CI: 0.3%-17.9%) and 9.8% (95%CI: 1.7%-18.5%), respectively, per SD increase in the cumulative effect. For the five grouping variables, compared with those in group 0 (reference), the HRs of new-onset MAFLD in the Q_3 and Q_4 groups for the equally weighted cumulative effects of ehALT were 1.651 (95%CI: 1.199-2.273) and 1.535 (95%CI: 1.119-2.106), respectively. For the unequally weighted cumulative effects of ehALT, the HRs of new-onset MAFLD in the Q_3 and Q_4 groups were 1.616 (95%CI: 1.162-2.246) and 1.580 (95%CI: 1.155-2.162), respectively; Q_1 and Q_2 were not significantly different from those in the reference group. Additionally, the trend test indicated that the risk of new-onset MAFLD showed an increasing trend for all cumulative occurrences of ehALT; and (3) Single ehALT (control group): Compared with those of the reference group, the univariate and multivariate models did not differ significantly for the continuous or categorical variable of a single ehALT occurrence. Additionally, the trend test showed that the risk of new-onset MAFLD did not increase (Table 3).

Figure 5 shows the dose-response relationship between the cumulative effects of ehALT and the risk of new-onset MAFLD using restricted cubic splines. After adjustment for sex, WC, SBP, DBP, BMI, SUA, TG, HDL-C, LDL-C, FPG, and HbA1c, the equally and unequally weighted cumulative effects of ehALT had a positive nonlinear relationship with the risk of new-onset MAFLD in approximately 95% of the enrolled participants (*i.e.*, total association $P = 0.005$ and nonlinearity $P = 0.011$; total association $P = 0.006$ and nonlinearity $P = 0.029$). Specifically, for the equally weighted cumulative effects of ehALT (Figure 5A), the HR increased rapidly to 1.5 with a cumulative effect of 10 U/L and then maintained a steady value thereafter, to a cumulative effect of 38.5 U/L. For the unequally weighted cumulative effects of ehALT (Figure 5B), the HR increased rapidly to 1.5 with a cumulative effect of 20 U/L and then maintained a steady value thereafter, with a cumulative effect of 82.0 U/L.

DISCUSSION

In this study, a large-scale, longitudinal population-based cohort of 7817 participants in China had a prevalence of MAFLD of 34.27%, and 83.13% of participants with MAFLD had normal ALT levels. The optimal ALT cut-off point for

Table 3 Risks of new-onset metabolic dysfunction-associated fatty liver disease stratified by different categories of cumulative effects of excess high-normal alanine aminotransferase level

Categories	Univariate, HR (95%CI)	P value	Sex-, WC-, SBP-, DBP- and BMI-adjusted ¹ , HR (95%CI)	P value	Multivariate-adjusted ² , HR (95%CI)	P value
Cumulative occurrences of ehALT (<i>n</i> = 3553)						
0 (101/1738)	1.000		1.000		1.000	
1 (82/851)	1.672 (1.249-2.237)	0.001	1.312 (0.975-1.765)	0.073	1.261 (0.935-1.699)	0.128
2 (70/512)	2.325 (1.714-3.154)	< 0.001	1.495 (1.091-2.049)	0.012	1.443 (1.050-1.982)	0.024
3 (83/452)	2.963 (2.215-3.962)	< 0.001	1.559 (1.142-2.128)	0.005	1.551 (1.135-2.119)	0.006
P value for trend ³	< 0.001		0.005		0.005	
Equally weighted cumulative effects of ehALT (<i>n</i> = 3553)						
Increase per SD ⁴	1.227 (1.149-1.310)	< 0.001	1.090 (1.006-1.182)	0.036	1.088 (1.003-1.179)	0.041
0 (reference)	1.000		1.000		1.000	
Q ₁ (0.01-3.00 U/L)	1.541 (1.069-2.225)	0.020	1.139 (0.785-1.652)	0.493	1.083 (0.744-1.575)	0.678
Q ₂ (3.01-8.50 U/L)	1.770 (1.261-2.485)	0.001	1.329 (0.941-1.876)	0.106	1.297 (0.916-1.837)	0.916
Q ₃ (8.51-20.50 U/L)	2.620 (1.926-3.564)	< 0.001	1.671 (1.214-2.300)	0.002	1.651 (1.199-2.273)	0.002
Q ₄ (≥ 20.51 U/L)	2.852 (2.120-3.838)	< 0.001	1.578 (1.151-2.162)	0.005	1.535 (1.119-2.106)	0.008
P value for trend ³	< 0.001		0.005		0.007	
Unequally weighted cumulative effects of ehALT (<i>n</i> = 3553)						
Increase per SD ⁴	1.231 (1.154-1.312)	< 0.001	1.102 (1.020-1.191)	0.014	1.098 (1.017-1.185)	0.016
0 (reference)	1.000		1.000		1.000	
Q ₁ (0.01-5.00 U/L)	1.709 (1.209-2.416)	0.002	1.190 (0.835-1.697)	0.336	1.114 (0.779-1.591)	0.555
Q ₂ (5.01-15.00 U/L)	1.749 (1.234-2.478)	0.002	1.283 (0.900-1.830)	0.168	1.278 (0.895-1.826)	0.177
Q ₃ (15.01-40.50 U/L)	2.319 (1.688-3.187)	< 0.001	1.636 (1.178-2.273)	0.003	1.616 (1.162-2.246)	0.004
Q ₄ (≥ 40.51 U/L)	2.996 (2.236-4.015)	< 0.001	1.626 (1.191-2.220)	0.002	1.580 (1.155-2.162)	0.004
P value for trend ³	< 0.001		0.003		0.004	
Single ehALT occurrence (control group, <i>n</i> = 1969)						
Increase per SD ⁴	1.055 (0.866-1.286)	0.594	1.010 (0.819-1.245)	0.929	1.012 (0.811-1.262)	0.917
0 (reference)	1.000		1.000		1.000	
Q ₁ (0.01-1.50 U/L)	0.688 (0.218-2.171)	0.524	0.497 (0.150-1.528)	0.214	0.557 (0.174-1.783)	0.324
Q ₂ (1.51-3.50 U/L)	0.612 (0.151-2.485)	0.492	0.462 (0.113-1.886)	0.282	0.352 (0.085-1.462)	0.151
Q ₃ (3.51-8.50 U/L)	1.250 (0.460-3.397)	0.662	1.326 (0.486-3.620)	0.582	1.043 (0.374-2.908)	0.936
Q ₄ (≥ 8.51 U/L)	1.990 (0.808-4.898)	0.134	1.688 (0.675-4.221)	0.263	1.828 (0.772-4.593)	0.199
P value for trend ³	0.142		0.259		0.264	

¹Adjusted for sex, waist circumference, systolic blood pressure, diastolic blood pressure, and body mass index.²Adjusted for sex, waist circumference, systolic blood pressure, diastolic blood pressure, body mass index, serum uric acid, triglycerides, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, fasting plasma glucose, and glycated haemoglobin.³For the trend test, Cox proportional hazards regression models were used with group medians in each group instead of grouping variables (*e.g.*, 0, Q₁, Q₂, Q₃, and Q₄). The cumulative excess high-normal alanine aminotransferase (ehALT) level concentrations were 12 U/L, 15 U/L, 20 U/L, and 26 U/L in the four groups, and the equally weighted cumulative effects of ehALT were 0, 1.5 U/L, 5.5 U/L, 13 U/L, and 33.5 U/L in the five groups. The unequally weighted cumulative effects of ehALT were 0, 2.5 U/L, 10.5 U/L, 24.5 U/L, and 69 U/L in the five groups, and the single ehALT occurrence (control group) were 0, 0.5 U/L, 2.5 U/L, 5.5 U/L, and 17.5 U/L in the five groups.⁴The SD of the equally weighted cumulative effects of excess high-normal alanine aminotransferase (ehALT) level was 15.93, the SD of the unequally weighted cumulative effects of ehALT was 32.73, and the SD of a single ehALT occurrence (control group, only 2019 ALT > 18.5 U/L) was 3.76.

WC: Waist circumference; SBP: Systolic blood pressure; DBP: Diastolic blood pressure; BMI: Body mass index; ehALT: Excess high-normal alanine

aminotransferase; HR: Hazard ratio; CI: Confidence interval.




Figure 3 Optimal alanine aminotransferase cut-off points for the diagnosis of metabolic dysfunction-associated fatty liver disease. The optimal alanine aminotransferase (ALT) cut-off point was 18.5 U/L based on receiver-operating characteristic curve and Youden index, and the cut-off point for high-normal ALT was 18.6-40 U/L. AUC: Area under the curve.




Figure 4 Incidence rate of metabolic dysfunction-associated fatty liver disease under the cumulative effects of excess high-normal alanine aminotransferase levels and a single excess high-normal alanine aminotransferase level occurrence in different groups. The incidence rate of metabolic dysfunction-associated fatty liver disease showed a linear increasing trend in the cumulative excess high-normal alanine aminotransferase (ehALT) levels groups, while a single ehALT occurrence did not show a linear increasing trend in different groups. MAFLD: Metabolic dysfunction-associated fatty liver disease; ehALT: Excess high-normal alanine aminotransferase.




Figure 5 Dose-response relationship between the cumulative effects of excess high-normal alanine aminotransferase levels and the risk of new-onset metabolic dysfunction-associated fatty liver disease. A: Equally weighted cumulative effects of excess high-normal alanine aminotransferase (ehALT) levels; B: Unequally weighted cumulative effects of ehALT. The equally and unequally weighted cumulative effects of ehALT had a positive nonlinear relationship with the risk of new-onset metabolic dysfunction-associated fatty liver disease in approximately 95% of the enrolled participants. Gray histogram: The probability distribution of the population corresponding to the accumulation of ehALT. Orange solid line: The restricted cubic splines [hazard ratios (HRs)]. Black dashed curves: The 95% confidence interval of the restricted cubic splines (HRs). Green dashed line: The reference line of the restricted cubic splines (HR = 1). Blue dashed lines: The dividing lines indicating population proportions of less than 5%, 3%, and 1%, respectively. MAFLD: Metabolic dysfunction-associated fatty liver disease; ehALT: Excess high-normal alanine aminotransferase; HR: Hazard ratio.

the diagnosis of MAFLD was 18.5 U/L. Our findings indicate that MAFLD has become one of the most common chronic liver diseases and is a growing public health problem. According to a systematic review and meta-analysis, NAFLD and normal ALT levels are closely related to diabetes, hypertension, and metabolic syndrome[18]. Additionally, the highest ALT cut-off points among 526641 participants without excessive alcohol consumption or known liver disease were 32 U/L, 37 U/L for men; 31 U/L for women; 39 U/L for overweight people; and 36 U/L for patients with diabetes, all of which were lower than the upper limit for ALT (40 U/L)[19]. According to the Liver-Bible-2020 cohort study, the best ALT cut-off for steatosis detection was 35 U/L in males and 22 U/L in females, and the best cut-off for fibrosis detection was 27 U/L in males[20]. Wahlang *et al*[21] reported that elevated ALT within the normal range was a substitute biomarker of NAFLD. However, there is no scientific evidence for whether the long-term, dynamic, or continuous accumulation of ehALT affects new-onset MAFLD (NAFLD accompanied by metabolic disorders).

The incidence rate of MAFLD was significantly different and showed a linear growth trend with the cumulative effects of ehALT in the different groups, whereas there was no such relationship for a single ehALT occurrence (control group). After adjustment for confounding factors, compared with those in the 1ALT group, the cumulative ehALT level was significantly associated with new-onset MAFLD according to three factors: The cumulative frequency of ehALT and both the equally and unequally weighted cumulative effects of ehALT. Moreover, there was no such relationship for a single ehALT occurrence (control group). This finding suggested that the cumulative effects of ehALT within the long-term normal range will significantly increase the risk of new-onset MAFLD. A prospective study conducted by Gawrieh *et al* [22] revealed that the histological characteristics of NAFLD, advanced fibrosis, and the frequency and severity of NASH gradually increase when ALT levels increase gradually from < 20 U/L to 20-39 U/L, within the normal range. Our previous study showed that, compared with those in the stable low-ALT subgroup (13.10-13.92 U/L for 3 consecutive years), the stable middle-ALT (22.65-24.08 U/L) and stable high-ALT (32.50-39.78 U/L) groups had a significantly increased risk of MAFLD among men and women in the general population[14]. Thus, long-term hALT intake can increase the risk of developing MAFLD and aggravate the severity of NASH and advanced fibrosis. However, in-depth studies are presently lacking regarding the cumulative effects of and dynamic changes in hALT and the risk of new-onset MAFLD.

With lifestyle changes, a growing number of patients with MAFLD (dominated by NAFLD with metabolic disorders) who were physically asymptomatic had fluctuating ALT levels mostly within the normal range, although liver biopsy results revealed marked inflammation or fibrosis in some patients[23]. ALT is an enzyme that exists widely in the cytoplasm of liver cells. Once hepatocyte apoptosis and damage occur, the serum ALT concentration increases significantly. Therefore, ALT is the most sensitive detection indicator reflecting liver function damage and liver inflammation and is an important marker for detecting steatosis, diagnosing NASH, evaluating NASH-related fibrosis stages, and detecting liver cirrhosis[24]. Clinically, most physicians assess the hepatic risk of NAFLD based on changes in ALT levels, which are often overlooked in patients with MAFLD who have long-standing normal ALT levels, leading to aggravation of the degree of hepatic inflammatory response with insulin resistance and multi-hit pathogenesis, and further resulting in NASH, liver fibrosis, and eventually the development of cirrhosis and even hepatocellular carcinoma [25].

This study used a population-based cohort to explore the cumulative effects of ehALT and the risk of new-onset MAFLD. The main strengths of this study were the determination of the optimal ALT cut-off points and the use of different measurement methods for the cumulative effects of ehALT from three perspectives (*i.e.*, cumulative number of ehALT measurements and equally and unequally weighted cumulative effects). In contrast, previous studies have investigated the association between a single ALT measurement or the ALT trajectory and neglected the effects of new-onset MAFLD by considering the quantitative cumulative effects of ALT on long-term dynamic changes *via* a lifespan approach.

This study has several limitations. First, the cohort's follow-up time was short, and the proportion of participants with aALT was low, which led to a decrease in the dose-response relationship and made it difficult to accurately detect a significant difference between the aALT group (> 40 U/L) and the hALT group (18.6–40 U/L) for the cumulative effects on the risk of new-onset MAFLD. Additionally, randomized controlled trials with different lifestyle interventions (including weight loss through diet and physical exercise) will be conducted to explore whether those interventions can improve long-term ALT levels in individuals who are high-normal and ultimately prevent MAFLD.

CONCLUSION

In conclusion, a large-scale population-based study in Henan Province indicated that a high-normal ALT level was 18.6–40 U/L, that a normal ALT level was common in patients with MAFLD, and that a long-term change in the ALT level had cumulative effects on the risk of new-onset MAFLD. We recommend that individuals in this population, especially those in young adulthood, maintain long-term ALT levels within the normal range.

ARTICLE HIGHLIGHTS

Research background

Additional evidence is needed regarding the association between repeated high-normal alanine transaminase (ALT) measurements and metabolic dysfunction-associated fatty liver disease (MAFLD), and only a few cross-sectional studies have shown that ALT trajectory is associated with the risk of MAFLD. In light of the public health burden of MAFLD in China, we investigated the association between persistently elevated high-normal ALT levels and new-onset MAFLD using an ambispective cohort from a health examination population.

Research motivation

MAFLD has become a growing public health problem and affects up to one-third of the global population, with a heavy disease burden. MAFLD can occur in the presence of normal ALT levels, and a trajectory within the normal range can increase the risk of MAFLD. However, the link between repeated high-normal ALT measurements and new-onset MAFLD has not been well studied.

Research objectives

We investigated the optimal ALT cut-off points for diagnosing MAFLD and the association between repeated high-normal ALT measurements and the risk of new-onset MAFLD in a health examination population in China.

Research methods

This study used data from an ambispective cohort of individuals from a health examination population in China. Repeated high-normal ALT measurements were assessed by considering equally and unequally weighted cumulative effects of excess high-normal ALT (ehALT), and participants were categorized into quartile groups. We performed multivariable Cox proportional hazards regression analysis to evaluate the association between cumulative ehALT and the risk of new-onset MAFLD and calculated the hazard ratios (HRs) and 95% confidence intervals.

Research results

A total of 83.13% of participants with MAFLD had normal ALT levels. The HRs of new-onset MAFLD in the group of patients with 2 or 3 cumulative episodes of ehALT (Q_3 and Q_4 for the equally and unequally weighted cumulative effects of ehALT) were greater than those in the group with low-normal ALT levels from 2017 to 2019. Additionally, the dose-response relationship indicated that the equally and unequally weighted cumulative effects of ehALT had positive nonlinear relationships with the risk of new-onset MAFLD.

Research conclusions

A cohort study of the Chinese adult population revealed that persistently elevated high-normal ALT levels were associated with a dose-dependent increase in the risk of new-onset MAFLD in all participants. The identification and management of high-normal ALT levels for several years may play a crucial role in preventing MAFLD.

Research perspectives

Long-term prospective cohort or randomized controlled trials are needed to confirm the relationship between repeated

high-normal ALT measurements and new-onset MAFLD. Future studies should focus on whether a healthy lifestyle can improve ALT levels and prevent MAFLD.

FOOTNOTES

Author contributions: Chen JF and Ding SY designed the study and acquired funding; Chen JF and Wu ZQ were responsible for developing the methodology; Yan S and Wang YX participated in the formal analysis and investigation; Chen JF wrote the original draft; Chen JF, Wu ZQ, Liu HS, Yan S, Wang YX, Xing M, Song XQ, and Ding SY participated in the review and editing.

Supported by National Natural Science Foundation of China, No. 72101236; China Postdoctoral Science Foundation, No. 2022M722900; Collaborative Innovation Project of Zhengzhou City, No. XTCX2023006; and Nursing Team Project of the First Affiliated Hospital of Zhengzhou University, No. HLKY2023005.

Institutional review board statement: The study was reviewed and approved by the Institutional Review Board of the First Affiliated Hospital of Zhengzhou University (Approval No. 2020-KY-381).

Informed consent statement: Signed informed consent was obtained from all participants.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: No additional data are available.

STROBE statement: The authors have read the STROBE Statement-checklist of items, and the manuscript was prepared and revised according to the STROBE Statement-checklist of items.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Jing-Feng Chen 0009-0005-2898-1490; Zhuo-Qing Wu 0000-0003-2415-5202; Hao-Shuang Liu 0000-0002-3108-7001; Su Yan 0009-0007-3959-2989; You-Xiang Wang 0009-0002-6797-876X; Miao Xing 0009-0003-3440-3430; Xiao-Qin Song 0000-0002-1788-3933; Su-Ying Ding 0000-0002-2649-8332.

S-Editor: Wang JJ

L-Editor: A

P-Editor: Yuan YY

REFERENCES

- 1 Lim GEH, Tang A, Ng CH, Chin YH, Lim WH, Tan DJH, Yong JN, Xiao J, Lee CW, Chan M, Chew NW, Xuan Tan EX, Siddiqui MS, Huang D, Nouredin M, Sanyal AJ, Muthiah MD. An Observational Data Meta-analysis on the Differences in Prevalence and Risk Factors Between MAFLD vs NAFLD. *Clin Gastroenterol Hepatol* 2023; **21**: 619-629.e7 [PMID: 34871813 DOI: 10.1016/j.cgh.2021.11.038]
- 2 Eslam M, Sanyal AJ, George J; International Consensus Panel. MAFLD: A Consensus-Driven Proposed Nomenclature for Metabolic Associated Fatty Liver Disease. *Gastroenterology* 2020; **158**: 1999-2014.e1 [PMID: 32044314 DOI: 10.1053/j.gastro.2019.11.312]
- 3 Yang J, Luo S, Li R, Ju J, Zhang Z, Shen J, Sun M, Fan J, Xia M, Zhu W, Liu Y. Sleep Factors in Relation to Metabolic Dysfunction-Associated Fatty Liver Disease in Middle-Aged and Elderly Chinese. *J Clin Endocrinol Metab* 2022; **107**: 2874-2882 [PMID: 35900115 DOI: 10.1210/clinem/dgac428]
- 4 Eslam M, El-Serag HB, Francque S, Sarin SK, Wei L, Bugianesi E, George J. Metabolic (dysfunction)-associated fatty liver disease in individuals of normal weight. *Nat Rev Gastroenterol Hepatol* 2022; **19**: 638-651 [PMID: 35710982 DOI: 10.1038/s41575-022-00635-5]
- 5 Liu J, Ayada I, Zhang X, Wang L, Li Y, Wen T, Ma Z, Bruno MJ, de Knecht RJ, Cao W, Peppelenbosch MP, Ghanbari M, Li Z, Pan Q. Estimating Global Prevalence of Metabolic Dysfunction-Associated Fatty Liver Disease in Overweight or Obese Adults. *Clin Gastroenterol Hepatol* 2022; **20**: e573-e582 [PMID: 33618024 DOI: 10.1016/j.cgh.2021.02.030]
- 6 Huang YP, Zhang S, Zhang M, Wang Y, Wang WH, Li J, Li C, Lin JN. Gender-specific prevalence of metabolic-associated fatty liver disease among government employees in Tianjin, China: a cross-sectional study. *BMJ Open* 2021; **11**: e056260 [PMID: 34911725 DOI: 10.1136/bmjopen-2021-056260]
- 7 Kim M, Yoon EL, Cho S, Lee CM, Kang BK, Park H, Jun DW, Nah EH. Prevalence of advanced hepatic fibrosis and comorbidity in metabolic dysfunction-associated fatty liver disease in Korea. *Liver Int* 2022; **42**: 1536-1544 [PMID: 35338555 DOI: 10.1111/liv.15259]
- 8 Quek J, Ng CH, Tang ASP, Chew N, Chan M, Khoo CM, Wei CP, Chin YH, Tay P, Lim G, Tan DJH, Lim WH, Chan KE, Teng M, Tan E, Tamaki N, Huang DQ, Siddiqui MS, Young DY, Nouredin M, Muthiah MD. Metabolic Associated Fatty Liver Disease Increases the Risk of Systemic Complications and Mortality. A Meta-Analysis and Systematic Review of 12 620 736 Individuals. *Endocr Pract* 2022; **28**: 667-672 [PMID: 35364328 DOI: 10.1016/j.eprac.2022.03.016]
- 9 Lee GB, Huh Y, Lee SH, Han B, Kim YH, Kim DH, Kim SM, Choi YS, Cho KH, Nam GE. Association of low muscle strength with

- metabolic dysfunction-associated fatty liver disease: A nationwide study. *World J Gastroenterol* 2023; **29**: 5962-5973 [PMID: [38131000](#) DOI: [10.3748/wjg.v29.i45.5962](#)]
- 10 **Wang M**, Wang M, Zhang R, Zhang L, Ding Y, Tang Z, Fan H, Wang H, Zhang W, Chen Y, Wang J. A combined association of serum uric acid, alanine aminotransferase and waist circumference with non-alcoholic fatty liver disease: a community-based study. *PeerJ* 2022; **10**: e13022 [PMID: [35265397](#) DOI: [10.7717/peerj.13022](#)]
 - 11 **Kang Y**, Park S, Kim S, Koh H. Normal serum alanine aminotransferase and non-alcoholic fatty liver disease among Korean adolescents: a cross-sectional study using data from KNHANES 2010-2015. *BMC Pediatr* 2018; **18**: 215 [PMID: [29976192](#) DOI: [10.1186/s12887-018-1202-z](#)]
 - 12 **Seko Y**, Sumida Y, Tanaka S, Mori K, Taketani H, Ishiba H, Hara T, Okajima A, Yamaguchi K, Moriguchi M, Mitsuyoshi H, Kanemasa K, Yasui K, Minami M, Imai S, Itoh Y. Serum alanine aminotransferase predicts the histological course of non-alcoholic steatohepatitis in Japanese patients. *Hepatol Res* 2015; **45**: E53-E61 [PMID: [25429984](#) DOI: [10.1111/hepr.12456](#)]
 - 13 **Wu Y**, Yang X, Morris HL, Gurka MJ, Shenkman EA, Cusi K, Bril F, Donahoo WT. Noninvasive Diagnosis of Nonalcoholic Steatohepatitis and Advanced Liver Fibrosis Using Machine Learning Methods: Comparative Study With Existing Quantitative Risk Scores. *JMIR Med Inform* 2022; **10**: e36997 [PMID: [35666557](#) DOI: [10.2196/36997](#)]
 - 14 **Chen JF**, Qin Q, Wu ZQ, Yan S, Song XQ, Ding SY. A cohort study on the correlation between alanine aminotransferase trajectories and new-onset metabolic fatty liver disease. *Zhonghua Liu Xing Bing Xue Za Zhi* 2022; **43**: 234-240 [PMID: [35184490](#) DOI: [10.3760/cma.j.cn112338-20210809-00621](#)]
 - 15 **Zhang YB**, Yang G, Bu Y, Lei P, Zhang W, Zhang DY. Development of a machine learning-based model for predicting risk of early postoperative recurrence of hepatocellular carcinoma. *World J Gastroenterol* 2023; **29**: 5804-5817 [PMID: [38074914](#) DOI: [10.3748/wjg.v29.i45.5804](#)]
 - 16 **Rong Y**, Chun-Yan N, Hong-Xin Z, Lu Y, Wen W, Yu T. Association of Adolescent Obesity with Nonalcoholic Fatty Liver Disease and Related Risk Factors in Xi 'an, China. *Ann Hepatol* 2018; **17**: 85-91 [PMID: [29311392](#) DOI: [10.5604/01.3001.0010.7538](#)]
 - 17 **Li X**, Jansen L, Chang-Claude J, Hoffmeister M, Brenner H. Risk of Colorectal Cancer Associated With Lifetime Excess Weight. *JAMA Oncol* 2022; **8**: 730-737 [PMID: [35297997](#) DOI: [10.1001/jamaoncol.2022.0064](#)]
 - 18 **Ma X**, Liu S, Zhang J, Dong M, Wang Y, Wang M, Xin Y. Proportion of NAFLD patients with normal ALT value in overall NAFLD patients: a systematic review and meta-analysis. *BMC Gastroenterol* 2020; **20**: 10 [PMID: [31937252](#) DOI: [10.1186/s12876-020-1165-z](#)]
 - 19 **Huang DQ**, Yeo YH, Tan E, Takahashi H, Yasuda S, Saruwatari J, Tanaka K, Oniki K, Kam LY, Muthiah MD, Hyogo H, Ono M, Barnett SD, Li J, Zou B, Fung J, Lee TY, Wong VW, Yuen MF, Dan YY, Lim SG, Cheung R, Toyoda H, Eguchi Y, Nguyen MH. ALT Levels for Asians With Metabolic Diseases: A Meta-analysis of 86 Studies With Individual Patient Data Validation. *Hepatol Commun* 2020; **4**: 1624-1636 [PMID: [33163833](#) DOI: [10.1002/hep4.1593](#)]
 - 20 **Valenti L**, Pelusi S, Bianco C, Ceriotti F, Berzuini A, Iogna Prat L, Trotti R, Malvestiti F, D'Ambrosio R, Lampertico P, Colli A, Colombo M, Tsochatzis EA, Fraquelli M, Prati D. Definition of Healthy Ranges for Alanine Aminotransferase Levels: A 2021 Update. *Hepatol Commun* 2021; **5**: 1824-1832 [PMID: [34520121](#) DOI: [10.1002/hep4.1794](#)]
 - 21 **Wahlang B**, Appana S, Falkner KC, McClain CJ, Brock G, Cave MC. Insecticide and metal exposures are associated with a surrogate biomarker for non-alcoholic fatty liver disease in the National Health and Nutrition Examination Survey 2003-2004. *Environ Sci Pollut Res Int* 2020; **27**: 6476-6487 [PMID: [31873887](#) DOI: [10.1007/s11356-019-07066-x](#)]
 - 22 **Gawrieh S**, Wilson LA, Cummings OW, Clark JM, Loomba R, Hameed B, Abdelmalek MF, Dasarathy S, Neuschwander-Tetri BA, Kowdley K, Kleiner D, Doo E, Tonascia J, Sanyal A, Chalasani N; NASH Clinical Research Network. Histologic Findings of Advanced Fibrosis and Cirrhosis in Patients With Nonalcoholic Fatty Liver Disease Who Have Normal Aminotransferase Levels. *Am J Gastroenterol* 2019; **114**: 1626-1635 [PMID: [31517638](#) DOI: [10.14309/ajg.0000000000000388](#)]
 - 23 **Sun DQ**, Zheng KI, Xu G, Ma HL, Zhang HY, Pan XY, Zhu PW, Wang XD, Targher G, Byrne CD, Chen YP, Yuan WJ, Zheng MH. PNPLA3 rs738409 is associated with renal glomerular and tubular injury in NAFLD patients with persistently normal ALT levels. *Liver Int* 2020; **40**: 107-119 [PMID: [31519069](#) DOI: [10.1111/liv.14251](#)]
 - 24 **Kim WR**, Flamm SL, Di Bisceglie AM, Bodenheimer HC; Public Policy Committee of the American Association for the Study of Liver Disease. Serum activity of alanine aminotransferase (ALT) as an indicator of health and disease. *Hepatology* 2008; **47**: 1363-1370 [PMID: [18366115](#) DOI: [10.1002/hep.22109](#)]
 - 25 **Shim JJ**, Kim JW, Oh CH, Lee YR, Lee JS, Park SY, Kim BH, Oh IH. Serum alanine aminotransferase level and liver-related mortality in patients with chronic hepatitis B: A large national cohort study. *Liver Int* 2018; **38**: 1751-1759 [PMID: [29377574](#) DOI: [10.1111/liv.13705](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

