

World Journal of *Gastroenterology*

World J Gastroenterol 2024 March 21; 30(11): 1470-1643



EDITORIAL

- 1470 MicroRNAs in hepatocellular carcinoma treatment: Charting the path forward
Lin HT, Castaneda AFA, Krishna SG, Mumtaz K
- 1475 Innovative pathways allow safe discharge of mild acute pancreatitis from the emergency room
Kothari DJ, Sheth SG
- 1480 Current remarks and future directions on the interactions between metabolic dysfunction-associated fatty liver disease and COVID-19
Brilakis L, Theofilogiannakou E, Lykoudis PM
- 1488 Routine utilization of machine perfusion in liver transplantation: Ready for prime time?
Parente A, Sun K, Dutkowski P, Shapiro AJ, Schlegel A
- 1494 Advancements in Barrett's esophagus detection: The role of artificial intelligence and its implications
Massironi S

REVIEW

- 1497 MicroRNAs: A novel signature in the metastasis of esophageal squamous cell carcinoma
Wei QY, Jin F, Wang ZY, Li BJ, Cao WB, Sun ZY, Mo SJ

MINIREVIEWS

- 1524 Morphological and biochemical characteristics associated with autophagy in gastrointestinal diseases
Chang YF, Li JJ, Liu T, Wei CQ, Ma LW, Nikolenko VN, Chang WL

ORIGINAL ARTICLE

Retrospective Study

- 1533 Efficacy of radiofrequency ablation combined with sorafenib for treating liver cancer complicated with portal hypertension and prognostic factors
Yang LM, Wang HJ, Li SL, Gan GH, Deng WW, Chang YS, Zhang LF

Clinical Trials Study

- 1545 Effect of *Aspergillus niger* prolyl endopeptidase in patients with celiac disease on a long-term gluten-free diet
Stefanolo JP, Segura V, Grizzuti M, Heredia A, Comino I, Costa AF, Puebla R, Temprano MP, Niveloni SI, de Diego G, Oregui ME, Smecuol EG, de Marzi MC, Verdú EF, Sousa C, Bai JC
- 1556 Effects of *Lactobacillus paracasei* N1115 on gut microbial imbalance and liver function in patients with hepatitis B-related cirrhosis
Hu YC, Ding XC, Liu HJ, Ma WL, Feng XY, Ma LN

Prospective Study

- 1572** Washed microbiota transplantation for Crohn's disease: A metagenomic, metatranscriptomic, and metabolomic-based study

Chen SJ, Zhang DY, Wu X, Zhang FM, Cui BT, Huang YH, Zhang ZL, Wang R, Bai FH

Basic Study

- 1588** Silent information regulator sirtuin 1 ameliorates acute liver failure *via* the p53/glutathione peroxidase 4/gasdermin D axis

Zhou XN, Zhang Q, Peng H, Qin YJ, Liu YH, Wang L, Cheng ML, Luo XH, Li H

- 1609** Identification of an immune-related gene signature for predicting prognosis and immunotherapy efficacy in liver cancer *via* cell-cell communication

Li JT, Zhang HM, Wang W, Wei DQ

META-ANALYSIS

- 1621** Effects of neoadjuvant chemotherapy *vs* chemoradiotherapy in the treatment of esophageal adenocarcinoma: A systematic review and meta-analysis

Csontos A, Fazekas A, Szakó L, Farkas N, Papp C, Ferenczi S, Bellyei S, Hegyi P, Papp A

CASE REPORT

- 1636** Myocardial metastasis from ZEB1- and TWIST-positive spindle cell carcinoma of the esophagus: A case report

Shibata Y, Ohmura H, Komatsu K, Sagara K, Matsuyama A, Nakano R, Baba E

ABOUT COVER

Editorial Board of *World Journal of Gastroenterology*, David L Morris, MD, FRCS (Ed), Professor, Department of Surgery, University of New South Wales, Sydney 2217, New South Wales, Australia. david.morris@unsw.edu.au

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (WJG, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now abstracted and indexed in Science Citation Index Expanded (SCIE), MEDLINE, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJG as 4.3; Quartile category: Q2. The WJG's CiteScore for 2021 is 8.3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yi-Xuan Cai; Production Department Director: Xu Guo; Cover Editor: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski

EXECUTIVE ASSOCIATE EDITORS-IN-CHIEF

Xian-Jun Yu (Pancreatic Oncology), Jian-Gao Fan (Chronic Liver Disease), Hou-Bao Liu (Biliary Tract Disease)

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

March 21, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

PUBLISHING PARTNER

Shanghai Pancreatic Cancer Institute and Pancreatic Cancer Institute, Fudan University
Biliary Tract Disease Institute, Fudan University

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/gerinfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/gerinfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

POLICY OF CO-AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/310>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/gerinfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

PUBLISHING PARTNER's OFFICIAL WEBSITE

<https://www.shca.org.cn>
<https://www.zs-hospital.sh.cn>



Routine utilization of machine perfusion in liver transplantation: Ready for prime time?

Alessandro Parente, Keyue Sun, Philipp Dutkowski, AM James Shapiro, Andrea Schlegel

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): A
Grade B (Very good): 0
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Kalinowski P, Poland

Received: January 3, 2024

Peer-review started: January 3, 2024

First decision: January 19, 2024

Revised: January 30, 2024

Accepted: February 28, 2024

Article in press: February 28, 2024

Published online: March 21, 2024



Alessandro Parente, AM James Shapiro, Department of Surgery, Division of Transplantation, University of Alberta Hospital, Edmonton T6G 2B7, Canada

Keyue Sun, Immunity and Inflammation, Lerner Research Institute, Cleveland, OH 44195, United States

Philipp Dutkowski, Swiss Hepato-Pancreato-Biliary Center, Department of Visceral and Transplantation Surgery, University Hospital Zurich, Zurich 8091, Switzerland

Andrea Schlegel, Immunity and Inflammation, Digestive Disease and Surgery Institute, Cleveland, OH 44195, United States

Corresponding author: Alessandro Parente, FEBS, MD, PhD, Surgeon, Department of Surgery, Division of Transplantation, University of Alberta Hospital, No. 8440 112 St NW, Edmonton T6G 2B7, Canada. aleparen@gmail.com

Abstract

The last decade has been notable for increasing high-quality research and dramatic improvement in outcomes with dynamic liver preservation. Robust evidence from numerous randomized controlled trials has been pooled by meta-analyses, providing the highest available evidence on the protective effect of machine perfusion (MP) over static cold storage in liver transplantation (LT). Based on a protective effect with less complications and improved graft survival, the field has seen a paradigm shift in organ preservation. This editorial focuses on the role of MP in LT and how it could become the new “gold standard”. Strong collaborative efforts are needed to explore its effects on long-term outcomes.

Key Words: Liver transplantation; Machine perfusion; Viability assessment; Hypothermic oxygenated perfusion; Normothermic machine perfusion

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Machine perfusion (MP) has garnered the interest of the transplant community given its proven beneficial effects on the clinical outcomes after liver transplantation (LT). Herein, we discuss the historical background of MP in LT and the available clinical evidence. Furthermore, we highlighted the obstacles and the need for future research, in particular with respect to viability assessment and prolonged preservation times.

Citation: Parente A, Sun K, Dutkowski P, Shapiro AJ, Schlegel A. Routine utilization of machine perfusion in liver transplantation: Ready for prime time? *World J Gastroenterol* 2024; 30(11): 1488-1493

URL: <https://www.wjgnet.com/1007-9327/full/v30/i11/1488.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v30.i11.1488>

INTRODUCTION

Machine perfusion (MP) technology is an old concept first introduced by Alexis Carrell and Charles Lindberg in the 1920's, then utilized first in clinical kidney transplantation in 1968[1], however static cold storage (SCS) appeared simpler and more practical, effectively hindering the progress in MP technology development for more than 3 decades. Recently, MP has emerged as one of the most promising approaches to improve post-transplant outcomes after liver transplantation (LT).

Two main ex-situ liver perfusion and one in-situ donor approach are increasingly used in clinical practice today[2]. The first ex-situ technique is known as hypothermic oxygenated perfusion (HOPE) using a highly oxygenated (pO_2 : > 60 kPa) artificial solution at hypothermic temperatures (8-12 °C)[2]. The second perfusion technique, normothermic MP (NMP), uses a blood-based perfusates at 37 °C, thus recreating a near-physiologic environment. Both HOPE and NMP are mainly applied after SCS, however NMP-preservation is also started upfront after minimal SCS of 2-3 hrs or as described with "ischemia-free organ transplantation (IFOT)", as a technique avoiding all SCS[2]. In contrast, during in-situ normothermic regional perfusion (NRP) a veno-arterial extracorporeal membrane oxygenation technique is used to recirculate and oxygenate donor blood immediately after circulatory arrest in donation after circulatory death (DCD) donors[2].

The importance of oxygen was understood early with the use of HOPE in kidney transplants in 1968, however the earlier perfusion concepts in LT were simply hypothermic (HMP). Guarrera *et al*[3] presented the first clinical study in LT in 2010 with HMP using a homegrown device (Figure 1). NRP was used first in uncontrolled DCD in 2007[4] and NMP started later off in clinics in 2016[5].

The very first randomized controlled trial (RCT) with NMP and the Organox-Metra® device was published in 2018 by Nasralla *et al*[6]. The authors demonstrated less early allograft dysfunction (EAD) and lower peak aspartate aminotransferase levels during the first week after transplant in the recipient of NMP-treated grafts compared to SCS[6]. Only one year later, similar results in terms of EAD were illustrated by another single center RCT, however with much smaller number of patients[7].

Next, in 2021, another two important RCTs were published. The first was focused on HOPE in DCD livers and demonstrated a four times lower risk of non-anastomotic biliary strictures compared to SCS, together with lower EAD-rates, postreperfusion syndrome and retransplantation[8]. And the second RCT paralleled such results with HOPE and showed less 90-d post-operative complications, shorter intensive care and hospital stay compared to SCS in extended criteria donor livers after brain death (DBD)[9]. The following 2 years, the findings of these studies with HOPE were supported by further RCTs. The Bologna team showed that patients receiving HOPE-treated extended criteria DBD livers had significantly lower EAD-rates, better 1-year graft survival, less post-LT complications and lower hospital readmission rates compared to SCS[10]. Authors from the United States published the results of another multicenter RCT using the portable OCS™ device provided by Transmedics for upfront NMP[11]. The NMP group in this study had significantly lower EAD-rates and less signs of ischemia-reperfusion injury on histopathology[11].

In the first half of 2023, two RCTs were published: The largest RCT ever conducted with HOPE and the first one with IFOT. In the former, the authors illustrated that, although the number of serious complications between HOPE and SCS was similar, a post hoc analysis revealed that liver-related serious complications occurred less frequently in the HOPE group compared to SCS. Notably, graft failure due to liver-related complications did not occur with HOPE but in 7% of SCS grafts (6 of 85)[12]. Subsequently, a Chinese group reported the results of the first RCT of DBD livers randomly assigned to SCS or IFOT, demonstrating significantly reduced EAD-rates, post-reperfusion syndrome, less non-anastomotic biliary stricture, and less cumulative post-operative complications at 12 months[13]. A group from Poland assessed the role of HOPE in their country. In low-risk DBD donors HOPE had no impact on outcomes, however authors demonstrated significantly lower EAD-rates and post-operative complications in donor livers with higher risk, *i.e.*, with a donor risk index of > 1.7[14]. Two additional RCTs were recently presented with NMP and HOPE in the United States. NMP with the Organox-Metra® device had a positive effect on the post-reperfusion syndrome[15], and HOPE showed lower primary non function, less EAD rates and reduced biliary strictures[16].

Such RCTs were recently combined with other retrospective studies in several meta-analyses[17], summarizing the beneficial effects of MP over SCS[18,19]. Currently, there are no available RCTs comparing HOPE and NMP, however results of ongoing RCT are awaited (Clinicaltrials.gov: NCT04644744).

While there are no RCT with NRP available yet, this technology appears to be beneficial to reduce biliary complications, improve graft survival and increase organ utilization if combined with limited SCS up to 6-7 hrs[20-22]. It seems




Figure 1 Overview on the history of organ and liver preservation. Key steps are highlighted for each time frame, with a final focus on the future research questions. HOPE: Hypothermic oxygenated perfusion; NRP: Normothermic regional perfusion; RCT: Randomized controlled trial; COR: Controlled oxygenated rewarming.

unlikely that a RCT would proceed as the clinical outcomes have been so positive to date—in fact it is currently mandated in France and other European countries that DCD donation should not proceed unless NRP is applied. Clinical studies demonstrated however several limitations, especially when donor warm ischemia time before or SCS after NRP are prolonged or when these grafts were used for retransplant candidates[23–26]. Perfusion technologies are not mutually exclusive, and indeed when looking at real-world data have been combined with satisfactory outcomes[27]. In particular, in Italy where the no-touch period following DCD donation is 20 min, combining NRP and HOPE has yielded good outcomes[28].

Despite its clinical benefits, MP is still not routinely used worldwide. There are several reasons behind this which have been identified as lack of funding, clarity as to what is research and what is accepted clinical care delivery, knowledge and availability of healthcare staff trained in the operation of this equipment[29,30]. In addition, it must be noted that the current available evidence is limited at one year follow-up. In fact, all RCTs and meta-analyses that have been published have reported their outcomes at one year. This appears to be one of the main limitations of the current available literature. As life expectancy of LT recipients continues to increase, exceeding 70% at 5 years[31], more evidence is needed to better understand the impact of MP on long-term outcomes after LT. Additionally, many RCTs have reported similar study endpoint outcome measures with however heterogeneous definitions often lacking certain endpoints[18]. Future research will need to endeavor in more homogeneous outcome reporting to provide replicable data worldwide.

Perfusing marginal organ with the aim to increase liver utilization rates necessitate reliable viability tests. Although there are several biochemical factors that have been explored both for HOPE and NMP[32,33], there are no clear guidelines regarding a systematic viability assessment during liver perfusion and available parameters often lack sensitivity and acceptable positive predictive value to discriminate between livers of metabolically good quality and others with too high risk for failure in the recipient. Most viability parameters are usually measured in the perfusate, with arbitrary timepoints and cutoffs, reflecting a post ischemia-reperfusion injury downstream of instigating mitochondria in hepatocytes, including perfusate lactate, transaminases, cytokine, and lactate dehydrogenase. Bile chemistry is also applied to assist identification of livers at risk of ischemic cholangiopathy. The early occurring cellular damage during MP was recently assessed with proteomics analysis of bile collected during sequential HOPE, rewarming and subsequent NMP[34]. On the other hand, mitochondrial transition pore opening with danger signal release appears to be a more upfront parameter of liver injury[33,35,36]. There is evidence that MP could be utilized to reprogram mitochondrial metabolism during HOPE before normothermic reperfusion. The slow metabolism of succinate and concomitant ATP reloading are two protective mechanisms described with HOPE. A recent study has demonstrated that the analysis of flavin mononucleotide from complex I, perfusate nicotinamide adenine dinucleotide hydrogen, and mitochondrial CO₂ production during HOPE allows a more objective viability assessment of liver quality on a subcellular level which seems to be more reliable when compared to donor derived data, in particular for high-risk organs such as DCD[33].

In the real world, MP is used also for logistic reasons and prolonged liver preservation is increasingly used to enable timely LT with both NMP and HOPE[6,11,15,37]. A human liver graft was preserved for 3 d and successfully transplanted[38] and some discarded livers were preserved with MP up to 7 d with satisfactory viability parameters[39]. Although this approach might be of interest in the future to enable the modification of cellular metabolisms during ex-situ MP, its daily applicability is still limited to small number of cases and challenges with difficult devices required particularly for prolonged NMP[40].

Next to a prolonged *ex-situ* perfusion, super cooling and cryopreservation are potential but as yet untested clinically as additional technique to prolong organ preservation. A recent study has explored the concept of cryopreservation which enabled to preserve experimental kidney grafts for 100 d through vitrification and nanowarming[41]. Despite being compelling novel strategies, such techniques are currently limited to animal models or small studies and their potential future applications remains unclear.

CONCLUSION

In summary, within the past 5 years, increasing evidence has demonstrated the clinical benefits of MP in the setting of LT (Figure 1). We believe that this progress with MP technology and the clear beneficial effects will soon outpace and replace the standard SCS preservation for the human liver grafts, and now necessitates both a paradigm shift and rapid change in clinical practice to capitalize on these advances. This will pave the way for a new era in organ transplantation, leading to the application of MP routinely in clinical practice, in particular for marginal organs. However, several challenges remain and will need to be addressed in future research.

FOOTNOTES

Author contributions: Parente A and Schlegel A designed research; Parente A and Schlegel A wrote the manuscript; Sun K, Shapiro AMJ and Dutkowski P revised the manuscript. All authors approved the final version of the manuscript.

Conflict-of-interest statement: All authors declare no conflict of interest.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Canada

ORCID number: Alessandro Parente 0000-0001-5506-224X; Philipp Dutkowski 0000-0002-3016-604X; AM James Shapiro 0000-0002-6215-0990.

S-Editor: Qu XL

L-Editor: A

P-Editor: Cai YX

REFERENCES

1. Belzer FO, Ashby BS, Gulyassy PF, Powell M. Successful seventeen-hour preservation and transplantation of human-cadaver kidney. *N Engl J Med* 1968; **278**: 608-610 [PMID: 4866541 DOI: 10.1056/NEJM196803142781108]
2. De Goeij FHC, De Meijer V, Mergental H, Guarrera JV, Asthana S, Ghinolfi D, Boteon YL, Selzner N, Kalisvaart M, Pulitano C, Sonnenday C, Martins PN, Berlakovich G, Schlegel A. Challenges With the Implementation of Machine Perfusion in Clinical Liver Transplantation. *Transplantation* 2023 [PMID: 38057969 DOI: 10.1097/TP.0000000000004872]
3. Guarrera JV, Henry SD, Samstein B, Odeh-Ramadan R, Kinkhabwala M, Goldstein MJ, Ratner LE, Renz JF, Lee HT, Brown RS Jr, Emond JC. Hypothermic machine preservation in human liver transplantation: the first clinical series. *Am J Transplant* 2010; **10**: 372-381 [PMID: 19958323 DOI: 10.1111/j.1600-6143.2009.02932.x]
4. Fondevila C, Hessheimer AJ, Ruiz A, Calatayud D, Ferrer J, Charco R, Fuster J, Navasa M, Rimola A, Taurá P, Ginés P, Manyalich M, García-Valdecasas JC. Liver transplant using donors after unexpected cardiac death: novel preservation protocol and acceptance criteria. *Am J Transplant* 2007; **7**: 1849-1855 [PMID: 17564639 DOI: 10.1111/j.1600-6143.2007.01846.x]
5. Ravikumar R, Jassem W, Mergental H, Heaton N, Mirza D, Perera MT, Quaglia A, Holroyd D, Vogel T, Coussios CC, Friend PJ. Liver Transplantation After Ex Vivo Normothermic Machine Preservation: A Phase I (First-in-Man) Clinical Trial. *Am J Transplant* 2016; **16**: 1779-1787 [PMID: 26752191 DOI: 10.1111/ajt.13708]
6. Nasralla D, Coussios CC, Mergental H, Akhtar MZ, Butler AJ, Ceresa CDL, Chiocchia V, Dutton SJ, García-Valdecasas JC, Heaton N, Imber C, Jassem W, Jochmans I, Karani J, Knight SR, Kocabayoglu P, Malagò M, Mirza D, Morris PJ, Pallan A, Paul A, Pavel M, Perera MTPR, Pirenne J, Ravikumar R, Russell L, Upponi S, Watson CJE, Weissenbacher A, Ploeg RJ, Friend PJ; Consortium for Organ Preservation in Europe. A randomized trial of normothermic preservation in liver transplantation. *Nature* 2018; **557**: 50-56 [PMID: 29670285 DOI: 10.1038/s41586-018-0047-9]
7. Ghinolfi D, Rreka E, De Tata V, Franzini M, Pezzati D, Fierabracci V, Masini M, Cacciatoinsilla A, Bindi ML, Marselli L, Mazzotti V, Morganti R, Marchetti P, Biancofiore G, Campani D, Paolicchi A, De Simone P. Pilot, Open, Randomized, Prospective Trial for Normothermic Machine Perfusion Evaluation in Liver Transplantation From Older Donors. *Liver Transpl* 2019; **25**: 436-449 [PMID: 30362649 DOI: 10.1002/lt.25362]
8. van Rijn R, Schurink IJ, de Vries Y, van den Berg AP, Cortes Cerisuelo M, Darwish Murad S, Erdmann JI, Gilbo N, de Haas RJ, Heaton N, van Hoek B, Huurman VAL, Jochmans I, van Leeuwen OB, de Meijer VE, Monbaliu D, Polak WG, Slangen JJG, Troisi RI, Vanlander A, de Jonge J, Porte RJ; DHOPE-DCD Trial Investigators. Hypothermic Machine Perfusion in Liver Transplantation - A Randomized Trial. *N Engl J*

- Med* 2021; **384**: 1391-1401 [PMID: 33626248 DOI: 10.1056/NEJMoa2031532]
- 9 **Czigany Z**, Pratschke J, Froněk J, Guba M, Schöning W, Raptis DA, Andrassy J, Kramer M, Strnad P, Tolba RH, Liu W, Keller T, Miller H, Pavicevic S, Uluk D, Kocik M, Lurje I, Trautwein C, Mehrabi A, Popescu I, Vondran FWR, Ju C, Tacke F, Neumann UP, Lurje G. Hypothermic Oxygenated Machine Perfusion Reduces Early Allograft Injury and Improves Post-transplant Outcomes in Extended Criteria Donation Liver Transplantation From Donation After Brain Death: Results From a Multicenter Randomized Controlled Trial (HOPE ECD-DBD). *Ann Surg* 2021; **274**: 705-712 [PMID: 34334635 DOI: 10.1097/SLA.0000000000005110]
 - 10 **Ravaioli M**, Germinario G, Dajti G, Sessa M, Vasuri F, Siniscalchi A, Morelli MC, Serenari M, Del Gaudio M, Zanfi C, Odaldi F, Bertuzzo VR, Maroni L, Laurenzi A, Cescon M. Hypothermic oxygenated perfusion in extended criteria donor liver transplantation-A randomized clinical trial. *Am J Transplant* 2022; **22**: 2401-2408 [PMID: 35671067 DOI: 10.1111/ajt.17115]
 - 11 **Markmann JF**, Abouljoud MS, Ghobrial RM, Bhati CS, Pelletier SJ, Lu AD, Ottmann S, Klair T, Eymard C, Roll GR, Magliocca J, Pruett TL, Reyes J, Black SM, Marsh CL, Schnickel G, Kinkhabwala M, Florman SS, Merani S, Demetris AJ, Kimura S, Rizzari M, Saharia A, Levy M, Agarwal A, Cigarroa FG, Eason JD, Syed S, Washburn WK, Parekh J, Moon J, Maskin A, Yeh H, Vagefi PA, MacConmara MP. Impact of Portable Normothermic Blood-Based Machine Perfusion on Outcomes of Liver Transplant: The OCS Liver PROTECT Randomized Clinical Trial. *JAMA Surg* 2022; **157**: 189-198 [PMID: 34985503 DOI: 10.1001/jamasurg.2021.6781]
 - 12 **Schlegel A**, Mueller M, Muller X, Eden J, Panconesi R, von Felten S, Steigmiller K, Sousa Da Silva RX, de Rougemont O, Mabrut JY, Lesurtel M, Cerisuelo MC, Heaton ND, Allard MA, Adam R, Monbaliu D, Jochmans I, Haring MPD, Porte RJ, Parente A, Muiesan P, Kron P, Attia M, Kollmann D, Berlakovitch G, Rogiers X, Petterson K, Kranich AL, Amberg S, Müllhaupt B, Clavien PA, Dutkowski P. A multicenter randomized-controlled trial of hypothermic oxygenated perfusion (HOPE) for human liver grafts before transplantation. *J Hepatol* 2023; **78**: 783-793 [PMID: 36681160 DOI: 10.1016/j.jhep.2022.12.030]
 - 13 **Guo Z**, Zhao Q, Jia Z, Huang C, Wang D, Ju W, Zhang J, Yang L, Huang S, Chen M, Zhu X, Hu A, Ma Y, Wu L, Chen Y, Han M, Tang Y, Wang G, Wang L, Li L, Xiong W, Zhang Z, Shen Y, Tang Z, Zhu C, Chen X, Hu X, Guo Y, Chen H, Zhang T, Zeng P, Lai S, Wang T, Chen Z, Gong J, Yu J, Sun C, Li C, Tan H, Liu Y, Dong Y, Liao B, Ren J, Zhou Z, Andrea S, Björn N, Cai C, Gong F, Rong J, Huang W, Guan X, Clavien PA, Stefan TG, Huang J, He X. A randomized-controlled trial of ischemia-free liver transplantation for end-stage liver disease. *J Hepatol* 2023; **79**: 394-402 [PMID: 37086919 DOI: 10.1016/j.jhep.2023.04.010]
 - 14 **Grat M**, Morawski M, Zhylo A, Rykowski P, Krasnodębski M, Wyporski A, Borkowski J, Lewandowski Z, Kobryn K, Stankiewicz R, Stypułkowski J, Hołówko W, Patkowski W, Mielczarek-Putka M, Struga M, Szczepankiewicz B, Górnica B, Krawczyk M. Routine End-ischemic Hypothermic Oxygenated Machine Perfusion in Liver Transplantation From Donors After Brain Death: A Randomized Controlled Trial. *Ann Surg* 2023; **278**: 662-668 [PMID: 37497636 DOI: 10.1097/SLA.0000000000006055]
 - 15 **Chapman WC**, Barbas AS, D'Alessandro AM, Vianna R, Kubal CA, Abt P, Sonnday C, Barth R, Alvarez-Casas J, Yersiz H, Eckhoff D, Cannon R, Genyk Y, Sher L, Singer A, Feng S, Roll G, Cohen A, Doyle MB, Sudan DL, Al-Adra D, Khan A, Subramanian V, Abraham N, Olthoff K, Tekin A, Berg L, Coussios C, Morris C, Randle L, Friend P, Knechtle SJ. Normothermic Machine Perfusion of Donor Livers for Transplantation in the United States: A Randomized Controlled Trial. *Ann Surg* 2023; **278**: e912-e921 [PMID: 37389552 DOI: 10.1097/SLA.0000000000005934]
 - 16 **Panayotova GG**, Lunsford KE, Quillin RC 3rd, Rana A, Agopian VG, Lee-Riddle GS, Markovic D, Paterno F, Griesemer AD, Amin A, Alonso D, Rocca JP, Borja-Cacho D, Hernandez-Alejandro R, Fung JJ, Pelletier SJ, Shah SA, Guarrera JV. Portable hypothermic oxygenated machine perfusion for organ preservation in liver transplantation: A randomized, open-label, clinical trial. *Hepatology* 2023 [PMID: 38090880 DOI: 10.1097/HEP.0000000000000715]
 - 17 **Parente A**, Dondossola D, Dutkowski P, Schlegel A. Current evidence on the beneficial HOPE-effect based on systematic reviews and meta-analyses in liver transplantation. *J Hepatol* 2024; **80**: e116-e119 [PMID: 37952765 DOI: 10.1016/j.jhep.2023.10.033]
 - 18 **Parente A**, Tirotta F, Pini A, Eden J, Dondossola D, Manzia TM, Dutkowski P, Schlegel A. Machine perfusion techniques for liver transplantation - A meta-analysis of the first seven randomized-controlled trials. *J Hepatol* 2023; **79**: 1201-1213 [PMID: 37302578 DOI: 10.1016/j.jhep.2023.05.027]
 - 19 **Tingle SJ**, Dobbins JJ, Thompson ER, Figueiredo RS, Mahendran B, Pandanaboyana S, Wilson C. Machine perfusion in liver transplantation. *Cochrane Database Syst Rev* 2023; **9**: CD014685 [PMID: 37698189 DOI: 10.1002/14651858.CD014685.pub2]
 - 20 **Hessheimer AJ**, de la Rosa G, Gastaca M, Ruiz P, Otero A, Gómez M, Alconchel F, Ramírez P, Bosca A, López-Andújar R, Atutxa L, Royo-Villanova M, Sánchez B, Santoyo J, Marín LM, Gómez-Bravo MÁ, Mosteiro F, Villegas Herrera MT, Villar Del Moral J, González-Abos C, Vidal B, López-Domínguez J, Lladó L, Roldán J, Justo I, Jiménez C, López-Monclús J, Sánchez-Turrión V, Rodríguez-Laíz G, Velasco Sánchez E, López-Baena JA, Caralt M, Charco R, Tomé S, Varo E, Martí-Cruchaga P, Rotellar F, Varona MA, Barrera M, Rodríguez-Sanjuan JC, Briceño J, López D, Blanco G, Nuño J, Pacheco D, Coll E, Domínguez-Gil B, Fondevila C. Abdominal normothermic regional perfusion in controlled donation after circulatory determination of death liver transplantation: Outcomes and risk factors for graft loss. *Am J Transplant* 2022; **22**: 1169-1181 [PMID: 34856070 DOI: 10.1111/ajt.16899]
 - 21 **Hessheimer AJ**, Coll E, Torres F, Ruiz P, Gastaca M, Rivas JI, Gómez M, Sánchez B, Santoyo J, Ramírez P, Parrilla P, Marín LM, Gómez-Bravo MÁ, García-Valdecasas JC, López-Monclús J, Boscá A, López-Andújar R, Fundora-Suárez J, Villar J, García-Sesma Á, Jiménez C, Rodríguez-Laíz G, Lladó L, Rodríguez JC, Barrera M, Charco R, López-Baena JA, Briceño J, Pardo F, Blanco G, Pacheco D, Domínguez-Gil B, Sánchez Turrión V, Fondevila C. Normothermic regional perfusion vs. super-rapid recovery in controlled donation after circulatory death liver transplantation. *J Hepatol* 2019; **70**: 658-665 [PMID: 30582980 DOI: 10.1016/j.jhep.2018.12.013]
 - 22 **Oniscu GC**, Mehew J, Butler AJ, Sutherland A, Gaurav R, Hogg R, Currie I, Jones M, Watson CJE. Improved Organ Utilization and Better Transplant Outcomes With In Situ Normothermic Regional Perfusion in Controlled Donation After Circulatory Death. *Transplantation* 2023; **107**: 438-448 [PMID: 35993664 DOI: 10.1097/TP.0000000000004280]
 - 23 **Richards JA**, Gaurav R, Upponi SS, Swift L, Fear C, Webb GJ, Allison MED, Watson CJE, Butler AJ. Outcomes of livers from donation after circulatory death donors with extended agonal phase and the adjunct of normothermic regional perfusion. *Br J Surg* 2023; **110**: 1112-1115 [PMID: 37079886 DOI: 10.1093/bjs/znad099]
 - 24 **Gaurav R**, Butler AJ, Kosmoliaptis V, Mumford L, Fear C, Swift L, Fedotovs A, Upponi S, Khwaja S, Richards J, Allison M, Watson CJE. Liver Transplantation Outcomes From Controlled Circulatory Death Donors: SCS vs in situ NRP vs ex situ NMP. *Ann Surg* 2022; **275**: 1156-1164 [PMID: 35258511 DOI: 10.1097/SLA.0000000000005428]
 - 25 **Antoine C**, Jasseron C, Dondero F, Savier E; French National Steering Committee of Donors After Circulatory Death. Liver Transplantation From Controlled Donors After Circulatory Death Using Normothermic Regional Perfusion: An Initial French Experience. *Liver Transpl* 2020; **26**: 1516-1521 [PMID: 32531132 DOI: 10.1002/lt.25818]
 - 26 **Savier E**, Lim C, Rayar M, Orlando F, Boudjema K, Mohkam K, Lesurtel M, Mabrut JY, Pittau G, Begdadi N, Cherqui D, Adam R, Dondero

- F, Sepulveda A, Soubrane O, Bucur P, Barbier L, Salame E, Jasseron C, Antoine C, Riou B, Scatton O. Favorable Outcomes of Liver Transplantation from Controlled Circulatory Death Donors Using Normothermic Regional Perfusion Compared to Brain Death Donors. *Transplantation* 2020; **104**: 1943-1951 [PMID: 32639402 DOI: 10.1097/TP.0000000000003372]
- 27 **van Leeuwen OB**, Bodewes SB, Lantinga VA, Haring MPD, Thorne AM, Brüggewirth IMA, van den Berg AP, de Boer MT, de Jong IEM, de Kleine RHJ, Lascaris B, Nijsten MWN, Reijntjens KMEM, de Meijer VE, Porte RJ. Sequential hypothermic and normothermic machine perfusion enables safe transplantation of high-risk donor livers. *Am J Transplant* 2022; **22**: 1658-1670 [PMID: 35286759 DOI: 10.1111/ajt.17022]
- 28 **De Carlis R**, Schlegel A, Frassoni S, Olivieri T, Ravaioli M, Camagni S, Patrono D, Bassi D, Pagano D, Di Sandro S, Lauterio A, Bagnardi V, Gruttadauria S, Cillo U, Romagnoli R, Colledan M, Cescon M, Di Benedetto F, Muiesan P, De Carlis L. How to Preserve Liver Grafts From Circulatory Death With Long Warm Ischemia? A Retrospective Italian Cohort Study With Normothermic Regional Perfusion and Hypothermic Oxygenated Perfusion. *Transplantation* 2021; **105**: 2385-2396 [PMID: 33617211 DOI: 10.1097/TP.0000000000003595]
- 29 **Parente A**, Flores Carvalho M, Panconesi R, Boteon YL, De Carlis R, Dutkowski P, Muiesan P, Dondossola D, Schlegel A. Trends and Obstacles to Implement Dynamic Perfusion Concepts for Clinical Liver Transplantation: Results from a Global Web-Based Survey. *J Clin Med* 2023; **12** [PMID: 37297960 DOI: 10.3390/jcm12113765]
- 30 **Flores Carvalho M**, Boteon YL, Guarrera JV, Modi PR, Lladó L, Lurje G, Kasahara M, Dutkowski P, Schlegel A. Obstacles to implement machine perfusion technology in routine clinical practice of transplantation: Why are we not there yet? *Hepatology* 2024; **79**: 713-730 [PMID: 37013926 DOI: 10.1097/HEP.0000000000000394]
- 31 OPTN/SRTR 2020 Annual Data Report: Liver. *HRSA*. Available from: https://srtr.transplant.hrsa.gov/annual_reports/2020/Liver.aspx
- 32 **Mergental H**, Laing RW, Kirkham AJ, Perera MTPR, Boteon YL, Attard J, Barton D, Curbishley S, Wilkhu M, Neil DAH, Hübscher SG, Muiesan P, Isaac JR, Roberts KJ, Abradelo M, Schlegel A, Ferguson J, Cilliers H, Bion J, Adams DH, Morris C, Friend PJ, Yap C, Afford SC, Mirza DF. Transplantation of discarded livers following viability testing with normothermic machine perfusion. *Nat Commun* 2020; **11**: 2939 [PMID: 32546694 DOI: 10.1038/s41467-020-16251-3]
- 33 **Eden J**, Breuer E, Birrer D, Müller M, Pfister M, Mayr H, Sun K, Widmer J, Huwyler F, Ungethüm U, Humar B, Gupta A, Schiess S, Wendt M, Immer F, Elmer A, Meierhofer D, Schlegel A, Dutkowski P. Screening for mitochondrial function before use-routine liver assessment during hypothermic oxygenated perfusion impacts liver utilization. *EBioMedicine* 2023; **98**: 104857 [PMID: 37918219 DOI: 10.1016/j.ebiom.2023.104857]
- 34 **Thorne AM**, Wolters JC, Lascaris B, Bodewes SB, Lantinga VA, van Leeuwen OB, de Jong IEM, Ustyantsev K, Berezikov E, Lisman T, Kuipers F, Porte RJ, de Meijer VE. Bile proteome reveals biliary regeneration during normothermic preservation of human donor livers. *Nat Commun* 2023; **14**: 7880 [PMID: 38036513 DOI: 10.1038/s41467-023-43368-y]
- 35 **Eden J**, Dutkowski P. Prolonging Preservation or Assessment of Organ Quality-What is Key? *Transpl Int* 2023; **36**: 12174 [PMID: 38020743 DOI: 10.3389/ti.2023.12174]
- 36 **Sampaziotis F**, Muraro D, Tysoe OC, Sawiak S, Beach TE, Godfrey EM, Upponi SS, Brevini T, Wesley BT, Garcia-Bernardo J, Mahbubani K, Canu G, Gieseck R 3rd, Berntsen NL, Mulcahy VL, Crick K, Fear C, Robinson S, Swift L, Gambardella L, Bargehr J, Ortmann D, Brown SE, Osnato A, Murphy MP, Corbett G, Gelson WTH, Mells GF, Humphreys P, Davies SE, Amin I, Gibbs P, Sinha S, Teichmann SA, Butler AJ, See TC, Melum E, Watson CJE, Saeb-Parsy K, Vallier L. Cholangiocyte organoids can repair bile ducts after transplantation in the human liver. *Science* 2021; **371**: 839-846 [PMID: 33602855 DOI: 10.1126/science.aaz6964]
- 37 **Brüggewirth IMA**, Mueller M, Lantinga VA, Camagni S, De Carlis R, De Carlis L, Colledan M, Dondossola D, Drefs M, Eden J, Ghinolfi D, Koliogiannis D, Lurje G, Manzia TM, Monbaliu D, Muiesan P, Patrono D, Pratschke J, Romagnoli R, Rayar M, Roma F, Schlegel A, Dutkowski P, Porte RJ, de Meijer VE. Prolonged preservation by hypothermic machine perfusion facilitates logistics in liver transplantation: A European observational cohort study. *Am J Transplant* 2022; **22**: 1842-1851 [PMID: 35315202 DOI: 10.1111/ajt.17037]
- 38 **Clavien PA**, Dutkowski P, Mueller M, Eshmuminov D, Bautista Borrego L, Weber A, Muellhaupt B, Sousa Da Silva RX, Burg BR, Rudolf von Rohr P, Schuler MJ, Becker D, Hefti M, Tibbitt MW. Transplantation of a human liver following 3 days of ex situ normothermic preservation. *Nat Biotechnol* 2022; **40**: 1610-1616 [PMID: 35641829 DOI: 10.1038/s41587-022-01354-7]
- 39 **Lau NS**, Ly M, Dennis C, Liu K, Kench J, Crawford M, Pulitano C. Long-term normothermic perfusion of human livers for longer than 12 days. *Artif Organs* 2022; **46**: 2504-2510 [PMID: 35929443 DOI: 10.1111/aor.14372]
- 40 **Eshmuminov D**, Becker D, Bautista Borrego L, Hefti M, Schuler MJ, Hagedorn C, Muller X, Mueller M, Onder C, Graf R, Weber A, Dutkowski P, Rudolf von Rohr P, Clavien PA. An integrated perfusion machine preserves injured human livers for 1 week. *Nat Biotechnol* 2020; **38**: 189-198 [PMID: 31932726 DOI: 10.1038/s41587-019-0374-x]
- 41 **Han Z**, Rao JS, Gangwar L, Namsrai BE, Pasek-Allen JL, Etheridge ML, Wolf SM, Pruett TL, Bischof JC, Finger EB. Vitrification and nanowarming enable long-term organ cryopreservation and life-sustaining kidney transplantation in a rat model. *Nat Commun* 2023; **14**: 3407 [PMID: 37296144 DOI: 10.1038/s41467-023-38824-8]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

