

World Journal of *Gastroenterology*

World J Gastroenterol 2024 April 21; 30(15): 2068-2178



EDITORIAL

- 2068** Combination treatment of inflammatory bowel disease: Present status and future perspectives
Triantafyllidis JK, Zografos CG, Konstadoulakis MM, Papalois AE
- 2081** Interplay between metabolic dysfunction-associated fatty liver disease and renal function: An intriguing pediatric perspective
Nardolillo M, Rescigno F, Bartiromo M, Piatto D, Guarino S, Marzuillo P, Miraglia del Giudice E, Di Sessa A
- 2087** Advancements in hemostatic strategies for managing upper gastrointestinal bleeding: A comprehensive review
Lee AY, Cho JY
- 2091** Understanding autoimmune pancreatitis: Clinical features, management challenges, and association with malignancies
Christodoulidis G, Kouliou MN, Koumarelas KE
- 2096** Probiotics: Shaping the gut immunological responses
Filidou E, Kandilogiannakis L, Shrewsbury A, Kolios G, Kotzampassi K

MINIREVIEWS

- 2109** Hepatocellular carcinoma and musculoskeletal system: A narrative literature review
Jadzic J, Djonic D

ORIGINAL ARTICLE

Case Control Study

- 2118** Urgent one-stage endoscopic treatment for choledocholithiasis related moderate to severe acute cholangitis: A propensity score-matched analysis
Zhou Y, Zhang YQ, Huang SJ, Liang Y, Liang X, Wali M, Feng YD

Retrospective Study

- 2128** Computed tomography-based radiomics to predict early recurrence of hepatocellular carcinoma post-hepatectomy in patients background on cirrhosis
Qian GX, Xu ZL, Li YH, Lu JL, Bu XY, Wei MT, Jia WD

Basic Study

- 2143** Taurine attenuates activation of hepatic stellate cells by inhibiting autophagy and inducing ferroptosis
Li S, Ren QJ, Xie CH, Cui Y, Xu LT, Wang YD, Li S, Liang XQ, Wen B, Liang MK, Zhao XF
- 2155** OSW-1 triggers necroptosis in colorectal cancer cells through the RIPK1/RIPK3/MLKL signaling pathway facilitated by the RIPK1-p62/SQSTM1 complex
Wang N, Li CY, Yao TF, Kang XD, Guo HS

LETTER TO THE EDITOR

- 2175** Circulating tumor DNA in liquid biopsy: Current diagnostic limitation

Liu SC

ABOUT COVER

Editorial Board Member of *World Journal of Gastroenterology*, Chen-Guo Ker, FACS, MD, PhD, Professor, HBP Surgeon, Department of General Surgery, E-Da Hospital, I-Shou University, Kaohsiung 824, Taiwan.
ed112739@edah.org.tw

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (WJG, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now abstracted and indexed in Science Citation Index Expanded (SCIE), MEDLINE, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJG as 4.3; Quartile category: Q2. The WJG's CiteScore for 2021 is 8.3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yi-Xuan Cai; **Production Department Director:** Xu Guo; **Cover Editor:** Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski

EXECUTIVE ASSOCIATE EDITORS-IN-CHIEF

Xian-Jun Yu (Pancreatic Oncology), Jian-Gao Fan (Chronic Liver Disease), Hou-Bao Liu (Biliary Tract Disease)

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

April 21, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

PUBLISHING PARTNER

Shanghai Pancreatic Cancer Institute and Pancreatic Cancer Institute, Fudan University
Biliary Tract Disease Institute, Fudan University

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

POLICY OF CO-AUTHORS

<https://www.wjgnet.com/bpg/GerInfo/310>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

PUBLISHING PARTNER's OFFICIAL WEBSITE

<https://www.shca.org.cn>
<https://www.zs-hospital.sh.cn>



Interplay between metabolic dysfunction-associated fatty liver disease and renal function: An intriguing pediatric perspective

Michele Nardolillo, Fabiola Rescigno, Mario Bartiromo, Dario Piatto, Stefano Guarino, Pierluigi Marzuillo, Emanuele Miraglia del Giudice, Anna Di Sessa

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B, B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Liu QQ, China

Received: December 27, 2023

Peer-review started: December 27, 2023

First decision: January 25, 2024

Revised: February 18, 2024

Accepted: March 26, 2024

Article in press: March 26, 2024

Published online: April 21, 2024



Michele Nardolillo, Fabiola Rescigno, Mario Bartiromo, Dario Piatto, Stefano Guarino, Pierluigi Marzuillo, Emanuele Miraglia del Giudice, Anna Di Sessa, Department of Woman, Child and of General and Specialized Surgery, Università degli Studi della Campania “Luigi Vanvitelli”, Naples 80138, Italy

Corresponding author: Anna Di Sessa, MD, PhD, Postdoc, Research Fellow, Department of Woman, Child and of General and Specialized Surgery, Università degli Studi della Campania “Luigi Vanvitelli”, Via de Crecchio 2, Naples 80138, Italy. anna.disessa@libero.it

Abstract

Over recent years, the nomenclature of non-alcoholic fatty liver disease has undergone significant changes. Indeed, in 2020, an expert consensus panel proposed the term “Metabolic (dysfunction) associated fatty liver disease” (MAFLD) to underscore the close association of fatty liver with metabolic abnormalities, thereby highlighting the cardiometabolic risks (such as metabolic syndrome, type 2 diabetes, insulin resistance, and cardiovascular disease) faced by these patients since childhood. More recently, this term has been further replaced with metabolic associated steatotic liver disease. It is worth noting that emerging evidence not only supports a close and independent association of MAFLD with chronic kidney disease in adults but also indicates its interplay with metabolic impairments. However, comparable pediatric data remain limited. Given the progressive and chronic nature of both diseases and their prognostic cardiometabolic implications, this editorial aims to provide a pediatric perspective on the intriguing relationship between MAFLD and renal function in childhood.

Key Words: Metabolic (dysfunction) associated fatty liver disease; Renal; Function; Children; Obesity

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Metabolic (dysfunction) associated fatty liver disease (MAFLD) has been closely linked to a wide spectrum of cardiometabolic consequences. Among these, accumulating data demonstrated an association between MAFLD and renal function in children with obesity. Worthy of note, a shared pathophysiology has been reported with a pivotal role of insulin-resistance in this dangerous interplay among obesity, renal hemodynamics, and metabolic derangements. Considering the relevant clinical and prognostic implications of this association, an increased awareness of this growing health concern for clinicians is needed.

Citation: Nardolillo M, Rescigno F, Bartiromo M, Piatto D, Guarino S, Marzuillo P, Miraglia del Giudice E, Di Sessa A. Interplay between metabolic dysfunction-associated fatty liver disease and renal function: An intriguing pediatric perspective. *World J Gastroenterol* 2024; 30(15): 2081-2086

URL: <https://www.wjgnet.com/1007-9327/full/v30/i15/2081.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v30.i15.2081>

INTRODUCTION

In 2020, an expert panel proposed replacing the nomenclature of non-alcoholic fatty liver disease (NAFLD) with the more inclusive term metabolic (dysfunction) associated fatty liver disease (MAFLD) to underscore the pathogenic link between the disease and metabolic derangements in both adults and children[1,2]. According to the definition[2], the acronym MAFLD was introduced to strengthen the association of pediatric hepatic steatosis (diagnosed by biopsy with histological evaluation, imaging, or blood biomarkers) with at least one of the following criteria: (1) Excess adiposity; (2) Prediabetes or type 2 diabetes (T2D); and (3) Evidence of metabolic dysregulation[2].

Similar to NAFLD, the prevalence of MAFLD has been increasing worldwide in parallel with the obesity epidemic since childhood[1-3]. Current estimates suggest that approximately a quarter of the global adult population presents with MAFLD, while pediatric data indicate a global prevalence of 34% among children and adolescents with obesity in the general population, and 45% among peers in the obesity clinical setting[3]. Given this, MAFLD represents a major health and economic concern.

Indeed, MAFLD has been closely linked to cardiovascular disease (CVD) risk, but recent evidence also supports its role as a predictor of atherosclerosis, heart failure, T2D, and cancer-related mortality[1,3]. Consistent with previous evidence for NAFLD[4,5], a higher incidence of chronic kidney disease (CKD) has also emerged in patients with MAFLD[6-10], underscoring the impact of the disease on cardiometabolic health. Notably, robust adult data document a stronger association of MAFLD with CKD than NAFLD[10,11]. However, similar evidence linking MAFLD to CKD is still limited in childhood[12,13], though the underlying complex interplay among liver, kidney, adiposity, and metabolic dysfunction has been highlighted in these young patients[14-17]. Therefore, in this editorial, we aim to discuss the available evidence in the field to provide a perspective on the relevant clinical and prognostic implications of the interplay between MAFLD and CKD in children with obesity.

THE PATHOPHYSIOLOGICAL COMPLEXITY OF THE INTERPLAY BETWEEN MAFLD AND KIDNEY

Although the interplay of several determinants such as genetics, lifestyle, and environmental factors has been well-documented in the development of MAFLD, its exact pathogenesis remains to be fully elucidated[1-3]. In particular, certain features of MAFLD, such as systemic inflammation, metabolic dysfunction, and vascular dysfunction, have been found to act as mediators in the tangled inter-organ crosstalk underlying the close association of MAFLD with extrahepatic diseases (*e.g.*, CVDs, cognitive impairment, thyroid dysfunction, and CKD)[1,3,6,8]. On this basis, the relationship between MAFLD and renal function is unsurprising given their shared pathogenic factors such as low-grade inflammation, oxidative stress, and insulin resistance (IR)[8,9,18] (Figure 1).

In this complex pathophysiological puzzle, the contribution of genetics has also been well-documented[8,9,19]. The risk variant rs738409 C>G of the Patatin-like phospholipase domain-containing 3 gene represents a major genetic determinant of MAFLD in both adults and children[19-21]. Additionally, other polymorphisms such as the transmembrane 6 superfamily member 2 loss-of-function variant rs58542926[22] and the rs641738 C>T variant in the membrane-bound O-acyltransferase 7 gene[23,24] have been found to increase susceptibility to MAFLD[8,19]. Notably, recent evidence also supports a multifaceted role of these polymorphisms in renal health[21,22,24,25].

Furthermore, a prominent pathogenic role for inflammation needs to be underlined[26], as demonstrated by the inclusion of C-reactive protein among MAFLD diagnostic criteria[1,2]. Indeed, both chronic low-grade inflammation and oxidative stress lead to hepatic fibrosis by exacerbating pro-inflammatory signaling pathway activation[18,26], which in turn promotes endothelial dysfunction[18]. Consequently, this deleterious cycle impairs renal function by increasing glomerular permeability and proteinuria[18,27]. Notably, this increase in systemic inflammation might also affect cardiovascular health[6,8,18].

To complicate matters, IR - as a shared pathophysiological factor - plays a central role in the development of both MAFLD and CKD[28,29]. In fact, certain processes mediated by IR, such as an increased release of free fatty acids and an altered secretion of adipokines, affect both the liver and kidney[18,30-32].




Figure 1 The complex pathophysiological interplay between metabolic (dysfunction) associated fatty liver disease and renal function.

T2D: Type 2 diabetes; FFA: Free fatty acids; MAFLD: Metabolic (dysfunction) associated fatty liver disease; CKD: Chronic kidney disease; ROS: Reactive oxygen species; RNS: Reactive nitrogen species; SCFAs: Short chain fatty acids; TMAOs: Trimethylamine-N-oxides; PNPLA3: Patatin-like phospholipase domain-containing 3; TM6SF2: Transmembrane 6 superfamily member 2; MBOAT7: Membrane-bound O-acyltransferase 7.

Interestingly, recent insights into the complex pathogenic interplay between MAFLD and CKD highlight a role for the gut-liver-kidney axis, suggesting an intriguing contribution of gut microbiota to the development and progression of both diseases[18,33,34]. This realizes a vicious circle among dysbiosis, IR, inflammation, and oxidative stress[18,32]. In particular, metabolites derived from the gut microbiota such as trimethylamine-N-oxides[34,35] exert systemic effects impairing both liver[33,34] and renal function[35]. Additionally, the crosstalk between the liver and kidney also affects the production of short-chain fatty acids[36], further amplifying the underlying pathophysiological processes (*e.g.*, inflammation, IR, and oxidative stress) of MAFLD and CKD[36-38].

MAFLD AND CARDIOMETABOLIC HEALTH: THE RENAL PERSPECTIVE

MAFLD has been widely recognized as a strong predictor of all-cause mortality including CVD-related mortality[39-42]. More interestingly, recent data have unveiled an intriguing correlation between MAFLD and renal impairments[43-45], particularly CKD[18,38,40].

Several shared pathophysiological factors such as metabolic abnormalities (*e.g.*, obesity and IR), inflammation, adipokines, gut dysbiosis, and genetics contribute to liver and renal damage development[4,18,19,38]. In particular, IR appears to play a central role in the tangled interplay between MAFLD and CKD through various molecular pathways[6, 8,18,38]. This exacerbates not only hypertension and atherogenic dyslipidemia but also activates renin-angiotensin system (RAS), which is closely related to endothelial dysfunction[6,8]. RAS activation further amplifies both proinflammatory and pro-coagulant states by releasing numerous mediators[6,8,18,38].

Moreover, evidence supports not only the intricate link between MAFLD and kidney function[8,43-45] but also the severity of hepatic damage with specific renal parameters in adults[46-48]. Indeed, recent studies have revealed associations between liver fibrosis, assessed *via* transient elastography, and kidney dysfunction parameters such as urinary albumin-to-creatinine ratio and estimated glomerular filtration rate (eGFR)[46-48]. However, research exploring the association between MAFLD and renal damage in pediatric populations remains limited[12,13].

Valentino *et al*[12] examined a cohort of Italian children with MAFLD and CKD during the initial coronavirus disease 2019 pandemic lockdown. After a six-month follow-up, children with MAFLD and CKD exhibited lower eGFR levels and an overall worse cardiometabolic profile compared to those without MAFLD[12]. De Groot *et al*[13] showed that children with a liver fat fraction > 2% and MAFLD presented with a worse cardiometabolic risk profile including higher blood pressure levels (as renal injury expression) compared to both children with a liver fat fraction < 2% and ≥ 2% liver fat without MAFLD. Of note, children with a liver fat fraction > 2% and MAFLD had an increased odds of clustering cardio-metabolic risk factor compared to those with liver fat fraction < 2% independently of MAFLD presence (odds ratio = 7.65, 95% confidence interval: 5.04-11.62)[13]. In line with adult findings[42-44], these results underscore an association between MAFLD and renal damage in childhood[12,13]. Nevertheless, further large-scale studies are warranted to

validate this intriguing link comprehensively.

CONCLUSION

Recently, several lines of evidence have supported not only a pathogenic link between renal health and metabolic dysfunction but also its prognostic impact, both in children[12,13,18] and adults[18,40-43]. A complex inter-organ crosstalk sustained by metabolic impairments (e.g., IR, visceral adiposity) has been supposed to be responsible for the relationship of MAFLD with extrahepatic diseases, including CKD[18]. More specifically, an intricate interplay among cardiometabolic risk factors, genetics, lipid nephrotoxicity, and hemodynamic changes has likely been implied in the relationship between MAFLD and renal impairment, but knowledge gaps in its exact pathophysiology still remain[8,18].

Compared to NAFLD, a better diagnostic and prognostic performance in identifying subjects at greater risk of hepatic and extra-hepatic complications has been demonstrated for the more inclusive MAFLD definition[8,18,41,48]. Based on these premises, children with MAFLD should receive more attention from clinicians through a multidisciplinary assessment that takes into account the intriguing MAFLD-associated multi-organ crosstalk.

Given the prognostic implications of the tangled relationship of MAFLD with extra-hepatic diseases[2,40-43], more scientific efforts are needed in this research area for a deeper understanding of the intricate interplay of molecular pathways contributing to hepatic and renal damage. On the other hand, this might also implement strategies for overall MAFLD management (including prevention, diagnosis, and treatment) to optimize patient outcomes since childhood.

FOOTNOTES

Author contributions: Nardolillo M and Di Sessa A wrote the manuscript; Marzuillo P, Miraglia del Giudice E, and Di Sessa A conceived the manuscript; Guarino S, Miraglia del Giudice E, and Di Sessa A supervised the manuscript drafting; Nardolillo M, Rescigno F, Bartiromo M, and Piatto D reviewed the literature data; Rescigno F prepared the figure. Each author contributed important intellectual content during manuscript drafting or revision.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Italy

ORCID number: Michele Nardolillo 0000-0002-1608-4777; Mario Bartiromo 0000-0002-5958-3314; Stefano Guarino 0000-0002-0551-5236; Pierluigi Marzuillo 0000-0003-4682-0170; Emanuele Miraglia del Giudice 0000-0002-1492-076X; Anna Di Sessa 0000-0002-5877-3757.

S-Editor: Wang JJ

L-Editor: A

P-Editor: Cai YX

REFERENCES

- 1 **Eslam M**, Newsome PN, Sarin SK, Anstee QM, Targher G, Romero-Gomez M, Zelber-Sagi S, Wai-Sun Wong V, Dufour JF, Schattenberg JM, Kawaguchi T, Arrese M, Valenti L, Shiha G, Tiribelli C, Yki-Järvinen H, Fan JG, Grønbaek H, Yilmaz Y, Cortez-Pinto H, Oliveira CP, Bedossa P, Adams LA, Zheng MH, Fouad Y, Chan WK, Mendez-Sanchez N, Ahn SH, Castera L, Bugianesi E, Ratziu V, George J. A new definition for metabolic dysfunction-associated fatty liver disease: An international expert consensus statement. *J Hepatol* 2020; **73**: 202-209 [PMID: 32278004 DOI: 10.1016/j.jhep.2020.03.039]
- 2 **Eslam M**, Alkhouri N, Vajro P, Baumann U, Weiss R, Socha P, Marcus C, Lee WS, Kelly D, Porta G, El-Guindi MA, Alisi A, Mann JP, Mouane N, Baur LA, Dhawan A, George J. Defining paediatric metabolic (dysfunction)-associated fatty liver disease: an international expert consensus statement. *Lancet Gastroenterol Hepatol* 2021; **6**: 864-873 [PMID: 34364544 DOI: 10.1016/S2468-1253(21)00183-7]
- 3 **Liu J**, Mu C, Li K, Luo H, Liu Y, Li Z. Estimating Global Prevalence of Metabolic Dysfunction-Associated Fatty Liver Disease in Overweight or Obese Children and Adolescents: Systematic Review and Meta-Analysis. *Int J Public Health* 2021; **66**: 1604371 [PMID: 34690666 DOI: 10.3389/ijph.2021.1604371]
- 4 **Mantovani A**, Scorletti E, Mosca A, Alisi A, Byrne CD, Targher G. Complications, morbidity and mortality of nonalcoholic fatty liver disease. *Metabolism* 2020; **111S**: 154170 [PMID: 32006558 DOI: 10.1016/j.metabol.2020.154170]
- 5 **Mantovani A**, Petracca G, Beatrice G, Csermely A, Lonardo A, Schattenberg JM, Tilg H, Byrne CD, Targher G. Non-alcoholic fatty liver disease and risk of incident chronic kidney disease: an updated meta-analysis. *Gut* 2022; **71**: 156-162 [PMID: 33303564 DOI: 10.1136/gutjnl-2020-323082]
- 6 **Pipitone RM**, Ciccioli C, Infantino G, La Mantia C, Parisi S, Tulone A, Pennisi G, Grimaudo S, Petta S. MAFLD: a multisystem disease. *Ther Adv Endocrinol Metab* 2023; **14**: 20420188221145549 [PMID: 36726391 DOI: 10.1177/20420188221145549]
- 7 **Kwon SY**, Park J, Park SH, Lee YB, Kim G, Hur KY, Koh J, Jee JH, Kim JH, Kang M, Jin SM. MAFLD and NAFLD in the prediction of

- incident chronic kidney disease. *Sci Rep* 2023; **13**: 1796 [PMID: 36720976 DOI: 10.1038/s41598-023-27762-6]
- 8 Mantovani A, Lombardi R, Cattazzo F, Zusi C, Cappelli D, Dalbeni A. MAFLD and CKD: An Updated Narrative Review. *Int J Mol Sci* 2022; **23** [PMID: 35806010 DOI: 10.3390/ijms23137007]
 - 9 Sun DQ, Jin Y, Wang TY, Zheng KI, Rios RS, Zhang HY, Targher G, Byrne CD, Yuan WJ, Zheng MH. MAFLD and risk of CKD. *Metabolism* 2021; **115**: 154433 [PMID: 33212070 DOI: 10.1016/j.metabol.2020.154433]
 - 10 Tanaka M, Mori K, Takahashi S, Higashiura Y, Ohnishi H, Hanawa N, Furuhashi M. Metabolic dysfunction-associated fatty liver disease predicts new onset of chronic kidney disease better than fatty liver or nonalcoholic fatty liver disease. *Nephrol Dial Transplant* 2023; **38**: 700-711 [PMID: 35612975 DOI: 10.1093/ndt/gfac188]
 - 11 Liang Y, Chen H, Liu Y, Hou X, Wei L, Bao Y, Yang C, Zong G, Wu J, Jia W. Association of MAFLD With Diabetes, Chronic Kidney Disease, and Cardiovascular Disease: A 4.6-Year Cohort Study in China. *J Clin Endocrinol Metab* 2022; **107**: 88-97 [PMID: 34508601 DOI: 10.1210/clinem/dgab641]
 - 12 Valentino MS, Marzuillo P, Esposito C, Bartiromo M, Nardolillo M, Villani AV, Maresca A, Furcolo G, Guarino S, Miraglia Del Giudice E, Di Sessa A. The Impact of COVID-19 Pandemic Lockdown on the Relationship between Pediatric MAFLD and Renal Function. *J Clin Med* 2023; **12** [PMID: 36902824 DOI: 10.3390/jcm12052037]
 - 13 de Groot J, Santos S, Geurtsen ML, Felix JF, Jaddoe VWV. Risk factors and cardio-metabolic outcomes associated with metabolic-associated fatty liver disease in childhood. *EClinicalMedicine* 2023; **65**: 102248 [PMID: 37855025 DOI: 10.1016/j.eclinm.2023.102248]
 - 14 Di Sessa A, Passaro AP, Colasante AM, Cioffi S, Guarino S, Umamo GR, Papparella A, Miraglia Del Giudice E, Marzuillo P. Kidney damage predictors in children with metabolically healthy and metabolically unhealthy obesity phenotype. *Int J Obes (Lond)* 2023; **47**: 1247-1255 [PMID: 37689826 DOI: 10.1038/s41366-023-01379-1]
 - 15 Sawyer A, Zeitler E, Trachtman H, Bjornstad P. Kidney Considerations in Pediatric Obesity. *Curr Obes Rep* 2023; **12**: 332-344 [PMID: 37558892 DOI: 10.1007/s13679-023-00522-3]
 - 16 Mangat G, Nair N, Barat O, Abboud B, Pais P, Bagga S, Raina R. Obesity-related glomerulopathy in children: connecting pathophysiology to clinical care. *Clin Kidney J* 2023; **16**: 611-618 [PMID: 37007691 DOI: 10.1093/ckj/sfac233]
 - 17 Di Sessa A, Guarino S, Umamo GR, Miraglia Del Giudice E, Marzuillo P. MASLD vs. NAFLD: A better definition for children with obesity at higher risk of kidney damage. *J Hepatol* 2024; **80**: e87-e89 [PMID: 37890724 DOI: 10.1016/j.jhep.2023.10.021]
 - 18 Theofilis P, Vordoni A, Kalaitzidis RG. Interplay between metabolic dysfunction-associated fatty liver disease and chronic kidney disease: Epidemiology, pathophysiologic mechanisms, and treatment considerations. *World J Gastroenterol* 2022; **28**: 5691-5706 [PMID: 36338895 DOI: 10.3748/wjg.v28.i39.5691]
 - 19 Lanzaro F, Guarino S, D'Addio E, Salvatori A, D'Anna JA, Marzuillo P, Miraglia Del Giudice E, Di Sessa A. Metabolic-associated fatty liver disease from childhood to adulthood: State of art and future directions. *World J Hepatol* 2022; **14**: 1087-1098 [PMID: 35978659 DOI: 10.4254/wjh.v14.i6.1087]
 - 20 Dongiovanni P, Donati B, Fares R, Lombardi R, Mancina RM, Romeo S, Valenti L. PNPLA3 I148M polymorphism and progressive liver disease. *World J Gastroenterol* 2013; **19**: 6969-6978 [PMID: 24222941 DOI: 10.3748/wjg.v19.i41.6969]
 - 21 Mantovani A, Pelusi S, Margarita S, Malvestiti F, Dell'Alma M, Bianco C, Ronzoni L, Prati D, Targher G, Valenti L. Adverse effect of PNPLA3 p.I148M genetic variant on kidney function in middle-aged individuals with metabolic dysfunction. *Aliment Pharmacol Ther* 2023; **57**: 1093-1102 [PMID: 36947711 DOI: 10.1111/apt.17477]
 - 22 Marzuillo P, Di Sessa A, Cirillo G, Umamo GR, Pedullà M, La Manna A, Guarino S, Miraglia Del Giudice E. Transmembrane 6 superfamily member 2 167K allele improves renal function in children with obesity. *Pediatr Res* 2020; **88**: 300-304 [PMID: 31923913 DOI: 10.1038/s41390-020-0753-5]
 - 23 Di Sessa A, Umamo GR, Cirillo G, Del Prete A, Iacomino R, Marzuillo P, Del Giudice EM. The Membrane-bound O-Acyltransferase7 rs641738 Variant in Pediatric Nonalcoholic Fatty Liver Disease. *J Pediatr Gastroenterol Nutr* 2018; **67**: 69-74 [PMID: 29601441 DOI: 10.1097/MPG.0000000000001979]
 - 24 Koo BK, An JN, Joo SK, Kim D, Lee S, Bae JM, Park JH, Kim JH, Chang MS, Kim W. Association Between a Polymorphism in MBOAT7 and Chronic Kidney Disease in Patients With Biopsy-Confirmed Nonalcoholic Fatty Liver Disease. *Clin Gastroenterol Hepatol* 2020; **18**: 2837-2839.e2 [PMID: 31546054 DOI: 10.1016/j.cgh.2019.09.017]
 - 25 Targher G, Mantovani A, Alisi A, Mosca A, Panera N, Byrne CD, Nobili V. Relationship Between PNPLA3 rs738409 Polymorphism and Decreased Kidney Function in Children With NAFLD. *Hepatology* 2019; **70**: 142-153 [PMID: 30912854 DOI: 10.1002/hep.30625]
 - 26 Masarone M, Rosato V, Dallio M, Gravina AG, Aglitti A, Loguerzio C, Federico A, Persico M. Role of Oxidative Stress in Pathophysiology of Nonalcoholic Fatty Liver Disease. *Oxid Med Cell Longev* 2018; **2018**: 9547613 [PMID: 29991976 DOI: 10.1155/2018/9547613]
 - 27 Stenvinkel P, Chertow GM, Devarajan P, Levin A, Andreoli SP, Bangalore S, Warady BA. Chronic Inflammation in Chronic Kidney Disease Progression: Role of Nrf2. *Kidney Int Rep* 2021; **6**: 1775-1787 [PMID: 34307974 DOI: 10.1016/j.ekir.2021.04.023]
 - 28 Sakurai Y, Kubota N, Yamauchi T, Kadowaki T. Role of Insulin Resistance in MAFLD. *Int J Mol Sci* 2021; **22** [PMID: 33923817 DOI: 10.3390/ijms22084156]
 - 29 Spoto B, Pisano A, Zoccali C. Insulin resistance in chronic kidney disease: a systematic review. *Am J Physiol Renal Physiol* 2016; **311**: F1087-F1108 [PMID: 27707707 DOI: 10.1152/ajprenal.00340.2016]
 - 30 Ma D, Feitosa MF, Wilk JB, Laramie JM, Yu K, Leidencker-Foster C, Myers RH, Province MA, Borecki IB. Leptin is associated with blood pressure and hypertension in women from the National Heart, Lung, and Blood Institute Family Heart Study. *Hypertension* 2009; **53**: 473-479 [PMID: 19204185 DOI: 10.1161/HYPERTENSIONAHA.108.118133]
 - 31 Zhao D, Zhu X, Jiang L, Huang X, Zhang Y, Wei X, Zhao X, Du Y. Advances in understanding the role of adiponectin in renal fibrosis. *Nephrology (Carlton)* 2021; **26**: 197-203 [PMID: 33073881 DOI: 10.1111/nep.13808]
 - 32 Li D, Lu Y, Yuan S, Cai X, He Y, Chen J, Wu Q, He D, Fang A, Bo Y, Song P, Bogaert D, Tsilidis K, Larsson SC, Yu H, Zhu H, Theodoratou E, Zhu Y, Li X. Gut microbiota-derived metabolite trimethylamine-N-oxide and multiple health outcomes: an umbrella review and updated meta-analysis. *Am J Clin Nutr* 2022; **116**: 230-243 [PMID: 35348578 DOI: 10.1093/ajcn/nqac074]
 - 33 Li X, Hong J, Wang Y, Pei M, Wang L, Gong Z. Trimethylamine-N-Oxide Pathway: A Potential Target for the Treatment of MAFLD. *Front Mol Biosci* 2021; **8**: 733507 [PMID: 34660695 DOI: 10.3389/fmolb.2021.733507]
 - 34 Barrea L, Muscogiuri G, Annunziata G, Laudisio D, de Alteriis G, Tenore GC, Colao A, Savastano S. A New Light on Vitamin D in Obesity: A Novel Association with Trimethylamine-N-Oxide (TMAO). *Nutrients* 2019; **11** [PMID: 31185686 DOI: 10.3390/nu11061310]
 - 35 Touroutzis T, Lioulis G, Fylaktou A, Moysidou E, Papagianni A, Stangou M. Microbiome in Chronic Kidney Disease. *Life (Basel)* 2022; **12** [PMID: 36294946 DOI: 10.3390/life12101513]

- 36 **Magliocca G**, Mone P, Di Iorio BR, Heidland A, Marzocco S. Short-Chain Fatty Acids in Chronic Kidney Disease: Focus on Inflammation and Oxidative Stress Regulation. *Int J Mol Sci* 2022; **23** [PMID: 35628164 DOI: 10.3390/ijms23105354]
- 37 **Marzocco S**, Fazeli G, Di Micco L, Autore G, Adesso S, Dal Piaz F, Heidland A, Di Iorio B. Supplementation of Short-Chain Fatty Acid, Sodium Propionate, in Patients on Maintenance Hemodialysis: Beneficial Effects on Inflammatory Parameters and Gut-Derived Uremic Toxins, A Pilot Study (PLAN Study). *J Clin Med* 2018; **7** [PMID: 30274359 DOI: 10.3390/jcm7100315]
- 38 **Wang TY**, Wang RF, Bu ZY, Targher G, Byrne CD, Sun DQ, Zheng MH. Association of metabolic dysfunction-associated fatty liver disease with kidney disease. *Nat Rev Nephrol* 2022; **18**: 259-268 [PMID: 35013596 DOI: 10.1038/s41581-021-00519-y]
- 39 **Kim H**, Lee CJ, Ahn SH, Lee KS, Lee BK, Baik SJ, Kim SU, Lee JI. MAFLD Predicts the Risk of Cardiovascular Disease Better than NAFLD in Asymptomatic Subjects with Health Check-Ups. *Dig Dis Sci* 2022; **67**: 4919-4928 [PMID: 35579799 DOI: 10.1007/s10620-022-07508-6]
- 40 **Zhou XD**, Cai J, Targher G, Byrne CD, Shapiro MD, Sung KC, Somers VK, Chahal CAA, George J, Chen LL, Zhou Y, Zheng MH; CHES-MAFLD consortium. Metabolic dysfunction-associated fatty liver disease and implications for cardiovascular risk and disease prevention. *Cardiovasc Diabetol* 2022; **21**: 270 [PMID: 36463192 DOI: 10.1186/s12933-022-01697-0]
- 41 **Wang Y**, Yu Y, Zhang H, Chen C, Wan H, Chen Y, Xia F, Yu S, Wang N, Ye L, Lu Y. Cardiovascular and renal burdens among patients with MAFLD and NAFLD in China. *Front Endocrinol (Lausanne)* 2022; **13**: 968766 [PMID: 36120461 DOI: 10.3389/fendo.2022.968766]
- 42 **Wei S**, Song J, Xie Y, Huang J, Yang J. The Role of Metabolic Dysfunction-Associated Fatty Liver Disease in Developing Chronic Kidney Disease: Longitudinal Cohort Study. *JMIR Public Health Surveill* 2023; **9**: e45050 [PMID: 37140958 DOI: 10.2196/45050]
- 43 **Chen S**, Pang J, Huang R, Xue H, Chen X. Association of MAFLD with end-stage kidney disease: a prospective study of 337,783 UK Biobank participants. *Hepatology* 2023; **17**: 595-605 [PMID: 36809487 DOI: 10.1007/s12072-023-10486-0]
- 44 **Liu Y**, Chai S, Zhang X. Effect of MAFLD on albuminuria and the interaction between MAFLD and diabetes on albuminuria. *J Diabetes* 2024; **16**: e13501 [PMID: 37974383 DOI: 10.1111/1753-0407.13501]
- 45 **Colasante AM**, Bartiromo M, Nardolillo M, Guarino S, Marzuillo P, Mangoni di S Stefano GSRC, Miraglia Del Giudice E, Di Sessa A. Tangled relationship between insulin resistance and microalbuminuria in children with obesity. *World J Clin Pediatr* 2022; **11**: 455-462 [PMID: 36439903 DOI: 10.5409/wjcp.v11.i6.455]
- 46 **Ciardullo S**, Ballabeni C, Trevisan R, Perseghin G. Liver Stiffness, Albuminuria and Chronic Kidney Disease in Patients with NAFLD: A Systematic Review and Meta-Analysis. *Biomolecules* 2022; **12** [PMID: 35053253 DOI: 10.3390/biom12010105]
- 47 **Freitas M**, Macedo Silva V, Xavier S, Magalhães J, Marinho C, Cotter J. Early Kidney Dysfunction in Metabolic-Associated Fatty Liver Disease: Is Transient Elastography Useful as a Screening Method? *Dig Dis* 2021; **39**: 653-662 [PMID: 33508843 DOI: 10.1159/000514811]
- 48 **Marc L**, Mihaescu A, Lupusoru R, Grosu I, Gadalean F, Bob F, Chisavu L, Olariu N, Tucicovschi V, Timar B, Sporea I, Timar R, Schiller A. Liver Steatosis: Better Predictor of CKD in MAFLD Than Liver Fibrosis as Determined by Transient Elastography With Controlled Attenuation Parameter. *Front Med (Lausanne)* 2021; **8**: 788881 [PMID: 35096879 DOI: 10.3389/fmed.2021.788881]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

