

World Journal of *Gastroenterology*

World J Gastroenterol 2024 May 7; 30(17): 2287-2373



EDITORIAL

- 2287 Quick and easy assessment of sarcopenia in cirrhosis: Can ultrasound be the solution?
Campani F, Li Cavoli TV, Arena U, Marra F, Lynch EN, Campani C
- 2294 Expanding indications for chronic hepatitis B treatment: Is it really desirable to treat everyone?
Di Dato F, Iorio R
- 2298 Surgical cystogastrostomy: Is it still worthwhile?
Au KP, Chok KSH
- 2302 Evaluation of urea breath test as a diagnostic tool for *Helicobacter pylori* infection in adult dyspeptic patients
Said ZNA, El-Nasser AM
- 2308 Chronic active and atrophic gastritis as significant contributing factor to the development of gastric cystica profunda
Papp V, Miheller P

MINIREVIEWS

- 2311 Contrast-enhanced guided endoscopic ultrasound procedures
Gheorghiu MI, Seicean A, Pojoga C, Hagiuc C, Seicean R, Sparchez Z

ORIGINAL ARTICLE

Retrospective Study

- 2321 Efficacy and safety of targeted therapy plus immunotherapy combined with hepatic artery infusion chemotherapy (FOLFOX) for unresectable hepatocarcinoma
Lin ZP, Hu XL, Chen D, Huang DB, Zou XG, Zhong H, Xu SX, Chen Y, Li XQ, Zhang J

Prospective Study

- 2332 Transanal eco-Doppler evaluation after hemorrhoidal artery embolization
Tutino R, Stecca T, Farneti F, Massani M, Santoro GA
- 2343 Diagnostic and prognostic performances of GALAD score in staging and 1-year mortality of hepatocellular carcinoma: A prospective study
Jitpraphawan O, Ruamtawee W, Treewatchareekorn M, Sethasine S

META-ANALYSIS

- 2354 Minocycline in the eradication of *Helicobacter pylori* infection: A systematic review and meta-analysis
Zhou K, Li CL, Zhang H, Suo BJ, Zhang YX, Ren XL, Wang YX, Mi CM, Ma LL, Zhou LY, Tian XL, Song ZQ

LETTER TO THE EDITOR

- 2369** Targeting therapy for hepatocellular carcinoma by delivering microRNAs as exosomal cargo
Suda T
- 2371** Metabolic dysfunction-associated fatty liver disease and low muscle strength: A comment
Karim MM, Butt AS

ABOUT COVER

Editorial Board Member of *World Journal of Gastroenterology*, Ming-Zhe Ma, MD, Assistant Professor, Department of Gastric Surgery, Fudan University Shanghai Cancer Center, Shanghai 200032, China. mmz666@163.com

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (WJG, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now abstracted and indexed in Science Citation Index Expanded (SCIE), MEDLINE, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJG as 4.3; Quartile category: Q2. The WJG's CiteScore for 2021 is 8.3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Hua-Ge Yu; Production Department Director: Xu Guo; Cover Editor: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski

EXECUTIVE ASSOCIATE EDITORS-IN-CHIEF

Xian-Jun Yu (Pancreatic Oncology), Jian-Gao Fan (Chronic Liver Disease), Hou-Bao Liu (Biliary Tract Disease)

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

May 7, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

PUBLISHING PARTNER

Shanghai Pancreatic Cancer Institute and Pancreatic Cancer Institute, Fudan University
Biliary Tract Disease Institute, Fudan University

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/gerinfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/gerinfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

POLICY OF CO-AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/310>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/gerinfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

PUBLISHING PARTNER's OFFICIAL WEBSITE

<https://www.shca.org.cn>
<https://www.zs-hospital.sh.cn>



Quick and easy assessment of sarcopenia in cirrhosis: Can ultrasound be the solution?

Francesca Campani, Tancredi Vincenzo Li Cavoli, Umberto Arena, Fabio Marra, Erica Nicola Lynch, Claudia Campani

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's classification

Scientific Quality: Grade B

Novelty: Grade B

Creativity or Innovation: Grade B

Scientific Significance: Grade B

P-Reviewer: Yibirin M, United States

Received: January 30, 2024

Revised: March 16, 2024

Accepted: April 15, 2024

Published online: May 7, 2024



Francesca Campani, Department of Health Science, University Hospital Careggi, University of Florence, Florence 50134, Italy

Tancredi Vincenzo Li Cavoli, Umberto Arena, Fabio Marra, Claudia Campani, Internal Medicine and Liver Unit, University Hospital Careggi, University of Florence, Florence 50134, Italy

Fabio Marra, Claudia Campani, Department of Experimental and Clinical Medicine, University of Florence, Florence 50134, Italy

Erica Nicola Lynch, Gastroenterology Research Unit, Department of Experimental and Clinical Biomedical Sciences "Mario Serio", University of Florence, Florence 50134, Italy

Erica Nicola Lynch, Department of Medical Biotechnologies, University of Siena, Siena 53100, Italy

Corresponding author: Claudia Campani, MD, PhD, Academic Fellow, Doctor, Research Fellow, Department of Experimental and Clinical Medicine, University of Florence, Largo Brambilla 3, Florence 50134, Italy. claudiacampani.cc@gmail.com

Abstract

Cirrhosis is frequently associated with sarcopenia, with reported rates of over 80% in patients with decompensated alcohol-related liver disease. Sarcopenia negatively impacts the prognosis of cirrhotic patients and affects the response to treatment of patients with hepatocellular carcinoma (HCC). For these reasons, identifying an easy-to-perform method to assess sarcopenia is a key element in the optimization of care in this patient population. Assessment of muscle mass by computed tomography is considered the standard of care for the diagnosis of sarcopenia, but exposure to radiation and high costs limit its application in this setting, especially for repeated assessments. We believe that ultrasound, a cheap and harmless technique also used for HCC screening in cirrhotic patients, could have an expanding role in the diagnosis and follow-up of sarcopenia in these patients.

Key Words: Sarcopenia; Ultrasound; Cirrhosis; Hepatocellular carcinoma; Computed tomography

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Cirrhosis is frequently associated with sarcopenia, which negatively impacts the prognosis of cirrhotic patients and affects the response to treatment of patients with hepatocellular carcinoma (HCC). For these reasons, identifying an easy-to-perform method to assess sarcopenia in is a key element in the optimization of care in this patient population. We believe that ultrasound, a cheap and harmless technique also used for HCC screening in cirrhotic patients, could have an expanding role in the diagnosis and follow-up of sarcopenia in these patients.

Citation: Campani F, Li Cavoli TV, Arena U, Marra F, Lynch EN, Campani C. Quick and easy assessment of sarcopenia in cirrhosis: Can ultrasound be the solution? *World J Gastroenterol* 2024; 30(17): 2287-2293

URL: <https://www.wjgnet.com/1007-9327/full/v30/i17/2287.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v30.i17.2287>

INTRODUCTION

Cirrhosis is frequently associated with sarcopenia, with reported rates of over 80% in patients with decompensated, alcohol-related cirrhosis[1]. Various pathogenetic mechanisms contribute to muscle wasting in these patients, such as altered protein metabolism, resulting in reduced levels of circulating branched chain amino acids[2] and decreased protein synthesis, increased autophagy, proteolysis, and mitochondrial oxidative dysfunction in the skeletal muscle due to hyperammonemia[3,4]. Chronic systemic inflammation[5], reduction in circulating testosterone levels[6,7], and physical inactivity[8,9] are other factors contributing to sarcopenia in patients with advanced liver disease. Sarcopenia negatively affects the prognosis of cirrhotic patients and the response to treatment in patients with hepatocellular carcinoma (HCC). For these reasons, identifying an easy-to-perform method to assess sarcopenia is a key element in the optimization of care in this patient population. Assessment of muscle mass by computed tomography (CT) is considered the standard of care for the diagnosis of sarcopenia, but exposure to radiation and high costs limit its application in this setting, especially for repeated assessments. We believe that ultrasound (US), a cheap and harmless technique also used for HCC screening in cirrhotic patients, may have an expanding role in the diagnosis and follow-up of sarcopenia in these patients.

DEFINITION OF SARCOPENIA

Sarcopenia is a progressive and generalized skeletal muscle disorder mainly defined by two parameters: Muscle mass and muscle strength. Low muscle strength is the key characteristic of probable sarcopenia, whereas a diagnosis of sarcopenia can be confirmed only after detection of low muscle quantity and quality[10]. Moreover, reduced physical performance is indicative of severe sarcopenia[10], which is associated with an increased likelihood of adverse outcomes including falls, fractures, disability, and mortality[10]. Loss of skeletal muscle mass and function commonly occurring with advancing age is classified as primary sarcopenia, but many other factors can cause or contribute to the development of secondary sarcopenia[10]. Systemic diseases, especially those characterized by inflammatory processes, are one of the leading causes of secondary sarcopenia[11]. Physical inactivity and inadequate energy or protein intake are also involved in the development of sarcopenia[11].

Sarcopenia is also common in overweight and obese patients[11,12], where the loss of muscle mass and function can be favored by chronic low-grade inflammation, increased oxidative stress, insulin resistance, sedentary lifestyle, and a higher incidence of comorbid chronic diseases that may negatively impact muscle metabolism[13]. Several lines of evidence show that sarcopenic obesity represents a strong and independent risk factor for frailty, comorbidities, and mortality, especially among the elderly[14,15].

PREVALENCE AND ROLE OF SARCOPENIA ACROSS LIVER DISEASES

Sarcopenia in metabolic dysfunction-associated steatotic liver disease patients

Sarcopenia is closely associated with metabolic dysfunction-associated steatotic liver disease (MASLD), the most common cause of chronic liver disease in Western countries[16,17]. Patients with sarcopenic MASLD are generally older and more frequently female[18]. Sarcopenia has been suggested to increase the risk of progression of liver fibrosis, and therefore its early recognition may play an important role in preventing the development of cirrhosis[19-21]. Petta *et al*[22] showed that MASLD-sarcopenic patients have more severe liver fibrosis compared with those without. Moreover, the co-occurrence of MASLD and sarcopenia is associated with higher mortality, suggesting that sarcopenia may play a role in increasing the risk of cardiovascular diseases, metabolic disorders, and physical disability in this group of patients[23,24].

Sarcopenia and cirrhosis

Sarcopenia affects between 30% to 70% of cirrhotic patients[25], with higher rates reported in men[26,27]. The etiology of cirrhosis plays a relevant role in the development of sarcopenia. The highest prevalence of sarcopenia can be found in

patients with alcohol-associated cirrhosis, with a prevalence of over 80% in alcohol-related decompensated cirrhosis[1, 28]. Alcohol consumption affects muscle mass leading to muscle autophagy, inhibition of proteasome activity and a decrease in insulin-like growth factor 1[29]. Sarcopenia can be both a cause and a consequence of complications of cirrhosis. Ascites may favor muscle loss through anorexia, reduced mobility, and frequent hospitalizations[30]. On the other hand, reduced muscle mass is an independent risk factor for hepatic encephalopathy[31,32] and is linked to an increased risk of decompensation[33].

In cirrhosis, sarcopenia also negatively impacts quality of life[34], increases the risk of infection[35], and prolongs the duration of hospitalizations[36]. Additionally, several studies show that a diagnosis of sarcopenia in cirrhotic patients is associated with an increased risk of falls, fractures, acute-on-chronic liver failure, and death[37-39]. Indeed, a recent systematic review and metaanalysis of 22 studies including 6965 cirrhotic patients showed that the risk of death was 2.6 times higher in patients with sarcopenia[27]. Low muscle density has been shown to predict mortality even in patients with compensated cirrhosis[33,40], and sarcopenic obesity is associated with a higher incidence of sepsis-related death [41]. The presence of sarcopenia prior to liver transplantation can significantly increase the length of hospital and intensive care unit (ICU) stay[42,43] and worsens the overall prognosis of these patients[44].

Sarcopenia and HCC

Up to 30%-40% of HCC patients are affected by sarcopenia at the time of diagnosis, at least partially because of the pro-inflammatory state triggered by the altered tumor microenvironment[45]. As sarcopenia influences the response to surgical, locoregional, and systemic treatments, its timely recognition is essential. In patients who undergo liver resection or liver transplantation, tackling sarcopenia reduces sepsis-related complications and length of ICU stay, and decreases patient mortality[46]. In patients treated with thermal ablation, sarcopenia has been linked to a reduced overall survival (OS) and to a higher risk of HCC recurrence[47]. A worse prognosis and high rate of progression has been also described for HCC patients treated with transarterial chemoembolization[48].

Sarcopenia also appears to impact the response to systemic treatments. Scheiner *et al*[49] showed that sarcopenia is associated with worse OS (6.5 months *vs* 20.9 months), progression-free survival (5.8 months *vs* 8.3 months) and objective response rate (22% *vs* 39%) in patients treated with atezolizumab-bevacizumab. Sarcopenic patients treated with sorafenib were subject to a higher drug exposure and increased dose-limiting toxicities *vs* non-sarcopenic patients[50]. In patients treated with lenvatinib, Dong *et al*[51] showed that sarcopenia is an independent prognostic factor of a shorter OS. Sarcopenia might also predict drug toxicity and poor tolerance to lenvatinib[52]. Based on the above findings, an adequate evaluation and diagnosis of sarcopenia in patients with HCC is likely to improve their prognosis.

CURRENT METHODS FOR SARCOPENIA DIAGNOSIS

Although the diagnosis of sarcopenia involves both a functional and quantitative assessment of muscle mass, current research is mainly directed at finding an objective and reproducible method to measure muscle mass. CT imaging currently represents the gold standard to quantify skeletal muscle. Muscle mass is conventionally reported as skeletal mass index (SMI), calculated as the total skeletal muscle area at the level of L3 normalized for height[26]. SMI is the only parameter for which cut-off values for the diagnosis of sarcopenia have been validated, < 50 cm for men and < 39 cm for women[10,26,53]. Alternatively, the psoas muscle index at L3 has been identified as an alternative to SMI, although it shows low accuracy in cirrhotic patients[54]. However, CT scan is not an adequate method to serially follow the improvement or deterioration of muscle mass over time, because of high radiation exposure[55]. For this reason, body composition is assessed with CT scans only when these are performed for other reasons, as in the setting of HCC.

Dual-energy x-ray absorptiometry (DXA), magnetic resonance imaging (MRI), and bioelectrical impedance analysis (BIA) are currently available alternatives, albeit various limitations should be considered. DXA is a costly, radiation-dependent technique influenced by body mass index and fluid retention. MRI is highly accurate but expensive and with restricted availability in most settings. BIA is population and device-dependent and is also affected by fluid retention. When technology-based devices (BIA, DXA, MRI or CT) are not available or feasible, anthropometric measures could be used to quantify skeletal muscle mass, at the expense of test sensitivity and reproducibility[56].

THE ROLE OF US IN SARCOPENIA DIAGNOSIS

US is an accurate and reliable technique, with high reproducibility for the assessment of muscle size[57,58]. Furthermore, abdominal US is used to screen cirrhotic patients for HCC semiannually, in accordance with guidelines of the European Association for the Study of the Liver and the American Association for the Study of Liver Diseases[59,60]. Therefore, the availability of US in virtually all cirrhotic patients, its non-invasiveness and independence of exposure to X-rays, make it an appealing tool for the initial diagnosis and follow-up of sarcopenia in cirrhosis[61], also in clinical studies.

The use of US in muscle assessment has been specifically explored in patients with cirrhosis. It must be noted that most studies included patients with a mild or moderate reduction in liver function (Child Pugh classes A and B) due to the high impact of ascites on the evaluation of psoas muscle by US[62-64]. Furthermore, HCC patients are generally excluded by these studies due to neoplastic cachexia which is considered a confounding factor.

The anatomical site that best represents total skeletal muscle mass has not yet been defined. The rectus femoris (RF) could be a possibility as it is exposed to an earlier age-related decline than other sites such as the biceps femoris[65]. Most

studies evaluating sarcopenia in cirrhotic patients through US used the measurement of large muscles in the upper and lower limbs because of their ease of identification and lesser susceptibility to fluid retention[63]. In fact, ascites can influence the sonic window, especially when examining the muscles of the abdominal wall or psoas[63]. The same issue can be encountered in patients with obesity, a condition that, due to the rising global prevalence, is going to be very frequent in patients with cirrhosis[63]. Another aspect that needs to be defined is the US parameter to be used in muscle mass assessment. Thickness and cross-sectional area of the muscle show similar results as those of DXA, CT, and MRI and may be used to confirm the presence of muscle mass depletion[66]. Echo intensity is a measure of muscle composition in terms of fatty infiltration and presence of fibrous tissue[67]. Indeed, US machines are increasingly equipped with software that could be useful in qualitative analysis of the muscle, defining its microvasculature or stiffness[68-70]. Two-dimensional shear wave elastography of the RF is another qualitative parameter proposed for the assessment of lean mass using US. The measurement of stiffness with this method was feasible in all patients and correlated with liver frailty index (LFI) in a study that involved 44 outpatients with cirrhosis. In addition, RF thickness inversely correlated with LFI [70].

Other key aspects that require standardization are the type of probe that should be employed, the anatomical sites of measurement, the patient's position during the examination, the probe direction and pressure exerted on the muscle, and the parameters that should be measured[68]. A linear probe with a frequency of 5-12 MHz is usually preferred, except for the psoas muscle, for which the use of a convex probe with a frequency of 3.5-5 MHz appears to be more adequate[63].

Despite the lack of standardization, there is growing evidence on the use of US to assess sarcopenia in cirrhotic patients. A recent review evaluating 17 studies assessed the role of US in the diagnosis of sarcopenia in older adults, and showed that US is accurate for the assessment of muscles size, especially when the evaluation is targeted at the quadriceps femoris[57]. In a prospective study including 159 cirrhotic outpatients, Tandon *et al*[71] demonstrated that the combination of body mass index and US-measured thigh muscle thickness was able to identify sarcopenic patients, in both genders, with the same efficacy as CT[71]. This implies an evident advantage in terms of increased screening feasibility and serial assessment to monitor the effectiveness of nutritional interventions[71]. Of note, even in the context of cirrhosis and obesity, the assessment of lean mass through US has been demonstrated to be well-correlated with SMI calculated from CT[72]. Similarly, when LFI or subjective global assessment were employed as references for the assessment of muscle function, a robust correlation with US measurements (*i.e.*, the antero-posterior diameter of the RF, rectus abdominis thickness) was found[70,73].

Besides demonstrating a strong correlation with the reference gold standard, the assessment of lean mass using US also correlates with various clinical outcomes. For example, rectus abdominis thickness predicts survival in a study that included a small group of cirrhotic patients, and both US-SMI and US-psoas to height ratio were significantly related to hospitalization in patients with decompensated liver cirrhosis[73,74].

CONCLUSION

Despite the above outlined limitations and the limited amount of data in large series, the wide availability of the instrument, its ease of application, and especially the possibility of repeated monitoring on the same patient makes US assessment of lean mass in patients with cirrhosis an attractive area of interest for future study.

FOOTNOTES

Author contributions: Campani F performed the bibliographic search; Campani F and Li Cavoli TV drafted the initial manuscript; Lynch EN and Campani C re-screened the search results; Lynch EN provided English language revision as a native speaker; Arena U, Marra F, Lynch EN, and Campani C revised the article critically for important intellectual content; and all authors approved the final version of the manuscript.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

ORCID number: Umberto Arena 0000-0002-8839-1506; Fabio Marra 0000-0001-8629-0878; Erica Nicola Lynch 0000-0002-2638-2559; Claudia Campani 0000-0003-3842-782X.

S-Editor: Wang JJ

L-Editor: A

P-Editor: Yu HG

REFERENCES

- 1 DiMartini A, Cruz RJ Jr, Dew MA, Myaskovsky L, Goodpaster B, Fox K, Kim KH, Fontes P. Muscle mass predicts outcomes following liver transplantation. *Liver Transpl* 2013; **19**: 1172-1180 [PMID: 23960026 DOI: 10.1002/lt.23724]
- 2 Montanari A, Simoni I, Vallisa D, Trifirò A, Colla R, Abbiati R, Borghi L, Novarini A. Free amino acids in plasma and skeletal muscle of patients with liver cirrhosis. *Hepatology* 1988; **8**: 1034-1039 [PMID: 3417224 DOI: 10.1002/hep.1840080509]
- 3 Chen HW, Dunn MA. Muscle at Risk: The Multiple Impacts of Ammonia on Sarcopenia and Frailty in Cirrhosis. *Clin Transl Gastroenterol* 2016; **7**: e170 [PMID: 27228401 DOI: 10.1038/ctg.2016.33]
- 4 Dasarathy S, Hatzoglou M. Hyperammonemia and proteostasis in cirrhosis. *Curr Opin Clin Nutr Metab Care* 2018; **21**: 30-36 [PMID: 29035972 DOI: 10.1097/MCO.0000000000000426]
- 5 Arroyo V, García-Martínez R, Salvatella X. Human serum albumin, systemic inflammation, and cirrhosis. *J Hepatol* 2014; **61**: 396-407 [PMID: 24751830 DOI: 10.1016/j.jhep.2014.04.012]
- 6 Moctezuma-Velázquez C, Low G, Mourtzakis M, Ma M, Burak KW, Tandon P, Montano-Loza AJ. Association between Low Testosterone Levels and Sarcopenia in Cirrhosis: A Cross-sectional Study. *Ann Hepatol* 2018; **17**: 615-623 [PMID: 29893704 DOI: 10.5604/01.3001.0012.0930]
- 7 Sinclair M, Grossmann M, Hoermann R, Angus PW, Gow PJ. Testosterone therapy increases muscle mass in men with cirrhosis and low testosterone: A randomised controlled trial. *J Hepatol* 2016; **65**: 906-913 [PMID: 27312945 DOI: 10.1016/j.jhep.2016.06.007]
- 8 Hiraoka A, Michitaka K, Kiguchi D, Izumoto H, Ueki H, Kaneto M, Kitahata S, Aibiki T, Okudaira T, Tomida H, Miyamoto Y, Yamago H, Suga Y, Iwasaki R, Mori K, Miyata H, Tsubouchi E, Kishida M, Ninomiya T, Kohgami S, Hirooka M, Tokumoto Y, Abe M, Matsuura B, Hiasa Y. Efficacy of branched-chain amino acid supplementation and walking exercise for preventing sarcopenia in patients with liver cirrhosis. *Eur J Gastroenterol Hepatol* 2017; **29**: 1416-1423 [PMID: 29016470 DOI: 10.1097/MEG.0000000000000986]
- 9 Román E, Torrades MT, Nadal MJ, Cárdenas G, Nieto JC, Vidal S, Bascañana H, Juárez C, Guarner C, Córdoba J, Soriano G. Randomized pilot study: effects of an exercise programme and leucine supplementation in patients with cirrhosis. *Dig Dis Sci* 2014; **59**: 1966-1975 [PMID: 24599772 DOI: 10.1007/s10620-014-3086-6]
- 10 Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T, Cooper C, Landi F, Rolland Y, Sayer AA, Schneider SM, Sieber CC, Topinkova E, Vandewoude M, Visser M, Zamboni M; Writing Group for the European Working Group on Sarcopenia in Older People 2 (EWGSOP2), and the Extended Group for EWGSOP2. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing* 2019; **48**: 16-31 [PMID: 30312372 DOI: 10.1093/ageing/afy169]
- 11 Mijnders DM, Koster A, Schols JM, Meijers JM, Halfens RJ, Gudnason V, Eiriksdottir G, Siggeirsdottir K, Sigurdsson S, Jónsson PV, Meirelles O, Harris T. Physical activity and incidence of sarcopenia: the population-based AGES-Reykjavik Study. *Age Ageing* 2016; **45**: 614-620 [PMID: 27189729 DOI: 10.1093/ageing/afw090]
- 12 Barazzoni R, Bischoff SC, Boirie Y, Busetto L, Cederholm T, Dicker D, Toplak H, Van Gossum A, Yumuk V, Vettor R. Sarcopenic obesity: Time to meet the challenge. *Clin Nutr* 2018; **37**: 1787-1793 [PMID: 29857921 DOI: 10.1016/j.clnu.2018.04.018]
- 13 Cleasby ME, Jamieson PM, Atherton PJ. Insulin resistance and sarcopenia: mechanistic links between common co-morbidities. *J Endocrinol* 2016; **229**: R67-R81 [PMID: 26931135 DOI: 10.1530/JOE-15-0533]
- 14 Peng TC, Chen WL, Chen YY, Chao YP, Wu LW, Kao TW. Associations between different measurements of sarcopenic obesity and health outcomes among non-frail community-dwelling older adults in Taiwan. *Br J Nutr* 2021; **126**: 1749-1757 [PMID: 33849669 DOI: 10.1017/S0007114521001288]
- 15 Atkins JL, Wannamethee SG. Sarcopenic obesity in ageing: cardiovascular outcomes and mortality. *Br J Nutr* 2020; **124**: 1102-1113 [PMID: 32616084 DOI: 10.1017/S0007114520002172]
- 16 Younossi ZM, Golabi P, Paik JM, Henry A, Van Dongen C, Henry L. The global epidemiology of nonalcoholic fatty liver disease (NAFLD) and nonalcoholic steatohepatitis (NASH): a systematic review. *Hepatology* 2023; **77**: 1335-1347 [PMID: 36626630 DOI: 10.1097/HEP.0000000000000004]
- 17 Sinn DH, Kang D, Kang M, Guallar E, Hong YS, Lee KH, Park J, Cho J, Gwak GY. Nonalcoholic fatty liver disease and accelerated loss of skeletal muscle mass: A longitudinal cohort study. *Hepatology* 2022; **76**: 1746-1754 [PMID: 35588190 DOI: 10.1002/hep.32578]
- 18 Golabi P, Gerber L, Paik JM, Deshpande R, de Avila L, Younossi ZM. Contribution of sarcopenia and physical inactivity to mortality in people with non-alcoholic fatty liver disease. *JHEP Rep* 2020; **2**: 100171 [PMID: 32964202 DOI: 10.1016/j.jhepr.2020.100171]
- 19 Joo SK, Kim W. Interaction between sarcopenia and nonalcoholic fatty liver disease. *Clin Mol Hepatol* 2023; **29**: S68-S78 [PMID: 36472051 DOI: 10.3350/cmh.2022.0358]
- 20 Petermann-Rocha F, Gray SR, Forrest E, Welsh P, Sattar N, Celis-Morales C, Ho FK, Pell JP. Associations of muscle mass and grip strength with severe NAFLD: A prospective study of 333,295 UK Biobank participants. *J Hepatol* 2022; **76**: 1021-1029 [PMID: 35085594 DOI: 10.1016/j.jhep.2022.01.010]
- 21 Kim G, Lee SE, Lee YB, Jun JE, Ahn J, Bae JC, Jin SM, Hur KY, Jee JH, Lee MK, Kim JH. Relationship Between Relative Skeletal Muscle Mass and Nonalcoholic Fatty Liver Disease: A 7-Year Longitudinal Study. *Hepatology* 2018; **68**: 1755-1768 [PMID: 29679374 DOI: 10.1002/hep.30049]
- 22 Petta S, Ciminnisi S, Di Marco V, Cabibi D, Cammà C, Licata A, Marchesini G, Craxi A. Sarcopenia is associated with severe liver fibrosis in patients with non-alcoholic fatty liver disease. *Aliment Pharmacol Ther* 2017; **45**: 510-518 [PMID: 28028821 DOI: 10.1111/apt.13889]
- 23 Iwaki M, Kobayashi T, Nogami A, Saito S, Nakajima A, Yoneda M. Impact of Sarcopenia on Non-Alcoholic Fatty Liver Disease. *Nutrients* 2023; **15** [PMID: 36839249 DOI: 10.3390/nu15040891]
- 24 Chung HH. Enhanced muscle function in cause-specific mortality among patients with NAFLD-related sarcopenia. *Liver Int* 2023; **43**: 1837-1838 [PMID: 37303293 DOI: 10.1111/liv.15617]
- 25 Mazurak VC, Tandon P, Montano-Loza AJ. Nutrition and the transplant candidate. *Liver Transpl* 2017; **23**: 1451-1464 [PMID: 29072825 DOI: 10.1002/lt.24848]
- 26 Carey EJ, Lai JC, Wang CW, Dasarathy S, Lobach I, Montano-Loza AJ, Dunn MA; Fitness, Life Enhancement, and Exercise in Liver Transplantation Consortium. A multicenter study to define sarcopenia in patients with end-stage liver disease. *Liver Transpl* 2017; **23**: 625-633 [PMID: 28240805 DOI: 10.1002/lt.24750]
- 27 Tantai X, Liu Y, Yeo YH, Praktikno M, Mauro E, Hamaguchi Y, Engelmann C, Zhang P, Jeong JY, van Vugt JLA, Xiao H, Deng H, Gao X, Ye Q, Zhang J, Yang L, Cai Y, Liu N, Li Z, Han T, Kaido T, Sohn JH, Strassburg C, Berg T, Trebicka J, Hsu YC, IJzermans JNM, Wang J, Su

- GL, Ji F, Nguyen MH. Effect of sarcopenia on survival in patients with cirrhosis: A meta-analysis. *J Hepatol* 2022; **76**: 588-599 [PMID: 34785325 DOI: 10.1016/j.jhep.2021.11.006]
- 28 **Ebadi M**, Burra P, Zanetto A, Montano-Loza AJ. Current treatment strategies and future possibilities for sarcopenia in cirrhosis. *J Hepatol* 2023; **78**: 889-892 [PMID: 36774981 DOI: 10.1016/j.jhep.2023.01.031]
- 29 **Thapaliya S**, Runkana A, McMullen MR, Nagy LE, McDonald C, Naga Prasad SV, Dasarathy S. Alcohol-induced autophagy contributes to loss in skeletal muscle mass. *Autophagy* 2014; **10**: 677-690 [PMID: 24492484 DOI: 10.4161/auto.27918]
- 30 **Aqel BA**, Scolapio JS, Dickson RC, Burton DD, Bouras EP. Contribution of ascites to impaired gastric function and nutritional intake in patients with cirrhosis and ascites. *Clin Gastroenterol Hepatol* 2005; **3**: 1095-1100 [PMID: 16271340 DOI: 10.1016/s1542-3565(05)00531-8]
- 31 **Bhanji RA**, Moctezuma-Velazquez C, Duarte-Rojo A, Ebadi M, Ghosh S, Rose C, Montano-Loza AJ. Myosteatosis and sarcopenia are associated with hepatic encephalopathy in patients with cirrhosis. *Hepatol Int* 2018; **12**: 377-386 [PMID: 29881992 DOI: 10.1007/s12072-018-9875-9]
- 32 **Tranah TH**, Ballester MP, Carbonell-Asins JA, Ampuero J, Alexandrino G, Caracostea A, Sánchez-Torrijos Y, Thomsen KL, Kerbert AJC, Capilla-Lozano M, Romero-Gómez M, Escudero-García D, Montoliu C, Jalan R, Shawcross DL. Plasma ammonia levels predict hospitalisation with liver-related complications and mortality in clinically stable outpatients with cirrhosis. *J Hepatol* 2022; **77**: 1554-1563 [PMID: 35872326 DOI: 10.1016/j.jhep.2022.07.014]
- 33 **Tapper EB**, Zhang P, Garg R, Nault T, Leary K, Krishnamurthy V, Su GL. Body composition predicts mortality and decompensation in compensated cirrhosis patients: A prospective cohort study. *JHEP Rep* 2020; **2**: 100061 [PMID: 32039402 DOI: 10.1016/j.jhepr.2019.11.005]
- 34 **Ando Y**, Ishigami M, Ito T, Ishizu Y, Kuzuya T, Honda T, Ishikawa T, Fujishiro M. Sarcopenia impairs health-related quality of life in cirrhotic patients. *Eur J Gastroenterol Hepatol* 2019; **31**: 1550-1556 [PMID: 31206408 DOI: 10.1097/MEG.0000000000001472]
- 35 **Montano-Loza AJ**, Meza-Junco J, Prado CM, Lieffers JR, Baracos VE, Bain VG, Sawyer MB. Muscle wasting is associated with mortality in patients with cirrhosis. *Clin Gastroenterol Hepatol* 2012; **10**: 166-173, 173.e1 [PMID: 21893129 DOI: 10.1016/j.cgh.2011.08.028]
- 36 **Du Y**, Karvellas CJ, Baracos V, Williams DC, Khadaroo RG; Acute Care and Emergency Surgery (ACES) Group. Sarcopenia is a predictor of outcomes in very elderly patients undergoing emergency surgery. *Surgery* 2014; **156**: 521-527 [PMID: 24929435 DOI: 10.1016/j.surg.2014.04.027]
- 37 **Tandon P**, Montano-Loza AJ, Lai JC, Dasarathy S, Merli M. Sarcopenia and frailty in decompensated cirrhosis. *J Hepatol* 2021; **75** Suppl 1: S147-S162 [PMID: 34039486 DOI: 10.1016/j.jhep.2021.01.025]
- 38 **Marasco G**, Dajti E, Ravaioli F, Brocchi S, Rossini B, Alemanni LV, Peta G, Bartalena L, Golfieri R, Festi D, Colecchia A, Renzulli M. Clinical impact of sarcopenia assessment in patients with liver cirrhosis. *Expert Rev Gastroenterol Hepatol* 2021; **15**: 377-388 [PMID: 33196344 DOI: 10.1080/17474124.2021.1848542]
- 39 **Praktiknjo M**, Book M, Luetkens J, Pohlmann A, Meyer C, Thomas D, Jansen C, Feist A, Chang J, Grimm J, Lehmann J, Strassburg CP, Abalde JG, Kukuk G, Trebicka J. Fat-free muscle mass in magnetic resonance imaging predicts acute-on-chronic liver failure and survival in decompensated cirrhosis. *Hepatology* 2018; **67**: 1014-1026 [PMID: 29059469 DOI: 10.1002/hep.29602]
- 40 **Beer L**, Bastati N, Ba-Ssalamah A, Pötter-Lang S, Lampichler K, Bican Y, Lauber D, Hodge J, Binter T, Pomej K, Simbrunner B, Semmler G, Trauner M, Mandorfer M, Reiberger T. MRI-defined sarcopenia predicts mortality in patients with chronic liver disease. *Liver Int* 2020; **40**: 2797-2807 [PMID: 32816394 DOI: 10.1111/liv.14648]
- 41 **Montano-Loza AJ**, Angulo P, Meza-Junco J, Prado CM, Sawyer MB, Beaumont C, Esfandiari N, Ma M, Baracos VE. Sarcopenic obesity and myosteatosis are associated with higher mortality in patients with cirrhosis. *J Cachexia Sarcopenia Muscle* 2016; **7**: 126-135 [PMID: 27493866 DOI: 10.1002/jcsm.12039]
- 42 **Montano-Loza AJ**, Meza-Junco J, Baracos VE, Prado CM, Ma M, Meeberg G, Beaumont C, Tandon P, Esfandiari N, Sawyer MB, Kneteman N. Severe muscle depletion predicts postoperative length of stay but is not associated with survival after liver transplantation. *Liver Transpl* 2014; **20**: 640-648 [PMID: 24678005 DOI: 10.1002/lt.23863]
- 43 **Kalafateli M**, Mantzoukis K, Choi Yau Y, Mohammad AO, Arora S, Rodrigues S, de Vos M, Papadimitriou K, Thorburn D, O'Beirne J, Patch D, Pinzani M, Morgan MY, Agarwal B, Yu D, Burroughs AK, Tsochatzis EA. Malnutrition and sarcopenia predict post-liver transplantation outcomes independently of the Model for End-stage Liver Disease score. *J Cachexia Sarcopenia Muscle* 2017; **8**: 113-121 [PMID: 27239424 DOI: 10.1002/jcsm.12095]
- 44 **Kaido T**, Ogawa K, Fujimoto Y, Ogura Y, Hata K, Ito T, Tomiyama K, Yagi S, Mori A, Uemoto S. Impact of sarcopenia on survival in patients undergoing living donor liver transplantation. *Am J Transplant* 2013; **13**: 1549-1556 [PMID: 23601159 DOI: 10.1111/ajt.12221]
- 45 **Jiang C**, Wang Y, Fu W, Zhang G, Feng X, Wang X, Wang F, Zhang L, Deng Y. Association between sarcopenia and prognosis of hepatocellular carcinoma: A systematic review and meta-analysis. *Front Nutr* 2022; **9**: 978110 [PMID: 36590214 DOI: 10.3389/fnut.2022.978110]
- 46 **Perisetti A**, Goyal H, Yendala R, Chandan S, Tharian B, Thandassery RB. Sarcopenia in hepatocellular carcinoma: Current knowledge and future directions. *World J Gastroenterol* 2022; **28**: 432-448 [PMID: 35125828 DOI: 10.3748/wjg.v28.i4.432]
- 47 **Fujiwara N**, Nakagawa H, Kudo Y, Tateishi R, Taguri M, Watadani T, Nakagomi R, Kondo M, Nakatsuka T, Minami T, Sato M, Uchino K, Enooku K, Kondo Y, Asaoka Y, Tanaka Y, Ohtomo K, Shiina S, Koike K. Sarcopenia, intramuscular fat deposition, and visceral adiposity independently predict the outcomes of hepatocellular carcinoma. *J Hepatol* 2015; **63**: 131-140 [PMID: 25724366 DOI: 10.1016/j.jhep.2015.02.031]
- 48 **Roth G**, Teyssier Y, Benhamou M, Abousalihac M, Caruso S, Sengel C, Seror O, Ghelfi J, Seigneurin A, Ganne-Carrie N, Gigante E, Blaise L, Sutter O, Decaens T, Nault JC. Impact of sarcopenia on tumor response and survival outcomes in patients with hepatocellular carcinoma treated by trans-arterial (chemo)-embolization. *World J Gastroenterol* 2022; **28**: 5324-5337 [PMID: 36185630 DOI: 10.3748/wjg.v28.i36.5324]
- 49 **Scheiner B**, Lampichler K, Pomej K, Beer L, Balcar L, Sartoris R, Bouattour M, Sidali S, Trauner M, Mandorfer M, Reiberger T, Scharitzer M, Tamandl D, Pinato DJ, Ronot M, Pinter M. Transversal psoas muscle thickness measurement is associated with response and survival in patients with HCC undergoing immunotherapy. *Hepatol Commun* 2023; **7** [PMID: 37708441 DOI: 10.1097/HCC.0000000000000261]
- 50 **Mir O**, Coriat R, Blanchet B, Durand JP, Boudou-Rouquette P, Michels J, Ropert S, Vidal M, Pol S, Chaussade S, Goldwasser F. Sarcopenia predicts early dose-limiting toxicities and pharmacokinetics of sorafenib in patients with hepatocellular carcinoma. *PLoS One* 2012; **7**: e37563 [PMID: 22666367 DOI: 10.1371/journal.pone.0037563]
- 51 **Dong D**, Shi JY, Shang X, Liu B, Xu WL, Cui GZ, Wang NY. Prognostic significance of sarcopenia in patients with hepatocellular carcinoma treated with lenvatinib: A retrospective analysis. *Medicine (Baltimore)* 2022; **101**: e28680 [PMID: 35119010 DOI: 10.1097/MD.00000000000028680]
- 52 **Uojima H**, Chuma M, Tanaka Y, Hidaka H, Nakazawa T, Iwabuchi S, Kobayashi S, Hattori N, Ogushi K, Morimoto M, Kagawa T, Tanaka K,

- Kako M, Koizumi W. Skeletal Muscle Mass Influences Tolerability and Prognosis in Hepatocellular Carcinoma Patients Treated with Lenvatinib. *Liver Cancer* 2020; **9**: 193-206 [PMID: 32399433 DOI: 10.1159/000504604]
- 53 **European Association for the Study of the Liver.** EASL Clinical Practice Guidelines on nutrition in chronic liver disease. *J Hepatol* 2019; **70**: 172-193 [PMID: 30144956 DOI: 10.1016/j.jhep.2018.06.024]
- 54 **Bhanji RA,** Montano-Loza AJ, Watt KD. Sarcopenia in Cirrhosis: Looking Beyond the Skeletal Muscle Loss to See the Systemic Disease. *Hepatology* 2019; **70**: 2193-2203 [PMID: 31034656 DOI: 10.1002/hep.30686]
- 55 **Prado CM,** Landi F, Chew STH, Atherton PJ, Molinger J, Ruck T, Gonzalez MC. Advances in muscle health and nutrition: A toolkit for healthcare professionals. *Clin Nutr* 2022; **41**: 2244-2263 [PMID: 36081299 DOI: 10.1016/j.clnu.2022.07.041]
- 56 **Barazzoni R,** Jensen GL, Correia MITD, Gonzalez MC, Higashiguchi T, Shi HP, Bischoff SC, Boirie Y, Carrasco F, Cruz-Jentoft A, Fuchs-Tarlovsky V, Fukushima R, Heymsfield S, Mourtzakis M, Muscaritoli M, Norman K, Nyulasi I, Pisprasert V, Prado C, de van der Schuren M, Yoshida S, Yu Y, Cederholm T, Compher C. Guidance for assessment of the muscle mass phenotypic criterion for the Global Leadership Initiative on Malnutrition (GLIM) diagnosis of malnutrition. *Clin Nutr* 2022; **41**: 1425-1433 [PMID: 35450768 DOI: 10.1016/j.clnu.2022.02.001]
- 57 **Nijholt W,** Scafoglieri A, Jager-Wittenaar H, Hobbelen JSM, van der Schans CP. The reliability and validity of ultrasound to quantify muscles in older adults: a systematic review. *J Cachexia Sarcopenia Muscle* 2017; **8**: 702-712 [PMID: 28703496 DOI: 10.1002/jcsm.12210]
- 58 **English C,** Fisher L, Thoires K. Reliability of real-time ultrasound for measuring skeletal muscle size in human limbs in vivo: a systematic review. *Clin Rehabil* 2012; **26**: 934-944 [PMID: 22324054 DOI: 10.1177/0269215511434994]
- 59 **European Association for the Study of the Liver.** EASL Clinical Practice Guidelines: Management of hepatocellular carcinoma. *J Hepatol* 2018; **69**: 182-236 [PMID: 29628281 DOI: 10.1016/j.jhep.2018.03.019]
- 60 **Singal AG,** Llovet JM, Yarchoan M, Mehta N, Heimbach JK, Dawson LA, Jou JH, Kulik LM, Agopian VG, Marrero JA, Mendiratta-Lala M, Brown DB, Rilling WS, Goyal L, Wei AC, Taddei TH. AASLD Practice Guidance on prevention, diagnosis, and treatment of hepatocellular carcinoma. *Hepatology* 2023; **78**: 1922-1965 [PMID: 37199193 DOI: 10.1097/HEP.0000000000000466]
- 61 **Pillen S,** van Alfen N. Skeletal muscle ultrasound. *Neurol Res* 2011; **33**: 1016-1024 [PMID: 22196753 DOI: 10.1179/1743132811Y.0000000010]
- 62 **Woodward AJ,** Wallen MP, Ryan J, Ward LC, Coombes JS, Macdonald GA. Evaluation of techniques used to assess skeletal muscle quantity in patients with cirrhosis. *Clin Nutr ESPEN* 2021; **44**: 287-296 [PMID: 34330481 DOI: 10.1016/j.clnesp.2021.05.029]
- 63 **Becchetti C,** Berzigotti A. Ultrasonography as a diagnostic tool for sarcopenia in patients with cirrhosis: Examining the pros and cons. *Eur J Intern Med* 2023; **116**: 27-33 [PMID: 37385916 DOI: 10.1016/j.ejim.2023.06.019]
- 64 **Zenith L,** Meena N, Ramadi A, Yavari M, Harvey A, Carbonneau M, Ma M, Abraldes JG, Paterson I, Haykowsky MJ, Tandon P. Eight weeks of exercise training increases aerobic capacity and muscle mass and reduces fatigue in patients with cirrhosis. *Clin Gastroenterol Hepatol* 2014; **12**: 1920-6.e2 [PMID: 24768811 DOI: 10.1016/j.cgh.2014.04.016]
- 65 **Abe T,** Kawakami Y, Kondo M, Fukunaga T. Comparison of ultrasound-measured age-related, site-specific muscle loss between healthy Japanese and German men. *Clin Physiol Funct Imaging* 2011; **31**: 320-325 [PMID: 21672141 DOI: 10.1111/j.1475-097X.2011.01021.x]
- 66 **Mirón Mombiela R,** Vucetic J, Rossi F, Tagliafico AS. Ultrasound Biomarkers for Sarcopenia: What Can We Tell So Far? *Semin Musculoskelet Radiol* 2020; **24**: 181-193 [PMID: 32438444 DOI: 10.1055/s-0039-3402745]
- 67 **Stock MS,** Thompson BJ. Echo intensity as an indicator of skeletal muscle quality: applications, methodology, and future directions. *Eur J Appl Physiol* 2021; **121**: 369-380 [PMID: 33221942 DOI: 10.1007/s00421-020-04556-6]
- 68 **Ticinesi A,** Meschi T, Narici MV, Lauretani F, Maggio M. Muscle Ultrasound and Sarcopenia in Older Individuals: A Clinical Perspective. *J Am Med Dir Assoc* 2017; **18**: 290-300 [PMID: 28202349 DOI: 10.1016/j.jamda.2016.11.013]
- 69 **Mitchell WK,** Phillips BE, Williams JP, Rankin D, Smith K, Lund JN, Atherton PJ. Development of a new Sonovue™ contrast-enhanced ultrasound approach reveals temporal and age-related features of muscle microvascular responses to feeding. *Physiol Rep* 2013; **1**: e00119 [PMID: 24303186 DOI: 10.1002/phy2.119]
- 70 **Alfuraih AM,** Tan AL, O'Connor P, Emery P, Wakefield RJ. The effect of ageing on shear wave elastography muscle stiffness in adults. *Aging Clin Exp Res* 2019; **31**: 1755-1763 [PMID: 30762201 DOI: 10.1007/s40520-019-01139-0]
- 71 **Tandon P,** Low G, Mourtzakis M, Zenith L, Myers RP, Abraldes JG, Shaheen AA, Qamar H, Mansoor N, Carbonneau M, Ismond K, Mann S, Alaboudy A, Ma M. A Model to Identify Sarcopenia in Patients With Cirrhosis. *Clin Gastroenterol Hepatol* 2016; **14**: 1473-1480.e3 [PMID: 27189915 DOI: 10.1016/j.cgh.2016.04.040]
- 72 **Dhariwal S,** Roy A, Taneja S, Bansal A, Gorski U, Singh S, De A, Verma N, Premkumar M, Duseja A, Dhiman R, Singh V. Assessment of Sarcopenia Using Muscle Ultrasound in Patients With Cirrhosis and Sarcopenic Obesity (AMUSE STUDY). *J Clin Gastroenterol* 2023; **57**: 841-847 [PMID: 35943413 DOI: 10.1097/MCG.0000000000001745]
- 73 **Ciocirlan M,** Mănuș M, Dicușescu M, Ciocirlan M. Is rectus abdominis thickness associated with survival among patients with liver cirrhosis? A prospective cohort study. *Sao Paulo Med J* 2019; **137**: 401-406 [PMID: 31721939 DOI: 10.1590/1516-3180.2019.000406082019]
- 74 **Hari A,** Berzigotti A, Štabuc B, Cagleviç N. Muscle psoas indices measured by ultrasound in cirrhosis - Preliminary evaluation of sarcopenia assessment and prediction of liver decompensation and mortality. *Dig Liver Dis* 2019; **51**: 1502-1507 [PMID: 31547952 DOI: 10.1016/j.dld.2019.08.017]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

